U. S. DEPARTMENT OF COMMERCE

RESEARCH PAPER RP815

Part of Journal of Research of the National Bureau of Standards, Volume 15, August 1935

COLOR AND SPECTRAL TRANSMITTANCE OF VEGETABLE OILS¹

By Harry I. McNicholas²

ABSTRACT

A spectrophotometric analysis of 125 vegetable oils, 111 of which are from the cotton seed, has been made as a rational physical basis for the discussion and development of improved methods for the color grading of oils. The lightness and chromaticity of all the oils, and their spectral transmittance from 440 to 720 $m\mu$, are indicated in tabular and graphical form. Consideration is given to some difficulties inherent in the color-grading problem, arising chiefly from the independent variation of the concentrations of several different pigments in the oils. Color grading in terms of the Lovibond glass standards is discussed along with other abridged methods of colorimetry suggested by the data.

CONTENTS

I.	Introduction	99
II.	Source and description of oils	100
III.	Spectral transmittance of oils	100
IV.	Colorimetric properties	106
V.	Discussion of data on oils and Lovibond glasses	113
VI.	Some abridged color-grading methods	117
VII.	Summary and conclusions	120

I. INTRODUCTION

The color of crude and refined vegetable oils is an important contributing factor in the determination of their market value. For many years the commercial color grading of these oils has been based entirely on the Lovibond system of glass standards. Many difficulties in the use of this system have arisen in practice, involving inherent properties of the system itself or as the result of the methods of application. Attempts to substitute colorimetric methods based on other material color standards have likewise failed to provide a satisfactory solution to the problem.

Paga

¹ The spectrophotometric measurements of the oils were made between July 1915 and July 1920 under the immediate supervision of I. G. Priest, with the cooperation and financial support of the Society of Cotton Products Analysts and the Interstate Cotton Seed Crushers Association. The author was employed by the above societies and detailed to the National Bureau of Standards as research associate from July 1915 to July 1916. He was then appointed to the Bureau staff. Reports involving some phases of this work have been published by Priest in the Cotton Oil Press, as follows: **3**, 86 (1919-20) no. **3**; **3**, 37 (1919-20) no. 9; **4**, 95 (1920-21) no. 3. ^a The author acknowledges the assistance of W. B. Emerson and M. K. Frehafer on some of the early spectrophotometric work. Throughout the recent analysis and preparation of data (1934-36) he is indebted to H. J. Keegan for assistance in computational work and to K. S. Gibson, D. B. Judd, and G. W. Haupt for many supplementary data supplied for the purposes of this paper. These data include all spectrophotometric or colorimetric data on the spectrum and the Lovibond glasses.

100 Journal of Research of the National Bureau of Standards [Vol. 15

As a rational physical basis for the analysis and discussion of the problem, and for the ultimate establishment of a more desirable procedure in color grading, an extensive investigation was undertaken of the spectral-transmissive and colorimetric properties of vegetable oils, covering a wide range of color variation in cottonseed oils from different geographical sources, and a more limited range of variation in vegetable oils of different plant origin.

The present paper gives the results of this study and a discussion of some difficulties inherent in the problem. Inasmuch as the color of these oils is at present universally expressed in terms of Lovibond glasses, this color system is also extensively employed in the present paper for comparison purposes.

II. SOURCE AND DESCRIPTION OF OILS

The vegetable oils included in this investigation were generously supplied by various manufacturers, dealers, and individuals interested in the work.³ Information is generally unavailable on details of the methods of processing, the chemical composition, and the exact geographical origin of these oils. For purely colorimetric purposes, however, it is believed that collectively they are fairly representative of the majority of the edible vegetable oils of commerce, at least of the cottonseed oils. Such oils vary in color from a light greenish yellow through canary or amber yellow to brown or deep red, and contain varying proportions of several different coloring matters commonly found in the tissues of plants.

The oils included in this work consist of 111 samples of refined cottonseed oils, designated in the tables and figures as groups A, B, D, E, and R, and 14 vegetable oils of other plant origin, herein, designated as group F. The latter group⁴ includes samples of sesame, peanut, soybean, corn, rape, and olive oils. The A and B groups were received in 1915 and 1916 from 21 different refineries scattered widely throughout the country. The B group consists entirely of nearly prime wintered oils. The D group was collected in 1919 and 1920 from various individuals. The E group was obtained in 1920 through the Bureau of Chemistry and Soils of the U. S. Department of Agriculture from various sources in North Carolina, South Carolina, Georgia, Alabama, Tennessee, and Texas. The R group of refined oils was obtained in like manner along with samples of the crude oils from which the refinings were made.

III. SPECTRAL TRANSMITTANCE OF OILS

Complete information on the color of an oil may be derived when the transmission of light is specified for the oil at various wave lengths throughout the visible spectrum. Such measurements have been made for a 5¼ inch thickness of each oil and the results represented graphically by the curves in figures 1 to 5. In these figures the solid-line curves represent cottonseed oils and the broken-line

³ The donors include Armour and Co., Harris Abattoir (Toronto), Procter and Gamble, Peet Bros., Globe Soap Co., Refuge Cotton Oil Co., Swift and Co., The Fort Worth Laboratories, Southern Cotton Oil Co., Corn Products Co., Phoenix Cotton Oil Co., Portsmouth Cotton Oil Co., Gunns Limited (Toronto), Capitol Refning Co., Armstrong Packing Co., Trinity Cotton Oil Co., American Cotton Oil Co., Falkenburg and Co., Empire Cotton Oil Co., Temple Cotton Oil Co., Independent Cotton Oil Co., Sea Island Cotton Oil Co., Magic City Oil Co., Felix Paquin, T. C. Law, F. N. Smalley, David Wesson, and P. S. Tilson.

⁴ Spectrophotometric data for oils F 10 to F 14 were taken from the work of Wesson, Cotton Oil Press 6, no. 5 (September 1922).

Color of Vegetable Oils



FIGURE 2.—Spectral transmittance of vegetable oils.

[Thickness 51/4 inches]



FIGURE 4.—Spectral transmittance of vegetable oils. [Thickness 51/4 inches]

curves the other oils of the F group. Oils of the A and B groups, in a $\frac{3}{6}$ inch thickness, are represented in figure 6. All data were obtained with the König-Martens spectrophotometer ⁵ at room temperature. There is no evidence of any change of importance in the transmittance of vegetable oils when the temperature is varied.⁶

The wide variation in form of the spectral-transmittance curves for the various oils is proof of the presence of several different pigments in widely varying concentration. Little is known regarding the exact identity of these pigments. The well-defined absorption bands centered for most of the oils at wave lengths 670 and 610 m μ ,



FIGURE 5.—Spectral transmittance of vegetable oils. [Thickness 5¼ inches]

and the similar bands shifted somewhat from these positions in a few cases (fig. 5), show the presence undoubtedly of chlorophyll compounds, which, if present alone, would give the oils the characteristic green color of these materials in solution. It will be convenient to refer to these pigments collectively as the green group of coloring matters. It is evident also that one or more yellow pigments must be present, which have strong absorptive properties in the blue and violet regions of the spectrum, such as would result from the presence of one or several of the known yellow plant pigments. In addition some yellow, amber, or brown pigments must be present, which are characterized by a more gradual increase in absorption from the red through the yellow and green to the blue. These yellow, amber, and brown materials will be denoted collectively as the brown group of

⁴ BS J. Research 1, 793 (1928) RP30. The arrangement of light source and auxiliary equipment for th spectrophotometric analysis of vegetable oils was not the same as described in this paper, but the method of measurement was essentially the same. Brass cells with glass end-plates were used to contain the oils The measured transmission of these end-plates with a thin film of bleached oil between was found to ba 0.892 for all wave lengths. Each measured transmission value of the cell filled with oil was divided by this number to give the transmittance of the loi litself as represented in the figures and in table 1. ⁶ See Judd, BS J. Research 1, 859, footnote 18, (1928) RP31.

McNicholas]

104 Journal of Research of the National Bureau of Standards [Vol. 15

pigments and are of considerable importance in determining the color of the oil in a 5¼ inch thickness. The general trend of the spectral-transmissive properties for the brown group is represented by those curves in figures 1 and 7 which exhibit little trace of the chlorophyll bands. Figure 7, lower part, shows the spectral transmittance of some crude cottonseed oils, herein denoted as the C group. During refining the brown pigments are removed in greater proportion than the green, as illustrated particularly by the transition from C 3 to R 3. Another factor possibly affecting the transmittance by the residual turbidity of some of the oils. However, all noticeably turbid oils were filtered through absorbent cotton before measurements were made.



FIGURE 6.—Spectral transmittance of cottonseed oils. [Thickness 3% inch]

In figures 1 to 4 the oils have been roughly classified in accordance with the form of their transmittance curves. All the oils show a strong absorption in the violet and blue regions of the spectrum, which increases markedly with increasing yellowness or redness of the oils. Through the color transition we observe the existence of two principal spectral types. The "green" type (fig. 4) is characterized by the presence chiefly of the green group of pigments with their sharp bands in the orange and red and lesser bands usually accompanying a sharp decrease in transmittance from the yellow into the green. The "brown" type (fig. 1), on the other hand, shows no very pronounced absorption bands and exhibits a more gradual decrease in transmittance through the yellow and green. The two types of oils are observed to merge one into the other so that the observed color variations in a 5¼ inch thickness may be ascribed to a corresponding variation in the absolute and relative concentrations of the two groups of pigments. It is probable that for purely colori-

Color of Vegetable Oils

McNicholas]

metric purposes the color variation over the entire range of vegetable oils could be satisfactorily described in terms of the independent



FIGURE 7.—Spectral transmittance of crude cottonseed oils and refinings from same. Showing the effect of the refining process on the selective removal of pigment components in the oils, and the reduction in total pigment content. The thickness ratio of crude to refined oils, found necessary to obtain comparable transmittance data, shows (by Beer's law) that the reduction in total pigment content is roughly 93 percent.

variation in concentration of two hypothetical pigments representative of the brown and green groups, respectively. This point is further discussed in section VI.

106 Journal of Research of the National Bureau of Standards 100.15

In figure 5 two small groups of oils are shown, each differing slightly from the oils in the preceding figures, and represented, respectively, by solid and broken lines. The solid-line curves represent cottonseed oils showing a definite displacement of the principal absorption band from the position at 670 m μ exhibited by the majority of all vegetable oils. This demonstrates the presence in general of more than one chlorophyll compound. These oils also contain a large proportion of the brown group of pigments. The broken-line curves in the figure represent samples of peanut, soybean, rape, and olive oils characterized here by the relatively high transmittance in the region about 580 m μ . These oils show the maximum proportion of the green pigments. They are the purest of the green spectral type and among the lightest of yellow oils included in this investigation. (See also fig. 10.) In other oils of the predominantly green spectral type (fig. 4) the transmittance maximum at 580 m μ is considerably reduced by the superposed characteristic transmittance of the brown pigments present. It may be noted that one oil, F 11, combines both spectral characteristics exhibited in figure 5, namely, the high transmittance maximum at 580 m μ and the displacement of the principal chlorophyll band.

The classification of oils into two principal spectral types is further illustrated by the curves in figure 6 for a % inch thickness. The dashdot curves represent all oils of groups A and B included in figure 1, along with oils A 1, B 6, and B 9 of figure 5. They are the purest A and B representatives of the brown spectral type. Among the remaining oils, the strong absorption in the blue corresponds to the sharp cut-off near 510 m μ as observed in figures 3 and 4.

A comparison of the solid and broken-line curves in figures 1 to 5 shows the essential sameness in spectral-transmissive properties between cottonseed and other vegetable oils herein examined. For each of the other vegetable oils there is a cottonseed oil not differing essentially in spectral type. This fact has been pointed out by Priest ⁷ for one sample of soybean oil, and is further illustrated by Wesson⁸ for peanut, rape, olive, and cocoanut oils. The most significant difference herein exhibited between cottonseed oils and the rape, soybean, sesame, and olive oils is the relatively higher transmittance of the latter group near 580 mµ as compared with that near 630 m μ . This follows from the smaller proportion of brown pigment materials. Hence, we may regard the problem of color grading over the entire range of cottonseed oils as practically encompassing that for all vegetable oils.

IV. COLORIMETRIC PROPERTIES

Having the spectral transmittance of an oil (or Lovibond-glass combination) we may proceed by purely analytical or graphical methods to derive ⁹ various desired colorimetric specifications, with reference to a standard illuminant, observer, and coordinate system. In the present paper the standard illuminant is Abbot-Priest sun.¹⁰

⁷ Cotton Oil Press (January 1920).
⁸ Cotton Oil Press 6, no. 5 (September 1922). These spectrophotometric data on the peanut, rape, and olive oils are reproduced in the present paper.
⁹ For methods of computing colorimetric properties from spectral transmittance and other data, see J. Opt. Soc. Am. and Rev. Sci. Instr. 6, 527 (1922); J. Opt. Soc. Am. 23, 359 (1933).
¹⁰ J. Opt. Soc. Am. and Rev. Sci. Instr. 12, 479 (1926).

which serves also as the basic stimulus 11 for a trilinear coordinate system.¹² The standard observer and coordinate system are defined by the (extrapolated) OSA visual response functions.¹³ These ref-erence conditions are the same as have been employed in all standardization work on Lovibond glasses at the National Bureau of Standards.14

Assuming that a 5¹/₄-inch column of oil is illuminated by light from the standard illuminant, the luminous transmittance T_s and the trilinear coordinates (r, g, b) have been determined from the given properties of the source and observer and the measured values of spectral transmittance. These quantities are listed in columns 3, 4, 5, and 6 of table 1. They specify, respectively, the purely luminous and chromatic aspects of the vegetable-oil colors (lightness and chromaticity, respectively). Inasmuch as the trilinear coordinates sum to unity by definition, any two of them may be used in a graphical representation of chromaticity. This is done on the (r, g) diagram of figure 8, which shows the Lovibond 35Y + N''R locus¹⁵ and a part of the spectrum (λ) locus. The relation of chromaticities on the 35Y + N''R scale to the complete spectrum locus, the black-body locus, and the standard illuminant, is shown on the smallscale (r, g) diagram in the figure.

The chromaticities of all the oils have also been evaluated on the one-dimensional (35Y+N''R) and λ scales by projection of points representing the oils parallel to the system of cross lines drawn in figure 8, and then reading the projected points from the given scales. The results are recorded in columns 8 and 9 of table 1. To a sufficient approximation for present purposes the analytical and graphical procedure ¹⁶ involved is equivalent to the establishment of a nearest chromaticity setting between an oil and the series of Lovibond-glass combinations (or the series of spectrum colors), when compared in a two-part photometric field under the standard conditions of illuminant and observer herein prescribed.

A comparison is made in figure 9 between the N'' values herein assigned to the oils and the N values submitted by donors of the oils (column 7, table 1). The spread of values about the straight line shows that any difference between the two chromaticity scales is negligible compared with erraticisms in the assignment of grades.

¹¹ J. Guild, On the fixed points of a colormetric system, Discussion by Priest, Trans. Opt. Soc. 32, 25 (1930-31).

<sup>(1930-31).
&</sup>lt;sup>12</sup> J. Opt. Soc. Am. and Rev. Sci. Instr. 9, 506 (1924).
¹³ J. Opt. Soc. Am. and Rev. Sci. Instr. 10, 230 (1925).
¹⁴ BS J. Research 13, 433 (1934) RP718. Also Oil and Soap 11, 246 (1934). This paper contains complete references to previous work.
¹⁵ BS J. Research 13, 433 (1934) RP718. Also Oil and Soap 11, 246 (1934). The N"R scale is based on the new derivation of unit red in the Lovibond system, yielding a consistent additive scale when com-

bined with 35 yellow

bined with 35 yellow. ¹⁶ To determine the directions of the projection lines, the $35Y + N''\lambda$ and N loci were first transferred by a prescribed transformation of coordinates from the OSA colorimetric system of figure 8 to the uniform-scale system devised by Judd, J. Research NBS 14, 41 (1935) RP756; or J. Opt. Soc. Am. 25, 24 (1935). In this new system, the nearest chromaticity match to be found for a given color out of any series of colors (such as the 35Y + N''R series, or the series of spectrum colors) is determined by drawing the shortest line (the perpendicular) from the point representing the given color to the locus representing the series. Accordingly, sets of perpendiculars were drawn both to the 35Y + N''R locus and to the spectrum locus; but since these loci in the uniform-scale system are nearly straight, parallel lines, these sets of perpendiculars are almost identical. Hence, one set will serve to determine both N'' and λ values of any color represented by points in the neighborhood of these loci. The series of cross lines shown in figure 8 corresponds to this set of perpendiculars. perpendiculars.

The luminous transmissions ¹⁷ of the cell-plus-oil and Lovibondglass combinations are compared in the upper part of figure 8. Values for combinations of 2, 3, and 4 glasses are shown.



FIG. 8.—Colorimetric properties of vegetable oils.

In figures 10 to 16 the spectral-transmittance curves for all the oils are replotted in a new classification, this time in accordance with their color grade. Again solid lines represent cottonseed oils and broken lines the other vegetable oils. The filled circles show the

¹⁷ Luminous transmissions differ from the luminous transmittances in column 3 of table 1 by a factor 0.892, as explained in footnote 5. Luminous transmissions given in figure 8 for a combination of two Lovibond glasses are multiplied by 0.901 and (0.901)², respectively, to obtain the transmissions for a combination of 3 or 4 glasses. Multiple reflections between glass surfaces are disregarded. The number 0.901 is the transmission of a single clear glass.

Color of Vegetable Oils

McNicholas]



PRIEST - GIBSON SCALE (35 YELLOW + N" RED)

FIGURE 9.—Comparison of Lovibond grades assigned by donors of oils with those derived in the present investigation.



FIGURE 10.—Spectral-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N'' between 0 and 5

109



FIGURE 11.—Spectral-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N'' between 5 and 6



FIGURE 12.—Special-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N'' between 6 and 7



FIGURE 13.—Spectral-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N' between 7 and 8.



FIGURE 14.—Spectral-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N'' between 8 and 10.



FIGURE 15.—Spectral-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N'' between 11 and 13.



FIGURE 16.—Spectral-transmissive properties of oils and Lovibond-glass combinations (filled circles) in the range of N'' between 13 and 20.

course of the spectral-transmittance¹⁸ curves for Lovibond-glass combinations on the scale of 35Y+N''R. In each figure, the assigned color grades of the group of oils therein represented lie between the N'' values indicated in the figure.

V. DISCUSSION OF DATA ON OILS AND LOVIBOND GLASSES

The wide variation in spectral-transmissive properties within the group of vegetable oils is illustrated by the classification in figures 1 to 5. A like variation between oils of nearly the same chromaticity is also illustrated in figures 10 to 16, along with the divergence of the oils, taken collectively, from the Lovibond-glass combinations in the same chromaticity range. These variations in spectral-transmissive properties are all translated into colorimetric terms and indicated by the spread of points in the upper and lower diagrams of figure 8.

The regular variation in spectral transmittance, lightness, and chromaticity for the sequence of color standards along the one-dimensional N'' scale follows from the regular variation in concentration of coloring material in the glass combinations. On the constantyellow basis, this variation resides solely in one component (red) of a two-component system (yellow+red). The lateral spread of points representing lightness and chromaticity of the vegetable oils follows from the irregular variations in form of the spectral-transmittance curves, and, assuming for simplicity that these variations may be ascribed likewise to variations in a two-component system (hypothetical brown and green pigments), we find the important difference exhibited in this case that the relative and absolute concentrations of the oil-pigment components may vary independently of one another. It is implied, therefore, that no one-dimensional chromaticity scale is possible on which all vegetable oils or all cottonseed oils can be accurately represented.

It may be noted at this point that the situation in this respect might be improved somewhat if it were feasible to modify or control the refining processes so as to confine the variation in pigment composition more nearly to one group of pigments, thus approaching the simpler condition of a one-component system. Inasmuch as information is not available on methods and conditions employed in the processing of these oils, no conclusive evidence on this matter can be gleaned from the present investigation. It is observed, however, that in refining the crude oils represented in figure 7 the brown pigments are more selectively removed than the green, when the latter are present in notable proportion. A continuation of this process would yield oils of a light-yellow or olive-oil hue, as illustrated in figures 5 and 10 by the broken-line curves, and in figure 8 (lower part) by the crossed circles for oils F 10 to F 14. Oil F 13, having perhaps the highest proportion of green pigments, is also of lowest Lovibond grade.

811-35-2

¹⁸ Given spectral-transmission curves for combinations of 2 glasses were divided by $(0.901)^2$ to derive the values of transmittance. When transmission curves for integral values of N'' were not available, they were found for present purposes by interpolation.

114 Journal of Research of the National Bureau of Standards [Vol. 15

It is known that the green pigments are the most fugitive to the bleaching action of light and chemical agents. Wesson ¹⁹ has shown for two samples of cottonseed oils that the bands at 670 and 610 m μ are nearly completely removed by the treatment with fuller's earth, and unpublished experiments of the author show the same effect after exposure to sunlight. When the brown pigments are present in considerable proportion, the removal of the green pigments does not have a large effect on the color. The absence of the green pigments would tend to confine the observed variation in chromaticity more closely to a one-dimensional range determined solely by the varying concentration of the brown pigments.

The fixed amount of yellow coloring material used in the combination of 35Y + N''R leads to the fact illustrated in figure 8 that none of the oils included in this investigation fall exactly on the established N'' scale, hence, none of these oils are accurately matched in chromaticity by any possible combination of a fixed 35-yellow with variable red. The position of the locus of glass combinations could be brought more nearly into coincidence with the general trend of points for the vegetable oils by use of a glass of yellow numeral higher than 35. This was corroborated by some experimental work with the B group of cottonseed oils. A color comparator was used, provided with means for equalizing brightnesses, and the chromaticities of the oils and Lovibond glasses were matched as closely as possible first with 35-yellow and variable red. Then, with the red value kept constant, more vellow was added until the best chromaticity match was obtained. It was found on the average that 80-vellow was required for these oils. This shift in the locus of glass combinations would be less suitable, however, for very light cottonseed oils; for, as recorded in table 1 and figure 9, the oils between N'' values of 1 and 3 were graded experimentally on a low-yellow basis in the neighborhood of 20Y.

It is evident that an improved one-dimensional Lovibond scale, which would coincide with the general trend of points for all vegetable oils, would require a fixation of the yellow numeral in some functional relation to the red. For the most accurate color grading, however, a two-dimensional Lovibond scale, with independent variation of both yellow and red, is necessary to cope completely with the independent variation in pigment concentrations within the group of vegetable oils.

A comparison of the luminous transmissions of the oils and Lovibond glasses in the upper part of figure 8 shows a general agreement in lightness, but the spread of the oils in this respect is greater than can be matched by any convenient choice of the number of clear and colored glasses used in the combination while maintaining the best chromaticity match. This condition introduces serious uncertainties in the present practice of color grading, because of the residual brightness difference between the fields of the color comparator when the final chromaticity setting is made.

It is evident from the data herein presented that the accurate color grading of vegetable oils can be based only on a two-dimensional chromaticity scale, which must be embodied in a colorimeter provided also with means for the independent equalization of brightnesses in the sample and comparison fields. Many such two-dimensional

¹⁹ Cotton Oil Press 6, no. 5 (September 1922).

scales could be devised and employed, provided there is no objection to the added complication in color grading. For limited ranges of color variation it is sometimes satisfactory for practical purposes to fix one of the scale variables at a constant value, or restrain its variation in some fixed relation to the other dimension, so that a onedimensional scale is effectively established. Illustrations of these possibilities have been given for the Lovibond system of yellow and red glasses and further examples are discussed in the following section.

TABLE 1.-Colorimetric properties of vegetable oils for a thickness of 5¼ inches

NBS lab. no.	Kind of oil or source			Lo	vibond	l grade					
		Sun- light trans- mit- tance	n- ht ns- it- nce				donor	By NBS on 35Y basis	Wave length	Spectral trans- mittance shown in	
		т.	r	g	b	Y	R	R	λ	Fig- ure	Fig- ure
		3	4	4 5	6	7		8	9	10	11
A1 A2 A3 A4 A5		0. 183 .070 .034 .521 .213	$\begin{array}{r} 0.\ 6093\\ .\ 6477\\ .\ 7140\\ .\ 5262\\ .\ 5934 \end{array}$	$\begin{array}{r} 0.\ 3822\\ .\ 3461\\ .\ 2832\\ .\ 4260\\ .\ 3976 \end{array}$	0.0085 .0062 .0028 .0477 .0090	35 35 35 20 35	$\begin{array}{ c c c } 7.6 \\ 12.0 \\ 20.0 \\ 2.5 \\ 7.6 \end{array}$	$7.6 \\ 11.3 \\ 20.4 \\ 2.2 \\ 6.2$	$587.0 \\ 591.3 \\ 599.5 \\ 579.8 \\ 585.3$	5 1 2 1 5	$ \begin{array}{r} 13 \\ 15 \\ 16 \\ 10 \\ 12 \end{array} $
A6 A7 A8 A9 A10		.514 .249 .142 .098 .234	.5255 .6195 .6540 .7153 .6269	.4236 .3726 .3409 .2818 .3659	0509 0079 0051 0029 0072	20 35 35 35 35 35	$ \begin{array}{c} 2.9\\ 7.4\\ 12.2\\ 20.2\\ 7.4 \end{array} $	$2.3 \\ 8.5 \\ 12.0 \\ 20.6 \\ 9.2$	$579.9 \\ 588.1 \\ 592.0 \\ 599.7 \\ 588.9$	$ \begin{array}{c} 1 \\ 2 \\ 5 \\ 2 \\ 2 \end{array} $	10 14 15 16 14
A11 A12 A13 A14 A15		$\begin{array}{c c} .154\\ .227\\ .640\\ .286\\ .203\end{array}$. 6863 . 6200 . 5002 . 5894 . 6330	.3098 .3736 .4339 .3996 .3596	.0040 .0064 .0659 .0110 .0075	35 35 24 35 35	12. 27. 02. 37. 612. 0	$ \begin{array}{c} 16.0 \\ 8.5 \\ 0.9 \\ 5.9 \\ 9.8 \end{array} $	596.1 588.1 578.3 585.0 589.6	$2 \\ 3 \\ 1 \\ 5 \\ 1$	16 14 10 11 14
A16 A17 A18 A19 A20		.338 .504 .305 .213 .082	. 5809 . 7233 . 5919 . 6116 . 7314	.4078 .2736 .3928 .3771 .2657	.0113 .0032 .0153 .0113 .0029	35 35 35 35 35	$\begin{array}{c} 7.1 \\ 20.2 \\ 7.1 \\ 7.6 \\ 20.0 \end{array}$	5.2 22.3 6.4 7.9 24.0	584. 1 600. 7 585. 5 587. 5 601. 7	5 1 1 1 1	11 16 12 13 16
A21 A22 A23 A24 A25	Refined cottonseed oils, re- ceived June 1915 from vari- ous manu ^f acturers	.170 .629 .404 .084 .188	.6612 .5108 .6010 .7151 .6569	.3326 .4314 .3908 .2821 .3388	.0062 .0578 .0083 .0028 .0043	35 20 35 35 35	$12.0 \\ 2.5 \\ 7.8 \\ 20.0 \\ 12.0$	$ \begin{array}{c} 12.9\\ 1.4\\ 6.8\\ 20.6\\ 12.2 \end{array} $	592. 9 578. 3 586. 1 599. 7 592. 3	1 1 3 2 3	15 10 12 16 15
A26 A27 A28 A29 A30		.690 .434 .682 .262 .346	.5244 .6065 .5246 .6496 .6128	.4351 .3855 .4313 .3449 .3789	.0405 .0080 .0441 .0055 .0083	20 35 20 35 35	$\begin{array}{c} 2.2 \\ 7.1 \\ 2.5 \\ 12.0 \\ 7.2 \end{array}$	1.8 7.3 2.0 11.5 7.9	579. 0 586. 7 579. 3 591. 5 587. 4	1 2 1 2 2	10 13 10 15 13
A31 A32 A33 A34 A35		$\begin{array}{c} .234\\ .405\\ .352\\ .166\\ .151\end{array}$. 6568 . 6075 . 6100 . 6583 . 6609	.3379 .3852 .3827 .3362 .3362 .3337	.0053 .0073 .0073 .0055 .0054	35 35 35 35 35 55	$ \begin{array}{c} 12.0\\ 7.6\\ 7.6\\ 12.0\\ 12.0\\ 12.0 \end{array} $	$ \begin{array}{c} 12.3 \\ 7.4 \\ 7.6 \\ 12.5 \\ 12.8 \end{array} $	592. 3 586. 8 587. 0 592. 5 592. 8	2 4 4 2 2	15. 13. 13. 15. 15.
A36 A37 A38 A39		.624 .417 .437 .378	.5320 .6023 .5999 .6024	.4323 .3892 .3912 .3893	. 0358 . 0085 . 0089 . 0083	20 35 35 35	2.5 7.1 7.1 7.1 7.1	$2.2 \\ 6.9 \\ 6.8 \\ 7.0$	579. 7 586. 3 586. 0 586. 3	1 2 2 3	10 12 12 12 12, 13
A41 A42 A43 A45 A46		.158 .118 .139 .279 .123	.6238 .7448 .6124 .6107 .6687	.3716 .2531 .3821 .3825 .3263	.0046 .0021 .0055 .0068 .0050	35 35 35 35 35 35	$ \begin{array}{c c} 8.9 \\ 24.0 \\ 7.9 \\ 7.6 \\ 12.7 \end{array} $	8.8 26.8 7.7 7.6 13.7	588.4 603.2 587.2 587.1 593.8	4 4 4 4 2	14 16 13 13 16

	Kind of oil or source			Lo	vibond	grade					
NBS lab. no.		Sun- light trans- mit- tance	Triline	By donor		By NBS on 35Y basis	Wave length	Spectral trans- mittance shown in			
		т.	r	g	b	Y	R	R	λ	Fig- ure	Fig- ure
1	2	3	4	5	6		7	8	9	10	11
B1 B2 B3 B4 B5		$(\begin{smallmatrix} 0.\ 175 \\ .\ 325 \\ .\ 324 \\ .\ 330 \\ .\ 225 \end{smallmatrix})$	0.6609 .6100 .6094 .6004 .5999	0. 3334 . 3803 . 3827 . 3913 . 3871	0.0057 .0097 .0079 .0084 .0131	35 35 35 35 35 35	7.8 7.0 7.2 6.7 6.3	12.8 7.7 7.6 6.8 7.0	592. 9 587. 2 587. 0 586. 1 586. 3	5 1 3 4 1	15 13 13 12 12
B6 B7 B8 B9 B10	Prime winter cottonseed oils	. 289 . 292 . 337 . 301 . 239	.5976 .6232 .6096 .6118 .6194	.3921 .3698 .3805 .3792 .3732	.0104 .0070 .0099 .0090 .0074	35 45 35 35 35	$\begin{array}{c} 6.5 \\ 7.2 \\ 7.0 \\ 6.3 \\ 7.1 \end{array}$	6.6 8.8 7.7 7.8 8.5	585.9 588.5 587.2 587.4 588.1	5 2 1 5 3	12, 13 14 13 13 14
B11 B12		.323	.6117 .6029	.3806 .3867	.0077	35 35	7.3 6.8	7.8 7.1	587.3 586.4	3 1	13 13
D1 D2 D3 D4 D5	Cottonseed oils	$\left(\begin{array}{c} .\ 009\\ .\ 151\\ .\ 386\\ .\ 422\\ .\ 365\end{array}\right)$.8388 .6063 .5884 .5822 .6115	$\begin{array}{r} .1612\\ .3854\\ .3957\\ .4072\\ .3810\end{array}$.0000 .0084 .0159 .0106 .0075	55 35 35 35	5.5 5.5 4.8 7.1	7.3 6.1 5.3 7.7	586.7 585.2 584.2 587.2	1 3 1 4 3	16 13 12 11 13
D6 D7 D8		$\left \begin{array}{c} .314\\ .288\\ .220\end{array}\right $.5967 .6176 .6135	. 3948 . 3753 . 3783	.0085 .0071 .0082	35 35	6.0 6.2	6.5 8.3 8.0	585.7 587.9 587.5	4 3 2	12 14 13, 14
E1 E2 E3 E4 E5	Ga Tenn Ala Ala Ala	.493 .050 .224 .410 .214	.5834 .7146 .6554 .6283 .6675	. 4065 . 2826 . 3407 . 3661 . 3287	.0101 .0028 .0039 .0057 .0038	35 35 35 35 35	5.423.012.79.314.1	5.420.512.09.213.5	584.3 599.6 592.1 589.0 593.6	3 3 4 4 3	$ \begin{array}{c} 11 \\ 16 \\ 15 \\ 14 \\ 16 \end{array} $
E6 E7 E8 E9 E10	N. C Ala Ala S. C Ga	.426 .299 .308 .439 .421	.5945 .6290 .6269 .5963 .5954	.3971 .3647 .3667 .3949 .3958	.0085 .0063 .0064 .0088 .0088	35 35 35 35 35	6.2 6.1	$\begin{array}{c} 6.3\\ 9.4\\ 9.2\\ 6.4\\ 6.4 \end{array}$	585.4 589.1 588.9 585.6 585.5	4 3 3 3 3	12 14 14 12 12
E11 E12 E13 E14 E15	Ga S. C S. C N. C Tex	.368 .411 .238 .297 .392	. 6030 . 6128 . 6316 . 6207 . 6003	.3894 .3806 .3617 .3725 .3914	.0076 .0066 .0067 .0068 .0084	35 35 35 35 35	7.57.58.08.17.6	$\begin{array}{c} 7.0 \\ 7.8 \\ 9.6 \\ 8.6 \\ 6.8 \end{array}$	586.3 587.3 589.4 588.2 586.1	$ \begin{array}{c} 4 \\ 3 \\ 2 \\ 3 \\ 3 \end{array} $	$12, 13 \\ 13 \\ 14 \\ 14 \\ 12$
E16 E17 E18 E19 E20	Ala N. C N. C Ga S. C	.336 .416 .396 .406 .396	.6072 .5916 .6002 .6018 .5944	.3847 .3994 .3915 .3903 .3969	.0081 .0091 .0083 .0080 .0088	35 35 35 35 35 35	5.7 7.5 7.0 6.0	$\begin{array}{c} 7.4 \\ 6.0 \\ 6.8 \\ 6.9 \\ 6.3 \end{array}$	586.8 585.2 586.0 586.2 585.4	333343 3	13 12 12 12 12
E21 E22 E23 E24	N. C Ga S. C Ala	.429 .374 .408 .336	. 5965 . 5946 . 5917 . 6073	. 3952 . 3970 . 3995 . 3844	. 0083 . 0085 . 0089 . 0083	35 35 35 35	$\begin{array}{c} 7.0 \\ 6.0 \\ 5.6 \\ 6.4 \end{array}$	$\begin{array}{c} 6.4 \\ 6.3 \\ 6.0 \\ 7.4 \end{array}$	585.6 585.4 585.1 586.8	4 4 4 2	12 12 12 13
E25 E26 E27 E28 E29 E30	Ala Ga Ga Ga S. C N. C	.295 .390 .337 .339 .240 .496	$\begin{array}{r} .\ 6163\\ .\ 5903\\ .\ 6038\\ .\ 6113\\ .\ 6295\\ .\ 5818\end{array}$.3764 .4014 .3879 .3814 .3637 .4080	.0074 .0083 .0083 .0073 .0068 .0102	35 35 35 35 35 35 35	7.4 6.2 7.2 8.0 5.5	$8.2 \\ 5.9 \\ 7.1 \\ 7.7 \\ 9.4 \\ 5.2$	587.7 585.0 586.4 587.2 589.2 584.1	$2 \\ 4 \\ 3 \\ 2 \\ 4$	14 11 13 13 14 11
E31 E32 E33 E34 E35	Ga S. C N. C Tex N. C	.486 .502 .460 .380 .369	. 5852 . 5856 . 5882 . 5957 . 6028	.4050 .4043 .4025 .3960 .3898	.0098 .0102 .0093 .0083 .0075	35 35 35 35 35	$5.2 \\ 5.5 \\ 5.5 \\ 6.8 \\ 7.2$	$5.5 \\ 5.6 \\ 5.8 \\ 6.4 \\ 6.9$	584. 5 584. 5 584. 8 585. 5 586. 3	4 2 4 4 4	11 11 11 12 12

TABLE 1.—Colorimetric properties of vegetable oils for a thickness of 5¼ inches— Continued

Journal of Research of the National Bureau of Standards [Vol. 15

116

NBS lab. no.	Kind of oil or sourse	and the second			Lo	ovibono	l grade	Wave length	Spectral trans- mittance shown in		
		Sun- light trans- mit- tance	Trilin	By donor		By NBS on 35Y basis					
		T.	r	g	b	Y	R	R	λ	Fig- ure	Fig- ure
		3	4	5	6		7	8	9	10	11
E36 E37 E38	Tex Ga Ala	0. 298 . 287 . 454	$\begin{array}{c} 0.\ 6096 \\ .\ 6135 \\ .\ 5918 \end{array}$	0.3827 .3790 .3993	0.0077 .0075 .0090	35 35 35	7.5 7.6 6.0	7.6 7.9 6.1	587.0 587.4 585.2	3 2 3	$ \begin{array}{c} 13 \\ 13 \\ 12 \end{array} $
R1 R2 R3 R4 R5	Cottonseed oils	$\left\{\begin{array}{c} .\ 232\\ .\ 405\\ .\ 398\\ .\ 289\\ .\ 261\end{array}\right.$	$\begin{array}{r} .\ 6473 \\ .\ 6032 \\ .\ 5896 \\ .\ 6124 \\ .\ 6008 \end{array}$.3468 .3880 .4022 .3794 .3887	$\begin{array}{r} .\ 0059\\ .\ 0088\\ .\ 0082\\ .\ 0082\\ .\ 0106\end{array}$			$ \begin{array}{r} 11.3\\ 7.0\\ 5.8\\ 7.9\\ 6.9 \end{array} $	$591. 2 \\586. 4 \\584. 9 \\587. 4 \\586. 2$	$2 \\ 1 \\ 4 \\ 1 \\ 1 \\ 1$	$ \begin{array}{c} 15 \\ 13 \\ 11 \\ 13 \\ 12 \end{array} $
R6 R7 R8		$\left \begin{array}{c} .484 \\ .376 \\ .340 \end{array} \right $. 6009 . 5970 . 6025	. 3902 . 3938 . 3887	.0089 .0092 .0088			6.8 6.5 7.0	$586.1 \\ 585.7 \\ 586.3$	$1 \\ 2 \\ 2$	$12 \\ 12 \\ 12, 13$
F1 F2 F3 F4 F5	Sesamedo Peanutdo Soybean	.103 .157 .544 .638 .230	$\begin{array}{r} .\ 6463 \\ .\ 6102 \\ .\ 5347 \\ .\ 5015 \\ .\ 6083 \end{array}$.3486 .3828 .4485 .4396 .3854	.0051 .0070 .0168 .0589 .0063	$\begin{array}{c} 60 \\ 52 \\ 25 \\ 17 \\ 35 \end{array}$	$10.7 \\ 7.0 \\ 1.5 \\ 1.0 \\ 7.0$	$11.\ 1 \\ 7.\ 6 \\ 1.\ 6 \\ 0.\ 7 \\ 7.\ 4$	$591.0 \\ 587.0 \\ 578.8 \\ 577.0 \\ 586.8 \\$	4 4 5 1 4	$15 \\ 13 \\ 10 \\ 10 \\ 13$
F6 F7 F8 F9 F10	do	.128 .082 .024 .439 .584	$\begin{array}{r} .\ 6308\\ .\ 6189\\ .\ 6344\\ .\ 5899\\ .\ 5525\end{array}$.3643 .3763 .3613 .3968 .4245	$\begin{array}{r} .\ 0049\\ .\ 0048\\ .\ 0042\\ .\ 0133\\ .\ 0230\end{array}$	65 90 	9.0 8.0	9.58.39.86.13.4	$589. 2 \\ 587. 9 \\ 589. 6 \\ 585. 2 \\ 581. 5$	$\begin{array}{c} 4\\ 4\\ 4\\ 1\\ 2\end{array}$	$14\\14\\14\\12\\10$
F11 F12 F13 F14	Rape S. olive C. olive Soybean	.423 .442 .100 .550	.5514 .5377 .5120 .5687	.4364 .4455 .4729 .4198	.0122 .0168 .0151 .0115			2.8 1.9 -0.2 4.2	580.7 579.2 575.1 582.7	5555	10 10 10 10

TABLE 1.—Colorimetric properties of vegetable oils for a thickness of 5¼ inches— Continued

VI. SOME ABRIDGED COLOR-GRADING METHODS

It may be recalled that the specifications of lightness and chromaticity for the vegetable oils, as recorded in table 1 and in figure 8, are all referred to a standard illuminant and colorimetric method, and the properties of the observer are rigidly specified. The method is based on a knowledge of the complete spectral-transmissive properties of the sample and requires extensive equipment and laborious computations to arrive at the desired results. It is essentially a method for the standardizing research laboratory and serves here as the fundamental basis for the discussion and design of abridged colorgrading methods for routine application in the field. In the design of such methods, it is generally desirable to avoid the use of a large number of material color standards and to eliminate, if possible, or reduce the effect of the individual-observer variation. Whereas the illuminant and colorimetric method may be readily controlled and standardized, in general the observer must necessarily vary, and his individual color-vision characteristics may introduce serious complications in the colorimetric procedure, depending on the degree of difference existing between the spectral-transmissive properties of the samples to be graded and the color standards defining the chromaticity This difference may exist regardless of the degree of agreescale.

ment in lightness or chromaticity obtainable between the samples and the standards of color.

It is beyond the scope of the present paper to give a detailed discussion of the application of the Lovibond method-or other abridged colorimetric methods-in the color grading of vegetable oils. The following remarks are included, however, on methods that have been investigated to some extent in connection with the present work, or that are obviously suggested by a consideration of the fundamental data herein presented. The former methods are one-dimensional by design. The latter methods are essentially two-dimensional, but may be reduced to one dimension as previously described, if the simplification is in conformity with practical requirements.

In an attempt to avoid all material color standards and effectively eliminate the properties of the observer, efforts were made during the early part of this work to discover a suitable one-number index to the chromaticity of the oils which could be expressed as a simple functional relationship between transmittances at selected wave lengths in the visible spectrum. An important condition imposed on the problem was the possibility of embodying the method in a practical oil-grading instrument measuring only transmittances at the selected wave lengths. In general, it was found that such abridged spectrophotometric methods were not wholly satisfactory, chiefly because they did not deal adequately with the independent variation of the concentrations of more than one important pigment in the oils. Unavoidable complexities in the experimental and computational aspects of the method made it doubtful if sufficient practical advantage was gained to warrant its routine application in the field.

An attempt was made by Priest 20 and the author to use a small number of selected samples of cottonseed oils as material color standards against which given cottonseed oils might be matched in chromaticity by the proper choice of their thickness alone. The thickness of the oil, or some suitable function thereof, could be used then as a one-number index to chromaticity. A color comparator was constructed in which the sample field was illuminated by a standard illuminant through a controlled variable thickness of the given sample of oil. The comparison field was illuminated by the same source through a fixed thickness of one of the oil standards. Several standards of different spectral type or thickness could be employed if desired. It was found that vacuum-sealed oils devoid of the green pigments were more stable in color than others, but none of the oils investigated proved sufficiently stable over a period of several months to serve as satisfactory color standards. The use of various yellow solutions instead of the oils, as color standards in this method, is suggested by the work of Arny, Kish, and Newmark,²¹ but no conclusive investigation has yet been made of the possibilities of such solutions in connection with this variation-in-thickness method.

The classification of the oils into two principal spectral types suggests a color-grading method based on the use of two standard solutions of fixed concentration and of proper spectral type to represent, respectively, the brown and green groups of pigment components in the oils. Light transmitted by variable thicknesses of the solutions

 ²⁰ Report of Proceedings of the Sixth Annual Convention of the Society of Cotton Products Analysts, Birmingham, p. 67 (May 1915); Cotton Oil Press p. 87 (July 1919).
 ²¹ J. Ind. Eng. Chem. **11**, 950 (1919).

(in series) would be compared with light transmitted by a fixed thickness of the oil. Then, by suitable choice of a fixed thickness of one solution with a variable thickness of the other, or by varying the relative and absolute thicknesses of both, either one- or two-dimensional chromaticity scales could be established, as desired. The latter variation corresponds precisely to the concentration variation in the vegetable oils. In view of the fact that all vegetable oils may be matched in chromaticity by suitable choice of yellow and red Lovibond glasses, it is certain that the double series of glass standards could be replaced by the two solution standards, which, because of a closer relation to the oil pigments in spectral-transmissive properties, would not only provide the same chromaticity matches but would also reduce the disturbing effect of the color-vision characteristics of the observer. Further investigation is required to discover suitable solutions for these standards which would be satisfactory with respect to stability, reproducibility, and spectral transmittance.

An inspection of the (r,g) diagram in figure 8 shows that a convenient and reproducible one-dimensional chromaticity scale is provided by the regular sequence of spectrum colors. From the reddest vegetable oils through the range of prime cottonseed oils the spectrum colors and the oil colors are nearly equal in chromaticity. This close agreement holds also for some very light oils of the F group. The evaluation of the oil colors on this wave-length scale has been described in section IV, and the λ values recorded in column 9 of table 1, for the standard observer on the OSA colorimetric system. Unfortunately, any attempt to incorporate this simple chromaticity scale into a practical color-grading method must seriously involve the color-vision characteristics of each different observer assigned to the color measurements. This condition follows from the wide difference in spectral-energy distribution existing between the light transmitted by the oils and that of a narrow wave-length band centered at the λ value required to obtain the nearest chromaticity match with the oils. It is known that under such conditions normal observers would vary in their individual λ values by amounts equivalent to 2 or 3 red units on the scale of 35Y + N''R. Hence, the use of this method would require a separate calibration of each observer making the color measurements. This would call for perhaps several glass or liquid filters standardized for the purpose and having spectral-transmissive properties approximating closely those of the oils. The added complication in the colorimetric procedure nullifies the apparent advantages of this simple basis for color grading.

It is also shown in figure 8 that a system of standard colors slightly less saturated than the pure spectrum colors is required to obtain exact chromaticity matches with all the oils. This condition could be realized by an extension of the method just described in which the pure spectrum colors would be diluted slightly by admixture with a variable amount of "white" light from the standard illuminant. A two-dimensional chromaticity scale is thus established by the independent variation of both wave length and purity, but the method is subject to the same pratical objections as the simpler method just described.

Another extension of the simple spectrum method is possible which would largely eliminate the variation in scale settings obtained by different observers. Instead of using a narrow band of wave lengths, alone or admixed with white, the chromaticity of an oil could

120 Journal of Research of the National Bureau of Standards [Vol 15

be accurately matched by use of a single wide band of wave lengths, covering roughly the same wave-length range as is transmitted by the oil itself.

In a colorimeter based on this principle, the comparison field would be illuminated by light from the standard illuminant after transmission through a selective light filter consisting essentially of the slitlens-prism arrangement of an ordinary monochromator used with exceptionally wide slits. A control of the widths, position, and form of the slits would provide a wide variation in form of the curve of spectral transmission for this filter. In particular, all wave lengths in the red could be transmitted, as is done by the oils, and all wave lengths shorter than a certain value cut off completely. The cut-off could be made abrupt or gradual, as desired, to simulate most closely the spectral-transmissive properties of the oils. It may be recalled that this is the condition required to reduce or eliminate from the method the variable color-vision characteristics of the observer, and may be satisfied in this method perhaps even better than is done by the use of Lovibond-glass combinations. The sample field of the colorimeter would be illuminated by the standard illuminant through a fixed thickness of oil, and independent means provided for equalizing the brightnesses of the sample and comparison fields.

The chief advantage of this method lies in the avoidance of all reference to material color standards. Again, as in other color-grading methods herein considered, one- or two-dimensional chromaticity scales could be established, as desired, the scale variables being related to instrumental parameters involving the position, width, or form of the slits. Considerable research and development work would be necessary to determine the best procedure in the practical application of the method.

In regard to all color-grading methods, it may be noted that scale variables in one method are always convertible into corresponding scale variables in another method. Such conversions are illustrated in the present paper by the transformation of scale values in the OSA system to values on the special N'' scale and the λ scale. In accordance with the single or double dimensionality of the scales, we deal respectively with nearest or exact chromaticity matches. Hence, it is always possible, upon the adoption of any improved method for the color grading of vegetable oils, to supply conversion tables along with the method, or perhaps to calibrate the scale of the colorimeter itself, so that results could be given in Lovibond terms until it is deemed desirable to use only the terms suited to the revised grading system itself.

VII. SUMMARY AND CONCLUSIONS

1. The spectral-transmittance and colorimetric properties of 125 vegetable oils are presented in tabular and graphical form. The investigation includes 111 cottonseed, 3 peanut, 5 soybean, 2 sesame, 1 rape, 1 corn, and 2 olive oils.

2. The oils in a 5¼ inch thickness, may be classified roughly into two general spectral types, determined, respectively, by the varying concentrations of two principal groups of pigment components. One group includes the green materials derived from chlorophyll; the other group includes perhaps several yellow or amber-colored materials of unknown identity. 3. With respect to spectral-transmissive and colorimetric properties, there is no essential difference between the cottonseed and the other vegetable oils included in this investigation. Hence the problem of color grading over the range of cottonseed oils apparently encompasses that for all vegetable oils.

4. The spread in lightness over the group of vegetable oils is greater than can be conveniently matched in the present method of color grading, in which the number of clear and colored glasses used with the Lovibond tintometer is varied while maintaining the best chromaticity match in the fields of the instrument. An improved color comparator is desirable, providing a more suitable arrangement for equalizing brightnesses of the sample and comparison fields.

5. The independent variation in relative and absolute concentrations of the several pigments present in the oils, makes it impossible to represent all oils accurately on any one-dimensional chromaticity scale. For accurate color grading in the Lovibond system, this situation demands the introduction into trade practice of a twodimensional Lovibond scale established by the independent variation of both yellow and red. It is shown that the best yellow numeral varies approximately from 20 for very light cottonseed oils to 80 or above for the prime yellow oils. An improved one-dimensional Lovibond scale would be provided by the fixation of yellow in some definite relation to red.

6. A brief discussion is given of some color-grading methods based on the following principles:

(1) The measurement of transmittances at a few selected wave lengths and the combination of these quantities in suitable relation to one another to yield a satisfactory one-number index to chromaticity.

(2) The grading in terms of a variable thickness of a sample oil required to obtain a nearest chromaticity match with a fixed thickness of a standard oil or standard solution.

(3) The grading of a fixed thickness of the sample oil in terms of the variable thicknesses of two standard solutions of fixed concentration (in series), representing, respectively, the brown- and green-pigment groups in the oils.

(4) The use of a variable light filter constructed on the principle of the well-known monochromator, but provided with special control of the position, width, and form of the slits, so that the spectral transmission of the filter could be controlled and made to conform as closely as possible to the spectral transmission of the oils to be graded.

7. Some experimental work is described on the application of methods (1) and (2) to cottonseed oils. It was found, however, that the oils themselves were not sufficiently stable in color to serve as satisfactory color standards. Other possible standards were not investigated. Both methods were deemed unsuitable for routine commercial application, however, chiefly because of complications involving the independent variation in concentration of more than one important pigment in the oils. Each of these methods was designed to establish a practical one-dimensional chromaticity scale. Methods (3) and (4) provide the possibility of either one or twodimensional scales for all oils, but no experimental work has been done on the methods. They are suggested by the fundamental data presented in the paper.

WASHINGTON, May 10, 1935.