

Nuclear Optical Model Analysis of Neutron Elastic Scattering for Calcium*

Randall S. Caswell

(May 14, 1962)

A nuclear optical model calculation of neutron elastic scattering using five parameters has been made. Appropriate estimates of the effect of compound elastic scattering at low energies are included. Comparisons have been made with experimental data, and results for the parameters adopted are given for elastic scattering in the energy range from 0.734 to 18 Mev. Although reasonably good agreement between the calculation and experimental results have been obtained, there is a discrepancy between experiments below 2 Mev, and experimental data is needed between 6 and 14 Mev. Both angular distributions and Legendre expansion coefficients for the total elastic scattering are given.

1. Introduction

The nuclear optical model has been outstandingly successful in describing the elastic scattering of neutrons and other nuclear particles above the energy of perhaps 6 Mev, where compound elastic scattering processes are not important. Below these energies it is necessary to include some estimate of compound elastic scattering. At neutron energies below the first excited state of the target nucleus, inelastic scattering is not possible. In this energy range it is possible to describe the compound elastic scattering by use of the fluctuation cross section [Feshbach, 1960]. Above this energy the angular distribution of the compound elastic scattering may be obtained from the compound statistical model [Hauser and Feshbach, 1952].

Initially in the present work the experimental data was analyzed using the six-parameter nuclear optical model of Bjorklund and Fernbach [1958]. However, it was found that at least as good fits to the experimental data could be obtained with a five-parameter model, using derivative surface absorption instead of an adjustable parameter to determine the width of the imaginary (absorptive) well. This latter model, which contains a Saxon real potential, a Thomas spin-orbit potential, and a surface imaginary potential, was used in this calculation.

This calculation is the first of a series which will provide data in a form suitable for input to neutron penetration calculations with digital computers. The present calculations are for calcium, an important component of concrete which is widely used for neutron shielding. Comparisons with experimental elastic scattering angular distributions are included. Angular distributions and Legendre coefficients for the calculated angular distributions are both presented. The calculations have been programmed in Fortran for the IBM 7090 computer at the National Bureau of Standards.

*This work was sponsored by Defense Atomic Support Agency.

2. Description of the Calculation

2.1. Solution of the Wave Equation

Although the exact phase shift analysis carried out here is standard in many respects, it seems desirable to record the specific method of calculation, including the Legendre polynomial analysis.

We start with the radial part of the non-relativistic Schrödinger wave equation:

$$\frac{d^2 U_l(r)}{dr^2} + \frac{2m}{\hbar^2} \left[E - \frac{l(l+1)\hbar^2}{2mr^2} - (V(r) + iW(r)) \right] \times U_l(r) = 0. \quad (1)$$

The potential is described by $V(r) + iW(r)$ where $V(r)$ is the real part of the potential and $W(r)$ is the imaginary part. The wave function $U_l(r)$ will also have both real and imaginary parts:

$$U_l(r) = X_l(r) + iY_l(r). \quad (2)$$

Equation (1) becomes:

$$\frac{d^2 (X_l + iY_l)}{dr^2} + \frac{2m}{\hbar^2} \left[E - \frac{l(l+1)\hbar^2}{2mr^2} - V - iW \right] \times (X_l + iY_l) = 0. \quad (3)$$

Separating this equation into real and imaginary parts, we obtain two coupled real equations which may be solved on the computer.

$$\frac{d^2 X_l}{dr^2} + \frac{2m}{\hbar^2} \left[E - \frac{l(l+1)\hbar^2}{2mr^2} - V \right] X_l + \frac{2m}{\hbar^2} W Y_l = 0 \quad (4)$$

$$\frac{d^2 Y_l}{dr^2} + \frac{2m}{\hbar^2} \left[E - \frac{l(l+1)\hbar^2}{2mr^2} - V \right] Y_l - \frac{2m}{\hbar^2} W X_l = 0. \quad (5)$$

These equations are integrated step-by-step outward in radius by the second order Runge-Kutta method [see, for example, Zurmühl, 1961]. The number of

integration steps is typically 100 to 500. If the wave functions become large, they are automatically renormalized to smaller values by the code. The running time for 15 l 's and 200 steps is about 1 min per energy including all angular distributions. The mass appearing in these equations is the reduced mass of the neutron in the neutron-nucleus center-of-mass system. The energy, E , is that of the neutron in the center-of-mass system. When the complex potential ($V+iW$) approaches a non-vanishing constant at the origin, then

$$U_l(r) \rightarrow C_l r^{l+1} \text{ as } r \rightarrow 0 \quad (6)$$

where C_l is a complex number independent of r [Amster, Leshan, and Walt, 1960]. In the presence of the spin-orbit term (see below) there is a $1/r$ behavior of the potential energy for small values of r ; however, eq (6) still holds since this term is dominated by the $1/r^2$ dependence of the centrifugal potential [see Bohm, 1951]. Since the cross sections depend only on the logarithmic derivative of the wave function, the normalization is arbitrary and C_l can be taken to be any non-zero value. We have chosen $k(1+i)$ for C_l where k is a small number chosen to prevent underflow or overflow problems. Provision is made in the code to calculate for values of l up to $l=29$.

For the optical model parameters we have used a potential whose real and imaginary parts are:

$$V(r) = V_c \rho(r) - \alpha V_c \left(\frac{\hbar}{2Mc} \right)^2 (\vec{l} \cdot \vec{\sigma}) \frac{1}{r} \frac{d\rho}{dr}, \quad (7)$$

$$W(r) = W_s \left(4a \frac{d\rho}{dr} \right) = W_s \left(4\rho^2 \exp \left[\frac{r-R_0}{a} \right] \right) \quad (8)$$

where

$$\rho(r) = \frac{1}{1 + \exp \left[\frac{r-R_0}{a} \right]}$$

the usual "Saxon" potential; α is the strength of the spin-orbit interaction relative to the "Thomas" term for a nucleon; M is the neutron mass; V_c is the central real potential; W_s is the lowest value of the imaginary surface potential; $\frac{\hbar}{Mc}$ is the Compton wavelength for a nucleon; \vec{l} is the orbital angular momentum of the neutron; $\vec{\sigma}$ is the Pauli spin operator of the incident neutron; R_0 is the nuclear radius; and a is the "diffuseness" parameter of the real potential. If the spin angular momentum of the incident neutron is parallel to the orbital angular momentum of the incident neutron, then $j=l+1/2$, and $\vec{\sigma} \cdot \vec{l} = l$. If the spin and orbital angular momentum are anti-parallel, then $j=l-1/2$ and $\vec{\sigma} \cdot \vec{l} = -(l+1)$. This leads to two real potentials, $V^+(r)$ and $V^-(r)$ corresponding to the parallel and anti-parallel $\vec{\sigma} \cdot \vec{l}$ interactions. The wave equation is solved for each of these. To obtain the desired

cross sections, we need the phase shifts η_l^+ and η_l^- which measure the effect of the complex potential on the neutron partial waves. These phase shifts are obtained by a calculation which compares the value for logarithmic derivative f_l obtained by numerical integration in the presence of the nuclear potential with the corresponding quantity ($\Delta_l + iS_l$) for the zero phase shift spherical Hankel function solutions obtained in the absence of the nuclear potential.

The required expression for η_l is given by Blatt and Weisskopf [1952]:

$$\eta_l = \frac{f_l - \Delta_l + iS_l}{f_l - \Delta_l - iS_l} \exp(2i\xi_l). \quad (9)$$

For neutrons

$$\exp(2i\xi_l) = \frac{G_l(R) - iF_l(R)}{G_l(R) + iF_l(R)}, \quad (10)$$

where $G_l(R)$ and $F_l(R)$ are solutions of eq (1) with $(V(r) + iW(r)) = 0$. In terms of spherical Bessel functions, $F_l(x) = xj_l(x)$ and $G_l(x) = xn_l(x)$ where $x = kR$.

2.2. Evaluation of the Cross Sections Directly From the Nuclear Optical Model

The total cross section predicted by the nuclear optical model is made up of a shape-elastic scattering cross section and a compound nucleus formation cross section [Feshbach, Porter, and Weisskopf, 1954]:

$$\sigma_t = \sigma_e + \sigma_c.$$

However, the compound nucleus may decay into the entrance channel producing "compound elastic" scattering or it may produce inelastic scattering or charged particle reactions:

$$\sigma_c = \sigma_{ce} + \sigma_r.$$

These equations may now be rearranged to separate the elastic scattering (σ_{se} and σ_{ce} are not distinguishable experimentally) from the reaction cross section:

$$\sigma_t = (\sigma_{se} + \sigma_{ce}) + \sigma_r = \sigma_e + \sigma_r.$$

In the present calculation σ_{ce} is estimated by the "fluctuation" cross section, σ_{fl} , which is important below energies of about 6 Mev:

$$\sigma_t = \sigma_{se} + \sigma_{fl} + \sigma_r.$$

When inelastic scattering and reactions are zero or negligible, the entire cross section is elastic:

$$\sigma_t = \sigma_{se} + \sigma_{fl}.$$

In this case the fluctuation cross section may be obtained very simply from the phase shifts of the optical model alone (see Feshbach [1960] for a discussion of the fluctuation cross section).

We now proceed to evaluate the various cross sections from the calculated phase shifts (η_i 's). For both incident particles and the nucleus without spin, the expressions for the incident plane wave, the total wave (which shows the effect of the potential of the nucleus on the incident wave), and the scattered wave are:

$$e^{ikz} = \sum_{l=0}^{\infty} i^l \frac{2l+1}{2} [h_l^*(kr) + h_l(kr)] P_l(\cos \theta), \quad (11)$$

$$\psi_{\text{tot}}(r) = \sum_{l=0}^{\infty} i^l \frac{2l+1}{2} [h_l^*(kr) + \eta_l h_l(kr)] P_l(\cos \theta), \quad (12)$$

and

$$\begin{aligned} \psi_{\text{scatt}}(r) &= \psi_{\text{tot}}(r) - e^{ikz} \\ &= \sum_{l=0}^{\infty} i^l \frac{2l+1}{2} (\eta_l - 1) h_l(kr) P_l(\cos \theta). \end{aligned} \quad (13)$$

The incident neutrons which have spin 1/2 may be represented by a plane wave:

$$\psi_{\text{inc}} = e^{ikz} \chi_{\text{inc}} \quad (14)$$

where χ_{inc} describes the spin wave function of the neutron. $\chi_{\text{inc}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is spin "up," $\chi_{\text{inc}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ corresponds to spin "down." In the presence of the nuclear potential the asymptotic form of the wave function of the neutron is:

$$\psi \xrightarrow{r \rightarrow \infty} \psi_{\text{inc}} + \frac{1}{r} e^{ikr} F(\theta) \chi_{\text{inc}} \quad (15)$$

where $F(\theta)$ is the scattering matrix.

$$F(\theta) = A(\theta) + B(\theta) \vec{\sigma} \cdot \vec{n} \quad (16)$$

where \vec{n} is a unit vector perpendicular to the plane of scattering. According to the Basel convention [Huber and Meyer, 1961], the positive direction of polarization, given by the unit vector \vec{n} , is:

$$\vec{n} = \frac{\vec{k}_i \times \vec{k}_f}{|\vec{k}_i \times \vec{k}_f|} \quad (17)$$

where \vec{k}_i and \vec{k}_f are the wave vectors of the incident and scattered neutrons.

In the equations above, $h_l^* = j_l - in_l$ and represents an incoming spherical wave and $h_l = j_l + in_l$ and represents an outgoing spherical wave. To extend this treatment to the case of neutrons with spin incident on spin-zero nuclei [Lepore, 1950], we introduce the operators π_l^+ and π_l^- which select the states for which $j = l + \frac{1}{2}$ and $j = l - \frac{1}{2}$, respectively. These operators are

$$\pi_l^+ = \frac{l+1 + \vec{\sigma} \cdot \vec{l}}{2l+1} \quad \text{and} \quad \pi_l^- = \frac{l - \vec{\sigma} \cdot \vec{l}}{2l+1}. \quad (17a)$$

If $j = l + \frac{1}{2}$, $\pi^+ = 1$, $\pi^- = 0$. If $j = l - \frac{1}{2}$, $\pi^+ = 0$, $\pi^- = 1$. Applying our operators to the total wave function, eq (16) we have

$$\begin{aligned} \psi_{\text{tot}} &= \sum_{l=0}^{\infty} \frac{2l+1}{2} i^l [\pi^+ + \pi^-] [h_l^*(kr) + \eta_l h_l(kr)] \\ &\quad \times P_l(\cos \theta) \chi_{\text{inc}}. \end{aligned} \quad (17b)$$

Let η_l^+ and η_l^- be phase shifts corresponding to the parallel and anti-parallel spin-orbit orientations for the neutron. At large radii we use the asymptotic form of the Hankel functions,

$$\begin{aligned} \psi_{\text{tot}} &= \sum_{l=0}^{\infty} \frac{(2l+1)i^l}{2kr} \left\{ \pi_l^+ \left(\eta_l^+ e^{i \left[kr - (l+1) \frac{\pi}{2} \right]} \right. \right. \\ &\quad \left. \left. + e^{-i \left[kr - (l+1) \frac{\pi}{2} \right]} \right) + \pi_l^- \left(\eta_l^- e^{i \left[kr - (l+1) \frac{\pi}{2} \right]} \right. \right. \\ &\quad \left. \left. + e^{-i \left[kr - (l+1) \frac{\pi}{2} \right]} \right) \right\} \times P_l(\cos \theta) \chi_{\text{inc}}. \end{aligned} \quad (17c)$$

If we now subtract the incident wave from the total wave to obtain the scattered wave, we find

$$\begin{aligned} \psi_{\text{scatt}} &= \sum_{l=0}^{\infty} \frac{e^{ikr}}{2ikr} \left[(l+1)(\eta_l^+ - 1) + l(\eta_l^- - 1) \right. \\ &\quad \left. + (\eta_l^+ - \eta_l^-) \vec{\sigma} \cdot \vec{l} \right] P_l(\cos \theta) \chi_{\text{inc}}, \end{aligned} \quad (18)$$

where we have used the expressions (17a) for the operators and the condition that when $\eta_l^+ = \eta_l^- = 1$, ψ_{scatt} must equal zero. The right-left asymmetry in the scattering of a polarized beam is introduced through the $(\vec{\sigma} \cdot \vec{l})$ operator:

$$(\vec{\sigma} \cdot \vec{l}) P_l(\cos \theta) = \vec{\sigma} \cdot \left[\vec{n} \left(-i \frac{\partial}{\partial \theta} \right) P_l(\cos \theta) \right]. \quad (19)$$

However, $\frac{\partial}{\partial \theta} = -\sin \theta \frac{\partial}{\partial(\cos \theta)}$. Finally,

$$(\vec{\sigma} \cdot \vec{l}) P_l(\cos \theta) = (\vec{\sigma} \cdot \vec{n}) \left[i \sin \theta \frac{\partial}{\partial(\cos \theta)} P_l(\cos \theta) \right], \quad (20)$$

$$(\vec{\sigma} \cdot \vec{l}) P_l(\cos \theta) = (\vec{\sigma} \cdot \vec{n}) i P_l^i(\cos \theta). \quad (21)$$

where $P_l^i(\cos \theta) = \sin \theta \frac{\partial}{\partial(\cos \theta)} P_l(\cos \theta)$. Note: a $(-1)^m$ term is sometimes included in the definition of the associated Legendre function $P_l^m(\cos \theta)$, in which case our function has the opposite sign, and a minus sign appears on the right side of eq (21). $A(\theta)$ and $B(\theta)$ appearing in the scattering matrix, eq (16) are:

$$A(\theta) = \frac{1}{2ik} \sum_{l=0}^{\infty} [(l+1)(\eta_l^+ - 1) + l(\eta_l^- - 1)] P_l(\cos \theta) \quad (22)$$

$$B(\theta) = \frac{1}{2k} \sum_{l=1}^{\infty} [\eta_l^+ - \eta_l^-] P_l^1(\cos \theta). \quad (23)$$

The differential scattering cross section is

$$\frac{d\sigma}{d\Omega} = \frac{1}{2} \text{tr} F^+ F = |A|^2 + |B|^2. \quad (24)$$

The polarization is given by

$$P = \frac{2 \text{Re}(A^* B)}{|A|^2 + |B|^2}. \quad (25)$$

For a completely polarized incident beam with spin "up," $P = (L - R)/(L + R)$, where L and R are the measured left and right detector counting rates.

The total, compound nucleus formation, and shape elastic cross sections (integrated over angle) are:

$$\sigma_i = \frac{\pi}{k^2} \sum_{l=0}^{\infty} [(l+1) 2 \text{Re}(1 - \eta_l^+) + l 2 \text{Re}(1 - \eta_l^-)] \quad (26)$$

$$\sigma_c = \frac{\pi}{k^2} \sum_{l=0}^{\infty} [(l+1)(1 - |\eta_l^+|^2) + l(1 - |\eta_l^-|^2)] \quad (27)$$

$$\sigma_{se} = \frac{\pi}{k^2} \sum_{l=0}^{\infty} [(l+1)|1 - \eta_l^+|^2 + l|1 - \eta_l^-|^2]. \quad (28)$$

For use in inelastic scattering calculations and in our estimate of compound elastic scattering, the "transmission coefficients" are also calculated:

$$T_l^+ = (1 - |\eta_l^+|^2) \text{ and } T_l^- = (1 - |\eta_l^-|^2). \quad (29)$$

2.3. Compound Elastic Scattering

Whereas shape elastic scattering is given directly by the nuclear optical model, our estimate of compound elastic scattering is treated in four different ways depending upon the incident neutron energy. Below 3.35 Mev, it may be obtained from the optical model phase shifts, with a small correction for the




FIGURE 1. Low-lying energy levels of Ca^{40} .

Spins and parities in parentheses are taken from Troubetzkoy, Kalos, Lustig, Ray, and Trupin [1961]. Levels taken as a continuum above 5 Mev.

observed (n, p) cross section. In the energy range from 3.35 to 5.0 Mev, a Hauser-Feshbach [1952] hand calculation using discrete energy levels of the target nucleus (see fig. 1) was made. From 5.0 to 5.7 Mev a Hauser-Feshbach calculation using a statistical-model residual nucleus was made. In this case the compound elastic scattering is isotropic because of the randomness of the spins and parities of the levels assumed in the statistical model. The angular isotropy has been proved by Wolfenstein [1951] and by Hauser and Feshbach. Above 6 Mev compound elastic scattering is neglected.

The fluctuation cross section, in the absence of inelastic scattering and reactions, may be obtained from the phase shifts of the optical model [Feshbach, 1960]:

$$|A|_{j_l}^2 = \frac{1}{4k^2} \sum_{l=0}^{\infty} \{ (l+1)^2 (1 - |\eta_l^+|^2)^2 + l^2 (1 - |\eta_l^-|^2)^2 \} \times [P_l(\cos \theta)]^2$$

$$|B|_{j_l}^2 = \frac{1}{4k^2} \sum_{l=1}^{\infty} \{ (1 - |\eta_l^+|^2) + (1 - |\eta_l^-|^2) \} [P_l^1(\cos \theta)]^2. \quad (30)$$

At energies below the energy of the first excited state in the target nucleus (3.35 Mev for Ca^{40}), we have used eq (30) to calculate the compound elastic scattering. Note that this leads not to isotropic scattering but to an angular distribution which goes predominantly as $[P_l(\cos \theta)]^2$ for neutrons of angular momentum l .

The cross section for inelastic scattering of a neutron of incident angular momentum l , final angular momentum l' , from a nucleus of initial angular momentum i and final angular momentum i' is given by Hauser and Feshbach as

$$\sigma(l, i|l', i') = \frac{1}{2(2i+1)} \sum_{\alpha, \beta} \sigma(l, j_\alpha|l', j'_\beta|\theta) \quad (31)$$

where

$$\sigma(l, j|l', j'|\theta) = \frac{\pi}{k^2} (2l+1) T_l(E) \sum_{p, q, r} \frac{A_J(l, j|l', j'|\theta)}{1 + \sum_{p, q, r} T_p(E'_q)/T_l(E')}, \quad (32)$$

and is the cross section for production of neutrons of energy E' of angular momentum l' , channel spin j' , moving in a direction θ . The r index refers to possible channel spins, p to possible final neutron angular momenta, E'_q to final neutron energies. J is the total angular momentum of the compound nucleus state. The prime on the summation indicates that we omit the term referring to the final neutron E', l', j' we are considering. As we shall see the denominator of eq (32) is essentially the branching ratio of the specific neutron transition under consideration versus all other possibilities. If we immediately specialize to the case of compound elastic scattering, ($E = E', l = l', j = j'$), consider spin zero nuclei only ($j = j' = \frac{1}{2}, i = 0$, and $J = l \pm \frac{1}{2}$), and T_l 's as defined in eq (29), the angular distribution for the l th partial wave of compound elastically scattered

neutrons becomes:

$$\left(\frac{d\sigma}{d\Omega}\right)_{l_{CE}} = \frac{\pi}{2k^2} \left\{ \frac{(l+1)T_l^+(E)}{1 + \sum_{p,q,r}' T_p^+(E'_q)/T_l^+(E)} A_{l+\frac{1}{2}} \left(l, \frac{1}{2} \middle| l, \frac{1}{2} \middle| \theta \right) + \frac{lT_l^-(E)}{1 + \sum_{p,q,r}' T_p^+(E'_q)/T_l^-(E)} A_{l-\frac{1}{2}} \left(l, \frac{1}{2} \middle| l, \frac{1}{2} \middle| \theta \right) \right\}. \quad (33)$$

The A -functions that occur in eq (33) are easily expressed in terms of Z -coefficients used by Blatt and Biedenharn [1952] and tabulated by Biedenharn [1953]:

$$A_J(l, j|l, j|\theta) = \frac{1}{4\pi(2l+1)} \sum_L [Z(lJlJ; jL)]^2 P_L(\cos \theta). \quad (34)$$

The question of phases does not enter since the Z -coefficient appears only when squared. Note that these results appear naturally in a Legendre series expansion which may be easily combined with the Legendre coefficient expansion as determined by the computer for the shape elastic scattering. Equations (33) and (34) are used to calculate the region of discrete states of the target nucleus (3.35 to 5 Mev). A correction to the branching ratio appearing in eq (33) is applied to allow for experimentally observed reaction cross sections. From 5 to 5.7 Mev, the residual nucleus is considered as statistical, and the compound elastic scattering is isotropic.

2.4. Legendre Coefficient Analysis of the Elastic Scattering

The elastic scattering angular distribution is represented in the following way:

$$\frac{d\sigma}{d\Omega}(E, \theta) = \frac{\sigma_{\text{total elastic}}}{4\pi} \sum_l (2l+1) S_l(E) P_l(\cos \theta) \quad (35)$$

where $S_0(E) = 1$.

The differential cross section as obtained from eq (24) appears as a sum of the products of two series:

$$\frac{d\sigma}{d\Omega} = \sum_l A_l P_l(\cos \theta) \sum_m A_m P_m(\cos \theta) + \sum_l B_l P_l(\cos \theta) \sum_m B_m P_m^1(\cos \theta). \quad (36)$$

The first term in eq (36) may be represented by a single series of Legendre coefficients

$$\frac{d\sigma}{d\Omega} = \sum_k a_k P_k(\cos \theta), \quad (37)$$

where a_k is given by

$$a_k = \sum_{l=0} \sum_{m=0} A_l B_m \frac{C_{\frac{1}{2}(l-m+k)} C_{\frac{1}{2}(l+m-k)} C_{\frac{1}{2}(m-l+k)}}{C_{\frac{1}{2}(l+m+k)}} \times \frac{2k+1}{l+m+k+1}, \quad (38)$$

where $C_j = 1.3.5 \dots (2j-1)/j!$; l, m are positive integers; $(l+m+k)$ is even; and $|l-m| \leq k \leq l+m$, that is, the three vectors satisfy the "triangle condition" [Whittaker and Watson, 1958].

The product of the two series in $P_l^i(\cos \theta)$ can be reduced to product of two series in $P_l(\cos \theta)$ which can then be solved by eq (38). First we express the $P_l^i(\cos \theta)$'s in terms of polynomial expansions in $P_l(\cos \theta)$ [see Morse and Feshbach, 1953]:

$$P_l^1(\cos \theta) = \frac{l(l+1)}{(2l+1) \sin \theta} [P_{l-1}(\cos \theta) - P_{l+1}(\cos \theta)] \quad (39)$$

$$P_m^1(\cos \theta) = \sin \theta [(2m-1)P_{m-1}(\cos \theta) + (2m-5)P_{m-3}(\cos \theta) + (2m-9)P_{m-5}(\cos \theta) + \dots] \quad (40)$$

Any product term $P_l^1(\cos \theta)P_m^1(\cos \theta)$ resulting from the series product in eqs (36) may be expressed in terms of a product of expressions from eqs (39) and (40). Note that in the product $\sin \theta$ cancels out, leaving only terms in Legendre polynomials. Relations (38 to 40) were programmed for the computer so that a Legendre coefficient analysis of the form of eq (35) could be directly obtained from the calculated phase shifts.

The calculation could equally well be done through use of vector-coupling coefficients [Edmonds, 1957]. It is expected that the problem for the computer would be about the same either way. The C_j coefficients used here are closely related to the Clebsch-Gordan coefficient ($ab00|abc0$) as given in eq (5) of Blatt, Biedenharn, and Rose [1952].

3. Choice of Optical Model Parameters and Comparison With Experiment

Calcium is composed of 96.97 percent Ca^{40} with 2.06 percent of Ca^{44} , and smaller amounts of Ca^{42} , Ca^{43} , Ca^{45} , and Ca^{46} . In this calculation it is considered as pure Ca^{40} except that a slightly larger value of R_0 is used to approximate the effect of the other isotopes.

In the initial attempts to fit calcium neutron cross section data, the model of Bjorklund and Fernbach [1958] was used. The real potential is as described by eq (7), and the imaginary potential is a Gaussian absorption located at the surface:

$$W = W_s \exp \left[- \left(\frac{r-R_0}{b} \right)^2 \right]. \quad (41)$$

This model, which has been quite successful in fitting neutron elastic scattering data at 4.1, 7, and 14 Mev, involves six parameters: R_0 , V_c , α , a , W_s , and b . In attempting to fit this data, R_0 was held constant at $1.25 A^{1/3}$ fermis, b at 0.980 fermis, α at 35. The average atomic weight was used for A . The other three parameters were varied. A problem of the optical model is the large number of arbitrary parameters. In an attempt to reduce the number of parameters, the width of the surface absorption, b , was eliminated as an independent parameter and derivative surface absorption used instead as given in eq (8). Somewhat surprisingly the agreement with the experimental data was considerably improved at 14.6 Mev, and was equally good at the energies of 2 Mev and below where data existed which could be directly compared with optical model predictions using the fluctuation cross section. The improved agreement at 14.6 Mev may be stated in another way; namely, the value of b (0.980 fermis) is too small at this energy, and a thicker shell of surface absorption of neutrons should be used. This is given automatically by the use of a derivative surface absorption.

Using the derivative surface absorption model, R_0 was held constant at $1.25A^{1/3}$ fermis, the value indicated by the work of Bjorklund and Fernbach, and also by the optical model analyses of proton polarization reported by Gursky and Stewart [1961]. Actually the parameters V_c and R_0 act very much like a single parameter $V_c R_0^n$ where n is equal to 2 at low energies and increases slightly with energy. Our choice is to hold R_0 fixed and vary V_c . The strength of the spin-orbit coupling, α , was also held at the value of 35 used by Bjorklund and Fernbach because it was felt that changes or improvements in this value should come from the analysis of polarization data to which α is more sensitive, rather than angular distribution data which is considered here. Three parameters were allowed to vary, V_c , W_s , and a . Although a could have been held fixed, independent of energy, at a value of about 0.580 fermis to obtain fairly good fits to the experimental data everywhere, it was found that better fits were obtained if a was taken as 0.600 fermis at 14.6 Mev and at 0.550 fermis below 2 Mev. Smooth curves drawn through the data were used to obtain parameters for the optical model calculations, even though local variations sometimes gave better fits (see fig. 2). This apparent jumping around of the experimental data is due at the low energies to the presence of resonances in the cross section, which the optical model averages over.

It should also be mentioned that a calculation which predicts the energy variation of the parameters of the optical model by use of a nonlocal potential has been made by Perey and Buck [1961]. The predictions of Perey and Buck have not been used here since it seemed unlikely that they would improve the cross section data.

Experimental data used for the parameter determination at 14.6 Mev were taken from Cross and Jarvis [1960]; below 2 Mev, the data of Seagrave, Cranberg, and Simmons [1958] and Lane, Langs-

dorf, Monahan, and Elwyn [1960] as collected by Howerton [1961] were used. Comparison of the calculation to experimental data is shown in figures 3 through 12. Comparison is also made to data at




FIGURE 2. Optical model parameters versus energy as adopted for the calculation.




FIGURE 3. Elastic scattering at 14.6 Mev.

Experimental data are those of Cross and Jarvis [1960]. The curves in this and succeeding figures are from this calculation using the optical model parameters of figure 2.




FIGURE 4. *Elastic scattering at 6.0 Mev.*

Experimental data are from Seagrave, Cranberg, and Simmons [1958].

4.1 Mev of Vincent, Morgan, and Prud'homme [1960] and to an angular distribution of Seagrave, Cranberg, and Simmons [1958] at 6.0 Mev. Neither of these distributions was used to obtain the optical model fit because the Hauser-Feshbach hand calculation involved made parameter fitting quite difficult (it later turned out that at 6.0 Mev the compound elastic scattering could be neglected). Nevertheless the agreement between calculation and experiment is about the same here as at the other energies. We may therefore be reasonably confident that the calculation predicts the cross sections quite well where they have not been measured.

It would not be possible for these calculations to agree with both the Seagrave et al. and the Lane et al. experimental measurements, since these disagree with each other. In general the angular distributions of Lane et al. are more isotropic, while the distributions of Seagrave et al. have a larger cross section in the forward direction. Somewhat more weight was given to the Seagrave et al. experiment, since the geometry appears to be somewhat cleaner and less subject to multiple scattering effects which would tend to make the distributions too isotropic.

This optical model underestimates the magnitude of the non-elastic (or compound nucleus formation) cross section at 14.5 Mev. The calculation gives




FIGURE 5. *Elastic scattering at 4.02 Mev.*

The calculation is compared to the 4.1 Mev experimental data of Vincent, Morgan, and Prud'homme [1960].




FIGURE 6. *Elastic scattering at 2.0 Mev.*

Experimental data are from Seagrave, Cranberg, and Simmons (1958).




FIGURE 7. Elastic scattering at 1.730 Mev.

Experimental data are from Lane, Lane, Langsdorf, Monahan, and Elwyn [1960].




FIGURE 9. Elastic scattering at 1.250 Mev.

Experimental data are from Lane, Langsdorf, Monahan, and Elwyn [1960].




FIGURE 8. Elastic scattering at 1.570 Mev.

Experimental data are from Lane, Langsdorf, Monahan, and Elwyn [1960].




FIGURE 10. Elastic scattering at 1.005 Mev.

Experimental data are from Lane, Langsdorf, Monahan, and Elwyn [1960] at 1.01 ± 0.03 Mev (crosses) and from Seagrave, Cranberg, and Simmons [1958] at 1.0 Mev (vertical lines).

1.227 barns at 14.6 Mev, whereas the experimental measurement at 14.5 Mev of Flerov and Talyzin [1956] gives 1.36 ± 0.02 barns. This leads to a corresponding error in the total cross section, but does not affect the elastic cross section values. The total elastic cross section at 14.6 Mev is 0.803 barns, in agreement with the experimental value of 0.82 ± 0.07 barns of Cross and Jarvis.

4. Results of the Calculation

The calculated angular distributions for total elastic scattering are given in table 1 and the Legendre coefficient expansions are given in table 2.




FIGURE 11. Elastic scattering at 0.930 Mev.

Experimental data are from Lane, Langsdorf, Monahan, and Elwyn [1960].




FIGURE 12. Elastic scattering at 0.770 Mev.

Experimental data are from Lane, Langsdorf, Monahan, and Elwyn [1960].

TABLE I. Angular distributions of neutron elastic scattering for calcium (barns/steradian)

Angle	18. 000 Mev	17. 100 Mev	16. 300 Mev	15. 500 Mev	14. 750 Mev	14. 000 Mev	13. 300 Mev
deg							
0.	2. 2740	2. 1727	1. 9827	1. 8448	1. 7511	1. 6825	1. 5868
5. 0	2. 1104	2. 0223	1. 8512	1. 7277	1. 6449	1. 5857	1. 5013
10. 0	1. 6774	1. 6223	1. 4995	1. 4131	1. 3584	1. 3234	1. 2685
15. 0	1. 1189	1. 1011	1. 0364	0. 9947	0. 9740	0. 9678	0. 9495
20. 0	0. 5980	0. 6068	0. 5893	. 5842	. 5908	. 6074	. 6203
25. 0	. 2290	. 2467	. 2544	. 2684	. 2884	. 3147	. 3451
30. 0	. 0439	. 0558	. 0677	. 0837	. 1029	. 1262	. 1585
35. 0	. 0031	. 0037	. 0084	. 0161	. 0261	. 0384	. 0614
40. 0	. 0349	. 0270	. 0232	. 0212	. 0208	. 0212	. 0310
45. 0	. 0755	. 0657	. 0582	. 0507	. 0437	. 0366	. 0354
50. 0	. 0914	. 0847	. 0787	. 0712	. 0632	. 0545	. 0479
55. 0	. 0792	. 0772	. 0753	. 0712	. 0660	. 0600	. 0533
60. 0	. 0527	. 0540	. 0556	. 0553	. 0541	. 0523	. 0484
65. 0	. 0281	. 0302	. 0330	. 0350	. 0366	. 0377	. 0371
70. 0	. 0142	. 0152	. 0175	. 0197	. 0220	. 0237	. 0251
75. 0	. 0112	. 0106	. 0117	. 0123	. 0142	. 0147	. 0162
80. 0	. 0141	. 0127	. 0129	. 0128	. 0127	. 0113	. 0118
85. 0	. 0174	. 0161	. 0161	. 0154	. 0143	. 0117	. 0111
90. 0	. 0178	. 0173	. 0179	. 0174	. 0164	. 0139	. 0130
95. 0	. 0152	. 0156	. 0170	. 0175	. 0175	. 0162	. 0162
100. 0	. 0110	. 0121	. 0143	. 0160	. 0174	. 0179	. 0194
105. 0	. 0072	. 0086	. 0113	. 0139	. 0166	. 0184	. 0217
110. 0	. 0049	. 0061	. 0088	. 0119	. 0152	. 0175	. 0219
115. 0	. 0040	. 0048	. 0072	. 0099	. 0130	. 0150	. 0196
120. 0	. 0039	. 0042	. 0059	. 0079	. 0100	. 0113	. 0150
125. 0	. 0038	. 0038	. 0047	. 0057	. 0067	. 0071	. 0093
130. 0	. 0033	. 0032	. 0036	. 0038	. 0041	. 0039	. 0047
135. 0	. 0028	. 0028	. 0031	. 0031	. 0030	. 0027	. 0028
140. 0	. 0026	. 0029	. 0035	. 0039	. 0041	. 0039	. 0043
145. 0	. 0029	. 0035	. 0048	. 0059	. 0069	. 0071	. 0085
150. 0	. 0035	. 0044	. 0064	. 0083	. 0101	. 0105	. 0131
155. 0	. 0043	. 0052	. 0076	. 0099	. 0121	. 0126	. 0161
160. 0	. 0048	. 0053	. 0075	. 0097	. 0117	. 0122	. 0157
165. 0	. 0049	. 0048	. 0063	. 0077	. 0090	. 0093	. 0120
170. 0	. 0045	. 0039	. 0043	. 0047	. 0051	. 0051	. 0067
175. 0	. 0040	. 0030	. 0026	. 0021	. 0017	. 0015	. 0020
180. 0	. 0038	. 0026	. 0019	. 0010	. 0004	. 0000	. 0001

TABLE I. Angular distributions of neutron elastic scattering for calcium (barns/steradian)—Continued

Angle	12. 700 Mev	12. 100 Mev	11. 500 Mev	10. 900 Mev	10. 400 Mev	9. 890 Mev	9. 410 Mev
deg							
0.	1. 5352	1. 4533	1. 4105	1. 3773	1. 3298	1. 3199	1. 3476
5. 0	1. 4571	1. 3852	1. 3492	1. 3219	1. 2816	1. 2756	1. 3061
10. 0	1. 2434	1. 1979	1. 1794	1. 1679	1. 1470	1. 1511	1. 1891
15. 0	0. 9480	0. 9357	0. 9393	0. 9476	0. 9521	0. 9692	1. 0165
20. 0	. 6382	. 6555	. 6781	. 7039	. 7321	. 7609	0. 8159
25. 0	. 3726	. 4078	. 4410	. 4768	. 5212	. 5571	. 6152
30. 0	. 1848	. 2238	. 2577	. 2945	. 3445	. 3815	. 4370
35. 0	. 0792	. 1109	. 1378	. 1684	. 2143	. 2470	. 2946
40. 0	. 0382	. 0573	. 0739	. 0947	. 1302	. 1553	. 1916
45. 0	. 0345	. 0416	. 0489	. 0599	. 0832	. 0997	. 1240
50. 0	. 0434	. 0428	. 0439	. 0480	. 0604	. 0695	. 0830
55. 0	. 0491	. 0459	. 0446	. 0448	. 0497	. 0535	. 0588
60. 0	. 0466	. 0441	. 0428	. 0417	. 0426	. 0431	. 0433
65. 0	. 0373	. 0368	. 0362	. 0352	. 0345	. 0334	. 0312
70. 0	. 0259	. 0268	. 0264	. 0256	. 0247	. 0232	. 0205
75. 0	. 0163	. 0171	. 0164	. 0155	. 0148	. 0134	. 0113
80. 0	. 0105	. 0105	. 0091	. 0079	. 0073	. 0063	. 0051
85. 0	. 0089	. 0079	. 0061	. 0046	. 0042	. 0035	. 0032
90. 0	. 0108	. 0096	. 0078	. 0063	. 0062	. 0057	. 0059
95. 0	. 0148	. 0144	. 0131	. 0119	. 0123	. 0117	. 0121
100. 0	. 0193	. 0205	. 0198	. 0189	. 0202	. 0193	. 0196
105. 0	. 0226	. 0255	. 0254	. 0247	. 0269	. 0257	. 0257
110. 0	. 0233	. 0274	. 0276	. 0271	. 0298	. 0285	. 0284
115. 0	. 0209	. 0252	. 0255	. 0259	. 0281	. 0270	. 0269
120. 0	. 0158	. 0194	. 0197	. 0197	. 0223	. 0218	. 0221
125. 0	. 0097	. 0121	. 0124	. 0127	. 0149	. 0151	. 0158
130. 0	. 0047	. 0059	. 0063	. 0069	. 0086	. 0095	. 0106
135. 0	. 0027	. 0032	. 0035	. 0042	. 0057	. 0067	. 0081
140. 0	. 0042	. 0048	. 0049	. 0053	. 0068	. 0076	. 0090
145. 0	. 0083	. 0096	. 0094	. 0093	. 0109	. 0111	. 0122
150. 0	. 0130	. 0152	. 0146	. 0139	. 0158	. 0152	. 0159
155. 0	. 0159	. 0188	. 0179	. 0168	. 0190	. 0178	. 0182
160. 0	. 0156	. 0186	. 0178	. 0168	. 0188	. 0176	. 0179
165. 0	. 0121	. 0146	. 0143	. 0137	. 0154	. 0145	. 0149
170. 0	. 0070	. 0086	. 0089	. 0090	. 0103	. 0100	. 0106
175. 0	. 0025	. 0034	. 0042	. 0048	. 0058	. 0061	. 0069
180. 0	. 0007	. 0013	. 0023	. 0032	. 0040	. 0046	. 0055

TABLE I. Angular distributions of neutron elastic scattering for calcium (barns/steradian)—Continued

Angle	8.950 Mev.	8.510 Mev.	8.100 Mev.	7.700 Mev.	7.330 Mev.	6.970 Mev.	6.663 Mev.
0.	1.3531	1.3565	1.3728	1.4119	1.4324	1.4620	1.4826
5.0	1.3142	1.3211	1.3389	1.3795	1.4010	1.4320	1.4533
10.0	1.2039	1.2203	1.2422	1.2867	1.3108	1.3456	1.3687
15.0	1.0400	1.0692	1.0962	1.1457	1.1732	1.2131	1.2385
20.0	0.8472	0.8887	0.9205	0.9741	1.0046	1.0491	1.0766
25.0	.6513	.7014	.7358	.7911	0.8231	0.8704	0.8990
30.0	.4736	.5265	.5608	.6141	.6457	.6930	.7211
35.0	.3276	.3775	.4087	.4565	.4853	.5294	.5554
40.0	.2182	.2603	.2861	.3257	.3500	.3882	.4106
45.0	.1429	.1746	.1939	.2237	.2426	.2731	.2910
50.0	.0946	.1156	.1284	.1485	.1617	.1839	.1972
55.9	.0645	.0763	.0835	.0952	.1034	.1180	.1270
60.0	.0450	.0500	.0531	.0585	.0628	.0713	.0768
65.0	.0308	.0316	.0323	.0338	.0356	.0397	.0427
70.0	.0193	.0183	.0179	.0175	.0180	.0196	.0211
75.0	.0102	.0090	.0085	.0078	.0078	.0084	.0091
80.0	.0045	.0039	.0036	.0034	.0035	.0039	.0043
85.0	.0030	.0032	.0032	.0037	.0038	.0046	.0048
90.0	.0058	.0065	.0065	.0075	.0076	.0087	.0087
95.0	.0118	.0125	.0123	.0135	.0132	.0145	.0142
100.0	.0188	.0193	.0186	.0196	.0190	.0203	.0197
105.0	.0245	.0247	.0236	.0243	.0235	.0247	.0239
110.0	.0270	.0271	.0259	.0264	.0255	.0267	.0261
115.0	.0258	.0260	.0251	.0257	.0251	.0262	.0252
120.0	.0215	.0223	.0220	.0228	.0229	.0245	.0248
125.0	.0161	.0174	.0179	.0191	.0199	.0219	.0229
130.0	.0116	.0132	.0144	.0161	.0174	.0198	.0213
135.0	.0094	.0112	.0126	.0146	.0162	.0189	.0207
140.0	.0100	.0116	.0129	.0150	.0165	.0193	.0212
145.0	.0127	.0137	.0146	.0165	.0177	.0204	.0222
150.0	.0157	.0162	.0164	.0181	.0189	.0216	.0231
155.0	.0174	.0175	.0173	.0187	.0192	.0218	.0232
160.0	.0168	.0169	.0165	.0178	.0182	.0208	.0223
165.0	.0141	.0143	.0140	.0155	.0160	.0188	.0204
170.0	.0103	.0108	.0109	.0125	.0133	.0163	.0183
175.0	.0071	.0079	.0083	.0101	.0111	.0143	.0165
180.0	.0058	.0067	.0073	.0091	.0103	.0135	.0159

Angle	6.300 Mev.	6.000 Mev.	5.700 Mev.	5.430 Mev.	5.160 Mev.	4.910 Mev.	4.670 Mev.
0.	1.5013	1.5253	1.5377	1.5673	1.5719	1.6092	1.6033
5.0	1.4726	1.4980	1.5109	1.5411	1.5464	1.5837	1.5786
10.0	1.3897	1.4188	1.4335	1.4651	1.4723	1.5096	1.5064
15.0	1.2617	1.2958	1.3128	1.3464	1.3563	1.3935	1.3932
20.0	1.1018	1.1409	1.1604	1.1956	1.2087	1.2453	1.2483
25.0	0.9254	0.9681	0.9894	1.0255	1.0414	1.0770	1.0832
30.0	.7473	.7910	.8134	0.8489	0.8669	0.9006	0.9097
35.0	.5799	.6219	.6439	.6774	.6966	.7279	.7367
40.0	.4321	.4697	.4902	.5202	.5395	.5671	.5792
45.0	.3086	.3398	.3580	.3835	.4019	.4254	.4377
50.0	.2105	.2345	.2497	.2704	.2872	.3065	.3181
55.0	.1363	.1531	.1654	.1813	.1961	.2113	.2218
60.0	.0828	.0935	.1031	.1149	.1276	.1392	.1481
65.0	.0463	.0524	.0597	.0683	.0791	.0877	.0951
70.0	.0231	.0262	.0318	.0383	.0474	.0538	.0598
75.0	.0102	.0117	.0160	.0213	.0290	.0340	.0386
80.0	.0049	.0058	.0093	.0139	.0206	.0247	.0282
85.0	.0050	.0060	.0089	.0132	.0191	.0227	.0252
90.0	.0086	.0098	.0122	.0166	.0218	.0250	.0268
95.0	.0138	.0152	.0173	.0217	.0265	.0295	.0306
100.0	.0190	.0207	.0225	.0271	.0317	.0344	.0352
105.0	.0231	.0249	.0268	.0315	.0361	.0388	.0395
110.0	.0255	.0274	.0295	.0343	.0393	.0422	.0430
115.0	.0260	.0282	.0307	.0359	.0412	.0447	.0459
120.0	.0252	.0277	.0309	.0364	.0423	.0468	.0486
125.0	.0239	.0267	.0305	.0363	.0429	.0489	.0515
130.0	.0229	.0259	.0302	.0363	.0435	.0515	.0551
135.0	.0227	.0256	.0304	.0368	.0446	.0551	.0597
140.0	.0233	.0261	.0311	.0378	.0460	.0595	.0653
145.0	.0242	.0270	.0322	.0390	.0477	.0649	.0719
150.0	.0251	.0278	.0332	.0403	.0494	.0707	.0791
155.0	.0252	.0280	.0337	.0411	.0508	.0767	.0864
160.0	.0244	.0276	.0336	.0414	.0516	.0824	.0934
165.0	.0229	.0265	.0329	.0411	.0519	.0872	.0994
170.0	.0210	.0252	.0319	.0406	.0518	.0910	.1042
175.0	.0196	.0241	.0311	.0401	.0517	.0933	.1072
180.0	.0190	.0237	.0308	.0399	.0516	.0941	.1082

TABLE I. Angular distributions of neutron elastic scattering for calcium (barns/steradian)—Continued

Angle	4.440 Mev	4.230 Mev	4.020 Mev	3.820 Mev	3.640 Mev	3.460 Mev	3.290 Mev
0.	1.7219	1.6126	1.5996	1.6072	1.3986	1.5570	1.5703
5.0	1.6038	1.5889	1.5765	1.5836	1.3787	1.5360	1.5497
10.0	1.6118	1.5197	1.5093	1.5150	1.3208	1.4746	1.4896
15.0	1.4832	1.4109	1.4034	1.4077	1.2298	1.3776	1.3944
20.0	1.3190	1.2710	1.2672	1.2712	1.1132	1.2522	1.2711
25.0	1.1321	1.1107	1.1109	1.1161	0.9799	1.1074	1.1283
30.0	0.9366	0.9408	0.9449	0.9528	.8394	0.9525	0.9751
35.0	.7455	.7719	.7794	.7907	.7006	.7966	.8206
40.0	.5696	.6126	.6229	.6370	.5708	.6475	.6726
45.0	.4170	.4698	.4818	.4971	.4559	.5115	.5375
50.0	.2923	.3475	.3602	.3749	.3592	.3927	.4195
55.0	.1972	.2477	.2604	.2727	.2824	.2936	.3208
60.0	.1304	.1703	.1823	.1915	.2250	.2148	.2421
65.0	.0884	.1138	.1247	.1312	.1854	.1552	.1822
70.0	.0665	.0754	.0850	.0900	.1607	.1129	.1390
75.0	.0589	.0519	.0600	.0651	.1475	.0850	.1098
80.0	.0600	.0396	.0464	.0524	.1421	.0885	.0917
85.0	.0648	.0352	.0408	.0479	.1409	.0602	.0817
90.0	.0691	.0358	.0404	.0476	.1410	.0574	.0774
95.0	.0702	.0390	.0428	.0491	.1401	.0579	.0768
100.0	.0668	.0432	.0464	.0511	.1366	.0601	.0783
105.0	.0590	.0475	.0504	.0534	.1301	.0634	.0813
110.0	.0481	.0514	.0545	.0587	.1204	.0675	.0855
115.0	.0363	.0552	.0587	.0616	.1083	.0725	.0912
120.0	.0261	.0591	.0636	.0684	.0948	.0791	.0990
125.0	.0201	.0637	.0694	.0769	.0808	.0877	.1096
130.0	.0201	.0693	.0765	.0865	.0677	.0987	.1239
135.0	.0277	.0762	.0852	.0968	.0562	.1126	.1423
140.0	.0430	.0846	.0956	.1078	.0474	.1293	.1647
145.0	.0654	.0942	.1075	.1198	.0417	.1485	.1907
150.0	.0933	.1047	.1206	.1335	.0392	.1696	.2191
155.0	.1243	.1156	.1341	.1491	.0397	.1915	.2484
160.0	.1555	.1261	.1473	.1662	.0425	.2128	.2765
165.0	.1839	.1354	.1592	.1834	.0464	.2319	.3013
170.0	.2066	.1428	.1686	.1982	.0505	.2471	.3207
175.0	.2212	.1475	.1747	.2085	.0534	.2568	.3331
180.0	.2262	.1491	.1768	.2121	.0545	.2602	.3374

Angle	2.970 Mev	2.690 Mev	2.440 Mev	2.210 Mev	2.000 Mev	1.810 Mev	1.630 Mev
0.	1.4990	1.3767	1.2546	1.1456	1.0433	0.9309	0.8244
5.0	1.4793	1.3603	1.2405	1.1334	1.0327	.9221	.8173
10.0	1.4218	1.3123	1.1991	1.0975	1.0017	.8963	.7963
15.0	1.3311	1.2361	1.1332	1.0403	0.9522	.8552	.7628
20.0	1.2143	1.1369	1.0473	0.9656	.8873	.8013	.7187
25.0	1.0797	1.0211	0.9466	.8779	.8111	.7378	.6663
30.0	0.9364	0.8957	.8374	.7822	.7278	.6881	.6083
35.0	.7927	.7679	.7253	.6838	.6417	.5958	.5476
40.0	.6558	.6440	.6160	.5873	.5569	.5241	.4867
45.0	.5311	.5290	.5138	.4964	.4767	.4554	.4280
50.0	.4219	.4269	.4220	.4142	.4036	.3919	.3731
55.0	.3298	.3396	.3426	.3425	.3391	.3349	.3232
60.0	.2550	.2682	.2764	.2818	.2839	.2851	.2789
65.0	.1962	.2119	.2230	.2321	.2380	.2424	.2403
70.0	.1520	.1694	.1814	.1924	.2005	.2065	.2071
75.0	.1201	.1387	.1500	.1614	.1704	.1767	.1787
80.0	.0988	.1175	.1268	.1376	.1465	.1520	.1544
85.0	.0858	.1036	.1102	.1196	.1276	.1315	.1337
90.0	.0795	.0949	.0986	.1061	.1127	.1148	.1163
95.0	.0782	.0900	.0908	.0962	.1013	.1015	.1019
100.0	.0803	.0879	.0862	.0896	.0933	.0916	.0909
105.0	.0848	.0882	.0846	.0864	.0889	.0856	.0837
110.0	.0908	.0910	.0863	.0870	.0885	.0841	.0811
115.0	.0981	.0968	.0917	.0920	.0930	.0878	.0839
120.0	.1068	.1063	.1016	.1021	.1032	.0974	.0926
125.0	.1176	.1202	.1168	.1181	.1196	.1133	.1078
130.0	.1316	.1391	.1375	.1403	.1426	.1358	.1296
135.0	.1498	.1632	.1640	.1686	.1720	.1645	.1575
140.0	.1727	.1922	.1957	.2025	.2071	.1986	.1905
145.0	.2006	.2252	.2315	.2407	.2466	.2367	.2274
150.0	.2326	.2607	.2698	.2814	.2885	.2770	.2661
155.0	.2671	.2967	.3084	.3223	.3305	.3172	.3044
160.0	.3015	.3309	.3448	.3608	.3699	.3550	.3400
165.0	.3329	.3608	.3765	.3943	.4041	.3877	.3706
170.0	.3580	.3841	.4011	.4202	.4304	.4130	.3941
175.0	.3743	.3988					

TABLE 1. Angular distribution of neutron elastic scattering for calcium (barns/steradian)—Continued

Angle	1.480 Mev	1.340 Mev	1.210 Mev	1.096 Mev	0.991 Mev	0.853 Mev	0.734 Mev
deg							
0.	0.4400	0.6742	0.3484	0.3193	0.2848	0.4682	0.4299
5.0	.4365	.6691	.3459	.3172	.2831	.4656	.4277
10.0	.4263	.6541	.3388	.3111	.2780	.4577	.4211
15.0	.4097	.6301	.3273	.3013	.2697	.4450	.4104
20.0	.3879	.5982	.3121	.2883	.2586	.4280	.3960
25.0	.3619	.5602	.2938	.2726	.2453	.4074	.3786
30.0	.3330	.5178	.2733	.2550	.2302	.3842	.3588
35.0	.3025	.4730	.2515	.2360	.2140	.3590	.3372
40.0	.2717	.4275	.2292	.2165	.1972	.3328	.3146
45.0	.2416	.3829	.2071	.1969	.1802	.3063	.2915
50.0	.2131	.3404	.1857	.1778	.1636	.2801	.2685
55.0	.1868	.3007	.1655	.1596	.1477	.2548	.2460
60.0	.1630	.2646	.1467	.1424	.1325	.2306	.2244
65.0	.1418	.2319	.1295	.1264	.1184	.2078	.2038
70.0	.1230	.2028	.1139	.1117	.1053	.1865	.1844
75.0	.1066	.1769	.0998	.0983	.0932	.1669	.1664
80.0	.0922	.1540	.0871	.0861	.0822	.1489	.1498
85.0	.0797	.1338	.0758	.0752	.0723	.1326	.1348
90.0	.0689	.1163	.0659	.0656	.0636	.1182	.1215
95.0	.0600	.1016	.0576	.0575	.0562	.1060	.1100

TABLE 1.—Angular distribution of neutron elastic scattering for calcium (barns/steradian)—Continued

Angle	1.480 Mev	1.340 Mev	1.210 Mev	1.096 Mev	0.991 Mev	0.853 Mev	0.734 Mev
deg							
100.0	.0530	.0902	.0511	.0511	.0503	.0961	.1007
105.0	.0484	.0824	.0465	.0465	.0461	.0889	.0937
110.0	.0464	.0790	.0443	.0441	.0437	.0847	.0894
115.0	.0476	.0805	.0447	.0441	.0435	.0839	.0878
120.0	.0521	.0875	.0480	.0467	.0456	.0866	.0893
125.0	.0603	.1003	.0542	.0519	.0499	.0930	.0937
130.0	.0720	.1188	.0634	.0598	.0565	.1028	.1010
135.0	.0872	.1426	.0752	.0699	.0652	.1157	.1108
140.0	.1051	.1709	.0893	.0820	.0755	.1312	.1227
145.0	.1250	.2023	.1049	.0955	.0869	.1484	.1360
150.0	.1459	.2352	.1212	.1095	.0989	.1665	.1501
155.0	.1666	.2677	.1373	.1234	.1108	.1844	.1640
160.0	.1858	.2978	.1523	.1363	.1218	.2010	.1768
165.0	.2022	.3236	.1651	.1473	.1312	.2152	.1879
170.0	.2148	.3435	.1749	.1557	.1384	.2261	.1963
175.0	.2228	.3560	.1810	.1610	.1429	.2329	.2016
180.0	.2255	.3602	.1831	.1628	.1445	.2352	.2034

TABLE 2. Legendre coefficients for calcium elastic scattering

Energy.....Mev	18.000	17.100	16.300	15.500	14.750	14.000	13.300
Total elastic.....barns	0.836	0.826	0.806	0.797	0.800	0.804	0.827
$S_l(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1	0.80315	0.79852	0.77512	0.75673	0.74259	0.74200	0.72305
2	.63456	.63283	.61087	.59713	.58931	.59324	.58335
3	.52042	.52075	.50234	.49179	.48678	.49052	.48014
4	.44965	.44827	.43098	.41843	.40920	.40525	.38564
5	.38692	.38273	.36377	.34726	.33268	.32239	.29412
6	.33349	.32659	.30864	.29195	.27627	.26353	.23572
7	.27457	.26459	.24661	.22811	.20948	.19201	.16411
8	.19745	.18289	.16280	.14236	.12222	.10318	.07932
9	.11379	.10103	.08647	.07360	.06212	.05073	.04016
10	.04970	.04153	.03233	.02535	.02005	.01572	.01149
11	.01833	.01441	.01021	.00735	.00540	.00401	.00267
12	.00600	.00448	.00294	.00197	.00136	.00096	.00059
13	.00176	.00125	.00076	.00048	.00031	.00021	.00012
14	.00048	.00033	.00019	.00011	.00007	.00004	.00002
15	.00012	.00008	.00004	.00002	.00001	.00001	
Energy.....Mev	12.700	12.100	11.500	10.900	10.400	9.900	9.410
Total elastic.....barns	0.841	0.882	0.912	0.951	1.030	1.082	1.177
$S_l(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1	0.72659	0.71210	0.72014	0.72907	0.72052	0.72939	0.73597
2	.58770	.57698	.58101	.58442	.57153	.57326	.57503
3	.47996	.46317	.45735	.44935	.42450	.41510	.40540
4	.37632	.34809	.33140	.31266	.27692	.25958	.24144
5	.27959	.24550	.22536	.20430	.16755	.15023	.13010
6	.22017	.18950	.16944	.14854	.11804	.10130	.08231
7	.14672	.11953	.10163	.08462	.06413	.05288	.04149
8	.06418	.04400	.03236	.02305	.01255	.00863	.00521
9	.03164	.02328	.01665	.01139	.00748	.00496	.00330
10	.00883	.00607	.00426	.00285	.00179	.00117	.00076
11	.00197	.00123	.00084	.00054	.00031	.00020	.00012
12	.00042	.00024	.00016	.00010	.00005	.00003	.00002
13	.00008	.00004	.00003	.00002			
14	.00002						
Energy.....Mev	8.950	8.510	8.100	7.700	7.330	6.970	6.663
Total elastic.....barns	1.238	1.341	1.407	1.523	1.591	1.709	1.777
$S_l(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1	0.74212	0.74199	0.74508	0.74309	0.74353	0.73635	0.73453
2	.57477	.56609	.56447	.55856	.55588	.54614	.54279
3	.39604	.37741	.36886	.35637	.34828	.33352	.32556
4	.22696	.20325	.19213	.17775	.16886	.15474	.14743
5	.11689	.09579	.08670	.07400	.06757	.05690	.05233
6	.06991	.05333	.04520	.03526	.02998	.02295	.01961
7	.03407	.02518	.02073	.01597	.01319	.01015	.00839
8	.00360	.00168	.00119	.00055	.00043	.00013	.00013
9	.00215	.00126	.00081	.00050	.00033	.00019	.00012
10	.00049	.00027	.00017	.00011	.00007	.00004	.00002
11	.00008	.00004	.00003	.00002	.00001		
12	.00001						
Energy.....Mev	6.300	6.000	5.700	5.430	5.160	4.910	4.670
Total elastic.....barns	1.844	1.965	2.057	2.196	2.313	2.472	2.538
$S_l(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1	0.73165	0.72296	0.70672	0.68561	0.65982	0.62793	0.61459
2	.53906	.52817	.51546	.49904	.48011	.47289	.46532
3	.31721	.30341	.28995	.27561	.25893	.24041	.22903
4	.14042	.12812	.11992	.11103	.10245	.09839	.09343
5	.04818	.03950	.03571	.03080	.02742	.02337	.02113
6	.01680	.01218	.01029	.00798	.00669	.00581	.00506
7	.00692	.00491	.00397	.00300	.00240	.00181	.00147
8	.00014	.00003	.00004	.00002	.00003	.00022	.00023
9	.00008	.00004	.00003	.00002	.00001		
10	.00002	.00001					

TABLE 2. Legendre coefficients for calcium elastic scattering—Continued

Energy.....Mev	4.440	4.230	4.020	3.820	3.640	3.460	3.290
Total elastic.....barns	2.669	2.761	2.858	2.965	3.043	3.149	3.481
$S_1(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1.....	0.59352	0.57163	0.54777	0.52606	0.50422	0.47943	0.42278
2.....	.45321	.44185	.43012	.42050	.26083	.39623	.36924
3.....	.21764	.20350	.18836	.17691	.16300	.14962	.12522
4.....	.14034	.08320	.07840	.07496	.07244	.06756	.06291
5.....	.01825	.01618	.01416	.01204	.01051	.00885	.00701
6.....	.00408	.00359	.00330	.00279	.00291	.00241	.00238
7.....	.00109	.00087	.00069	.00050	.00040	.00030	.00022
8.....	.00024	.00026	.00037	.00247	.00037	.00031	
Energy.....Mev	2.970	2.690	2.440	2.210	2.000	1.810	1.630
Total elastic.....barns	3.502	3.551	3.476	3.432	3.371	3.228	3.065
$S_1(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1.....	0.39586	0.36179	0.34086	0.31364	0.28852	0.27864	0.26563
2.....	.35509	.33340	.31893	.30155	.28452	.26741	.25148
3.....	.10885	.08494	.06471	.04677	.03169	.01740	.00559
4.....	.05815	.05179	.04688	.04314	.03973	.03473	.03123
5.....	.00512	.00366	.00261	.00177	.00119	.00078	.00049
6.....	.00537	.00135	.00109	.00084	.00065	.00051	
7.....	.00012	.00007	.00005				
Energy.....Mev	1.480	1.340	1.210	1.096	0.991	0.853	0.734
Total elastic.....barns	1.727	2.779	1.500	1.426	1.322	2.303	2.224
$S_1(E), l=0$	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
1.....	0.26421	0.25235	0.25450	0.25701	0.25074	0.23850	0.23522
2.....	.23576	.22312	.20742	.19271	.17951	.15989	.13935
3.....	-.00221	-.00704	-.01040	-.01179	-.01216	-.01144	-.00941
4.....	.02695	.02480	.02103	.01784	.01589	.01330	.01028
5.....	.00031	.00020	.00012	.00008			

5. Conclusions

Reasonably satisfactory average cross section predictions for calcium at neutron energies below 18 Mev can be made with the nuclear optical model using compound elastic scattering cross sections as calculated from the optical model in the absence of inelastic scattering, and using a Hauser-Feshbach calculation if inelastic scattering is present. In order to achieve the goal of calculation of any neutron cross section as required, Hauser-Feshbach and direct interaction inelastic scattering codes for inelastic scattering, and nuclear reaction codes will be required, for which the present type of calculation provides input data.

More experimental data in the region from 6 to 14 Mev is needed. Also the difference between the Seagrave et al. and the Lane et al. data at low energies needs to be resolved.

6. References

Amster, H. J., Leshan, E. J., and Walt, M., Computer techniques, Ch. IV. I., Fast Neutron Physics, Pt I, J. B. Marion and J. L. Fowler, Ed. (1960).
 Biedenharn, L. C., U. S. Atomic Energy Commission Report ORNL-1501 (1953).
 Biedenharn, L. C., Blatt, J. M., and Rose, M. E., Rev. Mod. Phys. **24**, 249 (1952).
 Bjorklund, F., and Fernbach, S., Phys. Rev. **109**, 1295 (1958).
 Blatt, J. M., and Biedenharn, L. C., Rev. Mod. Phys. **24**, 258 (1952).
 Blatt, J. M., and Weisskopf, V. F., Theoretical Nuclear Physics, p. 333, Wiley (New York) (1952).
 Bohm, D., Quantum Theory, p. 344, Prentice-Hall (1951).
 Cross, W. G., and Jarvis, R. G., Nuclear Physics **15**, 155 (1960).

Edmonds, A. R., Angular Momentum in Quantum Mechanics, Princeton Univ. Press, Princeton, N. J., p. 63 (1957).
 Feshbach, H., The complex potential model, Ch. VI. of Nuclear Spectroscopy, Part B, Fay Ajzenberg-Selove, Ed., Academic Press (New York) (1960).
 Feshbach, H., Porter, C. E., and Weisskopf, V. F., Phys. Rev. **96**, 448 (1954).
 Flerov, N. N., and Talyzin, V. M., Atomnaya Energiya **1**, 155 (1956).
 Gursky, M. L., and Stewart, Leona, Bull. Am. Phys. Soc. II, **6**, 367 (1961).
 Hauser, W., and Feshbach, H., Phys. Rev. **87**, 366 (1952).
 Howerton, R. J., U. S. A.E.C. Report (UCRL-5573) (1961).
 Huber, P., and Meyer, K. P., Proc. Intern. Symposium on Polarization Phenomena of Nucleons, p. 436, Berkhauser Verlag (Basel and Stuttgart) (1961).
 Lane, R. O., Langsdorf, A. S. Jr., Monahan, J. B., and Elwyn, A. J., private communication (1960). (See Howerton, 1961.)
 Lepore, J., Phys. Rev. **79**, 137 (1950).
 Morse, P. M., and Feshbach, H., Methods of Theoretical Physics, Mc-Graw Hill, New York, vol. 2, p. 1325 (1953).
 Perey, F. G. J., and Buck, B., Neutron Physics Division Progress Report, ORNL-3193 (1961).
 Seagrave, J. D., Cranberg, L., and Simmons, J. E., Private communication (1958). (See Howerton, 1961.)
 Troubetzkoy, E. S., Kalos, M. H., Lustig, H., Ray, J. H., and Trupin, B. H., NDA Report 2133-4 (NDL-TR-5), Nuclear Development Corporation of America, White Plains, N.Y. (1961).
 Vincent, L. D., Morgan, I. L., Prud 'homme, J. T., WADD-TR-60-217 (1960).
 Whittaker, E. T., and Watson, G. N., Modern Analysis, p. 331, Cambridge (1958).
 Wolfenstein, L., Phys. Rev. **82**, 690 (1951).
 Zurmühl, R., Praktische Mathematik für Ingenieure und Physiker, p. 412, Springer-Verlag, (Berlin, Göttingen, Heidelberg) (1961).

(Paper 66A5-174)