JOURNAL OF RESEARCH of the National Bureau of Standards – A. Physics and Chemistry Vol. 71A, No. 6, November–December 1967

Lifetimes of Energy Levels in Neutral Iron

C. H. Corliss and J. L. Tech

Institute for Basic Standards, National Bureau of Standards, Washington, D.C. 20234

(August 15, 1967)

Mean radiative lifetimes for 408 energy levels of neutral iron are calculated from the known transition probabilities of 3288 lines of Fe I.

Key Words: Atomic spectra, energy levels, iron, iron-lifetimes in, lifetimes in Fe I, radiative lifetimes, spectroscopy.

1. Introduction

Let us first briefly review the concept of mean radiative lifetime. Even in the absence of de-excitation caused by such external factors as collisional processes or an ambient radiation field, an atom in an excited state E_i will eventually suffer a spontaneous transition to a state of lower energy E_j and will emit a photon of energy in accordance with Bohr's energy relation, $h\nu = E_i - E_j$. The probability per unit time, A_{ij} , associated with this type of spontaneous decay is a physical constant for the particular transition and was first described by Einstein [1917]. For a system of N_i atoms in an excited state of energy E_i , we can write the decay rate as

$$\frac{dN_i}{dt} = -\sum_j A_{ij} N_i$$

the summation being extended over all states E_j of energy lower than E_i . By setting $1/\tau_i = \sum_j A_{ij}$ and integrating, we obtain

$$N_i = N_i^0 e^{-t/\tau_i}$$

where N_i^{ρ} is the number of atoms in the excited state at time t=0. After a time interval equal to τ_i , therefore, the number of atoms remaining in the excited level will have been reduced by a factor 1/e. This time interval is defined as the mean radiative lifetime of the given energy state and can be measured by a variety of wellestablished experimental methods, as described in the well-known work by Mitchell and Zemansky [1934]. As a result of the connection between transition probabilities (*A*-values) and oscillator strengths (*f*-values), namely

$$A_{ij} = \frac{6.670}{\lambda^2} \frac{g_j}{g_i} f_{ji}$$

where g is the statistical weight, 2J+1, of the level, it is possible to calculate the lifetime of an energy level

$$\tau_i = 1/\Sigma_i A_{ij}$$

provided only that the *f*-values for all possible downward transitions are known. The practicability of this indirect method is seriously limited, however, because for levels other than those yielding simple resonance lines the method requires not only a rather complete knowledge of the energy-level structure of the given atom, but also *f*-values in quantities not usually available. In the case of iron, however, these requirements no longer are insurmountable.

2. Discussion of the Data

For a variety of reasons, the spectrum of iron has traditionally been the object of considerable attention. The term analysis is sufficiently advanced that it was possible for Corliss and Bozman [1962] to derive *f*-values for a large number of iron lines by a simple reduction of accurate arc emission intensities. Corliss and Warner [1964, 1966] later extended this work by adding new and fainter lines. For many of the lines, the evaluation of 19 other sets of measurements in the literature made it possible for them to arrive at a "best" value for *gf*.

In these previous compilations the calculation of gf-values included the application of an empirically determined normalization to put the relative gf-values on an absolute scale, the normalization being given as a function of upper energy level for the transition. This function, which was essentially a correction to level populations for departures from a Boltzmann distribution in the arc, reduces the value of gf calculated from line intensities measured in the free-burning arc below that which would have been calculated with a simple Boltzmann factor. The correction affected all lines that arise from upper energy levels lying above 46,000 cm⁻¹, the magnitude of the correction becoming progressively larger as the value of the upper level increased.

Recent investigations by Huber and Tobey [1967]. who measured oscillator strengths for Fe I by use of a shock tube, do not support these excitation corrections in the case of FeI. A similar conclusion may be drawn from the work of Cowley and Warner [1967], who have decided on the basis of a theoretical model for the arc that a single normalization function is not applicable to all spectra but that the function depends on both the ionization potential and the ionization stage of the atom under study. For iron, their work indicates that no correction at all should be applied. In view of these concordant results, we have recalculated gf-values for all lines to which the excitation correction had previously been applied. The recalculated gf-values, together with other supporting material for the present calculation of lifetimes, are being published separately in a monograph by Corliss and Tech [1967]. That monograph covers the spectral range from 2084 to 9900 angstroms. As demonstrated there, the best available evidence indicates that the data presented in this new compilation do not now suffer from any serious systematic error.

Using these revised data we have calculated a number of intensity-related quantities, including *A*-values, for over 3000 spectral lines, and from these have been able to derive mean radiative lifetimes for over 400 energy levels of neutral iron. In a work of this kind, it is essential that the spectral range considered be sufficiently extensive to contain most of the lines in which the energy of the system is radiated and that no systematic error affects the data over that spectral range. We feel that the homogeneity and the extensive wavelength coverage of the present material on iron meets these requirements.

It has thus become possible for the first time to present for a complex atom a fairly comprehensive picture of individual lifetimes for a large number of its energy states. Apart from their own intrinsic interest, these lifetimes can additionally serve as the basis for a computation of the radiation damping constant for any line given in the aforementioned monograph. These constants, required in any calculation of line absorption coefficients for pure radiation damping, are given simply by the sum of the reciprocal lifetimes of the upper and lower energy levels involved in the transition. Thus, for a line produced by the transition $E_i \rightarrow E_j$, the damping constant will be given by

$$\Gamma_{ij} = \frac{1}{\tau_i} + \frac{1}{\tau_j} = \Gamma_i + \Gamma_j.$$

Of course, in most actual sources, for example those at high temperatures or high particle densities, other factors affecting the lifetimes come into play in fixing the damping constants and must be included in any such calculation of absorption coefficients or line profiles. However, the above expression for the contribution owing to pure radiation damping is exact and determines the natural width of the line.

3. Results

The calculated lifetimes for energy levels in iron are given in table 1, which is arranged in six columns as follows:

- col. 1-electron configuration (with parent term)
- col. 2-term designation
- col. 3-J-value for the individual levels of the term
- col. 4-level value rounded to the nearest cm⁻¹
- col. 5-lifetime in nanoseconds given in most cases to three significant figures, although not
- more than two figures are meaningful. col. 6-the number of downward transitions contributing to the lifetime determination.

Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions	Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions
3d ⁶ 4s(a ⁶ D)4p	z ⁷ D°	5	19351	182000.	2	$3d^6 4s(a + D)4p$	y ⁵ P°	3	36767	4.09	10
		4	19562	73700. 75600	2			2	$37158 \\ 37410$	3.58 3.15	9
		$\frac{3}{2}$	19737 19913 20020	89200. 137000.	2 2 2	$3d^{7} (b^{2}F)4s$	d ³ F	2	36941	<4320.	2
2.16.4.(-6D)/4-	7129	6	22650	799000	1	3d7 (a + F)4n	av3D°	3	38175	9 09	16
$3d^{\circ} 4s(a^{\circ}D)4p$	z'F	5 4	22650 22846 22997	12000. 8370.	$\begin{array}{c}1\\2\\3\end{array}$	Su (a 1)+p	y D	$\frac{3}{2}$	38678 38996	2.92 2.43 2.09	10 14 16
		3	23111	10500.	2	$\frac{3}{64} \frac{64}{2} \frac{(a4D)}{4p}$	~ 5D°	4	30696	1.64	7
		$\frac{2}{1}$	23193 23245	9540. 15700.	4	5 <i>a~</i> 45(<i>a~</i> D)4p	x D	32	39020 39970 40231	1.35 1.23	13 13
$3d^{6} 4s(a \ ^{6}\mathrm{D})4p$	z ⁷ P°	4	23711	17200.	3			1	40405	1.00	9
		3	24181 24507	26900. 85100	4			0	40491	1.03	3
		2	24307	05100.	5	3d ⁵ 4s ² (a ⁶ S)4p	y ⁷ P°	2	40052	119.	6
$3d^{6} 4s(a \ ^{6}\mathrm{D})4p$	<i>z</i> ⁵ D°	4	25900	59.8	6			3	40207	21.8	7
		3	26140 26340	60.6 50.1	6			4	40422	33.1	4
		1	26479	54.4	5	$3d^{6}4s(a^{4}D)4p$	$x {}^{5}\mathrm{F}^{\circ}$	5	40257	2.65	3
		0	26550	54.4	2			4	40594	1.81	7
2.76.4 ($= 6D$) $4 =$	~ 5F°	5	26875	47.4	4			3	40842	1.49	. 9
$5a^{4}(a^{-}D)^{4}p$	2 1	4	27167	50.8	6			ĩ	41131	.86	5
		3	27395	45.8	6		500		10005	0.04	0
		2	27560 27666	$43.2 \\ 39.2$	6 5	$3d^{\circ} 4s(b^{+}P)4p$	z "5"	2	40895	8.04	9
	5 D.0	2	20057	02.4	7	$3d^6 4s(b + P)4p$	<i>x</i> ⁵ P°	3	42533	7.87	11
$3d^{\circ} 4s(a^{\circ}D)4p$	z "P	3 2	29050	23.4 23.7	9			1	42000	10.8	7
		ĩ	29733	19.7	6						
9.16.4 (1D) 4	3179	4	21207	020	7	$3d^{6} 4s(a + H)4p$	y ⁵G°	6	42784	5.63	1
$3d^{\circ} 4s(a^{\circ}D)4p$	zor	3	31807	230. 249.	10			4	43023	7.05	5
		2	32134	370.	7			3	43138	4.90	4
2.16.4.(-1D).4.5	3D°	2	21292	110	0			2	43210	4.36	3
$5a^{-4}4s(a^{-1}D)4p$	2 0	2	31686	101.	7	$3d^{6} 4s(a^{6}D)5s$	<i>e</i> 7D	5	42816	3.15	7
		1	31937	98.5	6			4	43163	2.83	10
$3d^{7}(a+F)4n$	v ⁵ D°	4	33096	4.38	7.			2	43634	2.40	8
5 <i>a</i> (<i>a</i> 1)+ <i>p</i>	y D	3	33507	3.33	10			ī	43764	2.12	6
		2	33802	4.13	13	$3d^{6}4s(a^{4}H)4p$	z ⁵ H°	6	43321	10.9	1
		$ \begin{array}{c} 1\\ 0 \end{array} $	34017	2.58	4			5	42992	11.5	5
			01122		학생님 전문			43	43109	47.9	5
$3d^{7} (a {}^{4}\mathrm{F})4p$	y ⁵ F°	5	33695	4.69	5						
		3	34329	3.58	9	$3d^{6}4s(b^{4}P)4p$	w ⁵ D°	4	43500	2.02	8
		2	34547	3.58	8			3	43923	1.66	10
		1	34692	3.29	6			1	44184	1.05	14
$3d^{6} 4s(a {}^{4}D)4p$	z ³ P°	2	33947	14.4	14			0	44459	1.63	3
		1	34363	5.97	12						
		0	34556	31.7	4	$3d^6 4s(b + F) 4p$	w ⁵ F°	5	44244	7.21	4
$3d^{7}(a {}^{4}\mathrm{F})4p$	<i>z</i> ⁵ G°	6	34844	6.28	1			3	44166	8.98	9
		5 4	34782	5.80	12	성이 있는 것 같아요.		2	44285	7.54	8
		3	35612	4.68	11	상태 영화 영화 문		1	44378	7.37	4
		2	35856	4.29	7	2.16.4.1.1014	5D°	4	44415	5.54	6
$3d^{7}(a^{4}F)4p$	z ³ G°	5	35379	5.35	7	$3a^{-4s(b-F)4p}$	<i>U</i> *D	3	44551	4.87	6
		4	35768	4.80	12			2	44664	4.72	10
		3	36079	4.92	13			1	44761	3.14	5
$3d^{7}(a^{4}F)4p$	y ³ F°	4	36686	3.85	12			0	77021	1.55	-
		3	37163	3.82	13	$3d^{7} (a + P)4p$	y 5S°	2	44512	1.09	9
		2	37521	3.48	11						

 TABLE 1. Radiative lifetimes for energy levels in neutral iron

Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions	Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions
3d ⁶ 4s (a ⁶ D)5s	e ⁵ D	$ \begin{array}{c} 4 \\ 3 \\ 2 \\ 1 \\ 0 \end{array} $	$\begin{array}{r} 44677 \\ 45061 \\ 45334 \\ 45509 \\ 45595 \end{array}$	$\begin{array}{c} 4.50 \\ 4.44 \\ 3.80 \\ 3.63 \\ 3.29 \end{array}$	$13 \\ 15 \\ 14 \\ 8 \\ 3$	3d ⁶ 4s (a ⁴ G)4p	v ⁵ F°	$5 \\ 4 \\ 3 \\ 2 \\ 1$	47606 47930 48123 48239 48351	$\begin{array}{c} 4.25 \\ 2.09 \\ 2.22 \\ 1.50 \\ 1.50 \end{array}$	6 13 11 17 7
3d ⁶ 4s(b ⁴ P)4p	<i>x</i> ³ D°	$3 \\ 2 \\ 1$	$\begin{array}{c} 45221 \\ 45282 \\ 45552 \end{array}$	2.91 2.69 2.37	12 15 13	3d ⁶ 4s(b ⁴ F)4p	x ³G°	3 4 5	47834 47812 47835	$6.64 \\ 6.70 \\ 5.69$	12 15 10
3d ⁶ 4s (a ⁴ H)4p	у ³ G°	$5\\4\\3$	45295 45428 45563	8.05 8.20 18.6	9 14 12	3d7 (a 4F)5s	e ³ F	4 3 2	47961 48532 48928	4.74 3.99 3.73	19 20 9
3d ⁶ 4s (b ⁴ F)4p	x 5G°		45608 45726 45833 45914 45965	9.01 5.92 3.89 4.44 3.70	1 5 7 7 4	3 <i>d</i> ⁷ (<i>a</i> ⁴ P)4 <i>p</i>	v ⁵ P° w ³ G°	$ \begin{array}{c} 3 \\ 2 \\ 1 \\ 5 \end{array} $	47967 48163 48290 48231	.23 .68 .86 18.8	6 13 9 9
3d ⁶ 4s(a ⁴ H)4p	<i>z</i> ³ I°	2 7 6	45978 46027	22.5 28.7	1 3	2 d6 4 a (b 4 D) 4 p	3D°	4.3	48362 48476	15.7 6.14	89
$3d^5 4s^2(a\ {}^6\mathrm{S})4p$	w ⁵ P°	5 3 2	46136 46137 46314	.28	3 4 6	$3d^{3} 4s(0 + r) 4p$	х °Г	1 0 5	48516 48460 48383	1.33 1.46 1.41 3.07	19 14 7
		1	46410	.25	8	$3d^7 (a^2H)4p$	$z \cdot \Pi$ $v^{1}C^{\circ}$	4	40303	2 55	12
$3d^6 4s(b \ ^4\mathrm{P})4p$	z $^{3}S^{\circ}$	1	46601	.51	11		2°	2	49053	7.21	2
3d ⁷ (a ⁴ P)4p	у ^з Р°	$\begin{array}{c} 0 \\ 1 \\ 2 \end{array}$	$46673 \\ 46902 \\ 46727$	2.84 1.15 1.48	5 13 10	$3d^6 4s(b \ {}^4\mathrm{F})4p$	w ³ F°	4 3	49109 49243	1.49 1.45	11 14
3d7 (a 4P)4p	u ⁵ D°	$ \begin{array}{c} 4 \\ 3 \\ 2 \\ 1 \\ 0 \end{array} $	46721 46745 46889 47177 47172	1.58 .37 .56 .96 .63	$15 \\ 17 \\ 14 \\ 12 \\ 4$	3d ⁶ 4s(b ⁴ F)4p	v ³ D°	2 3 2 1	49433 49135 49243 49298	1.20 1.72 1.57 1.55	11 12 13
3d7 (a 2G)4p	$x {}^{3}\mathrm{F}^{\circ}$	$\frac{4}{3}$	46889 47093	$\begin{array}{c} 1.64 \\ 1.80 \end{array}$	$\frac{14}{17}$	$3d^{7}(a^{2}G)4p$	3° x ³ H°	3	49227 49434	30.9	1
3d ⁶ 4s(a ⁴ H)4p	z $^{3}\mathrm{H}^{\circ}$	2 6	47197 46982	2.40 5.19	14 7		<i>y</i> 11	5 4	49604 49727	1.01 1.07 .80	9 12
2J7(-4E)5	512	5 4	47008 47107	4.07 2.87	14 16	$3d^7 (a {}^2\mathrm{G})4p$	v ³G°	5 4 3	$\begin{array}{c} 49461 \\ 49628 \\ 49851 \end{array}$	$1.05 \\ 1.00 \\ .86$	12 16 11
<i>3a</i> ² (<i>a</i> ² F) <i>3s</i>	e °r	54	47006 47378 47756	4.24 3.99 3.58	17		$z {}^{1}\mathrm{D}^{\circ}$	2	49477	9.44	10
		2	48037	3.30 3.47 2.86	19 16 12	3d ⁶ 4s(a ⁶ D)5p	x ⁷ P°	3	49805	7.73	1
3d ⁷ (a ⁴ P)4p	w ³D°	$\frac{3}{2}$	47017 47136 47272	1.27 1.13 1.61	12 15 16 11	$3d^{7} (a^{2}D)4p$:	w ³ P°	$\begin{array}{c} 0 \\ 1 \\ 2 \end{array}$	49951 50043 50187	$1.09 \\ 1.29 \\ 1.50$	6 13 11
3d ⁶ 4s(a ⁴ G)4p	w ⁵ G°	6 5 4 3 2	47363 47420 47590 47693 47831	$5.90 \\ 3.29 \\ 5.36 \\ < 6.28 \\ 4.12$	5 7 10 8 7	3d ⁶ 4s (a ⁶ D)4d	<i>e</i> ⁷ F		50342 50833 51192 51149 51331 51208	0.63 .60 .55 .63 .83 .50	$3 \\ 9 \\ 11 \\ 12 \\ 10 \\ 6$
	1°	2	47420	3.43	13	3d ⁶ 4s(a ⁶ D)4d	f ⁷ D	5	50378	.72	4
$3d^{7} (a {}^{2}\text{G})4p$:	$z {}^{1}G^{\circ}$	4	47453	10.5	13			4 3 2	50808 50862 50999	.54 .59 .74	13 7 13

TABLE 1. Radiative lifetimes for energy levels in neutral iron-Continued

Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions	Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions
3d ⁶ 4s(a ⁶ D)4d	f⁵D	4	50423 50534	.80	14. 17		5°	3	51436	10.3	3
		2	50699	1.17	15	$3d^6 4s(a \ ^6\mathrm{D})4d$	e ⁷ S	3	51570	.50	11
			50981	1.25	6		6°	5	51630	3.47	8
3d ⁶ 4s(a ⁶ D)4d	<i>e</i> ⁷ P	4	50475	.86	8	$3d^{6}4s(a \ ^{6}\mathrm{D})5p$	u ⁵P°	3	51692	71.1	1
		$\frac{3}{2}$	50861	1.17	9	$3d^{7} (a^{2}P)4p$	y ¹D°	2	51708	1.34	11
$3d^{6} 4s(a \ ^{6}\mathrm{D})4d$	<i>e</i> ⁵ G	6	50523 50704	1.96	6		7°	2	51756	6.96	3
		5 4 2	50704	1.30	10		$x {}^{1}\mathrm{D}^{\circ}$	2	51762	5.07	6
		$\frac{3}{2}$	51219	.92 1.14	$11 \\ 12$	$3d^6 4s (a ^6\mathrm{D}) 4d$	e ⁵P	3	51837	2.44	13
3d7 (a 2G)4p:	z ¹ F°	3	50587	2.54	11			2	$52067 \\ 52020$	$2.65 \\ 2.71$	15 8
	$x {}^{1}G^{\circ}$	4	50614	3.90	7	$3d^7 (a {}^2\mathrm{P})4p$	u ³D°	3	51969	.81	12
$3d^6 4s(a \ ^6\mathrm{D})4d$	e 7G	7	50652	.74	1			$\frac{2}{1}$	$52297 \\ 52512$.47 .38	$\frac{11}{14}$
		5	50968 51229	.91 .89	4 9	$3d^7 (a^2D)4p$:	$t \ ^{3}\mathrm{D}^{\circ}$	1	52181	1.97	8
		4 3	$51335 \\ 51461$.79 .57	13 8			$\frac{2}{3}$	$52683 \\ 52213$	$\begin{array}{c} 1.17\\ 1.09\end{array}$	$\begin{array}{c} 11 \\ 10 \end{array}$
		$\frac{2}{1}$	$51540 \\ 51567$.58 .61	6 4	$3d^{7} (a^{2}H)4p$	w $^{3}\mathrm{H}^{\circ}$	6	52431	.96	5
3d ⁶ 4s(a ⁶ D)5p	u ⁵ F°	5	51017	7.44	2			5 4	$52613 \\ 52769$.84 1.06	6 7
		4 3	$\begin{array}{c} 51381 \\ 51619 \end{array}$	$6.09 \\ 12.3$	3	$3d^{7} (a^{2}H)4p$	y ³I°	7	52655	1.46	2
		2	51828	2.14	4			6 5	$52514 \\ 52899$	$1.24 \\ 1.22$	7 6
$3d^{6} 4s(a \ {}^{4}\mathrm{G})4p:$	<i>x</i> ³ H°	6 5	$51023 \\ 51069$	6.97 6.31	$ \begin{array}{c} 7\\ 12 \end{array} $	$3d^{7} (a^{2}P)4p$	x ³ S°	1	52858	.46	11
3d ⁶ 4s(a ⁶ D)5p	$t~^5\mathrm{D}^\circ$	4	51077	< 6.28	2	3d ⁷ (a ² P)4p:	$v~^3\mathrm{P^o}$	2	52916	.61	9
		2	51630	2.73	8		200	1	53808	.47	12
			51837 51942	5.08	1		s °D°	$\frac{3}{2}$	$52954 \\ 53275$	3.94 7.87	4. 2
$3d^6 4s(a \ ^6D)4d$	$f^{5}\mathbf{F}$	5	51103	1.28	9	$3d^7 (a \ {}^4\mathrm{F})4d$	g ${}^{5}\mathrm{F}$	5	53061	2.95	6
		4 3	51462 51604	1.12 2.09	11 13			4 3	53394 53831	3.72 4.21	11
		$\frac{2}{1}$	51705 51755	$2.12 \\ 3.45$	12 8			$\frac{2}{1}$	$54258 \\ 54386$	7.36 1.84	6 10
3d ⁶ 4s(a ⁶ D)4d	<i>e</i> ⁵ S	2	51149	1.05	8	$3d^{7} (a^{2}H)4p$	z 1 I $^{\circ}$	6	53094	1.06	4
3d ⁶ 4s(a ⁴ G)4p:	v ³ F°	2	51201	2.47	11	$3d^{7} (a {}^{4}\mathrm{F})4d$	h ⁵ D	4	53155 53546	4.39	6
		4	51305	2.05	13			2	53967	2.73	8
3d ⁶ 4s(a ⁴ D)5s	e ³D	3	51294	2.56	19	$2 J_{7} (-4E) A J$	(5D	1	59161	< 5.20	0
		1	51740 52040	2.05 2.64	19	5a' (a + f) + a	JP	3	53161 53569	< 4.78 2.09	4 12
3d ⁶ 4s(a ⁴ D)5s	g ⁵ D	4	51351	1.50	19			1	53925	2.62	11
		$\frac{3}{2}$	$51771 \\ 52050$	$1.57 \\ 1.91$	$\begin{array}{c} 23\\ 23\end{array}$	$\int 3d^7 (a {}^4\mathrm{F})4d$	$f^{5}G$	6 5	$53169 \\ 53282$	$4.15 \\ 1.82$	5 6
		$\begin{array}{c}1\\0\end{array}$	$52214 \\ 52257$	$1.52 \\ 1.48$	19 6			4 3	$53769 \\ 54161$	$2.97 \\ 2.02$	$12 \\ 12$
$3d^{7}(a^{2}\mathrm{H})4n$	u ³ C°	5	51374	111	11			2	54376	3.48	5
οα (α 11)+p	u U	4	51668 51826	.80	18 10	3d7 (a 2P)4p	z ¹ P°	1	53230	.57	11
	4°	4	51409	2.79	11	$3d^7 (a \ {}^4\mathrm{F})4d$	<i>e</i> ⁵ H	7 6	53275 53353	4.35	1

FABLE	1.	Radiative	lifetimes	for	energy	Tevels	in	neutral	iron-	Continued	ł
--------------	----	-----------	-----------	-----	--------	--------	----	---------	-------	-----------	---

Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions	Configuration	Term	J	Level cm ⁻¹	Lifetime ns	Number of tran- sitions
		5 4 3	53874 54237 54491	3.21 3.60 3.72	$3 \\ 3 \\ 1$		s ³ G°	5 4 3	55907 55906 56098	.74 .65 1.33	7 9 3
	9°	4	53329	3.35	5	3d ⁶ 4s(b ² H)4p:	<i>u</i> ³ H°	6 5	56334 56383	.89 45	4
3d ⁶ 4s (b ⁴ D)4p	t ⁵ P°	3 2	53389 54112	9.82 1.13	1 3	$2 d6 4_{2} (a 6D) E d$	1	4	56423	.46	9
	v ¹ F°	3	53661	1.98	9	$3d^6 4s(a^6\mathrm{D})5d$	2	4	56452	1.29	5
$3d^7 (a {}^2\mathrm{H})4p$	y ¹H°	5	53722	1.69	2	$3d^7 (a^2D)4p$:	<i>u</i> ³ F°	4	56593	.52	6
3d7 (a 4F)4d	<i>e</i> ³ G	5 4	53739 54067	2.28 2.79	12			3 2	56783 56859	.53 .44	7
		3	54379	3.71	3	3d ⁶ 4s(a ⁶ D)5d	3	4	56843	1.02	5
$3d^7 (a \ {}^4\mathrm{F})4d$	<i>f</i> ³ D	$\frac{3}{2}$	53748 54067	2.66 2.21	12 14	2d64c(h2H)4p	v ¹ G°	4 7	56951 57028	.42	5
	$x {}^{1}F^{\circ}$	1	54449 53763	.99	14	5 <i>a</i> +3(<i>b</i> 11)+ <i>p</i>	х I	6 5	57070 57104	.70 .73	3
3d ⁶ 4s(a ⁶ D)6s	g ⁷ D	5 4 3	53801 54125 54414	3.90 2.23	4 6		t ³ F°	4 3 2	57550 57641 57709	.51 1.26 .44	5 8 5
		$\frac{2}{1}$	54612 54748	< 3.98 3.30	4 3	$3d^6 4s(a \ ^4\mathrm{D})4d$	<i>i</i> 5D	43	57698 57814	.59 .56	7
3d7 (a 4F)4d	e ³ H	6 5 4	$53841 \\ 54267 \\ 54555$	3.51 19.6 2.90	2 1 4	3d ⁶ 4s(a ⁶ D)7s	h 7D	2 5	57974 57897	.49 2.75	9 2
$3d^{6}4s(b^{4}D)4d:$	10°(5D°?)	3	53892	1.18	9	$3d^{6}4s(a \ ^{4}\mathrm{D})4d$	g ⁵ G	6 5	$58002 \\ 58272$	$.50 \\ 1.80$	3 4
3d ⁶ 4s(a ⁴ G)4p:	t ³ G°	5 4 3	53983 54237 54600	.67 .78 .67	8 11 13			4 3 2	58520 58710 58825	$< \substack{2.17 \\ 8.82 \\ 2.10}$	5 2 3
	11°	3	54005	9.69	7	$3d^{6}4s(a \ ^{4}\mathrm{D})4d$	4	2	58213	.71	7
3d ⁶ 4s(b ⁴ D)4d:	12°(⁵ F°?)	5	54014	1.77	5		r ³G°	5 4	59927 60172	.37 .13	13
	13°	4	54301	2.05	9		. 9110	5	60365	.09	2
3d7 (a 4F)4d	$f{}^3\mathrm{F}$	$4 \\ 3 \\ 2$	54683 55125 55379	3.62 2.57 2.15	9 9 6		t ^s H ^o	5 4	60366 60549 60758	.17 .16 .18	4 4 5
	w ¹ G°	4	54811	2.94	4		q ³ G°	3	60807	.32	4
$3d^7 (a \ {}^4\mathrm{F})4d$	e ³ P		54880 55376 55727	3.85 2.47 1.08	5 5 2			3 4 3 2	53358 53610 53734 53749	3.88 7.73 8.45 2.88	7 3 6 6
	v ³ H°	4 5 6	55446 55430 55490	.88 .74 .66	6 9 5			3 4 3 3	53785 53882 54289 54357	6.62 7.13 4.58 2.30	3 4 7 7
	x ¹H°	5	55526	1.96	5 5			3	57565 60564	2.48 .93	4
3d ⁷ (a ² D)4p:	w ¹ D°	2	55754	.17	8			2	02081	1.14	3
	w ¹ F°	3	55791	.74	8	Fe II $(a \ ^{6}\mathrm{D}_{4\frac{1}{2}})$	Limit		63700		

TABLE 1. Radiative lifetimes for energy levels in neutral iron-Continued

In pure LS coupling the lowest three terms in table 1 would be metastable. This metastability is clearly exhibited in our results for the lifetimes of these levels. Though finite because of slight departures from LS coupling, the lifetimes obtained for these levels are nevertheless several orders of magnitude longer than those obtained for the nonmetastable levels. Another point of interest is the marked tendency throughout the table towards equality in the lifetimes of the individual levels of a single term, as expected.

Although the 3300 lines entering into these determinations represent more than three quarters of the classified lines observed in the laboratory, and the strongest ones at that, there are a few cases where the values stated for the lifetimes must be considered as an upper limit. For example, the $d {}^{3}F_{2}$ level at 36941 cm⁻¹, which is the lowest even level in our list, must make its strongest downward combinations in the infrared, beyond the range of our list of lines. For a few other levels, lines of significant intensity lacked reliable values of log gf. Where known, such values are marked with the "less than" (<) symbol.

Of the 408 lifetimes reported in table 1, three have been measured elsewhere by more direct methods. Karstensen and Richter [1965] measured lifetimes for $z \, {}^{3}F_{4}^{\circ}$ and $z \, {}^{3}D_{1}^{\circ}$ by a delayed coincidence method in a hollow cathode and Otten and Wagner [1967] measured the lifetime for $z \, {}^{5}F_{5}^{\circ}$ by a double resonance experiment in an atomic beam. The lifetimes in nanoseconds are compared below.

Level	This Paper	Elsewhere
z ${}^5\mathrm{F}_5^{\mathrm{o}}$	47	54
z $^3\mathrm{F}^{\mathrm{o}}_4$	238	220
z $^{3}\mathrm{D}_{1}^{\mathrm{o}}$	98	100

A detailed discussion of the accuracy of the absolute scale of the oscillator strengths from which our lifetimes are derived is given in section 5 of Corliss and Warner [1966]. From that discussion we conclude that our general scale of lifetimes may be small by about 30 ± 10 percent. In the comparison of values for the three levels shown in the preceding paragraph our scale is small by about 3 percent, which is well within the uncertainty of the individual values. This latter uncertainty cannot be precisely evaluated, but may be about 50 percent.

The work reported in this paper is based on the work originally conceived by the late W. F. Meggers and brought to fruition in his monumental "Tables of Spectral-Line Intensities," which was published in 1961 as NBS Monograph 32.

The present authors were privileged to be associated for many years with Dr. Meggers. His encouragement and good counsel were always a source of inspiration to us.

4. References

Corliss, C. H. and Bozman, W. R. (1962), Experimental transition probabilities for spectral lines of seventy elements, NBS Mono. 53 (July 20, 1962).

Corliss, C. H. and Tech, J. L. (1967), Oscillator strengths and transition probabilities for 3328 Lines of Fe I. NBS Mono. 108.

 Corliss, C. H. and Warner, B. (1964), Ap. J. Suppl. 8, 395, No. 83.
 Corliss, C. H. and Warner, B. (1966), J. Res. NBS 70A (Phys. and Chem.), No. 4, 325.

Einstein, A. (1917), Phys. Zeits. 18, 121.

Huber, M. and Tobey, F. L. (1967), Shock Tube Spectroscopy Laboratory, Scientific Report No. 16, Harvard College Observatory. Also Astrophys. J. (in press).

Karstensen, F. and Richter J. (1965), New Method of Measuring Transition Probabilities, AF EOAR Grant 63-110.

Mitchell, A. G. C. and Zemansky, M. W., Resonance Radiation and

Excited Atoms, p. 145 (Cambridge University Press, 1934). Otten, E. W. and Wagner, R. (1967), Physica **33**, 283.

Warner, B. and Cowley, C. R. (1967), J. Quant. Spectrosc. Radiative Transfer 7, 751.

(Paper 71A6-472)