


Reference

NBS
Publi-
cations



NOV 17 1983

NBSIR 83-2742 (R)

Photonuclear Data-Abstract Sheets
1955 - 1982
Volume III (Carbon)


U.S. DEPARTMENT OF COMMERCE
National Bureau of Standards
National Measurement Laboratory
Center for Radiation Research
Washington, D.C. 20234

July 1983



QC
100
.U56
83-2742
V. III
1983

DEPARTMENT OF COMMERCE
BUREAU OF STANDARDS



Ref.

QC100

.456

NO. 83-2742

V. 3

1983

NBSIR 83-2742

PHOTONUCLEAR DATA—ABSTRACT SHEETS
1955 - 1982
VOLUME III (CARBON)

E. G. Fuller, Henry Gerstenberg

U.S. DEPARTMENT OF COMMERCE
National Bureau of Standards
National Measurement Laboratory
Center for Radiation Research
Washington, D.C. 20234

July 1983

U.S. DEPARTMENT OF COMMERCE, Malcolm Baldrige, *Secretary*
NATIONAL BUREAU OF STANDARDS, Ernest Ambler, *Director*




TABLE OF CONTENTS

Table of Contents	i
Introduction.	1
Carbon (A=11)	3
Carbon (A=12)	11
Carbon (A=13)	459
Carbon (A=14)	505
Definition of Abbreviations and Symbols	515



Photonuclear Data-Abstract Sheets
1955-1982


I. Introduction

As used in connection with this collection of data-abstract sheets, the term photonuclear data is taken to mean any data leading to information on the electromagnetic matrix element between the ground state and excited states of a given nuclide. The most common types of reactions included in this compilation are: (e, e') , (γ, γ) , (γ, γ') , (γ, n) , (γ, p) , etc. as well as ground-state particle capture reactions, e.g. (α, γ_0) . Two reactions which fit the matrix element criterion are not included in the compilation because of their rather special nature. These are heavy particle Coulomb excitation and the thermal neutron capture reaction (n, γ_0) . While the energy region of particular interest extends from 0 to 150 MeV, papers are indexed which report measurements in the region from 150 MeV to 4 GeV. Most of the experiments listed are concerned with the excitation energy range from 8 to 30 MeV, the region of the photonuclear giant resonance.

The hierarchical grouping of the photonuclear data-abstract sheets within the file is by: 1. Target Element, 2. Target Isotope, and 3. by the Bibliographic Reference Code assigned to the paper from which the data on the sheet were abstracted. In this file, colored pages are used to mark the beginning and end of the sheets for each chemical element. A brief historical sketch of the element is given on the divider sheet marking the start of each section; the information for this sketch was derived from references such as the Encyclopaedia Britannica. In those cases where the sheets for a given element make up a major part of a volume, colored pages are also used to delineate sections pertaining to the individual isotopes of the element. Each of the sections of the file, as delineated by two colored divider sheets, represents a 27 year history of the study of electromagnetic interactions in either a specific nuclide or a specific element.

The data-abstract sheets are filed under the element and/or isotope in which the ground-state electromagnetic transition takes place. For example, the abstract sheet for a total neutron yield measurement for a naturally occurring copper sample would appear in the elemental section of the copper file. On the other hand, a measurement of the ^{62}Cu 9.73 minute positron activity produced in the same sample by photons with energies below the three-neutron separation energy for ^{65}Cu (28.68 MeV) would be filed with the sheets for ^{63}Cu . Similarly a measurement of the ground-state neutron capture cross section in ^{12}C would be filed under ^{13}C while the corresponding ground-state alpha-particle capture cross section would be filed under ^{16}O .

At the end of this volume there is a master list of the abbreviations that have been used in the index section of the abstract sheets. The listings are those used in the final published index, Photonuclear Data Index, 1973-1981, NBSIR 82-2543, issued in August 1982 by the U. S. Department of Commerce, National Bureau of Standards, Washington, DC 20234. In some cases two notations are entered for the same quantity. The second entry is the abbreviation that was used in one or more of the earlier published editions of the index.



CARBON

Z=6

Carbon exists in three allotropic forms; diamond, graphite, and amorphous carbon. Ancient Hindu writings indicate that diamonds were known at least as early as 1200 B. C. The word diamond is a corruption of the Greek word *adamas* "the invincible". Graphite was often confused with other minerals having a similar appearance; its first recognition is obscured in antiquity. C. W. Scheele demonstrated in 1779 that graphite oxidized to carbon dioxide. This was the first proof of the chemical constitution of this important gas. The name graphite comes from the Greek verb *graphein* "to write"; the name originated with A. G. Werner, the father of German geology, in 1789. Amorphous carbon is generally obtained by some type of thermal decomposition or partial combustion of natural organic matter such as coal, petroleum, gas, and timber. Historical references to these related carbon compounds abound in the literature. The inhabitants of the ancient city of Nineveh (founded not later than 6000 B. C.) used an asphalt based mortar for construction. Some translations of the Old Testament call this material "pitch" or "slime". Noah was told, when building his ark, to "pitch it within and without with pitch". When the Tower of Babel was built, Noah's descendants "had brick for stone, and slime had they for mortar".

C
A=11C
A=11

Elem. Sym.	A	Z
C	11	6
Ref. No. 61 Ja 2		JHH

Method 1.4 MeV Cockcroft-Walton generator; NaI

Reaction	E or ΔE	E ₀	Γ	∫σdE	J ^π	Notes
$B^{10}(p,\gamma)$	0.75- 1.30	1.14	$(2J+1)\Gamma_{\gamma} =$ 10ev		$5/2^{-}$	$E_{\gamma 0} = 3.74$ MeV; $W(\theta_{\gamma}) = 1 + (0.21 \pm 0.06)P_2(\cos\theta)$

Fig. 3. Decay scheme of the 9.74 MeV C^{11} level.

TABLE 3
Experimental partial decay widths

Decay energy (MeV)	$2J_i - 1$	$2J_f - 1$			LMM
		E1	M1	E2	
9.74	0	0.02	0.5	30	0.7
7.42	1	0.02	0.7	130	0.7
5.29	3	0.10	4.3	2700	0.7

Elem. Sym.	A	Z
C	11	6

Method Tandem van de Graaff; NaI

Ref. No. 62 Op 2
 JHH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
B ¹⁰ (p,γ ₀)	2.4-7.2	12.66				<p>σ(p,γ₀) data given in curves III in Figure 2.</p> <p>θ = 0° on (p,γ₀)</p>




Fig. 2 Yield curves of gamma rays from the B¹⁰(p,γ₀), B¹⁰(p,γ₁) and B¹⁰(p,γ₂) reactions within the energy range E₀ = 2.4-7.2 MeV. For clarity, only some of the experimental points are shown as the yield curves of the 40 and 717 keV gamma rays.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	11-24	D	2-17	NAI-D	3-24	DST
				(2.6 - 17.0)			253




Fig. 5. The coefficients A_0 , a_1 and a_2 for $^{10}\text{B}(p, \gamma)^{11}\text{C}$ obtained from least-squares fits to $W(\theta) = A_0(1 - \sum a_n P_n)$ with terms up to and including P_2 . The solid angle subtended by the detector at tenuates a_2 by about 3%.




Fig. 4. Angular distributions of $^{10}\text{B}(p, \gamma)^{11}\text{C}$ plotted in units of $1/A_0$, where A_0 is obtained from the least-squares fit to $W(\theta) = A_0[1 - \sum a_n P_n(\cos \theta)]$. The curves are the least-squares fits with terms up to and including P_2 .




Fig. 3. The 90° yield curve of $^{10}\text{B}(p, \gamma)^{11}\text{C}$. The arrows mark energies at which angular distributions were measured.




Fig. 4. The angular distribution for $^{10}\text{B}(p, \gamma)^{11}\text{C}$ with terms up to and including P_2 and P_4 .




Fig. 11. Comparison of (p, γ) reactions on ^{10}B , ^{11}B [ref. ¹²], ^{12}C [ref. ¹³], ^{14}N and ^{15}N [refs. ¹³, ¹⁴].

- ¹²J.F. Eichelberger, G.R. Grove and L.V. Jones, Mound Lab. progress report MLM 1100 (1963), Miamisburg, Ohio.
¹³C.J. Christensen, A. Nielson, A. Bahnsen, W.K. Brown, B.M. Rustad, Phys. Lett. **20B**, 11 (1967).
¹⁴F.E. Emery and T.A. Rason, Phys. Rev. **140**, A2039 (1965).
¹⁵R.J. Blin-Stoyle, to be published.

ELEM. SYM.	A	Z
C	11	6

METHOD	REF. NO.
	78 K1 5

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,N	ABX	19- 30	D	30	ACT-I		4PI


The cross section for the production of the ¹¹C activity by bombarding ¹²C with 30 MeV electrons has been measured absolutely. The result, 11.9 ± 0.2 μb, has a smaller absolute error than any previous measurement. This value corresponds to a (γ,n) cross section integrated to 30 MeV of 41.0 ± 0.6 McV mb, based on a distorted-wave Born-approximation electric dipole virtual photon spectrum.

TABLE I. The cross section for the production of ¹¹C by 30 MeV electrons.

Run	Cross section
I	12.15 ± 0.31
II	11.73 ± 0.47
III	11.34 ± 0.32
IV	11.99 ± 0.34
V	11.78 ± 0.32
Weighted average	11.92 ± 0.15

TABLE II. Comparison with other experiments.

σ (μb)	Projectile	Detector	Reference
12.7 ± 1.0	Electrons	Activity	7
10.1 ± 1.0	Photons	BF ₃	8
11.6 ± 1.2	Photons	Activity	9
13.9 ± 1.1	Photons	Activity	10
10.7 ± 0.9	Photons	BF ₃	11
11.9 ± 0.2	Electrons	Activity	This work



C
A=12

C
A=12

C
A=12

Elem. Sym.	A	Z
C	12	6

Method	Linac; neutron yield, radioactivity; Faraday Cup	Ref. No.	55 Ba 1	EGF
--------	--------------------------------------------------	----------	---------	-----

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C ¹² (γ,n)	Bremss. 18-260	22.5		$\int_0^{20} = 0.056 \pm 0.003$ MeV-b $\int_0^{250} = 0.080 \pm 0.01$ MeV-b		σ = 8.3 mb




FIG. 1. Yield curve for the reaction C¹²(γ,n)C¹¹. The data shown as dots were taken with a tantalum radiator of 169.77 mg/cm². However, the yield as expressed by Eq. (1) has been divided by N_e and by the thickness of the radiator in radiation lengths so that the ordinate scale represents the cross section for the reaction induced by the bremsstrahlung of one electron in a unit radiator.

TABLE I. Comparison of some parameters of the C¹²(γ,n)C¹¹ cross section as determined by different investigators.

Reference	Mean energy (MeV)	Energy at peak of cross section (MeV)	∫ ₀ ²⁵ σ(k)dk (MeV-barns)	∫ ₀ ²⁵⁰ σ(k)dk (MeV-barns)
This paper	34	22.5	0.032 ± 0.003	0.080 ± 0.01
4	---	21.3	0.029	...
5	---	22.0	0.027	...
6	27	---	---	0.090 ± 0.022
7	23-32	---	---	0.086 ± 0.02

Montalbetti, Katz, and Goldemberg, Phys. Rev. 91, 659 (1953).
 R. Nathans and J. Halpern, Phys. Rev. 93, 437 (1954).
 E. Strauch, Phys. Rev. 81, 973 (1951).
 L. Marshall, Phys. Rev. 83, 345 (1951).




FIG. 3. Cross-section curve for the reaction C¹²(γ,n)C¹¹. The shape of the curve was determined by the photon-difference analysis of the yield curve of Fig. 1, but the absolute value of the ordinate scale was determined from the data with a copper radiator. The smooth curve shown is consistent with the data, but because of the insensitivity of the method the detailed shape of the high-energy portion is uncertain.




FIG. 2. Yield curve for the reaction C¹²(γ,n)C¹¹ extended to 280 MeV. Absolute points taken with the tantalum radiator are shown as plain dots, while the points labeled with a subscript taken with a thick copper radiator, have been arbitrarily normalized. The ordinate scale is the same as for Fig. 1. The curves A to E are computed yield curves assuming various cross-section curves described in the text.

- "E" - Fig. 3 cross section cut off at 36 MeV.
- "A" - "E" + bump centered at 50 MeV and 12 MeV with magnitude to fit at 250 MeV.
- "B" - "E" + tail starting at 44 MeV with σ = 1.27 mb ~ (1/E²)
- "D" - "E" + tail (1/E²) fitted at 34 MeV with σ = 1.27 mb.
- "C" - tail + rise at meso C¹²(γ,n=0)C¹¹.

METHOD				REF. NO.		ANGLE
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR	
			TYPE	RANGE	TYPE	RANGE
Synchrotron; ion chamber monitor; $^{12}\text{C}(n,2n)$ threshold detector.				55 Ba 5		EGF
G, XN	ABY	30-200	C	150-250	THR-1	30-




FIG. 2. Angular distribution of photoneutrons with energies higher than 30 MeV. ● - C_{250} ; + - C_{200} ; ▲ - Be_{250} ; ○ - Al_{250} ; * - Pb_{250} ; ■ - data of work⁵.




FIG. 3. The dependence of the yield of photoneutrons with energies higher than 30 MeV in carbon (in units 10^{-27}cm^2 per eif. quantum) on the maximum energy of retarded radiation.




FIG. 4. The dependence of the yield of photoneutrons with energies higher than 30 MeV at an angle of 90° (in units 10^{-28}cm^2 per eif. quant steradian on the mass number A).

METHOD						REF. NO.		EGF
						55 Ca 2		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
G,3A	ABX	12-18	D	15.18	EMU-D	4-12	4PI	
				(14.8, 17.6)	ION-D	6-12	4PI	

Methane filled proportioned counter used to get absolute σ at 17.6 MeV.
 Emulsion data normalized at 17.6 MeV.

E 17.6 14.8 12.3 (MeV)
 σ 1.70 \pm 24 0.33 \pm 0.07 1.15 \pm 0.6 10^{-28} cm²


Elem. Sym.	A	Z
C	12	6

Method	Ref. No.
Synchrotron; neutron angular distribution; scintillator; ion chamber	55 Di 1 NVB

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C(\gamma, xn)$	70					Anisotropy suggests direct effects in light nuclei. Curves fitted to $a + b \sin^2 \theta$

TABLE II
 EXPERIMENTAL VALUES FOR b/a

Target	Correction factor for self-scattering	Corrected b/a
Lead	1.10	-0.08 ± 0.08
Tin	1.08	0.12 ± 0.17
Copper	1.48	0.23 ± 0.15
Iron	1.35	0.09 ± 0.25
Aluminum	1.17	0.36 ± 0.29
Carbon	1.8	1.6 ± 0.8
Beryllium (1)	2.6	
Beryllium (2)	1.35	1.2 ± 0.4



REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	4-10	D	80-187	MAG-D		DST

LEVELS 4.4, 7.7, 9.6

TABLE I. Cross section of elastic and inelastic scattering peaks in microbarns per steradian for various angles at an incident energy of 187 Mev. The radiative correction is included.

Lab angle	Elastic	4.4 Mev	7.68 Mev	9.61 Mev
35°	28.3 ± 1.4	2.35 ± 0.28	1.05 ± 0.22	0.49 ± 0.20
45°	68.7 ± 3.5	1.13 ± 0.19	0.24 ± 0.10	0.19 ± 0.09
		1.41 ± 0.11	0.44 ± 0.08	
50°	36.3 ± 1.1	1.12 ± 0.22	0.39 ± 0.09	0.24 ± 0.07
		1.46 ± 0.16	0.49 ± 0.07	0.39 ± 0.08
60°	0.42 ± 0.38	0.828 ± 0.083	0.114 ± 0.035	0.121 ± 0.030
		0.715 ± 0.086	0.180 ± 0.027	0.101 ± 0.021
70°	2.72 ± 0.11	0.981 ± 0.063	0.080 ± 0.025	0.133 ± 0.035
		0.500 ± 0.050	0.092 ± 0.017	0.127 ± 0.019
80°	0.711 ± 0.028	0.251 ± 0.025	0.048 ± 0.010	0.068 ± 0.012
		0.286 ± 0.034	0.051 ± 0.016	0.073 ± 0.013
90°	0.199 ± 0.014	0.179 ± 0.021	0.021 ± 0.005	0.043 ± 0.010
		0.457 ± 0.017	0.013 ± 0.006	0.053 ± 0.010

TABLE II. Ratio of cross section for excitation of 4.43-Mev level to point-nucleus scattering cross section for carbon.

Energy (Mev)	Lab angle	Ratio
80	90°	0.0035 ± 0.0008
150	90°	0.0119 ± 0.0015
150	80°	0.0155 ± 0.0017
150	70°	0.0145 ± 0.0016
187	90°	0.0092 ± 0.0021
		0.0178 ± 0.0030
187	80°	0.0167 ± 0.0017
		0.0190 ± 0.0023
187	70°	0.0179 ± 0.0023
		0.0184 ± 0.0019
187	60°	0.0157 ± 0.0016
		0.0135 ± 0.0016
187	50°	0.0103 ± 0.0020
		0.0131 ± 0.0014
187	45°	0.0066 ± 0.0011
		0.0081 ± 0.0010
187	35°	0.0048 ± 0.0006




FIG. 4. The angular distributions of the elastic and inelastic peaks at 187 Mev in carbon. The absolute cross sections have been obtained by the proton-normalization method discussed in the text.

(over)




FIG. 5. The angular distributions in carbon at 150 Mev of the elastic and inelastic peaks. The absolute cross sections have been obtained by the proton-normalization method discussed in the text.




FIG. 10. The cross sections for the elastic and inelastic peaks at 90° as a function of energy.

H. Glattli, E. Loepfe and P. Stoll
 Helv. Phys. Acta 28, 366 (1955)

C

12

6


METHOD

REF. NO.

55 G1 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.A	SPC	17	D	14.17	EMU	1-7	4PI

Li(p, γ) source

METHOD					REF. NO.		EGF
[Page 1 of 4]					55 Go 1		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,3A	SPC	THR-60	C	60	EMU-D		DST

E^* represents at excitation of Be^3
 All angles are in center of mass system

θ_1 = direction of first α

θ_c = angle between α_1 and break-up direction of Be^3

θ_{23} = direction of break-up relative to γ -beam





FIGURE 2. Angular distributions of α , in ground-state transitions: (a) $13 \text{ MeV} \leq E_\gamma < 15.6 \text{ MeV}$; (b) $15.6 \text{ MeV} \leq E_\gamma < 20 \text{ MeV}$; (c) $20 \text{ MeV} \leq E_\gamma < 26.4 \text{ MeV}$. The fitted, full-line curves have the general form defined by equation (A 2), and determine the parameter values (a) $k^2 = \infty$; (b) $k^2 = 0.9$, $2k \cos \eta = -0.45$; (c) $k^2 = 10$.

METHOD				REF. NO.		EGF	
				[Page 2 of 4] 55 Go 1			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




FIGURE 3. E^* histograms (three values for each star) for stars with $E_\gamma < 25$ MeV. (a) < 14.2 ; (b) 14.2 to 16.6 ; (c) 16.6 to 17.0 ; (d) 17.0 to 18.0 ; (e) 18.0 to 20.0 ; (f) 20.0 to 25.0 MeV. Stars which give an E^* value < 0.3 MeV are not included. Shading indicates stars which fail to give an E^* value between 1.0 and 5.0 MeV. The full-line curves correspond to the following values of the parameter b in equation (10): (a) 3.7 ; (b) 1.0 ; (c) 0 ; (d) 1.9 ; (e) 1.2 ; (f) 0.8 .

METHOD

REF. NO.

[Page 3 of 4]

55 Go 1

EGF

DETECTOR

ANGLE

TYPE

RANGE




FIGURE 4. Angular distributions and correlation in 2.05 MoV level transitions; $17.8 \text{ MeV} \leq E_\gamma < 18.6 \text{ MeV}$. The curves indicate possible theoretical results for a $J=2$ level, and $E1+E2$ (full lines) or $M1+E2$ (dashed lines) γ -ray interactions.




FIGURE 5. E^* histograms (three values for each star) for stars with (a) $25 \text{ MeV} \leq E_\gamma < 26.4 \text{ MeV}$; (b) $E_\gamma \geq 26.4 \text{ MeV}$. Stars which give an E^* value $< 0.3 \text{ MeV}$ are excluded from (a). In (b), shading indicates stars which fail to give an E^* value between 16.0 and 17.3 MeV.




FIGURE 6. Angular distributions and correlation in transitions to the 16.8 MoV level of ${}^9\text{Be}$; $E_\gamma \geq 26.4 \text{ MeV}$. The fitted curves assume $E1$ γ -ray interaction, and $J=2$ (full line) or $J=4$ (dashed line) for the ${}^9\text{Be}$ level.

METHOD

REF. NO.

[Page 4 of 4]

55 Go 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	

TABLE 2. RELATIVE INTENSITIES OF TRANSITIONS TO THE GROUND STATE AND EXCITED LEVELS OF ⁹Be

E_γ (MeV)	N	level excitation (MeV)						
		0.00	2.05	4.0	8 and 10	15	16.8	17.6
10.0 to 13.0	106	19	81	—	—	—	—	—
13.0 to 14.2	98	18	82	—	—	—	—	—
14.2 to 15.6	222	6	94	—	—	—	—	—
15.6 to 17.0	325	7	88	5	—	—	—	—
17.0 to 18.6	970	3	89	8	—	—	—	—
18.6 to 20.0	315	7	75	13	5	—	—	—
20.0 to 22.6	110	15	51	20	14	—	—	—
22.6 to 25.0	151	12	64	5	17	2	—	—
25.0 to 26.0	50	14	25	43	12	6	—	—
26.0 to 26.4	20	5	20	20	10	35	10	—
26.4 to 29.0	96	1	5	2	10	63	19	—
≥ 29	120	3	4	—	2	73	18	—

Column 2 gives the number (N) of stars analyzed. The intensities are expressed as percentages. About one in six of the stars assigned to column 3 of the table (13.5 MeV level) may involve the suspected 10.4 MeV level.

TABLE 3. γ -RAY ABSORPTION PROCESSES INITIATING THE ¹²C(γ , 3α) REACTION

E_γ (MeV)	E1 (%)	E2 (%)	M1 (%)	$r(E1)$	$r(E2)$
13.0 to 15.6	0	70	30	—	0.13
15.6 to 17.0	35	65	—	0.07	0.08
17.0 to 18.6	75	25	—	0.04	0.02
18.6 to 20.0	70	30	—	0.11	0.05
20.0 to 22.0	33	67	—	0.22	0.33
22.0 to 26.0	60	40	—	0.26	0.25
≥ 26.0	100	(5?)	—	?	?

At the lower γ -ray energies, for which there is no conclusive evidence of E1 absorption, $M1 + E2$ is assumed. This is consistent with the fact that $1^-(T=1)$ levels do not occur below about 16 or 17 MeV in ¹²C (Inglis 1953). In the last two columns, r = (ground-state/2.06 MeV level) branching ratio.

Elem. Sym.	A	Z
C	12	6
Method		Ref. No.
emulsions; bremsstrahlung		55 Ha 1
		EGF

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
----------	---------	----------------	---	------	----	-------

$C^{12}(\gamma, 3\alpha)$

27
33

Analysed data in terms of:
 $\gamma + C^{12} \rightarrow Be^{8*} + \alpha_1$
 $Be^{8*} \rightarrow 2\alpha$

Define α_1 are high energy α . Data for two bremsstrahlung energies are not separated. (also O^{16})

Measured 500 coplaner 3 prong stars; used only those for which a momentum balance to within $(1 \text{ MeV})^{1/2}$ was obtained; 485 satisfied this condition.




FIG. 1. Number of C^{12} disintegrations as a function of the sum of the kinetic energies of the three alpha particles $E_{\alpha\alpha}$. The lower abscissa scale refers to the excitation energy of Be^8 .




FIG. 2. (A) Number of stars vs Be^8 excitation energy for all C^{12} events for which $E_{\alpha\alpha} < 12.5$ Mev. (B) Distribution in energy of the first emitted alpha particle.




FIG. 3. (A) Distribution in energy of the first emitted alpha. (B) Number of stars vs Be^8 excitation energy for all C^{12} events for which $E_{\alpha\alpha} > 12.5$ Mev.




FIG. 4. (A) Angular distribution of the first emitted alpha for $E_{\alpha\alpha} < 12.5$ Mev and $E_{\alpha\alpha} > 12.5$ Mev. (B) Energy distribution of the first emitted alpha for $E_{\alpha\alpha} < 12.5$ Mev and $E_{\alpha\alpha} > 12.5$ Mev. (C) Angular distribution of the first emitted alpha for $E_{\alpha\alpha} < 12.5$ Mev and $E_{\alpha\alpha} > 12.5$ Mev.

METHOD				REF. NO.			
Synchrotron; ZnS counter; ion chamber				55 Jo 1		NVB	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	RLY	THR - 65	C	65	SCI-D	14 - +	DST




FIG. 5. The angular distributions of protons with an energy above 14 Mev.

TABLE I. Target thickness and the constants a and b in the angular distribution curve $a + (b \sin \theta + b \sin \theta \cos \theta)^2$.

Element	Target thickness mg/cm ²	a	b
Carbon	182	0.32	0.80
Aluminum	274	0.58	1.35
Nickel	352	0.94	1.45
Molybdenum	295	0.62	2.00




FIG. 10. The relative yield per mole for protons above 14 Mev as a function of the atomic number.

METHOD					REF. NO.		EGF
					55 Sp 2		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	RLY	THR - 20	C	17-20	ACT-I		4PI

BREAKS




FIG. 2. $C^{12}(\gamma,n)C^{11}$ yield curve near threshold. The data from Fig. 1 are shown by solid dots. The open circles represent the results of an earlier determination. The two arrows indicate the position of the threshold as erroneously quoted on two occasions.

TABLE I. Comparison of $C^{12}(\gamma,n)$ yield-curve breaks with those observed by Katz *et al.*

	Break energy (Mev)		Mev above threshold	
	Present results	Katz <i>et al.</i> *	Present results	Katz <i>et al.</i> *
Threshold (cal pt)	18.73 ± 0.03	18.73 ± 0.03	0.00 ± 0.04	0.00 ± 0.05
First break	19.10 ± 0.05	19.30 ± 0.05	0.37 ± 0.05	0.57 ± 0.05
Second break	19.55 ± 0.05	19.75 ± 0.05	0.82 ± 0.05	1.02 ± 0.05

* See reference 3.

Elem. Sym.	A	Z
C	12	6

Method 90° plates

Ref. No.	EGF
56 Co 1	

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes										
C ¹² (γ,p)	Bremss. 24			~ 60 MeV-mb		<p>Proton yield at 24 MeV Bremss was $12 \pm 3 \times 10^4$ p's/mole r. (1.97 mg/cm² polyethylene)</p> <p>Assuming ground state, resonances:</p> <table> <tr><td>23.1 MeV</td><td></td></tr> <tr><td>22.6 MeV</td><td>12 MeVmb</td></tr> <tr><td>21.5 MeV</td><td>9 MeV-mb</td></tr> <tr><td>20.8 MeV</td><td>6 MeV-mb</td></tr> <tr><td>17.8 MeV</td><td>2 MeV-mb</td></tr> </table> <p>Figure 9: "curved line" - detailed balance; Blair, Kington and Willard, Phys. Rev. 100, 21 (1955); Haines, Day - Phys. Rev. <u>91</u>, 599 (1953).</p>	23.1 MeV		22.6 MeV	12 MeVmb	21.5 MeV	9 MeV-mb	20.8 MeV	6 MeV-mb	17.8 MeV	2 MeV-mb
23.1 MeV																
22.6 MeV	12 MeVmb															
21.5 MeV	9 MeV-mb															
20.8 MeV	6 MeV-mb															
17.8 MeV	2 MeV-mb															



Fig. 9. Photon absorption cross-section curve for the electron photoprotons from carbon obtained from Fig. 5 on the assumption that the resultant B¹¹ nucleus is always left in its ground state. This curve gives an upper limit to the cross section for photon energies below about 20.5 Mev and a lower limit for photon energies above 23 Mev. The curve is uncertain above photon energies of 23 Mev because of statistical uncertainties and corrections. The solid curve is the cross section predicted from B¹¹(p,γ) data by detailed balancing. The point at 17.6 Mev is due to Mann and Titterton reference 10.

Elem. Sym.	A	Z
C	12	6
Ref. No. 56 Fa 1		EGF

Method ZnS paraffin; 23 MeV Bremss.

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n)$	Bremss. 23					<p>Angular distribution measured: $60^\circ - 160^\circ$</p> <p>$\sigma(\theta) = 1 + (1.55 \pm 0.88) \sin^2 \theta$; this about same as obtained for p's in this energy range.</p> <p>ZnS paraffin counter is insensitive to γ's; counts neutrons with $E_n > 0.3$ MeV.</p>




FIG. 2. Angular distribution of neutrons from carbon bombarded with bremsstrahlung of 23-MeV maximum energy

METHOD Linac				REF. NO. 56 Fr 1		NVB	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	0-35	D	187	MAG-D		DST

RMS radius of C^{12} : $(2.37 \pm 0.05) 10^{-13}$ cm.

J-PI

$$r_0 = (1.33 \pm 0.02) 10^{-13} \text{ cm}$$

9.61 MeV level 2^- or 0^-




FIG. 1. Scattering at 187 Mev at an angle of 110°. This is composite of two runs.




FIG. 3. Elastic and inelastic angular distributions at 187 Mev. The absolute cross sections have been obtained by the proton-normalization method discussed in the text.

METHOD

REF. NO.

56 Fr 1

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




FIG. 2. Scattering at 187 Mev at an angle of 80°.

TABLE I. Results. Differential cross sections in the laboratory system in units of 10^{-28} cm²/sterad for angles of 90° or less and in units of 10^{-28} cm²/sterad for angles greater than 90°. Errors are statistical only.

<i>E</i>	<i>θ_{lab}</i>	Elastic	4.43 Mev	7.65 Mev	9.61 Mev
187 Mev	35°	280 ± 14.0	2.32 ± 0.23	1.04 ± 0.22	0.481 ± 0.190
	45°	68.7 ± 2.3	1.41 ± 0.16 1.13 ± 0.19	0.440 ± 0.083 0.240 ± 0.096	0.185 ± 0.087
	50°	36.3 ± 1.5	1.46 ± 0.16 1.12 ± 0.23	0.490 ± 0.069 0.391 ± 0.090	0.391 ± 0.078 0.243 ± 0.070
	60°	9.53 ± 0.38	0.724 ± 0.087 0.834 ± 0.170	0.182 ± 0.027 0.146 ± 0.035	0.143 ± 0.024 0.122 ± 0.031
	70°	2.80 ± 0.11	0.514 ± 0.031 0.498 ± 0.065	0.095 ± 0.017 0.082 ± 0.027	0.131 ± 0.020 0.137 ± 0.036
	80°	0.748 ± 0.030	0.300 ± 0.036 0.263 ± 0.026	0.053 ± 0.017 0.040 ± 0.011	0.079 ± 0.014 0.072 ± 0.013
	90°	0.213 ± 0.015	0.168 ± 0.019 0.192 ± 0.023	0.014 ± 0.006 0.022 ± 0.006	0.062 ± 0.011 0.046 ± 0.011
	100°	46.6 ± 8.4 45.2 ± 2.7	83.6 ± 9.2 65.1 ± 4.6
	110°	9.84 ± 0.69	33.9 ± 2.0 35.8 ± 2.1	1.86 ± 0.93	12.6 ± 1.6
	120°	2.45 ± 0.30 1.90 ± 0.15	17.3 ± 1.21		
	130°	0.569 ± 0.069	8.83 ± 0.80
	138°	0.065 < 0.100	4.29 ± 0.30	< 0.305	2.78 ± 0.56 -1.11
150 Mev	70°	9.77 ± 0.39	0.610 ± 0.067	0.148 ± 0.025	0.103 ± 0.021
	80°	3.23 ± 0.13	0.323 ± 0.042	0.034 ± 0.017	0.081 ± 0.021
	90°	1.20 ± 0.05	0.170 ± 0.019	0.043 ± 0.011	0.047 ± 0.011
80 Mev	90°	3.48 ± 0.18	0.182 ± 0.040

Elem. Sym.	A	Z
C	12	6
Ref. No.		NVB
56 Li 1		

Method Synchrotron; $p + \alpha$ cross sections; nuclear emulsions

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, p+\alpha)$	Bremss. 70			$\int_{25}^{35} = 0.68 \text{ MeV-mb}$ $\int_{35}^{40} = 0.65 \text{ MeV-mb}$		




Fig. 4. Variation of cross-section with photon energy in the $p + \alpha$ reaction in carbon and nitrogen based on 32-stars and 21-stars respectively.

ELEM. SYM.	A	Z
C	12	6
REF. NO.	56 Ma 1	
	NVB	

METHOD				Van de Graaf; proton cross section; nuclear emulsion		REF. NO.		56 Ma 1		NVB	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE				
			TYPE	RANGE	TYPE	RANGE					
G,P	ABX	18	D	18	EMU-D			4 PI			
		(17.63)		(17.63)							
		(1.19±0.21) 10 ⁻²⁷ cm ²									

In Figure 5, "A, B, C, D and E" indicate five independent observations.

$$\frac{\sigma(\gamma, p)}{\sigma(\gamma, \alpha)} = 7.00 \pm 0.73$$

based on $\sigma(\gamma, 3\alpha)$ of Carver, Hay and Titterton, $\sigma(\gamma, p) = 1.19 \pm 0.21 \times 10^{-27} \text{ cm}^2$




Figure 2. Corrected number-energy histograms derived for the five sets of data by treating all tracks as protons are shown in A, B, C, D, and E. The sum of all the data (with a change of scale) is shown in F. The proton peaks corresponding to the ²⁷Al(γ, p)²⁶Mg reaction centre at 1.5 MeV.

METHOD						REF. NO.		
Self-absorption and activation curve analysis.						[Page 1 of 2] 56 Tz 1		EGF
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
G,N	RLY	22 - 24	C	22-24	ACT-I		PI	

ϕ = ratio of yields of detector placed before and after absorber.




Fig. 1.
 La courbe 1 représente $\phi(E)$. Elle comporte des mesures faites le 31 janvier 1956 et le 13 mars 1956.
 Les courbes 2 et 3 sont des courbes d'activation grossières faites dans des conditions différentes pour s'assurer de la correction de l'étalonnage en énergie vers 23 MeV.

METHOD

Self-absorption and activation curve analysis.

[Page 2 of 2]

REF. NO.

56 Tz 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 2.

Nous avons choisi arbitrairement une courbe de section efficace composée de pics étroits et de résonances larges. Leurs sections efficaces intégrées sont indiquées. La courbe d'activation a été obtenue à l'aide du spectre de Schiff. On constate que le domaine où la pente varie est très inférieur à la largeur de la résonance correspondante (de l'ordre 10 fois plus petit).

REF. C. Tzara
 J. Phys. Radium 17, 1001 (1956)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	
	56 Tz 2	EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	LFT	23	C	22 - 23	ACT-I		4PI

Self-absorption measurement to give limit on width of level responsible for break in neutron yield curve at 22.3 MeV.

WIDTH GREATER .4 MEV

METHOD	REF. NO.
	57 Ba 1 EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABY	30 - 260	C	120-260	THR-I	21	DST

$^{12}\text{C}(\text{N},2\text{N})^{11}\text{C}$ threshold detector.




FIG. 4. Angular distributions of neutrons of energy > 20 Mev from carbon and deuterium. O — yield of neutrons from carbon for $E_{\gamma m} = 225$ Mev; \times — the same for $E_{\gamma m} = 170$ Mev; \bullet — yield of neutrons from deuterium for $E_{\gamma m} = 255$ Mev. The solid curve gives the angular distribution of neutrons from deuterium for $E_{\gamma m} = 170$ Mev, obtained from the results, Refs. 4 — 10, transposed for neutrons.

TABLE I. Yield of photoneutrons of energy ≥ 20 Mev from deuterium and carbon

$E_{\gamma m}$ (Mev)	Yield from deuterium (without meson production) according to Refs. 4-10 $10^{24} \text{ cm}^2/Q$	Total yield from deuterium $\sigma_D \times 10^{24} \text{ cm}^2/Q$	Total cross section from carbon $\sigma_C \times 10^{24} \text{ cm}^2/Q$	σ_C/σ_D
170	0.82	0.84 ± 0.07	7.2 ± 0.25	8.58 ± 0.66
255	1.27	2.06 ± 0.06	18.9 ± 1.35	9.15 ± 1.41

Elem. Sym.	A	Z
C	12	6
Ref. No.		EH
57 Ca 2		

Method 35 MeV electron synchrotron; neutron yield; radioactivity;
 Cu⁶³(γ ,n) reaction. (Berman and Brown)

Reaction	E or ΔE	E ₀	Γ	$\int \sigma dE$	J π	Notes
C ¹² (γ ,n)	Bremss. 18-30	23	Γ 1/2: 4.2 MeV	$\int_{31}^{31} = 42 \pm 7$ MeV-mb		




Fig. 2.—The measured yield curve and the derived cross section curve for the reaction ¹²C(γ ,n).

TABLE 3
 PARAMETERS OF THE MEASURED (Y, N) REACTIONS

Reaction	Peak Energy (MeV)	Width at Half Maximum (MeV)	Integrated Cross Section to 31 MeV (MeV-mb)
¹² C(γ ,n)	23	4.2	42 \pm 7
¹³ C(γ ,n)	24	3.1	16 \pm 2
¹⁶ O(γ ,n)	—	8.3	290 \pm 50

METHOD


[Page 1 of 2]

REF. NO.


57 Ch 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	SPC	THR - 84	C	64,34	EMU-D	18-54	DST



a



b

FIG. 2 Angular distributions of the photoprotons. a) Be⁹: dots - $E_p > 20$ Mev, $E_{\gamma_{max}} = 68$ Mev; crosses - $E_p > 32$ Mev, $E_{\gamma_{max}} = 84$ Mev; b) C¹²: dots - $E_p > 18$ Mev, $E_{\gamma_{max}} = 64$ Mev; crosses - $E_p > 26$ Mev, $E_{\gamma_{max}} = 84$ Mev.

Elem. Sym.	A	Z
C	12	6
Ref. No.		EGF
57 Ch 1		

Method

[Page 2 of 2]

Reaction	Result	Excitation Energy	Source		Detector		Angle
			Type	Range	Type	Range	




FIG. 3. Energy distribution of photoprotons (yield given in relative units):
 a - Be⁹, E_{γmax} = 68 Mev; b - Be⁹, E_{γmax} = 84 Mev; c - C¹², E_{γmax} = 64 Mev; d - C¹²,
 E_{γmax} = 84 Mev.

Elem. Sym.	A	Z
C	12	6

Method ρ detector; N detector; activation

Ref. No.
 57 Co 1
 EGF

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	J^π	Notes
$C^{12}(\gamma, xn)$	$\sim 19-37$	22.8	3.5	$\int_{18}^{25} = 34 \text{ MeV}\cdot\text{mb}$ $\int_{18}^{38} = 64 \text{ MeV}\cdot\text{mb}$		$E_{th} = 18.72 \text{ MeV}; \sigma_{max} = 10.4 \text{ mb.}$




FIG. 13. The total photon absorption cross section for C^{12} and C^{13} .




FIG. 11. The cross sections for $C^{12}(\gamma, xn)$ and $C^{13}(\gamma, xn)$. The solid curve is the experimental cross section. The dotted curves for the cross sections $C^{12}(\gamma, n)C^{12}$ and $C^{13}(\gamma, pn)B^{11} - C^{13}(\gamma, 2n)C^{11}$ are based on reasonable assumptions about the neutron multiplicity. The curve $C^{13}(\gamma, pn)C^{12} - C^{13}(\gamma, pn)B^{11} - C^{13}(\gamma, 2n)C^{11}$ hypothetical is the assumed cross section for all processes involving neutrons. The assumptions made are discussed in the text.

Elem. Sym.	A	Z
C	12	6

Method	Betatron; photon scattering; absorption; NaI spectrometer	Ref. No.	57 Ha 1	NVB
--------	-----------------------------------------------------------	----------	---------	-----

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C(γ,γ)	Bremss. 19	15	$\Gamma_t = 79 \pm 16$ ev $\Gamma_{gs\gamma} =$ 54.5 ± 9.3 ev	1.90 ± 0.27 MeV-mb		Γ_t - total level width; $\Gamma_{gs\gamma}$ - ground state radiation width. Peak absorption cross section: $\sigma_p = 22.2 \pm 2.2$ b In Figure 2, $t = \left[\frac{\text{Doppler width}}{\text{level width}} \right]^2$




Fig. 4. The peak absorption cross section σ_p as a function of $t = \left[\frac{\text{Doppler width}}{\text{level width}} \right]^2$ is shown from the two curves. The points at $t = 0$ have been read from the curve of Fig. 2. The absorption cross section is a function of t for the excited state.
 $\sigma_p = \left[\frac{2\pi^2}{3} \frac{E_{\gamma}^2}{E_0^2} \right] \Gamma_{gs\gamma} E$
 The experimental value of the internal-conversion cross section σ_{ic} is 1.0 barn. It has been used to obtain the theoretical σ_p for $E_{\gamma} = 15.1$ MeV. The ratio σ_{ic}/σ_p is 0.045 for $E_{\gamma} = 15.1$ MeV.

REF.

D. L. Livesey
 Can. J. of Physics 35, 987 (1957)

ELEM. SIM.

C

12

6

METHOD

REF. NO.

[Page 1 of 2]

57 Li 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	SPC	THR - 35	C	35	EMU-D	5-12	DST

Group I $f(\theta) = A + B \sin \theta$ $A/B = 0.23 \pm 0.1$

II $f(\theta) = A + B(\sin \theta + p \sin \theta \cos \theta)^2$ $A/B = 0.5, p = 0.6$




Fig. 1. Energy distribution of 1940 proton tracks from propane exposed to 35 Mev. bremsstrahlung.

METHOD

REF. NO.

[Page 2 of 2]

57 Li 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




FIG. 2. Angular distributions of protons from propane in two energy groups: I, protons between 4.5 and 6.9 Mev.; II, protons above 6.9 Mev., with a dashed curve showing the function $f(\theta) = A - B(\sin \theta + \rho \sin \theta \cos \theta)^2$.




FIG. 3. Numbers of protons in Group I averaged over supplementary angles and plotted against $\sin^2 \theta$.

METHOD					REF. NO.		ANGLE
					57 Mu 1		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,3A	SPC	15, 18	D	15, 18	EMU-D	0 - 15	DST

(Li,p) γ -rays thick target E_0 500 keV.




Fig. 1. Energy-release histogram for $^{12}\text{C}(\gamma, 3\alpha)$.




Fig. 2. E^* histograms (three values for each star).




Fig. 4. Velocity vectors of $^{12}\text{C}(\gamma_1)\text{Be}^*(\alpha_1)\alpha_2$.




Fig. 3. Angular distributions and correlation in the 2.95 MeV transitions. The curves indicate possible theoretical results for a $J=2$ level, and $E1+E2$ (full lines) of $M1+E2$ (dotted lines) γ -ray interactions.

Elem. Sym.	A	Z
C	12	6

Method $C^{12}(n,2n)$ scintillator detector

Ref. No.	F/H
58 Ba 1	

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n!)$	Bremss. 170 255					<p>Figures 2 and 4: Solid curves are calculated for neutrons from Illinois and Cornell $d(\gamma, p)$ data.</p> <p>In Figure 2, carbon data has been normalized to the deuterium data assuming $\sigma_c = 9\sigma_d$.</p> <p>$E_{th} = 20.6$ MeV, for the neutron detector.</p>




FIG. 2. Dependence of yield of fast photoneutrons emitted from deuterium at 75° in the lab. on the peak bremsstrahlung energy. Solid curve—results from references 4-10 recalculated for neutrons. ●—fast neutron yield from deuterium; ○—fast neutron yield from carbon. The carbon neutron yields are normalized with respect to total cross section to the deuterium data assuming $\sigma_c = 9\sigma_d$.

TABLE I. Yield of >20-Mev photoneutrons from deuterium and carbon.

$E_{\gamma, \max}$ (MeV)	Deuterium yield (below meson threshold) according to data from the literature*	Total yield from deuterium $\sigma_D \times 10^{28}$ cm ² /Q	Total yield from carbon $\sigma_C \times 10^{28}$ cm ² /Q	σ_C/σ_D
170	0.82	0.84 ± 0.07	7.2 ± 0.25	3.58 ± 0.06
255	1.27	2.06 ± 0.06	18.9 ± 1.35	9.18 ± 1.44

* See references 4-10.

- * E. Whalin, Phys. Rev. 95, 1362 (1954).
- * J. C. Keck and R. M. Littauer, Phys. Rev. 93, 827 (1954).
- * L. Allen and A. Hanson, Phys. Rev. 95, 629 (1954).
- * L. Allen, Phys. Rev. 98, 705 (1955).
- * Tollestrup, Keck, and Smythe, Phys. Rev. 96, 850(A) (1954).
- * J. C. Keck and A. V. Tollestrup, Phys. Rev. 101, 360 (1956).
- * Whalin, Schriever, and Hanson, Phys. Rev. 101, 377 (1956).
- * Aleksandrov, Delone, Slovkhotov, Sokol, and Shtarkov, J. Exptl. Theoret. Phys. U.S.S.R. (to be published).




FIG. 4. Angular distributions of >20-Mev neutrons from carbon and deuterium at $E_{\gamma, \max} = 170$ and 255 Mev. ○—neutron yield from carbon at $E_{\gamma, \max} = 255$ Mev⁴ and $E_{\gamma, \max} = 170$ Mev⁵; ●—neutron yield from deuterium for $E_{\gamma, \max} = 255$ Mev. Solid curve—angular distribution of neutrons from deuterium for $E_{\gamma, \max} = 170$ Mev, obtained by recalculating the results of references 4-10 for neutrons. The absolute deuterium neutron yields are normalized with respect to total cross sections to the carbon data.

Ref. W.C. Barber
 Phys. Rev. 111, 1642 (1958)

Elem. Sym.	A	Z
C	12	6
Ref. No.		EH
58 Ba 2		

Method	Ref. No.
Activation	58 Ba 2
	EH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n)$						Ratio of yields for (γ, n) and $(e, e'n)$ reactions is consistent with mixture of intensities of 92% E1 to 8% E2.
$C^{12}(e, e'n)$						
$C^{12}(e, n)$						

METHOD					REF. NO.		EGF
					58 Ba 5		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.XN	RLY	THR - 23	C	18-23	BF ₃ -I		4PI

BREAKS




Fig. 1. — Courbe d'activation du carbone.
 a, voisinage du seuil; b, discontinuités importantes; c, zone de 22,8 MeV.

Discontinuités dans la courbe d'activation par réaction $^{12}\text{C}(\gamma, n)^{11}\text{C}$.

Katz et al. (¹).	Spicer et Penfold (²).	Janjik (³).	Basile et Gusakow.
19,3 ± 0,05 MeV	19,10 ± 0,05 MeV	19,11 ± 0,04 MeV	(19,1) MeV
19,8 ± 0,05 "	19,55 ± 0,05 "	19,60 ± 0,04 "	19,5 ± 0,05 "
20,1 ± 0,05 "			20,15 ± 0,05 "
20,5 ± 0,05 "			20,8 ± 0,05 "
20,7 ± 0,05 "			21,4 ± 0,05 "
21,1 ± 0,05 "			
21,6 ± 0,05 "			
22,4 ± 0,05 "			
22,8 ± 0,05 "			

Method emulsions ; brems. Ref. No. 58 Ch 1 EH

Reaction	E or ΔE	E ₀	Γ	∫σ _d E	Jπ	Notes
C ¹² (γ, p)	Brems. 30, 44					Angular distribution for proton energy groups 57, 7-10, 10-13, 13-16 ≥ 16 MeV.




Fig. 5. Angular distributions of proton groups from photodisintegration of C¹² by γ quanta with E_{γmax} = 44 Mev: ● - E_p = 5-7 Mev, ○ - E_p = 7-10 Mev, ▲ - E_p = 10-13 Mev, △ - E_p = 13-16 Mev, × - E_p ≥ 16 Mev.

METHOD				[Page 1 of 2]		REF. NO. 58 Ma 1		EGF
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
G,3A	ABX	12 - 40	C	150,250	EMU-D		4PI	
G,PA	ABX	25 - 30	C	150,250	EMU-D		4PI	
G,PT	ABX	27 - 70	C	150,250	EMU-D		4PI	

Some data on more complete breakup reaction, i.e. $^1\text{H}+^3\text{H}+2^4\text{He}$, etc.




FIG. 1. Dependence of the cross section for reaction (I) on γ -ray energy. The solid line gives our data, the dashed one the data of reference 3.

$\sigma(G,3A)$





FIG. 3. Dependence of the cross section for reaction (III) on γ -ray energy.

TABLE I. Integral cross sections for reactions (III) and (V) in mbn-Mev

E_γ , Mev	25-30	30-35	35-40	40-45
(III)	3.85 ± 0.20	1.78 ± 0.17	0.82 ± 0.13	0.38 ± 0.11
(V)	3.11 ± 0.21	1.64 ± 0.18	0.84 ± 0.14	0.16 ± 0.08



Method					Ref. No.		EGF	
					58 Ma 1			
Reaction	Result	Excitation Energy	Source		Detector		Angle	
			Type	Range	Type	Range		




FIG. 10. Angular distribution of protons in reaction (III):
 a - $E_\gamma < 50$ Mev, a' - $E_\gamma > 50$ Mev.

Reaction VI $^{12}\text{C} + \gamma \rightarrow \text{p} + \text{t} + 2\alpha - 27$ Mev.




FIG. 12. Angular distribution of protons in reaction (VI) for
 $E_\gamma < 70$ Mev.




FIG. 13. Dependence of the cross sections for several reactions on the energy of the γ -ray: 1-(I), 2-(III), 3-(VI), 4 - cross section for photoproduction of stars with two or more prongs in the interval $E_\gamma = 30$ to 80 Mev from reference 1, 5 - $\text{C}^{12}(\gamma, \text{n})\text{C}^{11}$ from reference 8.

Ref. V.K. Rasmussen, F.R. Metzger, C.P. Swann
 Phys. Rev. 110, 154 (1958)

Elem. Sym.	A	Z
C	12	6

Method Van de Graaf; photon scattering, absorption; NaI spectrometer

Ref. No.	NVB
58 Ra 1	

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C ¹² (γ,γ)	4.43	4.43				Mean life: $\tau = (6.5 \pm 1.2) 10^{-14}$ sec. Detector at 90°, 126°.

Ref. G. W. Tautfest
 Phys. Rev. 110, 708 (1958)

Elem. Sym.	A	Z
C	12	6

Method Li^9 delayed neutrons; BF_3 counters; Purdue University synchrotron;
 Ion chamber

Ref. No.	EH
58 Ta 1	

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$(\gamma, 3p)$	Bremss. 100-320			$\int_{100}^{320} = 0.31 \pm 0.05$ MeV-mb		




FIG. 9. Photon cross section for the reaction $\text{C}^{12}(\gamma, 3p)\text{Li}^9$ obtained by the photon difference method from the yield point of Fig. 7. The photon cross section below 100 Mev was chosen arbitrarily.

Elem. Sym.	A	Z
C	12	6
Ref. No.		EH
58 Wh 2		

Method Counter telescope; Bremss.; coinc. with n-counter.

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C(\gamma, p)$	Bremss. 40-110	37-78				Curves and tables giving energy and angular distribution of photoprotons having energies between 37-65 MeV produced by photons of 45-95 MeV; compare favorably with pseudo-deutror calculations of Dedrick, e.g.




Fig. 4. Photoproton energy spectra from carbon with 90-MeV bremsstrahlung.




Fig. 5. Angular distributions of photoprotons from carbon with 90-MeV bremsstrahlung.




Fig. 6. Angular distributions of photoprotons from carbon with 110-MeV bremsstrahlung.

Method

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
----------	-----------------	-------	----------	------------------	--------	-------




Fig. 6. Yield curves for 37-Mev photoneutrons from carbon.





Fig. 7. Cross sections for photoneutron production by 96-Mev photons on carbon. The curves shown are the quasi-neutron calculations by Dedrick.²





Fig. 9. Cross sections for the production of 37-Mev photoneutrons from carbon. The curves shown are the quasi-neutron calculations by Dedrick.²



Method NaI detectors; $T^3(p,\gamma)$ source.

Ref. No.
 58 Wo 1
 EH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	J^π	Notes
$C(\mu_e)$	20.3 - 20.8					σ_e varied 315 ± 5 to 324 ± 6 mb smoothly With resolution better than 70 keV, it should have been possible to detect resonances with integrated cross sections greater than 0.8 MeV-mb.

TABLE II. Comparison of expected total cross section with this experiment.

Component	Cross section (millibarns)*	
	20.3 Mev	20.8 Mev
Compton	179.6 ± 1	176.5 ± 1
Pair production	116.6 ± 0.5	117.6 ± 0.5
Triplet production	14.6 ± 1.5	14.8 ± 1.5
Nuclear (γ,n) and (γ,p)	9 ± 4	14 ± 4
Total:	320 ± 5	323 ± 5
This experiment	315 ± 5	324 ± 6

* Uncertainties are estimated.




FIG. 3. (A) Measured total cross section for carbon as a function of photon energy. (B) Various expectations of structure based on betatron (γ,n) yield curve breaks.

Ref. B. Ziegler
Z. Physik 152, 566 (1958)

Elem. Sym.	A	Z
C	12	6

Method 32 MeV betatron; pair spectrometer; absorption measurement;
ionization chamber

Ref. No.
58 Zi 1
EH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
C (μ_t)	Bremss. 30			30 ± 10 MeV-mb		




Fig. 6. a) σ_{π^+} for C; b) σ_{π^+} for $\pi^+ p$; c) σ_{π^+} for $\pi^+ n$.

- ¹ MONTALBETTI, R., L. KATZ u. J. GOLDBERG: Phys. Rev. 91, 659 (1953).
- ¹² HALPERN, J., u. A.K. MANN: Phys. Rev. 83, 370 (1951).
- ¹³ SPICER, B.M.: Phys. Rev. 99, 33 (1955).
- ¹⁴ BARBER, W.C., W. DE GEORGE u. D.D. REAGAN: Phys. Rev. 98, 73 (1955).

Elem. Sym.	A	Z
C	12	6
Ret. No.		EH
59 Ba 1		

Method Mark II linear acclerator; 18" spectrometer

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
C(e, e')	42.6	15.1 19 20		1.47 MeV-mb $\pm 20\%$	1 ⁺	






FIG. 1. Energy distribution of electrons scattered at 160° from carbon. The elastic peak is shown at the right with the ordinate scale indicated on the right margin. The dashed curve is the calculated tail of the elastic peak due to bremsstrahlung. The solid curves were drawn arbitrarily. The points without statistical errors indicated have errors comparable with those of neighboring points.

Ref. E.B. Bazhanov
 Zhur. Eksp. i Teoret. Fiz. 37, 374 (1959);
 Soviet Phys. JETP 10, 267 (1960)

Elem. Sym.	A	Z
C	12	6

Method
 Brems.; NaI-CsI telescope

Ref. No.	JH
59 Ba 2	

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes																																																							
(γ,p)	protons 17.8-52.2MEV					<p>Took difference of yield measured with 2 Brems. spectra 82-89 MeV.</p> <div style="display: flex; justify-content: space-around;"> <div style="width: 45%;">  <p>FIG. 2. Energy spectrum of protons ejected from C¹² when this nucleus is irradiated by the full bremsstrahlung spectrum having E_{γmax} = 89 Mev. Protons observed at 90°. ● - our data, ○ - data of Whitehead et al., x - data of Chuvilo and Shevchenko. Errors shown are statistical.</p> </div> <div style="width: 45%;">  <p>FIG. 3. Energy spectrum for protons from C¹² for rays in a narrow energy range; θ = 90°. Curves calculated from data of Dedrick; 1 - without taking into account scattering of protons in the nucleus, 2 - taking scattering into account. The errors shown are statistical.</p> </div> </div> <table border="1" style="width: 100%; margin-top: 10px;"> <thead> <tr> <th rowspan="2">Target</th> <th rowspan="2">Mean proton energy E_p, Mev</th> <th colspan="2">Proton yield, cm²10⁻³⁰ sterad·Mev·Q</th> <th rowspan="2">Target</th> <th rowspan="2">Mean proton energy E_p, Mev</th> <th colspan="2">Proton yield, cm²10⁻³⁰ sterad·Mev·Q</th> </tr> <tr> <th>E_{γmax} = 82 Mev</th> <th>E_{γmax} = 39 Mev</th> <th>E_{γmax} = 82 Mev</th> <th>E_{γmax} = 39 Mev</th> </tr> </thead> <tbody> <tr> <td rowspan="6">C¹²</td> <td>17.5</td> <td>4.03±0.040</td> <td>1.120±0.042</td> <td>10.1</td> <td>1.31±0.022</td> <td>0.360±0.010</td> <td>0.100±0.005</td> </tr> <tr> <td>19.4</td> <td>2.94±0.052</td> <td>0.834±0.035</td> <td>14.7</td> <td>0.71±0.013</td> <td>0.200±0.008</td> <td>0.050±0.003</td> </tr> <tr> <td>28.0</td> <td>1.94±0.034</td> <td>0.534±0.018</td> <td>20.7</td> <td>0.47±0.009</td> <td>0.130±0.006</td> <td>0.030±0.002</td> </tr> <tr> <td>31.4</td> <td>0.91±0.023</td> <td>0.252±0.011</td> <td>27.8</td> <td>0.27±0.006</td> <td>0.070±0.004</td> <td>0.020±0.001</td> </tr> <tr> <td>36.8</td> <td>0.45±0.010</td> <td>0.123±0.005</td> <td>38.7</td> <td>0.14±0.003</td> <td>0.030±0.002</td> <td>0.010±0.001</td> </tr> <tr> <td>44.0</td> <td>0.15±0.004</td> <td>0.040±0.002</td> <td>44.3</td> <td>0.05±0.001</td> <td>0.010±0.001</td> <td>0.003±0.001</td> </tr> </tbody> </table>	Target	Mean proton energy E _p , Mev	Proton yield, cm ² 10 ⁻³⁰ sterad·Mev·Q		Target	Mean proton energy E _p , Mev	Proton yield, cm ² 10 ⁻³⁰ sterad·Mev·Q		E _{γmax} = 82 Mev	E _{γmax} = 39 Mev	E _{γmax} = 82 Mev	E _{γmax} = 39 Mev	C ¹²	17.5	4.03±0.040	1.120±0.042	10.1	1.31±0.022	0.360±0.010	0.100±0.005	19.4	2.94±0.052	0.834±0.035	14.7	0.71±0.013	0.200±0.008	0.050±0.003	28.0	1.94±0.034	0.534±0.018	20.7	0.47±0.009	0.130±0.006	0.030±0.002	31.4	0.91±0.023	0.252±0.011	27.8	0.27±0.006	0.070±0.004	0.020±0.001	36.8	0.45±0.010	0.123±0.005	38.7	0.14±0.003	0.030±0.002	0.010±0.001	44.0	0.15±0.004	0.040±0.002	44.3	0.05±0.001	0.010±0.001	0.003±0.001
Target	Mean proton energy E _p , Mev	Proton yield, cm ² 10 ⁻³⁰ sterad·Mev·Q		Target	Mean proton energy E _p , Mev	Proton yield, cm ² 10 ⁻³⁰ sterad·Mev·Q																																																							
		E _{γmax} = 82 Mev	E _{γmax} = 39 Mev			E _{γmax} = 82 Mev	E _{γmax} = 39 Mev																																																						
C ¹²	17.5	4.03±0.040	1.120±0.042	10.1	1.31±0.022	0.360±0.010	0.100±0.005																																																						
	19.4	2.94±0.052	0.834±0.035	14.7	0.71±0.013	0.200±0.008	0.050±0.003																																																						
	28.0	1.94±0.034	0.534±0.018	20.7	0.47±0.009	0.130±0.006	0.030±0.002																																																						
	31.4	0.91±0.023	0.252±0.011	27.8	0.27±0.006	0.070±0.004	0.020±0.001																																																						
	36.8	0.45±0.010	0.123±0.005	38.7	0.14±0.003	0.030±0.002	0.010±0.001																																																						
	44.0	0.15±0.004	0.040±0.002	44.3	0.05±0.001	0.010±0.001	0.003±0.001																																																						

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,N	ABY	THR - 36	D	10 - 36	BF3-I		4PI




FIG. 5. Yield of neutrons per incident electron as a function of initial electron energy for the low-Z elements. The concrete target is a simple 3:1 sand:cement mixture. The numbers at the top right refer to the Cu-I curve at the indicated energies.

TABLE I. Thicknesses of the targets used in the experiment, with the exception of heavy water, all targets contained isotopes in their naturally-occurring proportions.

Target	Thickness (g/cm ²)	Thickness (radiation lengths)
Heavy water	0.698	"thin"
Be	0.559	0.00867
C-I	38.91	0.88
Al-I	24.10	1.00
Cu-A	1.372	0.108
Cu-I	13.26	1.04
Cu-II	26.56	2.08
Cu-III	39.86	3.13
Cu-IV	53.13	4.17
Ta-I	6.21	0.93
Pb-I	5.88	1.01
Pb-II	11.42	1.97
Pb-III	17.30	2.98
Pb-IV	22.89	3.94
Pb-VI	34.42	5.93
U-I	6.17	1.14
U-II	12.42	2.30
U-III	18.61	3.46
Concrete	23.5	1.19

¹³ A. I. Berman and K. L. Brown, Phys. Rev. 96, 83 (1954).




FIG. 14. Experimental and expected yields of neutrons per incident electron for 1-radiation-length targets at 34.3 Mev, as a function of atomic number Z. The experimental yields were obtained by dividing the measured yields from the targets labeled I by the actual target thicknesses listed in Table I. The expected yields were calculated from expression (8).

Ref. N.A. Burgov, G.V. Danilyan, B.S. Dolbilkin, L.E. Lazareva, F.A. Nilolaev
 Zhur. Eksp. i Teoret. Fiz. 37, 1811 (1959);
 Soviet Phys. JETP 10, 1278 (1960)

Elem. Sym.	A	Z
C	12	6
Ref. No.	59 Bu 1	JH

Method
 39 MeV Synchr.; magnetic pair spectrometer

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
total absorption	$\Delta E_\gamma = 22.2$ -23.5 Mev	~ 22.35 (from Fig.2)	$\lesssim 100$ keV	9 Mev-mb $\int_{22.2}^{22.5}$		Experimental peak width ~ 150 keV, but this includes smearing by the spectrometer resolution of ~ 100 keV.




FIG. 2

In Fig. 2 are shown the x-ray spectrum measured without absorber, $N_0(h\nu)$, and with absorber, $N(h\nu)$. All experimental points are referred to an equal γ dose, which was determined by means of a thin-walled ionization chamber and an integrator

ELEM. SYM.	A	Z
C	12	6
REF. NO.		EGF
59 Ch 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	RLY	THR - 30	C	30	TEL-D	15 - 45	90
G,XD	RLY	THR - 30	C	30	TEL-D	15 - 45	90

REL D TO P YLD




FIG. 3. Energy distributions: O - photoprotons and □ - photodeuterons from C¹³ with E_{γmax} = 30 Mev. The scale of the ordinate axis for deuterons is enlarged by a factor of 10.




FIG. 4. Ratio of the number of photodeuterons to the number of photoprotons with the same energy, as a function of energy: O - for C¹³ and ● - for Be⁹.




FIG. 7. Experimental and calculated ratios of the number of photodeuterons to the number of photoprotons from C¹³ as functions of particle energy: O - with E_{γmax} = 80 Mev and ● - with E_{γmax} = 300 Mev.

Ref. L.D. Cohen, W.E. Stephens
 Phys. Rev. Letters 2, 263 (1959)

Elem. Sym.	A	Z
C	12	6

Method Monochromatic capture γ 's; activation; NaI detector

Ref. No.	EH
59 Co 1	

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n)$	20.2-21					




FIG. 1. $C^{12}(\gamma, n)$ activation curve. Cross section in millibarns for the photoproduction of C^{11} radioactivity as a function of photon energy in Mev. The dashed figures indicate energy resolution vs energy deduced from tritium-zirconium target thickness measurements and Doppler angle calculations.

METHOD			REF. NO.				
			59 Ga 1				
			egf				
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	LFT	15	C	19-42	NAI-D		DST

Direct measurement of $\int \sigma_s dE = 2.11 \pm 0.31$.

15.1 MEV LEVEL

TABLE I. Properties of the 15.1-Mev, $T=1, J=1+$ level in C^{12} .

	This experiment	National Bureau of Standards experiment ^a
1. Peak absorption cross section	29.7 ± 1.1	22.2 ± 2.2 barns
2. Ratio of Doppler broadening to total level width	0.62 ± 0.10	...
3. Total level width	64.5 ± 10.4	79 ± 16 ev
4. Radiation width to the ground state of C^{12}	59.2 ± 9.7	54.5 ± 9.3 ev
5. Radiation width to the 4.43-Mev state of C^{12}	3.2 ± 2.5	≤ 5.5 ev
6. Alpha-particle width to the 2.9-Mev state of Be^8	2.1 ± 3.2	$(18.6 \leq \Gamma_\alpha \leq 24.5) \pm 8.2$ ev
7. Integrated elastic scattering cross section	2.33 ± 0.19	1.90 ± 0.27 mb-Mev

^a See reference 5.

⁵ E. Hayward & E. G. Fuller, Phys. Rev. 106, 991 (1957).




FIG. 4. The angular distribution of photons scattered from the 15.1-Mev level in C^{12} .

Elem. Sym.	A	Z
C	12	6

Method Cyclotron; gamma-ray yield; NaI(Tl) detector

Ref. No.
 59 Ge 1
 EH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$B^{11}(p,\gamma)$	$\sim 19.6-23$	22.5 ± 0.1				<p>$\sigma(\gamma,p) 90^\circ = 29 \pm 5$ mb.</p> <p>Two compound states at 21.4 and 22.1 MeV figured in absorption; data agrees with Barber's $C^{12}(e,e')$ data. Katz breaks not seen.</p> <p>Proton energy range 4 - 7.7 MeV.</p> <p>Yield at 90°</p>




Fig. 5 (a) The $B^{11}(p,\gamma)C^{12}$ yield curve with the data of Barber et al.¹⁰, Blaser et al.¹³ and Katz¹⁵ shown for comparison.

- Ref. 10: Blair, Kingston and Willard, Phys. Rev. 100, 21 (1955).
- Ref. 13: Blaser, Boehm, Marmier and Scherrer, Helv. Phys. Acta. 24, 465 (1951).
- Ref. 15: Katz, Washington Photoneuclear Conference (5/58).

Elem. Sym.	A	Z
C	12	6
Ref. No. 59 Go 1		EH

Method Counting γ 's.

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$B^{11}(p,\gamma)$	3-11					Measure ground state photon.




FIG. 1. The 90° yield of gamma rays leading to the ground state of C^{12} from the $B^{11}(p,\gamma)C^{12}$ reaction.

Elem. Sym.	A	Z
C	12	6

Method 30 MeV synchrotron; NaI(Tl) scintillator spectrometer; photon spectrum	Ref. No. 59 Ko 1	EH
----------------------------------------------------------------------------------	---------------------	----

Reaction	E or ΔE	E ₀	Γ	∫ σ _d E	Jπ	Notes
C(σ _t)	Bremss. 30.5	23	4.6	$\int_{10}^{27} = 0.12 \text{ MeV-mb}$		

TABLE 2

Experimentally determined giant resonance data; peak energy, E_{max}, width at half height, ΔE, and integrated cross section, $\int_{10 \text{ MeV}}^{27 \text{ MeV}} \sigma_d E$, and values of $\int_0^{\infty} \sigma_d E$ predicted by the dipole sum rule.

Absorber	E _{max} (MeV)	ΔE (MeV)	$\int_{10 \text{ MeV}}^{27 \text{ MeV}} \sigma_d E$	$\int_0^{\infty} \sigma_d E$
Carbon	23	4.6	0.12	0.18
Aluminium	21.5	7.8	0.24	0.40
Sulphur	21	6.6	0.30	0.48
Iron	19.5	9.4	0.76	0.84




Fig. 1. Experimentally determined photo-nuclear absorption cross sections (crosses). The open circles are the total absorption cross sections and the filled squares the dipole sum rule cross section where available. The solid circles give the total photo-nuclear absorption cross section as measured by Ziegler.

⁹⁾ B. Ziegler, Z. Physik 152 (1958) 566.

¹⁷⁾ J. Halpern and A. K. Mann, Phys. Rev. 83 (1951) 37.

¹⁸⁾ R. Montalbetti, L. Katz and J. Goldemberg, Phys. Rev. 91 (1953) 639.

¹⁹⁾ L. W. Jones and K. M. Terwilliger, Phys. Rev. 9 (1953) 699.

²⁰⁾ W. C. Barber, W. D. George and D. D. Reagar, Phys. Rev. 98 (1955) 73.

²¹⁾ F. Ferrero, R. Malvano, S. Menardi and O. Terracini, Nuclear Physics 9 (1958) 32.

²²⁾ L. Katz and A. S. Penfold, Phys. Rev. 81 (1951) 81.

²³⁾ S. A. E. Johansson, Phys. Rev. 97 (1955) 1136.

Ref. L.A. Kul'chitskii, V. Presperin
 Zhur. Eksp. i Teoret Fiz. 37, 1524 (1959);
 Soviet Phys, JETP 10, 1082 (1960)

Elem. Sym.	A	Z
C	12	6

Method
 88 MeV Synchr.; proton recoil telescopes

Ref. No.
 59 Ku 1 JH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n)$	Bremss: $E_{\gamma \max} = 88 \text{ Mev};$ $E_{\pi} > 10 \text{ Mev}$					Angular distribution; results consistent with quasi-deuteron model, but some direct resonance abs. may be present.




FIG. 3. Energy distribution of photoneutrons from C^{12} at laboratory angle 75° .




FIG. 4. Angular distribution of photoneutrons from C^{12} . Solid curve - converted data from reference 14; dashed curve - data calculated in reference 13.

METHOD

Betatron

REF. NO.

59 Oc 1

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, 2N	RLI	THR-100	C	THR-100	ACT-I		4PI

REL TO G, N

TABLE II. Relative integrated cross sections.

Element	(γ, n)	Position of the peak for (γ, n)	($\gamma, 2n$)	Position of the peak for ($\gamma, 2n$)
C ¹²	1	23 Mev	0.003	42 Mev
N ¹⁴	1	24 Mev	0.007*	
O ¹⁶	1	22 Mev	0.002	40 Mev
F ¹⁹	1	20 Mev	0.14	32 Mev
Na ²³	1	20 Mev	0.05	32 Mev
P ³¹	1	20 Mev	0.06 ($\gamma, 2p$)	45 Mev ($\gamma, 2p$)
			0.08 ($\gamma, 2pn$)	50 Mev ($\gamma, 2pn$)

* The (γ, n) integrated cross section was taken from reference 4.




FIG. 1. The energy dependence of several photoneuclear reactions. The relative yield scales of different graphs are independent.

METHOD

REF. NO.

59 Pa 3

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, G	ABX	17	D	15,18	NAI-D	17	90

Source Li(p,γ) with $E_p = 500$ keV.

TABELLE I

Die gemessenen totalen Streuquerschnitte in cm^2 unter Annahme von E1- und E2-Streuung

Element	eigene Werte	Fuller und Hayward ¹²⁾	Stearns ⁹⁾
Pb	$(5.8 \pm 1) \times 10^{-17}$	$(4-8) \times 10^{-17}$	$(5-9) \times 10^{-17}$
Al	$(2 \pm 1) \times 10^{-18}$	$(2-8) \times 10^{-18}$	
O	$(1-5) \times 10^{-18}$		
C	$(5.8 \pm 2) \times 10^{-20}$		

Durch die Wahl der Meßgeometrie ergibt sich für eine E1- wie eine E2-Winkelverteilung innerhalb der Fehlergrenzen der numerischen Rechnung der gleiche Wert für den totalen Streuquerschnitt. Die angegebenen Fehler enthalten nur den Fehler in der Bestimmung des primären γ -Flusses und die statistischen Fehler der Streuraten. Zum Vergleich sind die entsprechenden Ergebnisse von Fuller und Hayward ¹²⁾ und Stearns ⁹⁾ gegenübergestellt. Die angegebenen Werte gehören jeweils zu den Fehlergrenzen.

Elem. Sym.	A	Z
C	12	6

Method	β counting; self absorption measurement; bremsstrahlung	Ref. No.	59 Pe 2	EH
--------	---------------------------------------------------------------	----------	---------	----

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C(\mu_e)$	Bremss. 30.7					Carbon and oxygen detectors were used. The average value for the C absorption cross section is just that to be expected if σ_e is smooth.




FIG. 3. The experimental results (solid dots) compared to predicted results (crosses). The predicted results were obtained assuming completely "smooth" cross sections. $\langle \sigma \rangle_0$ is the total photon absorption cross section for oxygen averaged over the $(0^{+}, \pi^{+})$ cross section. $\langle \sigma \rangle_1$ is the total photon absorption cross section for oxygen averaged over the $(0^{+}, \pi^{-})$ cross section, and so on.

Method
 Proton counting; Bremsstrahlung, CsI spectrometer

Ref. No.
 59 Pe 4
 EH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
----------	---------	----------------	---	------	----	-------

C¹²(γ,p)

30-70

Ground state and first excited state of B¹¹ cross section measured. Angular distribution fit with:

$$\sum A_i \cos^i \theta \text{ and } \sum c_i P_i(\cos \theta)$$




Figure 8: Ground-state angular distributions at 5 photon energies. The solid curves are least-squares fits (see text). The dashed curve with the 22.1 MeV data is proportional to $2+3 \sin^2 \theta$, normalized to our data at 90° . The curves are labeled with the appropriate photon energy. Note that the cross section scales are not all equal. Uncertainties shown are statistical only.

TABLE III Angular distribution coefficients

Photon Energy, Mev	22.1	25.3	30.7	38.2	48.2
Ground state cross section	0.432 ± 0.021	0.589 ± 0.012	1.148 ± 0.007	1.090 ± 0.023	0.920 ± 0.016
Excited state cross section	0.184 ± 0.056	0.145 ± 0.041	0.112 ± 0.022	0.070 ± 0.027	0.054 ± 0.046
Excited state cross section at 90°	0.191 ± 0.142	0.130 ± 0.095	0.101 ± 0.051	0.013 ± 0.0161	0.042 ± 0.024
Excited state cross section at 180°	0.130 ± 0.087	0.101 ± 0.066	0.113 ± 0.040	0.015 ± 0.0120	0.015 ± 0.0179
Excited state cross section at 135°	0.149 ± 0.179	0.139 ± 0.125	0.106 ± 0.073	0.0310 ± 0.0211	0.057 ± 0.027
Excited state cross section at 45°	0.047 ± 0.010	0.307 ± 0.027	0.110 ± 0.004	0.045 ± 0.011	0.018 ± 0.016
Excited state cross section at 135°	0.061 ± 0.015	0.064 ± 0.011	0.068 ± 0.017	0.016 ± 0.010	0.025 ± 0.014
Excited state cross section at 135°	0.040 ± 0.025	0.171 ± 0.018	0.066 ± 0.013	0.020 ± 0.013	0.043 ± 0.027
Excited state cross section at 135°	0.116 ± 0.043	0.045 ± 0.016	0.045 ± 0.016	0.012 ± 0.008	0.027 ± 0.012
Excited state cross section at 135°	0.013 ± 0.041	0.066 ± 0.029	0.013 ± 0.017	0.011 ± 0.008	0.012 ± 0.005




Fig. 9. Cross section at 90° as a function of photon energy. Circles are for the ground state and squares are for the excited state. The solid curve is the least-squares fit to the data at 90° . The solid curve is the fit to the data at 90° and the dashed curve is the fit to the data at 135° .




Fig. 9. Coefficients c_i of Legendre polynomials $P_i(\cos \theta)$ in the least-squares fit shown in Fig. 8 as a function of energy. Uncertainties shown are statistical only. Where no uncertainty is shown it is smaller than the size of the symbol. The smooth curves are for illustrative purposes only.

METHOD					REF. NO.		JOC	
					59 Pe 5			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
G,G	ABX	19 - 61	C	19 - 61	NAI-D		135	




FIG. 5. The scattering cross section for C at 135°. The crosses designate points which can only be due to elastic scattering, while the dots designate points for mixed elastic and inelastic scattering. The mixing is defined by Eq. (15) in the text. The data were obtained from irradiations with bremsstrahlung energies of 27, 28, and 61 Mev. The solid curve is a predicted cross section (see text for details).

METHOD				REF. NO.		EGF	
				59 Sa 1		EGF	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	NOX	THR - 19	C	18-19	ACT-I		4PI
G,N	NOX	19 - 21	C	18-21	MOD-I		4PI


Résultats avec l'anthracène (MeV).

BREAKS

Katz (*).....	18,90	18,96	19,08
Présent travail.....	18,86	18,96	19,10
	$\pm 0,02$	$\pm 0,02$	$\pm 0,02$

Résultats obtenus avec la paraffine (MeV).

Spicer et Penfold.....	-	19,55	19,95	20,35 (*)	20,75 (*)
		$\pm 0,05$	$\pm 0,05$	$\pm 0,05$	$\pm 0,05$
Présent travail.....	19,31	19,56	19,95	20,37	20,75
	$\pm 0,02$	$\pm 0,02$	$\pm 0,02$	$\pm 0,02$	$\pm 0,02$
Katz (*).....	19,30	19,57	19,92	20,29	20,62



Elem. Sym.	A	Z
C	12	6
Ref. No.		JH
60 Ba 1		

Method: Brems.; from 100 MeV Syn.; NaI (Tl) telescopes

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C ¹² (γ,p)	protons					Units on ∫σdE 10 ⁻³⁰ in MeV/sr. Values for ∫σdE calculated according to Shklyarvskii, using harmonic oscillator potential, are 10.5 and 8.8, X 10 ⁻³⁰ cm ² MeV/sr for proton energy interval 18.6-24.2 and 24.2-29.9 respectively. θ = 57.5°
	18.6-24.2 MeV			135.2 ± 20.6	63 MeV 34 MeV	
	24.2-29.9 MeV			120.6 ± 18.9	70 MeV 42 MeV	
	29.9-38.7 MeV					




FIG. 1. Photoproton yield and cross section curves for C¹² with $\bar{E}_p = 21.4$ Mev. The right ordinate scale refers to curve a and the left to curves b and c and the shaded areas. The errors are statistical.




FIG. 2. The same as in Fig. 1, but for photoprotons with $\bar{E}_p = 27.0$ Mev.




FIG. 3. The same as in Fig. 1, but for photoprotons with $\bar{E}_p = 34.3$ Mev.

Elem. Sym.	A	Z
C	12	6

Method Stanford Mark II Linac; magnetic spectrometer; plastic scintillator counter telescope

Ref. No. 60 Ba 4
 JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(e^-, e^-)$	42.5	15.11	40^{+8}_{-6} eV	$2.0^{+0.4}_{-0.3}$ MeV-mb	$1+$	<p>$\int \sigma dE$ is average of 1.88 and 2.44 MeV-mb measured $\theta = 160^\circ$ and 132°, respectively.</p> <p>M1 transition.</p> <p>$\int \sigma dE$ average of 76 and 72 MeV-mb are 160° and 132°, respectively.</p> <p>Peaks at 17.3 ± 0.2 and 19.5 ± 0.2 MeV excitation energy superimposed on giant resonance.</p> <p>E1 transition.</p>




FIG. 2. Energy distribution of electrons, which were initially 42.5 Mev, after 160° scattering from a C target.




FIG. 4. Energy distribution of electrons, which were initially 42.5 Mev, after 132° scattering from a C target.

REF.

ELEM. SYM. A Z

A. Bussiere de Nercy and M. Langevin
J. Phys. Radium 21, 293 (1960)

(See 61 Bu 4)

C

12

6

METHOD

REF. NO.

60 Bu 3

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	LFT	15	C	23	NAI-D		DST

$$\Gamma_{\gamma} = 56 \pm 11 \text{ eV.}$$

$$\Gamma = 72 \pm 14 \text{ eV.}$$

Method Monochr. γ 's from $H^3(p,\gamma)He^4$ reaction, total abs.; NaI Ref. No. 60 Ca 1 JH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
H^3_t	20.0-21.2	20.15	165 ± 30 keV	1.1 ± 6 MeV-mb		σ_{max} (nuclear) = 6.5 ± 2 mb. σ_{max} (nuclear) = 7 ± 3 mb. σ_{max} (nuclear) = 22mb. Subtraction of calculated atomic cross-sections, using Borsellino for triplet.
		20.46	145 ± 30 keV	$1.0 \pm .75$ MeV-mb		
		20.92	225-350 keV	6.6 ± 1.2 MeV-mb		
				total: 21.20 8.7 ± 2.6 20.0		




FIG. 2. Total absorption in C^{12} . The points indicate the experimental results. The lines show the total atomic cross sections calculated as described in the text using the indicated triplet calculations. The curve marked Koch and Wyckoff includes a 25% addition to the pair production and the Borsellino triplet cross section.




FIG. 3. Total nuclear absorption in carbon-12. The cross section in millibarns is plotted as a function of photon energy.

Ref. V.P. Chizhov
 Zhur. Eksp. i Teoret. Fiz. 38, 809 (1960)
 Soviet Phys. JETP 11, 587 (1960)

Elem. Sym.	A	Z
C	12	6

Method 90 MeV brems.; scintillator counter telescope

Ref. No. 60 Ch 1
 JHH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C(γ,p) C(γ,d)	Bremss. 90					Ratio: σ(γ,d)/σ(γ,p), θ = 90° Energy range of E _p and E _d : 15.5 to 30 MeV

YIELD DATA TABLE:

It should be noted that the yield of photoprotons of the energy considered rises smoothly with Z for the elements plotted in Fig. 3, and that starting already with Al. no direct proportionality to Z is observed on account of the effect of the Coulomb barrier. For illustration, we give the yields of photoprotons Y(γ,p) per proton in the nucleus for several elements in relative units (the error in these measurements was estimated to be ≈ 10%):

	Li ⁶	Li ⁷	Be	C	Al	Cu
Y(γ,p) =	1.00	1.07	1.1	1.31	1.90	0.56




FIG. 6. Angular distributions of photoprotons of energies 15.5-30 Mev for Li⁶, Li⁷, Be, and C. The errors are statistical.

Table of σ(γ,d) to σ(γ,p) cross sections for photodeuteron energies 15.5-30 Mev as function of angle θ. The solid curve shows the results given by Eq. (2), arbitrarily normalized.




FIG. 7. Angular distributions of photodeuterons of energies 15.5-30 Mev for Li⁶, Li⁷, Be, and C. The errors are statistical. The solid curve gives the calculated results for photodeuterons from Be.

Method
 30 - MeV betatron; emulsions; proton recoil tracks

Ref. No.
 60 Em 2 JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
(γ, n)	Bremss.; $E_{\gamma \max} = 30$	(See last column Table III)				Neutron spectra at $30^\circ, 60^\circ, 120^\circ$ and 150° (Figure 1). These spectra, integrated over angle are compared with (γ, p) data of Cohen, et al, Phys. Rev. <u>104</u> , 108 (1956) in Figure 2. (Refer: 56 Co 1) For $E_n > 3 \text{ MeV}$, angular data fits $1 + 1.5 \sin^2 \theta$ (see Figure 3), expected from Wilkinson's indep-part. model for ejection from $l=1$ orbit.






Fig. 2. Energy spectra of C¹² at various angles and integrated from 0 to 30 MeV.






Fig. 3. Energy spectra of C¹² at various angles and integrated from 0 to 30 MeV.




Fig. 1. Energy spectra of C¹² at various angles and integrated from 0 to 30 MeV.

TABLE III. Levels in C¹² above 21 Mev.*

Peaks	Cohen et al. ^b	Ajzenberg and Lauritsen ^c	Jones ^d	Present work
b, b'	21.3 ± 0.2	21.34 21.80		21.4 ± 0.1
c, c'	22.2 ± 0.2	(22.1) 22.55 ± 0.1		22.2 ± 0.2
d, d'	23.0 ± 0.1	(22.8)	22.7 ± 0.2	22.9 ± 0.2
e		(24.3)	24.3 ± 0.2	25.6 ± 0.2
f			25.0 ± 0.2	24.3 ± 0.1
g				24.3 ± 0.2
h		(25.4)		25.6 ± 0.2

* Levels in parenthesis are given with uncertainty.
^b See reference 26.
^c See reference 32.
^d See reference 33.

M. Garnier, H. Gauvin, W. Sebaoun
 J. Phys. Radium 21, 893 (1960)

ELEM. SYM. A Z
 C 12 6

METHOD Van de Graaf; alpha cross section; nuclear emulsion;
 ion chamber. REF. NO. 60 Ga 2 NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, 5A	ABX	15, 18	D	15, 18	EMU-I		
		(14.8, 17.6)		(14.8, 17.6)			

$$\sigma(14.8) = 0.44 \pm 0.11 \times 10^{-28} \text{ cm}^2$$

$$\sigma(17.6) = 1.90 \pm 0.21 \times 10^{-28} \text{ cm}^2$$

Ref. K.N. Geller, J. Halpern, E.G. Muirhead
 Phys. Rev. 119, 716 (1960)

Elem. Sym.	A	Z
C	12	6

Method 25 MeV Betatron; activation of carbon in plastic scintillator

Ref. No. 60 Ge 2
 JH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(γ, n)	18.6-20.0					<p>E_γ (threshold) = 18.77 ± 0.03 MeV to be compared with Everling's neutron separation energy 18.721 ± 0.006 MeV value from mass data.</p> <p>Observed break(s) in yield curve (Figure 1) at:</p> <p>18.79 MeV 18.86 " 19.00 "</p>




FIG. 1. Activation curve for the reaction C¹²(γ, n)C¹¹.

METHOD			REF. NO.				
Betatron; neutron threshold; ion chamber			60 Ge 3		NVB		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	NØX	THR	C	THR	BF3-I		4 PI

THRESHOLD

TABLE I. Summary and comparison of neutron separation energies inferred from present threshold measurements with values predicted from mass data and reaction energies. All energies are expressed in the center-of-mass system in Mev.

Reaction	No. runs	Present results	Other results	Method	Reference
$C^{12}(\gamma, n)C^{11}$	1	15.115 ± 0.006 (calib)	15.116 ± 0.006	$B^{11}(d, n\gamma)C^{12}$	b

^b R. W. Kavanagh and C. A. Barnes, Phys. Rev. 112, 503 (1958)

METHOD					REF. NO.		
					60 Ha 2		egf
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	15	D	15,17	NAI-D		90

TABLE 2. VALUES OF $\Gamma_{\gamma_0} \sqrt{(\Gamma_{\gamma_0}/\Gamma)}$ (IN eV) OBTAINED IN THE PRESENT EXPERIMENT AT DIFFERENT PROTON BOMBARDING ENERGIES. THE ERRORS SHOWN INCLUDE ONLY THE STATISTICAL UNCERTAINTY

E_p (keV)	500	860	1125	aver.
series 1	53.4 ± 6.6	42.8 ± 4.5	52.9 ± 4.1	49.6 ± 3.0
series 2	34.6 ± 6.7	53.5 ± 4.6	44.3 ± 4.1	44.2 ± 3.0
aver.	44.0 ± 4.8	48.2 ± 3.3	48.6 ± 3.0	47.0 ± 2.1

TABLE 3. VALUES OF Γ_{γ_0} AND $\int \sigma_p dE$ OBTAINED BY VARIOUS INVESTIGATORS

investigation	Γ_{γ_0} (eV)	$\int \sigma_p dE$ (mb MeV)
Hayward & Fuller (1957)	54.5 ± 9.3	1.90 ± 0.27
Garwin (1959)	59.4 ± 9.7	2.33 ± 0.19
Barber & Gudden (1959)	—	$1.47 \pm 0.29^*$
present experiment†	50.5 ± 7.1	2.20 ± 0.31

* In more recent work (Barber, W. C., Berthold, F., Fricko, G. & Gudden, F. E. 1960 *Bull. Amer. Phys. Soc.* (2) 5, 271) it is reported that this result may indeed be too low.
 † From the results of Miller, Pixley & Segel (following paper).

Elem. Sym.	A	Z
C	12	6
Ref. No		JHH
60 Ja 1		

Method	Betatron; photon scattering; NaI
--------	----------------------------------

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, \gamma)$	Bremss. 25	15.1			1^+	Used polarized bremsstrahlung to show that transition is M1.




Fig. 9. Azimuthal angular distribution of elastically scattered 15 Mev gamma rays. This is a polar plot in which the radial scale is in counts per r. The inner circle corresponds to 10 counts/r; the outer circle corresponds to 20 counts/r. The average value of the 8 data points is 19.2 counts/r. Background has been subtracted and 1% detection efficiency corrections have been made to compensate for the position. The solid curve is given by: counts/r = $15.4(1 - 0.51 \cos^2 \theta)$. This corresponds to $P = 1.53; 0.51$ is appropriate rather than 0.53 due to the finite solid angle of the detector.




Fig. 10. Polarization as a function of angle. The experimental points shown are for 15 Mev bremsstrahlung gamma rays arising from 25-Mev electrons in a copper slit collimator. The circles are for the Al target with $\theta = 0.57$; the small circles are for 1% with $\theta = 0.64$. The statistical errors are shown. The solid curves are those given by the calculations discussed in the text.

Elem. Sym.	A	Z
C	12	6

Method: 320 MeV synchrotron; proton telescope; neutron counter

Ref. No. 60 St 1
 JHH

Reaction	E or ΔE	E ₀	Γ	$\int \sigma dE$	Jπ	Notes
$C^{12} (\gamma, np)$	Bremss. 320					$(\sigma/\sigma_{H^2}) = 2.2 \pm 0.3$ $[\sigma_{H^2} = 63 \mu b]$ Mean photon energy - 262 MeV Proton counter at 76°




FIG. 2. Relative cross sections per neutron-proton pair in the nucleus versus atomic number. The cross section of the element of interest is divided by the cross section for deuterium and by the factor NZ/A .




FIG. 3. The relative cross sections per neutron-proton pair, corrected for the probability of escape, is plotted against atomic number. The probability of escape factor is calculated using $r_0 = 1.50 \times 10^{-13}$ cm and $\lambda = 3.6 \times 10^{-12}$ cm. The probability of escape factor is given in expression (1). The data shown are those of Fig. 2 divided by $P(2R/\lambda)$.

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
Linac; total absorption; magnetic compton spectrometer; ion chamber	60 Ta 2
	NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G. MU-T	ABX	11-32	C	32	MAG-D		4PI

456

$$\int_{11}^{32} \sigma_{\text{nucl.}} dE = (200 \pm 35) \text{ MeV-mb}$$

$$\int_{11}^{32} \frac{\sigma}{E} dE = 8.7 \pm 1.5 \text{ mb}$$





FIG. 4. — Sections efficaces d'absorption de photons pour ¹²C.

Ref.

J.M. Wyckoff, H.W. Koch
 Phys. Rev. 117, 1261 (1960)


Elem. Sym.	A	Z
C	12	6

Method
 180 MeV Sychrotron; total absorption; NaI.

Ref. No.
 60 Wy 1

JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
σ_{total}	Bremss.: 35-90			0.267 $\left\{ \begin{array}{l} \gamma, \text{ meson} \\ \text{threshold} \end{array} \right.$ $\pm 15\%$ $\sim 15 \text{ MeV}$ MeV-barns		Substraction of $\sigma_{electronic}$ as base-line.



Elem. Sym.	A	Z
C	12	6
Ref. No. 60 Zi 1		JHH

Method 52 MeV betatron; magnetic pair spectrometer

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
(μ_e)	Bremss. 32 $E_x =$ 10-30					




Fig. 6. Kernabsorptionsquerschnitt von Kohlenstoff. a) Berechnet nach (3). Die strichpunktierte Gerade gibt die bei einer früheren Messung⁴⁾ (wobei nur der Gang mit der Energie, nicht der Absolutwert der Schwächung gemessen wurde) irrtümlich angenommene Lage der Nulllinie von σ_0 an, woraus ein zu kleiner Wert des maximalen Absorptionsquerschnittes resultierte. b) 1. $\sigma(\gamma, p)$ nach Cohen *et al.*²¹⁾; statt des dort angegebenen Wertes $4\pi(d\sigma/d\Omega)_{90^\circ}$ wurde entsprechend der Winkelverteilung von Penner und Leiss²²⁾ ein um 20% kleinerer Wert eingezeichnet.
2. $\sigma(\gamma, p)$ durch detailed balance aus $B^{11}(p, \gamma)$ gewonnen²¹⁾;
3. $\sigma(\gamma, p)$ nach Penner und Leiss²²⁾.
4. $\sigma(\gamma, n)$ nach Barber *et al.*²³⁾
5. Mittelwerte aus 1, 2, 3 addiert zu Kurve 4;
6. Experimentelle Kurve aus Fig. 6a.
7. Riesenresonanz von C^{12} nach G. R. Bishop und R. Wilson im Handbuch der Physik, Bd. 42, S. 337 (Springer Verlag 1957)
c) $\sigma(\gamma, 3z)$ nach Goward und Wilkins²⁴⁾, ohne Feinstruktur.

Ref 4: B. Ziegler, Z. Phys. 124, 506 (1958).
Ref 21: L. Cohen *et al.*, Phys. Rev. 10, 108 (1956)
Ref 22: Penner and Leiss - Phys. Rev. 114, 1101 (1959)
Ref 23: Barber, George and Reagan - Phys. Rev. 98, 73 (1955)
Ref 24: Goward and Wilkins - Proc. Roy. Soc. A 228, 376 (1955); Proc. Roy. Soc. A 217, 357 (1953).

METHOD				[Page 1 of 2]		REF. NO.		NVB
Linac						61 Bo 4		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
E.E./	SPC	0-164	D	194	MAG-D		135	




FIG. 3. — Spectre expérimental à 135° de diffusion d'électrons de 194 MeV sur une cible de graphite de 0,961 g/cm².




FIG. 4. — Spectre pris avec des cibles différentes et queues radiatives. La courbe la plus basse est la courbe n° 1.

METHOD				REF. NO.			
Linac				[Page 2 of 2]			
				61 Bo 4			
				NVB			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




FIG. 5. — Spectre de diffusion d'électrons de 194 MeV sur du carbone à 135° (après soustraction de la queue radiative).
 | Après soustraction de la QR n° 2.
 | Après soustraction de la QR n° 1.
 — Spectre théorique dans le cas d'un noyau à potentiel parabolique en supposant une répartition gaussienne de la charge dans le proton. (Le spectre a été normalisé pour que son sommet coïncide avec celui du spectre réel.)

TABLE I

ÉNERGIE D'EXCITATION MeV	$\frac{d\sigma}{d\Omega}$ MB/STÉRIAN	
	PROTON PONCTUEL	PROTON GAUSSIEN
15,3	6,904	2,16
30,6	29,36	11,16
45,9	31,57	13,16
61,2	24,38	10,95
76,5	16,42	7,78
91,8	9,38	4,65
107,1	3,89	2,03
122,4	1,07	0,59
137,7	0,19	0,12
153,0	0,02	0,01

REF. A. Bussiere de Nercy
 J. Phys. Radium 22, 535 (1961) (See 61Bu4)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		NVB
61 Bu 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	19-36	C	32	NAI-D		140




FIG. 2.

Method Synchrotron; Bremss; ~ monoergic photons by coincidence with electrons from donut target; proton telescope of 10 plastic scintillator separated by Cu absorbers.

Ref. No. 61 Ce 1
 JHH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C(γ,p)	245 ± 15					Figure 4 data fitted with two quasi-deuteron momentum distributions.




Fig. 4. Experimental results. Plot of the differential cross section per nucleus per selected photon for photoproton production from carbon at 60 deg by 245 ± 15 Mev photons. The curves are explained in the text.




Fig. 5. Plot of the ratio of the differential cross section to the total cross section for photoproton production from carbon at 60 deg by 245 Mev photons. The curves are explained in the text.

METHOD				REF. NO.		NVB	
Linac				61 Du 1			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	13-18	D	46-154	MAG-D		135




FIG. 2.




FIG. 3.




FIG. 4.

METHOD				REF. NO.			
				61 Du 1			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	

TABLEAU 1

E_0 MeV	$\left(\frac{d\sigma}{d\Omega}\right)_{Mott}$	$\left(\frac{d\sigma}{d\Omega}\right)_{el}$ expér.	$\left(\frac{d\sigma}{d\Omega}\right)_{Mott} F_{el}^2$	$\theta = 135^\circ$		$R_{16,1}$	$\left(\frac{d\sigma}{d\Omega}\right)_{15,1}$	$R_{16,1}$	$\left(\frac{d\sigma}{d\Omega}\right)_{16,1}$
				$10^{-29} \text{ cm}^2/\text{stéradian}$	10^{-3}				
46	1,76	1,3 ± 10 %	1,24	2,06 ± 10 %	2,56 ± 0,26				
63,5	0,92	0,42 ± 10 %	0,44	4,04 ± 10 %	1,78 ± 0,18	1,08 ± 32 %			0,43 ± 0,15
81	0,56	0,18 ± 10 %	0,16	6,66 ± 12 %	1,07 ± 0,13	3,75 ± 20 %			0,60 ± 0,12
82,5	0,54	0,15 ± 10 %	0,15	5,60 ± 12 %	0,84 ± 0,10	2,36 ± 20 %			0,36 ± 0,07
100,5	0,36	0,055 ± 10 %	0,054	7,45 ± 12 %	0,40 ± 0,05	9,45 ± 10 %			0,51 ± 0,05
119	0,26	0,016 ± 10 %	0,017	10,44 ± 57 %	0,18 ± 0,10	29,36 ± 20 %			0,50 ± 0,10
154	0,15	0,001 ± 10 %	0,0011			500 ± 40 %			0,55 ± 0,22




Fig. 5. — Calcul de Dalitz et Yennie. Noyau ponctuel. $\Gamma_\gamma = 55 \text{ eV}$.




Fig. 6. — Calcul de Dalitz et Yennie. Noyau ponctuel $\Gamma_\gamma = 1,6 \text{ eV}$ (normalisé pour $q^2 = 0,25 \text{ Fermi}^{-2}$)

Method Tandem accelerator; NaI

Ref. No. 61 Go 1
EH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
B ¹¹ (p,γ)C ¹²	2.6-11.4	22.5±0.02		∫ 4.7 X 10 ⁻⁴ MeV-b		<p>E_{P0} = 7.12 MeV</p> <p>σ(E_{P0} = 7.12 MeV) = 1.5X10⁻²⁸ cm².</p> <p>This corresponds, in the inverse reaction, to the peak in the giant resonance with E_{γ0} = 22.5±0.02 MeV.</p> <p>∫σ(γ,p)dE = 0.043 MeV-b by detailed balance method.</p> <p>The absolute cross section obtained here results from normalizing to the data of Huus and Day [Phys. Rev. <u>91</u>, 599 (1953)].</p>




Fig. 8. Partial energy level diagrams for C¹², Mg²⁴ and Na²³ showing the region of excitation energy (20-25 MeV) the positions of the first excited states and the energies at which each nucleus becomes unstable against neutron, proton and alpha particle emission. The approximate position of the peak of the giant resonance (GR) is also indicated.




Fig. 9. The yield of gamma rays leading to the ground and first excited state of C¹² resulting from proton capture in B¹¹ and measured at 90° to the incident beam. The different symbols on the γ₀ yield curve represent separate runs. The ordinates of the two yield curves are correct relative to each other. The abscissae are marked both in terms of the incident laboratory proton energy and in excitation energy in the compound nucleus C¹².




Fig. 10. Angular distributions of ground state gamma rays from the reaction B¹¹(p,γ)C¹² measured at four incident proton energies below and above the giant resonance in C¹². The proton energy of 7.12 MeV corresponds to the peak of the giant resonance. The solid lines are least square Legendre polynomial fits and the values of the coefficients are given in the expressions 1-a, P₁-b, P₂.

E _p (MeV)	Excitation energy (MeV in C ¹²)	a	b	Notes
7.12	22.5	-0.12 ± 0.03	-0.09 ± 0.05	Present work
21.1	21.1	0.00 ± 0.04	-0.26 ± 0.04	Present work
19.4	23.7	0.21 ± 0.06	0.37 ± 0.09	Present work
	23.3	0.27 ± 0.04	0.14 ± 0.06	Present work

Ret.

V. Presperin, L.A. Kul'chitskii
 Zhur. Eksp. i Teorat. Fiz. 41, 60 (1961)
 Soviet Phys. JETP 14, 46 (1962)

Elem. Sym. A Z

C 12 6

Method Cu^{63} (n,2n) threshold detectors; positrons counted

Ref. No. 61 Pr 1

JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$\text{C}^{12}(\gamma, n)$	Bremss.: 30 - 85					<p>In figure, curve 2 is calculated (using Penfold-Leiss) from curve 1, $\sigma(\gamma, n)$ for all $E_n > 11$ MeV; curve 3 is $\sigma(\gamma, n)$ for $E_n > 11-21$ MeV.</p> <p>Yields analysed in terms of $\text{Cu}^{63}(n, 2n)\text{Cu}^{62}$ cross section and assumed neutron spectrum of from E^{-n}; $n = 1.5$.</p>




FIG. 1

Elem. Sym.	A	Z
C	12	6

Method 24 MeV betatron; radioactivity; neutron yield; proportional counter;
 r chamber.

Ref. No.
 61 Ro 2
 EH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
$C^{12}(\gamma, n)$	Bremss. 25	23.0	3.2 MeV	$\int_{th}^{24} = 0.022 \text{ MeV-b}$		At E _n = 23.0, σ _{max} = 7.9 mb Relative yield curve normalized by standard Cu ⁶³ (γ, n) reaction [Roalsvi Haslam, McKenzie, Can. J. Phys. <u>37</u> , 607 (1959)].




Fig. 5. Absolute activation curve for the reaction $C^{12}(\gamma, n)C^{11}$.




Fig. 6. Cross-section curve for the reaction $C^{12}(\gamma, n)C^{11}$.

SEE PAGE 2 FOR TABLES

Method

Ref. No.

61 Ro 2

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
----------	---------	----------------	---	------	----	-------

TABLE II

References	E _{max} (MeV) energy for max σ ⁰	Yield at E _{max} (n/g-atom 100 r)	σ _{max} (mb)	Half width F _{1/2} (MeV)	E ₁ (MeV)	E ₂ (MeV)	Norm. integr. cross section (MeV)	As given by authors	Norm. to present yield (MeV bars)
Haslam et al. (1961)	22.4	—	11.6	4.2	0.000 (27)	0.000	2.41	—	—
Katz and Cameron (1961)	22.0	1.95 × 10 ⁶	12.1	2.8	0.000 (27)	0.000	2.28	—	—
Montalbetti et al. (1963)	21.4	2.05 × 10 ⁶	12.4	2.0	0.000 (24)	0.000	2.15	—	—
Nathans and Halpern (1964)	22.0	Activation curve not given in paper	8.6	2.0	0.000 (25)	0.000	2.02	—	—
Barber et al. (1965)	22.5	3.46 × 10 ⁶	8.2	4.2	0.002 (25)	0.007	2.20	—	—
Cook (1967)	22.0	2.75 × 10 ⁶	10.4	2.5	0.000 (25)	0.000	2.00	—	—
Present work	22.0	2.77 × 10 ⁶	7.9	2.2	0.000 (25)	0.002	2.25	0.000	0.000
Katz et al. (1964) ^a	—	—	—	—	—	—	—	0.000	0.000
Thorne (1967)	—	—	—	—	—	—	—	0.000	0.000
Average value ^b	22.4	2.65 × 10 ⁶	10.2	3.4	—	—	—	—	—
Standard deviation	0.6	0.3 × 10 ⁶	2.0	0.8	—	—	—	—	—

^aRedistribution of values of Haslam et al. (1961) using the photon-difference method. Values are normalized according to Montalbetti et al. (1963).
^bValues of Haslam et al. (1961) are not included directly, but indirectly under Katz and Cameron (1964).
 These values should be compared to the present value 0.000 MeV bars for the average yield curve.

TABLE I
Absolute yields of O¹⁶(γ,n)O¹⁶ and C¹²(γ,n)C¹² at 22 MeV

References	Absolute yield of O ¹⁶ (γ,n)O ¹⁶ (n/g-atom 100 r)	Absolute yield of C ¹² (γ,n)C ¹² (n/g-atom 100 r)
Price and Kavet (1960)	0.67 × 10 ⁶	0.62 × 10 ⁶
Haslam, Johns, and Horsley (1961)	—	0.75 × 10 ⁶
Johns, Horsley, Haslam, and Quanten (1961)	1.60 × 10 ⁶	—
Montalbetti et al. (1963)	2.7 × 10 ⁶	2.2 × 10 ⁶
Nathans and Halpern (1964)	—	2.5 × 10 ⁶
Barber et al. (1965)	—	2.3 × 10 ⁶
Cook (1967)	—	1.8 × 10 ⁶
Present work	2.37 × 10 ⁶	0.78 × 10 ⁶
		2.16 × 10 ⁶

^aAccording to the note added in proof the author mentions that this value should be reduced by 10%.
^bA redistribution of the absolute yield gave results in close agreement with this value.

Elem. Sym.	A	Z
C	12	6
Ref. No.		JHH
61 Se 2		

Method Cockraft-Walton generator; NaI

Reaction	E or ΔE	E _γ	Γ	∫σdE	Jπ	Notes
B ¹¹ (p,γ)	150-183 keV E ₀ = 163 keV	16.11	0.22±0.09 eV			Detector at 90° Pure E2 transition.




Fig. 1. Principal modes of decay of the 16.11-Mev state in C¹². The partial widths for these decays are given in Table I.




Fig. 2. Yield of gamma rays of energy 16.1 Mev from a natural boron target irradiated with 10¹² neutrons per second. The intensity units are arbitrary and the region is in arbitrary units. The arrow indicates the summing effect of most of the data were taken.

TABLE I. Parameters of the 16.11-Mev state in C¹². All widths are in eV.

	Reference 1	Present work
Γ _{γb}	7000	7300±500 ^a
Γ _p	5	69±15 ^b
Γ _{α0}	100	290±45
Γ _{α1}	5000	6300±500
Γ _{γ0}	2 ^c	0.22±0.09 ^c
Γ _{γ1}	70	6.8±1.1

^a Reference 7.
^b Using the gamma ray yield given in reference 4.
^c Taking for γ₀ = 0.3 ± 0.1% from reference 9.

Ref. 7: Hunt and Jones, Phys. Rev. 39, 1285 (1953).
 Ref. 4: Huus and Day, Phys. Rev. 91, 599 (1953).
 Ref. 9: Craig, Cross and Jarvis, Phys. Rev. 103, 1414 (1956).

TABLE II. Reduced widths in single-particle units for various radiations from the 16.11-Mev state in C¹². The basis for single-particle units is given in the text.

Radiation	η ²
p	0.8
α ₀	10 ⁻⁴
α ₁	2.5 × 10 ⁻³
γ ₀	0.15
γ ₁	0.20

METHOD			SOURCE		DETECTOR		REF. NO.	ANGLE
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE		
Van de Graaf; 3 alpha spectrum; nuclear emulsion							61 Se 3	NVB
G,3A	RLY	18	D	18 (17.6)	EMU-D			4 PI

E* peaks: E* is excitation energy in Be⁸.
 2.9 MeV
 8.7




FIG. 1. — Distribution de E*.




FIG. 2. — Distribution expérimentale et distributions calculées de E_α.

METHOD				REF. NO.			
Betatron; proton yield; radioactivity				61 Th 1			
				NVB			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.N	ABI	18-23	C	18-23	ACT-I		4PI

$$\int_{18}^{23} \sigma(E) dE = 14 \pm 3 \text{ MeV-mb}$$

18 BREAKS

Table 1. Positions of Breaks and the Integrated (γ, n) Cross Section of the Photon Absorption Levels corresponding to these Breaks in the $^{12}\text{C}(\gamma, n)^{11}\text{C}$ Reactions

(1)	(2)	(3)	(4)
18.73 Threshold			
18.90	0.0125		18.86
18.96	0.0155		18.96
19.08	0.0194	19.10	19.10
19.17	0.0236		
19.30	0.0378		
19.46	0.0583		
19.57	0.0708	19.55 \pm 0.05	19.56
19.76	0.0626		
19.92	0.127	19.95 \pm 0.05	19.95
20.13	0.206		
20.29	0.282	20.35 \pm 0.05	20.37
20.62	0.616	20.75 \pm 0.05	20.75
20.90	0.657		
21.08	0.761		
21.22	0.956		
21.58	3.13		
22.02	2.80		
22.88	4.12		

(1) Energy of break E_i (mev) (energies are to within ± 0.04 mev near threshold and ± 0.07 mev near 23 mev); (2) $\int (\sigma_{\gamma, n})_i dE_{\gamma}$ (mev mbn) (relative values are given to 3 significant figures, absolute values are probably known to within 20%); (3) Spicer and Penfold (1955) E_i (mev); (4) Sadeh (1959) E_i (mev) (Sadeh quotes his energies to within ± 0.02 mev; this seems rather optimistic).

Elem. Sym.	A	Z
C	12	6
Ref. No.		EGF
61 Va 1		

Method Stanford Mark II accelerator; 8 KI counters in focal plane of magnetic spectrometer

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(γ,p)	24	22.5		∫ ²⁴ = 41±9 MeV-mb		σ _{max} = 17.7±2.5 mb Assume f(θ) = 1+0.25 cosθ + 1.35 sin ² θ for ground state transitions. Integrals based on the measured angular distribution (1+1.5 sin ² θ); Pa. data is consistent if this distribution instead of isotropy is used.
(e,p)	40			∫ ⁴⁰ = 77±17 MeV-mb		




Fig. 2. Cross section of C¹²(γ, p) assuming 100% of transitions leave B¹¹ in ground state and cross section of C¹²(γ, n) from proton difference method.

In Figure 2, C¹²(γ, n) curve is from Barber et al, Phys. Rev. 98, 73, (1955).

ERRATA: In Figures 5 and 6 captions, (e, e'n) and (γ, n) should read (e, e'p) and (γ, p) respectively.




Fig. 5. Angular distributions for E₀ = 40 MeV of a proton energy group from C¹²(e, e'n) and C¹²(γ, n) reactions. The curves represent least-square fits to the data.
Circles: 2.36 [1 - 0.25 cos θ - 1.42 sin² θ - 0.09 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr.
Crosses: 2.49 [1 - 0.27 cos θ - 1.40 sin² θ - 0.04 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr




Fig. 6. Angular distribution for E₀ = 40 MeV of a proton energy group from C¹²(e, e'n) and C¹²(γ, n) reactions. The curves represent least square fits to the data.
Circles: 3.12 [1 - 0.24 cos θ - 1.30 sin² θ - 0.05 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr
Crosses: 2.83 [1 - 0.26 cos θ - 1.20 sin² θ - 0.04 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr.




Fig. 7. Angular distributions for E₀ = 40 MeV at different proton energy groups from C¹²(γ, n) reaction.
Circles: 1.57 [1 - 0.27 cos θ - 1.24 sin² θ - 0.04 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr
Crosses: 1.39 [1 - 0.27 cos θ - 1.24 sin² θ - 0.04 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr
Triangles: 1.03 [1 - 0.24 cos θ - 1.24 sin² θ - 0.04 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr
Squares: 0.83 [1 - 0.24 cos θ - 1.24 sin² θ - 0.04 sin⁴ θ cos θ] × 10⁻⁴¹ cm² MeVsr

METHOD			SOURCE		DETECTOR		ANGLE
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE	
G.G	ABX	40 - 120	C	132	SCI-D		DST

REF. NO.
61 W1 1 EGF




Fig. 7. The observed differential cross section at 90° against photon energy. For details of the solid line, see the text.




Fig. 8. The observed differential cross sections at 112° against photon energy. For details of the solid line, see the text.




Fig. 9. The observed differential cross sections at 135° against photon energy. For details of the solid line, see the text.




Fig. 10. The differential cross section for the scattering in the range 80-120 mev against $\cos \theta$. For details of the solid lines, see the text.

Elem. Sym.	A	Z
C	12	6

Method 100 MeV electron synchrotron; activation, NaI coinc. for annihilation radiation of β^+ from C^{11} .
 Ref. No. 62 Bo 3 JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n)$	Bremss. 30-90					




Fig. 3. The $C^{12}(\gamma, n)C^{11}$ yield, normalized to the data of ref. 6).





Fig. 4. The $C^{12}(\gamma, n)C^{11}$ cross section and integrated cross section.




Fig. 5. The comparison among the data of refs. 6, 8) and those of $C^{12}(\gamma, n)C^{11}$ of the present work.

Ref 6: Barber, George, Reagan - Phys. Rev. 98, 73 (1955)
 Ref 8: Cook - Phys. Rev. 106, 300 (1957)

Method: 90 MeV Synchrotron; magnetic spectrometer; emulsions; NaI counter telescope

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(γ,d)	Bremss. 90					
(γ,p)						




Fig. 4. The ratio of the yields of deuterons and protons with energies 15.5 to 30 MeV depending on the mass number of nuclei A for $E_{\gamma, \max} = 90$ MeV. The solid line stands for the normalized dependence.




Fig. 5. The relative deuteron and proton yields plotted against particle energy E for $E_{\gamma, \max} = 90$ MeV. a) Black circles denote Be and open circles Li; b) Black circles denote Li and open circles Li.




Fig. 7. Angular distributions of photodeuterons with the energies 15.5 to 30 MeV emitted in the photodisintegration by bremsstrahlung with $E_{\gamma, \max} = 90$ MeV. Crosses denote Li, black circles Li, open circles Be, and open squares C. Particle yields are given in arbitrary units.




Fig. 8. Angular distributions of photoprotons with the energies 15.5 to 30 MeV. See caption fig. 7.

Method Electrostatic generator, $H^3(p,\gamma)He^4$ reaction; activation of positron emitter; 2 NaI in coincidence.

Ref. No. 62 De 1
 JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
(γ, n)	20.1-21.2					$\sigma(\gamma, n) = 1.04 \pm 0.11$ mb at 20.48 MeV

1052




Figure 6: The $C^{12}(\gamma, n)C^{11}$ reaction cross section in millibarns as a function of photon energy in MeV.

Method Magnetic analysis of proton spectra produced by electron bombardment

Ref. No.	BG
62Dol	

Reaction	E or ΔE	E ₀	Γ	∫ σ _{dE}	Jπ	Notes
C ¹² (e,p)	30 24.5	6.7 7.2 (8.2) (8.9) 10.2 10.9	Giant resonance width=3.1	29.3 ∫ σ(γ,p)dE _γ = 20.3 50±8 MeV-mb		Correspondence between (e,pe') reaction and (γ,p) reaction. Assumed electron has associated with it a virtual photon spectrum. Electron production yields were analyzed by use of E1 virtual photon spectrum to obtain σ(γ,p). Angular distribution of proton measured; assumed of proton for A' + B' cosθ + C' sin ² θ + ... for incident photons.
				100% ground state transitions assumed.		

See page 2 for figures and tables

Elem. Sym.	A	Z
C	12	6

Method

Ref. No.
62 Do 1

Reaction	E or ΔE	E ₀	Γ	∫σdE	∫π
----------	---------	----------------	---	------	----

Notes




FIG. 21. Carbon (e,pe') cross section for E₀ = 30 MeV and θ = 76° under the assumption of 100% ground-state transitions.




FIG. 22. Carbon (e,pe') cross section for E₀ = 24.5 MeV and θ = 76° under the assumption of 100% ground-state transitions.




FIG. 36. Carbon (e,pe') proton angular distribution at E₀ = 24.5 MeV.




FIG. 37. Carbon (e,pe') proton angular distributions at E₀ = 24.5 MeV.

TABLE VIII. Comparison of the direct and inverse C¹²(e,pe') angular-distribution coefficients. The errors of this experiment are standard deviations.

Experiment	E _γ	C ₀	C ₁	C ₂	C ₃
C ¹² (e,pe') ^a	22-23	1	0.14 ± 0.02	-0.50 ± 0.03	...
C ¹² (e,pe') ^b	22.1	1 ± 0.02	0.09 ± 0.02	-0.56 ± 0.04	-0.03 ± 0.01
Bi ²⁰⁹ (e,pe') ^c	22.5	1	0.12 ± 0.03	-0.69 ± 0.05	...
This experiment	22.4	1 ± 0.05	0.16 ± 0.09	-0.61 ± 0.04	0.11 ± 0.02

^a See reference 56.
^b See reference 57.
^c See reference 59.

^d M. M. Hoffman and A. G. W. Cameron, Phys. Rev. 92, 11 (1953).




FIG. 38. Ratio of the even-parity terms in the angular distributions of protons from C.


Ref. R.D. Edge, G.A. Peterson
 Phys. Rev. 128, 2750 (1962)

Elem. Sym.	A	Z
C	12	6

Method Linac; counter telescope

Ref. No.	
62 Ed 1	BG

Reaction	E or ΔE	E_0	Γ_γ (eV)	$\int \sigma dE$ (MeV-mb) / π	Notes
$C^{12} (e, e')$	41.5	15.1	39	1.95 ± 0.2	<p>Nuclear states excited by 180° electron scattering; M1 transitions assumed.</p> <p>Inelastic electron scattering cross sections obtained by comparing inelastic peaks to e-p elastic scattering peak.</p> <p>Γ_γ from virtual photon theory.</p> <p>Limits not given for cross sections.</p> <p>$\Gamma_\gamma = 39$ eV corresponds to transition from 1^+ excited state to 0^+ ground state.</p> <p>Inelastic crosssection of 15.1 MeV state = $1.82 (\pm 10\%) \times 10^{-32}$ cm²/sr</p>



REF.

F. W. K. Firk, K. H. Lokan and E. M. Bowey
Proc. Padua Conf. 804 (1962)

ELEM. SYM. A C

C

12

6

METHOD

REF. NO.

62 Fi 2

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.N	RLX	21-29	C	25-32	TOF-D	2-15	




Fig. 2. The relative $O^{14}(\gamma, n)$ cross-section for $E_{\gamma, \text{max}} = 32.5$ and 26.5 MeV.

METHOD Stilbene scintillator; neutron spectrum

REF. NO.

62 Fu 5

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.N	ABX	21 - 31	C	31	SCI-D		90 (90±15)




Abb. 1. Ergebnisse für den C^{12} -Kern. 1 a und 1 b: C^{12} -Photo-neutronenspektren, gemessen bei zwei verschiedenen Einstellungen der Impulsformdiskrimination. Endenergie des Bremsstrahlspektrums $E_0=31$ MeV. Beobachtungsrichtung gegenüber dem γ -Strahl $\Theta=90^\circ \pm 15^\circ$. 1 c: Histogramm: Wirkungsquerschnitt der $C^{12}(\gamma, n_0)C^{11}$ -Reaktion, berechnet aus dem Spektrum 1 b unter der Annahme, daß der C^{11} -Kern stets im Grundzustand zurückbleibt. Die Fehlerbalken geben den statistischen Meßfehler an. Die Unsicherheit der angegebenen Absolutwerte beträgt etwa 25%. Kurve: Wirkungsquerschnitt der $C^{12}(\gamma, n)C^{11}$ -Reaktion nach BARBER et al.

REF.

K.O. Hermann, J.A. Scheer
Z. Physik 170, 162 (1962)

ELEM. SYM.	A	Z
C	12	6

METHOD

Betatron; proton spectrum; solid state

REF. NO.

62 He 1

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XP	SPC	19 - 27	C	31	SCD - D	3 - 9.4	




Fig. 1. Photoprotonen aus C^{14} . Die angegebenen γ -Energien E_γ gelten unter der Annahme, daß die Reaktionen zum Grundzustand des Folgekerns führen. Da der Detektor zur Vergrößerung der Zählrate möglichst nahe an das Target herangebracht worden war, war der von Elektronen und γ -Quanten herrührende Untergrund trotz der relativen Unempfindlichkeit des Detektors ziemlich stark.

Elem. Sym.	A	Z
C	12	6

Phys.Rev. 128, 2784 (1962)

Method: Synch-defining counter telescope, proton counter telescope - C analyzer

Ref. No. 62L11
BG

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(γ,p)	335					<p>Polarization of high energy photo-protons given in %.</p> <p>Mean proton production energy = 168 MeV</p> <p>$\theta_p = 45^\circ \quad +24 \pm 15$</p> <p>$\theta_p = 56^\circ \quad -11.8 \pm 15$</p> <p>$\theta_p = 90^\circ \quad -11.3 \pm 15$</p> <p>Postulates identical γ absorption process for each kind of nucleus (quasi-deuteron).</p> <p>Fig.2: combined results of Li⁷, Be⁹, B¹¹, C¹² at 3 angles of measurement. $\theta_{\text{plab}} = 90^\circ$ only for Carbon data.</p> <p>Fig. 2 A: Only E1 transitions considered. C-M1 transitions from $^3S_1 - ^1S_0$ also taken into account.</p> <p>Results are consistent both with a zero value for the polarization and also with theory.</p>

FIG. 2. Comparison of the experimental results with the predicted polarization.

Elem. Sym.	A	Z
C	12	6

Method Positron annihilation; neutron yield; 4π neutron; positron current

Ref. No. 62 Mi 2
 JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{12}(\gamma, n)$	18-26	22.2 23.3		$\int_{18}^{26} = 24 \text{ MeV}\cdot\text{mb}$		

Fig. 1. Section efficace de la réaction (γ, n) sur le carbone.

Tableau 1

Noyau Z	$\int_{E_{min}}^{E_{max}} (\sigma - \dots) dE$ (MeV - mb)	Energie des niveaux (MeV)	Largeur des niveaux (MeV)	Section efficace intégrée des niveaux discrets (MeV-mb)
C^{12}	24	22.2 23.3		
O^{16}	13	17.5	-1	0.6
		19.4		1.6
		21.1		1.4
		22.2		(4.5)
		23.2		(2.4)
Ca^{40}	53.5	(19.1) 19.3 (21.4)		

REF.

W.T. Morton, T.G. Walker
Phil. Mag. 7, 741 (1962)

C

12

6

METHOD

Synchrotron; p and α cross section; nuclear emulsion

REF. NO.


62 Mo 2

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, PA	ABI	25-120	C	120	EMU-D		4PI

$$\int_0^{120} \sigma dE = 1.9 \pm 0.4 \text{ MeV-mb}$$

Fig. 2

Energy dependence of $(\gamma, p + \alpha)$ reactions in ^{12}C , ^{14}N and ^{16}O .

Method
150 MeV linac - CsI(Tl)

Ref. No.
62Pal
B6

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(γ,p)						<p>Electrodisintegration and photo-disintegration cross section data (at 60°) given.</p> <p>Results compared with "quasi-deuteron" model and direct γ-p interaction.</p> <p>E_p = 33-98</p>




Fig. 5. Photoproton spectrum from carbon at 60° compared with the theoretical curves.

Method: Linac; NaI; detector at 90°. Ref. No. 62 Se 1 JHH

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C ¹² (γ,γ)	prems; 19	4.4 10.7 15.1			1+	0.31±0.07 ≤ 0.02 1.00 , relative numbers of γ-rays scattered at 90°.




FIG. 5. Pulse-height spectrum for γ rays scattered from a C sample. Sample-out background has been subtracted. The solid lines are KS response functions for 4.4- and 15.1-Mev γ rays. The dashed line is an estimate of the sample-associated background.

Elem. Sym.	A	Z
C	12	6
Ref. No.		JHH
63 Be 5		

Method Van de Graaff; γ yield; NaI

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$B^{11}(p,\gamma)$	6-13.2					
	~ 7.2	22.56				

Table I
Excited states of C^{12}

Configuration	Spin	Theory (%)	Expt.
$(1p_1)^{-1}(1p_3)^{-1}$	1^+	16.1	15.11 ¹⁾
$(1p_1)^{-1}(2s_1)^{-1}$	1^-	18.7	17.23 ²⁾
$(1p_1)^{-1}(1d_1)^{-1}$	1^+	22.2	22.25 ³⁾
$(1p_1)^{-1}(1d_3)^{-1}$	1^-	23.9	25.5 ⁴⁾
$(1s_1)^{-1}(1p_1)^{-1}$	1^-	34.3	$\sim 34^5)$
$(1p_1)^{-1}(1d_3)^{-1}$	2^-	18.1	16.57 ⁶⁾
$(1p_1)^{-1}(1d_1)^{-1}$	2^-	19.2	22.1 ⁷⁾
$(1p_1)^{-1}(2s_1)^{-1}$	2^-	22.9	23.6 ⁸⁾
$(1p_1)^{-1}(1d_3)^{-1}$	3^-	$\sim 26.4^9)$	26.9 ¹⁰⁾

¹⁾ This experiment.




Fig. 3. The relative yields at 90° of capture gamma radiation to the ground state (1+) and the excited state (2+) of C^{12} following proton capture in B^{11} . The incident proton energy range is from 6 to 13 MeV.

References

- Gove, Litherland and Batchelor, Nuclear Physics 16 (1960) 430
- Reay, Lee and Hintz, University of Minnesota Progress Report 1961 (unpublished)
- F. Auzenberg-Seiove and T. Lauritsen, Nuclear Physics 11 (1959) 1
- M. Elaine Tums, U. S. Naval Research Laboratory No. 13 (1960) (unpublished)
- J. Fujita, Prog. Theor. Phys. 16 (1956) 112
- Gammel, Morton and Titterton, Nuclear Physics 10 (1959) 25
- S. Yinn-Mau and G. E. Brown, Nuclear Physics 29 (1962) 49
- Bair, Kingston and Walard, Phys. Rev. 100 (1955) 21
- M. Nomoto, private communication

METHOD					REF. NO.		
Linac; electron scattering; magnetic spectrometer					63 Bo 3		NVB
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	4-25	D	200	MAG-D	20-190	135

E_0	$\frac{d^2\sigma}{d\Omega dE}$
4.43 MeV	$3.7 \pm 0.4 \mu\text{ barns/str.}$
16.1 MeV	$4.2 \pm 2 \mu\text{ barns/str.}$
19.8 MeV	$7 \pm 2 \mu\text{ barns/str.}$




FIG. 1. — Spectre des électrons de 198 MeV diffusés à 135° par une cible de graphite.




FIG. 3. — Spectre complet d'électrons de 198 MeV diffusés à 135° par une cible de graphite. Interprétation théorique par approximation de l'impulsion.

Ref.

N.A. Burgov, G.V. Danilyan, B.S. Dolbilkin, L.E. Lazareva,
F.A. Nikolaev
Izv. Akad. Nauk SSR 27, 866 (1963)

Elem. Sym.	A	Z
C	12	6

Method

250 MeV Synchrotron; pair spectrometer

Ref. No.

63 Bu 1

JHH

Reaction	E or ΔE	E ₀	Γ	∫σdE	∫π	Notes
$C^{12}(\mu)$		16.5 17.6 19.1 22.5 25.5				For relative intensities, see second column, Table 2, below.




Рис. 3. Аппаратурная кривая ядерного сечения поглощения γ -квантов C^{12} , исправленная на «функцию разрешения»

Таблица 2
Сравнение экспериментальных данных по C^{12} с результатами теоретических расчетов

E, MeV	Данная работа		[13]		[12]	
	I, %	E, MeV	I, %	E, MeV	I, %	E, MeV
16,5	5	17,03	12	17,03	12	17,03
17,6	6,5					
19,1	6	18,7	0,5	18,7	3	18,7
22,5	57	22,2 23,9	75 0,5	22,48	54	22,48
25,5	22	—	—	24,26	25	24,26
—	—	34,3	18	—	—	—

Таблица 1

Энергии уровней C^{12} (MeV) в интервале 16–27 MeV

По основному	По кривой выхода пик C^{12} (7)	По спектру фотопротонов (8)	По спектру фотонейтронов (11)	По сечению реакции (p, n) (10)	Согласно обзорной статье (5)
16,5	—	—	—	—	16,58 (2-?)
17,6	—	—	—	17,22	17,22 (1-?)
—	—	—	—	—	18,40
—	18,90	—	—	—	18,55
—	18,96	—	—	—	—
—	19,06	—	—	—	—
19,1	19,17	—	—	(19,2)	19,28 (1-?)
—	19,30	—	—	—	19,42
—	19,46	—	—	—	—
—	19,57	—	—	—	—
—	19,58	—	—	—	19,57
—	19,92	—	—	—	—
—	20,13	—	—	—	19,89
—	20,29	—	—	—	20,27
—	20,62	—	—	—	20,49
—	20,90	—	—	—	20,65
—	21,08	—	—	—	—
—	21,22	—	—	—	—
—	21,58	—	—	—	21,34
—	22,02	—	—	—	21,50
22,5	—	22,55	23,0	22,5	—
25,5	—	(25,5)	26,0	25,5	—
—	—	(27,5)	—	—	—

METHOD Synchrotron; total nuclear absorption; magnetic pair spectrometer; ion chamber REF. NO. 63 Bu 3 NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	13-27	C	220	MGP-D		4PI

Table I. C¹² energy levels in the region 16-28 MeV obtained by different methods

Present work	From the nuclear absorption cross section [2,4]		From the yield curve of C ¹² (γ,n) [9]		From photo-neutron spectrum [13]	From photo-proton spectrum [13]	From cross section for inverse reaction B ¹¹ (p, γ)C ¹² [11]	From cross section for reactions induced by charged particles [11]	
16.5								16.53	
17.6								17.23	
								17.77	
			18.00	(60)				18.40	
			18.96	(120)				18.85	
			19.08	(90)					
19.1		19.17				19.2	19.26		
20.15			19.30	(130)				19.42	
			19.46	(160)					
			19.57	(110)				19.67	
			19.76	(190)				19.83	
			19.92	(160)					
			20.13	(210)				20.27	
	20.46		20.27	(140)				20.49	
			20.62	(350)				20.65	
	20.92			20.90	(250)				
				21.08	(180)				
			21.22	(140)				21.34	
			21.53	(360)				21.80	
		22.02	(440)						
23.0	22.5			23.0	22.55	22.5			
25.6				26.0	25.5 27.5	25.5			

*Distances in keV between neighboring "breaks" are given in parentheses.




FIG. 2. Cross sections for the C¹²(γ,n) and C¹²(γ,p) reactions compared with the measured cross section for γ-ray absorption in carbon. 1—σ(γ,n), [22] with data above 25.5 MeV taken from [23]; 2—σ(γ,p); [13] 3—σ(γ,n) + σ(γ,p); 4—cross section for nuclear absorption in the 20-27 MeV region.

Nuclear absorption cross section & references on page 2

METHOD			REF. NO.				
			63 Bu 3				
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	

Table II. Parameters of C¹² giant resonance obtained by different methods

	E _{max} , MeV, of cross section	Cross section at maximum σ _{max} , mb	Γ, MeV	∫σdE, MeV-mb	Source
Absorption cross section	23	16	3.2	84(27)*	Present authors [1]
	23	17	4	100(27)	
Nuclear cross section of (γ,p) reaction	21.5	34	1.7	63(24)	[22]
	23	14.7**		46(24)	[7]
	22.1	3.1**	3.6		[24]
	23.5	2.7**			[10]
	23.3	12.7	3.3	57(27)	[12]
	22.3	12	3.1		[11]
Nuclear cross section of (γ,n) reaction	22.5	3.3	4.3	30(27)	[23]
	22.3	10.4	3.5	34(25)	[26]
	23.0	7.3	3.2	22(24)	[27]
	23.4	6.3	3.2	24(26)	[28]
	23.0	7.5	4	35(27)	[13]
	23	8	4.2	24(27)	[28]

*The upper integration limits (in MeV) are given in parentheses.
 **The data given here are the values corrected in [13] for anisotropic distribution.




FIG. 1. C¹² nuclear absorption cross section.

¹ B. Ziegler, Z. Physik 152, 566 (1958); Nuclear Phys. 17, 238 (1960).
² Burgov, Danilyan, Dolbilkin, Lazareva, and Nikolaev, JETP 37, 1811 (1959), Soviet Phys. JETP 10, 1278 (1960).
⁴ E. E. Carroll and W. E. Stephens, Phys. Rev. 118, 1256 (1960).
⁵ I. M. Thorson and L. Katz, Proc. Phys. Soc. (London) 77, 166 (1961).
⁷ Cohen, Mann, Patton, Reibel, Stephens, and Winhold, Phys. Rev. 104, 108 (1956).
¹⁰ Gemmel, Morton, and Titterton, Nuclear Phys. 10, 33 (1959).
¹¹ Gove, Litherland, and Batchelor, Nuclear Phys. 26, 480 (1961).
¹² V. J. Vanhuyse and W. C. Barber, Nuclear Phys. 26, 233 (1961).
¹³ W. R. Dodge and W. C. Barber, Phys. Rev. 127, 1746 (1962).
¹⁴ K. O. Hermann and J. A. Scheer, Z. Physik 170, 162 (1962).
¹⁵ Fuchs, Haeg, Lindenberger and Meyer-Berkhout, Z. Naturforsch. 17a, 439 (1962).
²⁰ Miller, Schuhl, Tamas, and Tzara, Phys. Lett. 2, 76 (1962).
²¹ F. Ajzenberg-Selove and T. Lauritsen, Nuclear Phys. 11, 1 (1959).
²² Bair, Kington, and Willard, Phys. Rev. 100, 21 (1955).
²³ J. Halpern and A. K. Mann, Phys. Rev. 33, 370 (1951).
²⁴ S. Penner and J. E. Leiss, Phys. Rev. 114, 110 (1959).
²⁵ Barber, George, and Reagan, Phys. Rev. 98, 70 (1955).
²⁶ B. C. Cook, Phys. Rev. 106, 300 (1957).
²⁷ Roalsvig, Gupta, and Haslam, Can. J. Phys. 39, 643 (1961).
²⁸ J. H. Carver and K. H. Lokan, Australian J. Phys. 10, 312 (1957).

Ref.

S. Costa, F. Ferrero, S. Ferroni, B. Minetti, C. Molino,
 R. Malvano
 Phys. Letters 6, 226 (1963)

Elem. Sym. A

Z

C

12

6

Method 100 MeV Synchrotron; 4π neutron detector; calculated integrated cross sections - fitted with polynomial of degree π

Ref. No.

63 Co 3

EGF

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
(γ, xn)						$\bar{v}_b = \int \frac{\sigma(E)}{E} dE$ <p>gets $\langle \bar{v}_p \cdot \bar{v}_n - \bar{v}_n \cdot \bar{v}_n \rangle$</p> $= (R_c^2 - R_p^2 - \frac{3}{\pi} \frac{trc}{e^2} \bar{v}_b \frac{A-1}{A^2}) \times \frac{2}{A-2}$ <p>See "Boron" for plots of this and $\int \sigma dE/60 \text{ NZ/A}$</p>



Figure 1: Photonuclear cross sections for several light elements versus γ -ray energy.

REF.

E. Finckh, R. Kosiek, K.H. Lindenberger, K. Maier,
 W. Meyer-Berkhout, M. Schechter, J. Zimmerer
 Z. Physik 174, 357 (1963)

ELEM. SYM.

A

Z

C

12

6

METHOD

Betatron; proton spectrum, cross section; CsI spectrometer

REF. NO.

63 Fi 4

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	SPC	19-30	C	31 (30.5)	SCI-D	3-14	90

$$\int_{18.5}^{30.5} \sigma(\gamma, p) dE = 55 \pm 8 \text{ MeV-mb}$$

assuming $W(\theta) = 1 + (1.2_{-0.5}^{+1}) \sin^2 \theta$ and ground state transition only.




Fig. 2 a, b. Zwei Energieverteilungen der Photo-neutronen aus C^{13} bei einer Endenergie von 0,5 MeV. Die Histogramme stellen die auf Energieverlust korrigierten Meßwerte dar, die ausserdem durchgezogen nach einem Mittelungsverfahren (gewogen) Nahere Erläuterungen im Text und in Tabelle 1




Fig. 3. Differentieller Wirkungsquerschnitt für Photoneutronenemission aus C^{13} unter 90° zum γ -Strahl unter der Annahme, daß nur Grundzustandsübergänge vorliegen (—). Differentieller Wirkungsquerschnitt für Photoneutronenemission aus C^{13} unter 90° nach FUCHS und HAAG* (---)

* FINK, F.W.K., and K.H. LOKAN: Phys. Rev. Letters 8, 321 (1962).

* FUCHS, H., u. D. HAAG: Z. Physik 171, 403 (1963).

METHOD Betatron; neutron spectrum, cross section; stilbene scintillator; ion chamber						REF. NO. 63 Fu 1		NVB
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
G, XN	ABX	20-29	C	31 (30.5)	SCI-D	2-11	90	






Fig. 3 a u. b. Spektren der Photoneutronen aus Kohlenstoff, unter 90° bei E₀ = (30,5 ± 1) MeV. a und b Ergebnisse zweier unabhängiger Messungen bei verschiedener Einstellung der Impulsformdiskrimination

Fig. 4 a u. b. Wirkungsquerschnitte für die Emission von Photopartikeln aus C¹². a C¹²(γ, p) B¹¹ Wirkungsquerschnitt von DODGE und BARBER²³, berechnet aus den Spektren der Protonen aus C¹²(α, p) B¹¹ unter der Annahme, daß 100% Übergänge zum Grundzustand von B¹¹ vorliegen. b C¹²(γ, n) C¹¹: Histogramm: Ergebnis dieser Arbeit für den Wirkungsquerschnitt von C¹²(γ, n) C¹¹ berechnet aus dem Neutronenspektrum in Fig. 3 b unter der Annahme, daß 100% Übergänge zum Grundzustand von C¹¹ vorliegen. Die rechte Skala gibt dσ/dΩ (90°) an und weist in ihren Absolutangaben einen Fehler von 20% auf, während die linke Skala σ angibt, berechnet aus dσ/dΩ (90°) unter der Annahme, daß die Winkelverteilung die Form A + B · sin² θ habe mit B/A = 1,5 ± 1 für alle Neutronenenergien. Der Fehler in den Absolutwerten ist dort etwa 25%. Die Fehlerbalken geben den statistischen Fehler an. Ausgezogene Kurve: σ für C¹²(γ, n) C¹¹ von BARBER et al.²³

²³ BARBER, W. C., W. D. GEORGE and D. D. REAGAN: Phys. Rev. 98, 73 (1955).

Elem. Sym.	A	Z
C	12	6
Ref. No.		JHH
63 Go 4		

Method
 Linac; electron scattering; magnetic spectrometer

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C ¹² (e,e')	41.5					Giant resonance ∫σ(γ abs.)dE inferred to be ~ 50 MeV-mb.




Fig. 3. Spectrum of inelastically scattered electrons from a styrofoam target (C₆H₈). The solid line is a visual fit to the data. Seen are the carbon elastic peak, the proton peak, the 15.1 MeV excited level in carbon and the giant resonance.




Fig. 4. The giant resonance in C¹² obtained from Fig. 3 after subtraction of the estimated radiation tail.

TABLE I
 Summary of experimental and theoretical inelastic electron scattering cross sections at 140

Element	Target	Density (g/cm³)	Thickness (mg/cm²)	Thickness (radiation length)	($\frac{d\sigma}{d\Omega}$) _{ex} measured	($\frac{d\sigma}{d\Omega}$) _{ex} calculated	($\frac{d\sigma}{d\Omega}$) _{ex} (calc.) maximum	($\frac{d\sigma}{d\Omega}$) _{ex} (calc.) average	($\frac{d\sigma}{d\Omega}$) _{ex} (measured) minimum
C ¹²	C ₆ H ₈ (Styrofoam)	0.033	466	0.0103	3.8 ± 0.8	16.7	5.6	4.4	3.6
Al ²⁷	Stack of 0.013 cm foils	2.70	380	0.016	4 ± 2	25	12.5	6.2	4.1
Fe	Gas	0.016	460	0.041	8 ± 4	16	9	4.5	1

Elem. Sym.	A	Z
C	12	6

Phys.Rev. 129, 1362 (1963)

Method 335 MeV Synch-scintillation counter telescope	Ref. No. 63Kil	B6
---------------------------------------------------------	-------------------	----

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(γ, p)	E _{γmax} = 335					<p>Data corrected for: nuclear absorption, γ attenuation, multiple scattering in target, multiple scattering in counter, pion contamination.</p> <p>Photoproton cross section found proportional to NZ/A (NZ probability of having np pair) over angular and energy region examined. (Z=3,4,5,6)</p> <p>Differential cross section data given.</p> <p>T_p = proton energy (Fig.1)</p>




Fig. 5. Differential cross sections for photoproton production from carbon.

Elem. Sym.	A	Z
C	12	6
Ref. No.		63Le1
		6-5

Phys.Rev.Letters 10, 493 (1963)

Method
60MeV linac - coincidence counter telescope

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
(e, e')	E _{in} = 55 E _{scatt} = 0 - 60	15.1 (18) (23)				Form factor versus q (momentum transfer) for giant dipole resonance is plotted. Fig. 1 is from erratum, Phys.Rev.Letters 11, 106 (1963).

ELECTROMAGNETIC STRUCTURE OF THE GIANT DIPOLE RESONANCE. F. H. Lewis, Jr., J. D. Walecka, J. Goldemberg, and W. C. Barber (Phys. Rev. Letters 10, 493 (1963)).

We found a slight numerical mistake in our calculations. The correct numerical results are as indicated in Fig. 1 (revised). All of the statements and arguments given in the paper are correct.




FIG. 2. Spectrum of 55-MeV electrons scattered at 130° by C¹²; the radiation tail was not unfolded in this curve.




FIG. 1 (revised). Form factors vs q for the giant dipole resonance in C¹². The dashed curve is discussed in reference 4. We have also indicated the value corresponding to the Thomas-Reiche-Kuhn sum rule.

Elem. Sym.	A	Z
C	12	6

Method 24 MeV betatron; silicon-diode charged particle detector

Ref. No. 63 Mu 1
 JHH

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
----------	-----------------	-------	----------	------------------	--------	-------

(γ, p)

Bremss.
 23.5

Small peak observed at 21.05 MeV ($E_p = 4.7$ MeV); also seen by Dodge and Barber [Phys. Rev. 127, 1746 (1962), our reference 62Dol].

Absorber used totall if other than protons are being detected; concludes there are no α 's seen; cross section assumes no excited state transitions.

Detector at 118°




FIG. 5. Pulse-height distributions from carbon. Curve A is the smoothed value of the data taken with no absorber between target and detector. Curve B is the smoothed value of the data taken with 7 mg cm^{-2} of aluminum between target and detector. The absorber data has been corrected at each point for the energy lost in the absorber by a proton of that energy.




FIG. 6. The differential cross section for the $C^{12}(\gamma, p)B^{11}$ reaction from 19 MeV to 22 MeV assuming 100% ground-state transitions.

Elem. Sym.	A	Z
C	12	6
Ref. No.		BG
63Rel		

Method
26 MeV Linac - NaI(Tl)

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
$B^{11}(p, \gamma)C^{12}$	$E_p = 15-25$					
(p, γ_0)				0.65 ± 0.16	4^0	Using data from this and other experiments - if angular distribution of Gove (Nuclear Phys. 26, 480 (1961) used ground state integrated sum reduced to 0.50 ± 0.12 MeV-mb while excited state sum essentially is unchanged.
(p, γ_1)				0.71 ± 0.24		
						Using reciprocity relationship, including spin statistical weight factors, the following is obtained:
$C^{12}(\gamma, p_0)B^{11}$				72 ± 17 MeV-mb		C^{12} in ground state
				13 ± 3	2^+	C^{12} in first excited state (Jπ not determined in this experiment) Isotropy in cross section assumed above. Correction for anisotropy would give 55 ± 13 for ground state transition.
$B^{11}(p, \gamma_0)C^{12}$		$E_\gamma = 34.4 \pm 0.5$		0.038 ± 0.01		
$C^{12}(\gamma, p_0)B^{11}$				4.1 ± 1		using reciprocity.




Fig. 4. Excitation function for the reaction $B^{11}(p, \gamma)C^{12}$. The solid curve indicates the γ_0 transitions, the dashed curve indicates the γ_1 transitions, the thin target no. 1 is indicated by \odot , the thick target is indicated by \square , and the thin target no. 2 is indicated by \dots




Fig. 5. Summary of data for the reaction $B^{11}(p, \gamma)C^{12}$.

METHOD			Li ⁷ (p,γ) source; photon scattering; NaI		REF. NO.	63 Sc 3	NVE
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	15 (15.1)	D	15, 18 (14.7, 17.6)	NAI-D		DST

G-WDTH

Tabelle. Parameter des 15,1 MeV I⁺; 1 Niveaus von C¹²

Autor	5	6	8	10	11	Vorliegende Messung
$\sigma_{\text{tot}}^{\text{max}}$	22.2 = 2.2	29.7 = 1.1		29 = +		28.9 = 3.3 barn
$\partial\Gamma$	< 1	0.62 = 0.10				> 0.55 < 1.20
Γ	79 = 16	64.5 = 10.4		60 = 8		44.7 = 10.3 eV
Γ_7^0	34.5 = 9.3	39.2 = 9.7	59.3 = 7.1	54 = 6	40 - 8 - 6	40.2 = 5.2 eV
$\Gamma_7^{4,4}$	≤ 5.5	3.2 = 2.5		6.5 = 3.0		eV
Γ_2	21.6 = 11.2	2.1 = 3.2	≤ 15 ¹¹			eV
$\int \sigma_S dE$	1.90 = 0.27	2.33 = 0.19	2.20 = 0.31	2.45 = 0.50	2.0 ± 0.4 ¹⁰ - 0.3	1.82 = 0.12 MeV·m

a) Übernommen von 9.

b) Der angegebene Wert ist der von ¹¹ gemessene integrierte Absorptionsquerschnitt. Zum Vergleich multipliziert man mit $\Gamma_7^0 \Gamma$.

⁹ MILLER, G.L., R.E. PINLEY and R.E. SEGEL: Proc. Roy. Soc. (London) A 259, 275 (1961).

¹⁰ BUSSIERE DE NERCY, A.: Ann. phys. 6, 1379 (1961).

¹¹ BARBER, W.C., F. BERTHOLD, G. FAICKE and F.E. GUDDEN: Phys. Rev. 120, 2081 (1960).

Method emulsions

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
C ¹² (γ, 3α)	Bremss. 70					<p>Q-value distribution in Figure 1 is based on 679 3-pronged stars.</p> <p>Shows that previous evidence for a transition to state at 4 MeV in Be⁸ by C¹²(γ, α)Be^{8*} is spurious. Result from resonant absorption at 15.11 MeV</p> <p>In Figure 1, shaded area contains all stars for which at least one of the possible Q-values for Be⁸ fell between 2.4 and 3.6 MeV (transition to 2.9 MeV state).</p> <p>In Table I ⊕ gives the spurious Q values in Be⁸, E_γ the corresponding photon absorption energy and E_σ the energies at which resonances are found in the cross section curve for the (γ, 3α) reaction.</p>

Table 1.
Spurious Q-values and corresponding photon absorption resonances.

Q (MeV)	E _γ (MeV)	E _σ (MeV)	E(., n)
3.6	14.1	-	
4.3	15.0	-	
5.3	16.3	15.6	
(6.2)	(17.5)	17.9	
6.9	18.5	-	
7.7	19.5	19.3	
(9.3)	(20.3)	21.0	
(9.5)	(22.3)	21.1	22.2
10.5	23.3	22.5	23.1




Fig. 1. Calculated Q-values.

Elem. Sym.	A	Z
C	12	6
Ref. No.		JHH
63 Wa 2		

Method Synchrotron; proton angular distribution; CsI

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
----------	---------	----------------	---	------	----	-------

C¹²(γ,p)

Bremss.
65

Angular distribution function is:
a+b sin²θ + A cosθ + B sin²θ cosθ




Fig. 2. The angular distributions for the proton energy bins 8-9 and 12-13 MeV. Vertical flags represent counting statistics and horizontal flags angular resolution. The curves are the least-squares fits of eq. (1).




Fig. 3. The angular distributions for the proton energy bins 16-17 and 20-21 MeV. The curves are the least-squares fits of eq. (1).




Fig. 4. The coefficients from the least-squares fitting of eq. (1) plotted as a function of proton energy. The flags represent the standard errors of the coefficients. The curves were drawn by eye to illustrate the trend of the coefficients.




Fig. 5. The b/a ratio as a function of proton energy. The crosses are from Vanhuyse and Barber's results.




Fig. 6. The energy spectrum at 75°. The number of counts are normalized to 13.5 mg/cm² target thickness and to the ion chamber monitor response.

1) V. J. Vanhuyse and W. C. Barber, Nuclear Physics 26 (1961) 233

Table 2
Coefficients of least-squares fit of $a + b \sin^2 \theta + A \cos \theta + B \sin^2 \theta \cos \theta$

Energy bin (MeV)	a (%)	b (%)	A (%)	B (%)
6.75-8	1.45 ± 0.02	0.52 ± 0.18	-0.17 ± 0.56	0.54 ± 0.71
8-9	0.49 ± 0.042	0.51 ± 0.32	0.21 ± 0.07	1.94 ± 0.08
9-10	0.28 ± 0.05	0.47 ± 0.32	1.61 ± 0.99	0.24 ± 0.25
10-11.2	0.29 ± 0.04	0.64 ± 0.41	0.21 ± 0.25	1.24 ± 0.21
11-12	0.52 ± 0.09	0.25 ± 0.09	0.20 ± 0.14	0.78 ± 0.11
12-13	0.80 ± 0.02	0.24 ± 0.19	0.67 ± 0.18	0.23 ± 0.15
13-14	0.76 ± 0.01	0.33 ± 0.27	1.94 ± 0.14	0.21 ± 0.06
14-15	0.90 ± 0.05	0.16 ± 0.29	0.74 ± 0.16	1.00 ± 0.08
15-16	0.44 ± 0.04	0.38 ± 0.27	0.89 ± 0.10	2.24 ± 0.27
16-17	1.12 ± 0.05	0.01 ± 0.10	0.74 ± 0.10	2.99 ± 0.14
17-18 (21.3 mu cm ⁻²)	0.74 ± 0.05	0.35 ± 0.24	-0.52 ± 0.09	1.01 ± 0.12
17-18 (19.1 mu cm ⁻²)	0.40 ± 0.01	0.55 ± 0.09	1.06 ± 0.12	1.84 ± 0.10
18-19	0.27 ± 0.02	0.71 ± 0.22	0.14 ± 0.28	1.71 ± 0.17
19-20	0.18 ± 0.05	0.77 ± 0.31	0.08 ± 0.40	0.80 ± 0.17
20-22	0.16 ± 0.04	0.72 ± 0.25	0.07 ± 0.37	0.69 ± 0.15

METHOD				REF. NO.			
$B^{11}(p,\gamma_0)C^{12}$ Van de Graaff				64 A1 2		JOC	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	19-29	D	4-14	NAI-D		DST




Fig. 6. The 90° yield curve in 50 keV steps for γ_0 .




Fig. 10. A sampling of angular distributions of γ_0 and of γ_1 . Angular distributions were actually taken about every 50 keV.

METHOD ${}^{11}\text{B}(p,\gamma_0){}^{12}\text{C}$ Van de Graaff [Page 2 of 2] REF. NO. 64 A1 2 JOC

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 11. The coefficients a_n obtained by expanding the angular distributions of γ_0 into a series of Legendre polynomials $W(\theta) = A_0(1 - \sum a_n P_n(\cos \theta))$. The finite solid angle subtended by the detector attenuates a_n by about 3%.




Fig. 13. Total cross section for the ${}^{11}\text{B}(p,\gamma_0){}^{12}\text{C}$ reaction as a function of energy. The data for $1 \text{ MeV} \leq E_p \leq 4 \text{ MeV}$, as well as the absolute scale, are from ref. ¹³. (R.G.Allas, S.S.Hanna and R.E.Segel, to be published.)

METHOD $Al^{27}(p,\gamma_0)Si^{28}$; tandem

REF. NO.

64 Al 3

JOC

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P.G	NOX	20 - 30	D	4 - 14	NAI-D		DST

Table I. Summary of experimental results on the (p,γ) giant resonance (references 9-11). The angular distributions that are quoted generally characterize the data to within 15% throughout the giant-resonance region.

Nucleus	Type of observation	Energy interval and range	Result
C^{12}	90° yield curve	50-keV steps 4.0-14 MeV	Broad ($\Gamma \approx 1$ MeV) overlapping levels; no correlation between γ_0 and γ_1 .
	Angular distribution ^a	50-keV steps 4.0-14 MeV	$W(\theta)_{\gamma_0} = 1 + 0.15P_1 - 0.6P_2$ $W(\theta)_{\gamma_1} = 1 + 0.15P_1$
Ne^{20}	90° yield curve	30-keV steps 4.3-9.1 MeV	Broad ($\Gamma \approx 400$ keV) levels usually well isolated, γ_0 and γ_1 well correlated.
	Angular distribution	100-keV steps 4.0-10.5 MeV	$W(\theta)_{\gamma_0} = 1 + 0.05P_1 - 0.7P_2$ $W(\theta)_{\gamma_1} = 1 + 0.05P_1 + 0.2P_2$
Si^{28}	90° yield curve	15-keV steps 4.0-12.5 MeV	Narrow ($\Gamma \approx 50$ keV) Ericson fluctuations, superimposed on intermediate structure; no correlation between γ_0 and γ_1 .
	Angular distribution	15-keV steps 4.0-4.32 MeV	$W(\theta)_{\gamma_0} = 1 + 0.07P_1$
		6.0-6.62 MeV	$W(\theta)_{\gamma_1} = 1 + 0.1P_1 - 0.45P_2 - 0.1P_3$
		8.0-8.54 MeV	
		10.0-10.28 MeV 11.54-11.53 MeV	

^aThe coefficients of P_1 are average values. They actually increase by about 0.03/MeV over the giant resonance.

METHOD

REF. NO.

64 A1 5

JOC


REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	NØX	THR-34	C	34	THR-I	6-	DST

TABLE I

Summary of present experimental data at 34 MeV bremsstrahlung

Element		$-\frac{a_2}{a_0}$	$\frac{a_1}{a_0}$
^4Be		0.43 ± 0.02	0.05 ± 0.01
^{12}C		0.61 ± 0.04	0.09 ± 0.02
^{13}Al		0.39 ± 0.03	0.05 ± 0.01
^{22}Ti		0.34 ± 0.02	0.06 ± 0.01
^{24}Cr	34 MeV	0.33 ± 0.02	0.02 ± 0.01
	22 MeV	0.13 ± 0.07	-0.02 ± 0.01
^{29}Cu		0.36 ± 0.02	0.10 ± 0.01
^{50}Sn		0.38 ± 0.02	0.11 ± 0.01
^{54}Ba		0.39 ± 0.03	0.11 ± 0.02
^{73}Ta	Before installation of iron shielding	0.26 ± 0.04	0.13 ± 0.02
	After installation of iron shielding	0.27 ± 0.02	0.12 ± 0.01
^{82}Pb	target diameter 3.0 cm	0.39 ± 0.03	0.15 ± 0.02
	target diameter 1.5 cm	0.40 ± 0.03	0.19 ± 0.02
^{83}Bi		0.42 ± 0.03	0.17 ± 0.02

$$Y = a_0 + a_1 \cos \theta + a_2 \cos^2 \theta$$



REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	RLX	THR - 150	D	550	MAG-D		51




FIG. 2. Electron-proton coincidence counting rate per 10^{11} equivalent quanta at 550 MeV as a function of the incident energy. The dashed lines indicate the contributions of the various shells and the background as explained in the text.

REF.

ELEM. SYM. A Z

W. Bertozzi, P. Demos, F. Hanser, S. Kowalski, C. Sargent,
 W. Turchinets, R. Fullwood and J. Russell
 Proc. Paris Conf. 1026 (1964)

C

12

6

METHOD

Polarization Data

REF. NO.

64 Be 8

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$^{12}\text{C}(n,\alpha)^9\text{B}$	NOX	THR-32	C	32	SCI-D		DST

$P_n = -.05 \pm .05$, a very small number. Neutron at 90° and 45°.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		JOC
64 B1 1		

METHOD			$\text{Be}^9(\text{He}^3, \gamma)\text{C}^{12}$		REF. NO.		JOC
					64 B1 1		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
HE3,G	ABX	28	D	2-5 (1.5-4.5)	NAI-D		90




Fig. 2(a). Spectrum of high-energy γ rays resulting from bombardment of Be^9 with a He^3 beam of energy 4 MeV. A plastic anticoincidence shield was employed to reduce cosmic ray background. γ rays to the ground state and first excited state of C^{12} can be clearly distinguished. The 17.6 MeV line is also shown, and is seen to be at least an order of magnitude more intense than the full-energy γ ray to the C^{12} ground state.

Fig. 2(b). Excitation function at 90° for the γ rays to the ground state and first-excited state of C^{12} . The ordinate represents the cross section measured in μb , assuming an isotropic angular distribution. The He^3 energy is the laboratory bombarding energy.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E, E/$	SPC	0-20	D	54	SCI-D		141




Fig. 1. — Electron scattering from a graphite foil ($d_{\text{eff}} = 0.30 \text{ g/cm}^2$).
 C^{12} level energies in MeV. For 15.1 MeV level, $q_f = 0.44 \text{ fm}^{-1}$.

REF. K.W. Chen, J.R. Dunning, Jr., J.R. Rees, W. Shlaer, J.K. Walker and R. Wilson
Phys. Rev. 135, 4B B1030 (1964)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		JOC
64 Ch 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,P	ABX		D	4 GEV	MAG-D	110-450	DST

TABLE I. Cross sections for production of protons by electrons.

Electron target	Energy angle (deg)	4 BeV proton energy (MeV)	$\frac{d^2\sigma}{d\Omega dE}$ $10^{-33} \text{ cm}^2/\text{sr MeV}$
H	59.8	374	3.8*
Li _a	59.8	448	0.082
Li _b	59.8	368	0.175
Li _c	59.8	332	0.280
C	59.8	374	0.425
Al	59.8	374	1.19
H	63.1	291	7.5*
Li _a	63.1	355	0.146
Li _b	63.1	319	0.204
Li _c	63.1	290	0.315
C	63.1	291	1.01
Al	63.1	291	2.42
H	67.1	208	16*
Li _a	67.1	226	0.6
Li _b	67.1	206	0.92
C	67.1	209	2.57
Al	67.1	209	6.4
H	72.1	124	46*
Li _a	72.1	166	1.20
Li _b	72.1	144	1.53
Li _c	72.1	124	2.46
Li _d	72.1	119	2.80
Li _e	72.1	109	2.90
C	72.1	123	6.6
Al	72.1	124	16.9
Cu	72.1	124	42.3
H	44.8	291	2.9*
Li ^o	44.8	337	0.16
Li ^o	44.8	293	0.29
C	44.8	291	0.76
Al	44.8	291	1.53
H	52.3	208	8.5*
Li ^o	52.3	200	0.91
C	52.3	208	2.13
Al	52.3	208	4.95
H	61.1	124	25*
C	61.1	145	3.95
C	61.1	124	5.75
C	61.1	115	6.43
Al	61.1	145	10.7
Al	61.1	124	16.0

* $d\sigma/d\Omega$ in mb/sr; inside a 5.5% momentum interval.




FIG. 4. Cross section, divided by A, as a function of A, for producing protons of 124 MeV from 4-BeV electrons at 72.1°.

METHOD						REF. NO.	
Linac						64 Cr 1	
						NVB	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,e/	ABX	4-10	D	250	MAG-D		DST

New method of obtaining widths.

WIDTHS

TABLE I. C¹² cross sections for elastic and inelastic scattering of electrons.

E ₀ MeV	θ	Elastic scattering (in 10 ⁻²¹ cm ² /sr)			4.43-MeV level (in 10 ⁻²¹ cm ² /sr)			7.66-MeV level (in 10 ⁻²¹ cm ² /sr)			9.64-MeV level (in 10 ⁻²¹ cm ² /sr)		
		dσ/dΩ	Area	Total	dσ/dΩ	Area	Total	dσ/dΩ	Area	Total	dσ/dΩ	Area	Total
187	84.0	3.76	0.05	0.12	1.86	0.03	0.06	2.96	0.15	0.17	5.83	0.58	0.61
	91.8	1.17	0.02	0.04	1.08	0.02	0.04	1.71	0.08	0.10	3.94	0.39	0.41
250	40.0	417.	5.	13.	26.6	1.0	1.2	63.1	3.9	4.3	59.3	7.4	7.6
	45.0	167.	2.	5.	16.4	0.4	0.7	42.2	1.7	2.1	47.4	2.8	3.1
	50.0	66.8	0.6	2.1	11.0	0.2	0.4	28.2	0.9	1.2	37.0	1.1	1.6
	55.0	27.0	0.3	0.9	7.01	0.14	0.25	15.8	0.8	0.9	26.2	0.9	1.2
	66.0	10.3	0.1	0.3	4.79	0.07	0.16	8.08	0.40	0.47	15.2	0.5	0.7
	65.0	4.04	0.05	0.13	3.26	0.05	0.11	4.65	0.23	0.27	11.0	0.4	0.5
	70.0	1.44	0.02	0.05	1.94	0.02	0.06	2.12	0.05	0.08	8.24	0.12	0.27
	75.0	0.420	0.008	0.016	1.28	0.01	0.04	1.30	0.04	0.06	5.57	0.08	0.18
	80.0	0.124	0.003	0.005	0.702	0.006	0.022	0.574	0.021	0.027	3.80	0.04	0.12
	85.0	0.0314	0.0010	0.0014	0.451	0.004	0.014	0.311	0.010	0.013	2.56	0.02	0.08
	90.0	0.0057	0.0003	0.0003	0.242	0.002	0.007	0.095	0.005	0.006	1.46	0.01	0.05
300	49.2	17.4	0.1	0.5	7.69	0.09	0.25	10.8	2.1	2.2	32.9	6.1	6.2
	54.8	4.30	0.04	0.14	4.12	0.05	0.13	4.90	0.49	0.51	19.5	2.0	2.0

TABLE II. Measured transition widths for excited states in C¹².

Level energy in MeV	λ	A _λ	Γ in eV
4.43	2	1.638±0.066	(11.2±1.2)×10 ⁻⁴
7.66	0	0.936±0.048	(6.5±0.7)×10 ⁻⁴
9.64	3	1.014±0.054	(3.6±0.4)×10 ⁻⁴

METHOD Linac		SOURCE		DETECTOR		ANGLE	
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE	ANGLE
E, E'	ABX	10-32	D	40-70	MAG-D		DST




FIG. 3. Cross section for inelastic scattering from carbon derived from the spectrum shown in Fig. 2.

130°




FIG. 11. The square of the form factor for the main part of the C¹² giant resonance plotted as a function of momentum transfer. The experimental point at 23 MeV is from work with photons. The other three experimental points are from 130° electron scattering experiments. The curves are calculated on the basis of different theories as explained in the text.

¹⁰ F. H. Lewis, Jr. (private communication).

TABLE II. Cross sections for excitation of the 18.1- and 19.5-MeV states observed in the Darmstadt carbon scattering experiments.

E ₀ (MeV)	deg	(dσ/dΩ) × 10 ²² (19.5-MeV state)	(dσ/dΩ) × 10 ²² (18.1-MeV state)	(dσ/dΩ) × 10 ²² (Both states)	⟨T ¹¹ ⟩²(q) ² (19.5-MeV state only)	⟨T ¹¹ ⟩²(q) ² (Both states)
56	152	1.3 ± 0.13	1.0 ± 0.3 ¹⁰	2.3 ± 0.3 ¹⁰	0.00053 ± 0.00005	0.00091 ± 0.00005 ¹⁰
50	152	0.88 ± 0.18	1.0 ± 0.2	1.9 ± 0.3	0.00026 ± 0.00005	0.00057 ± 0.00009
40	152	0.55 ± 0.3	0.5 ± 0.4	1.1 ± 0.5	0.00012 ± 0.00007	0.00023 ± 0.00011
55	128	1.4 ± 0.3	0.5 ± 0.4	1.9 ± 0.5	0.00034 ± 0.00007	0.00046 ± 0.00012

¹¹ F. Gudden, G. Fricke, H. G. Clerc, and P. Brix (unpublished).

FME




FIG. 12. Square of the form factor for the lowest 1⁻ state in C¹² plotted as a function of the momentum transfer. The two curves are from the shell-model calculation for two different assumptions about the two-particle interaction. The triangles are from the Stanford 180° scattering experiments. The circles are from the Darmstadt 152° scattering experiments. The solid circles are for the single peak at 19.5 MeV whereas the open circles include the broad 18-MeV peak as well. The square is derived by detailed balancing from B¹¹(p, γ) experiments which indicate a 1⁻ level at 17.2 MeV.

METHOD					REF. NO.		
Linac					64 Go 3		
					NVB		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	15	D	40-70	MAG-D		180

FMF

TABLE I. Data on M1 transitions.

	q MeV/c	Cross section*	Reference	$ \langle J_f T_1^{max}(q) J_i \rangle ^2 (1/q^2)$ (10^{-3})	$ \langle J_f T_1^{max}(q) J_i \rangle ^2$ [$10^{-14}(\text{MeV}/c)^{-2}$]
15.1-MeV level in C^{12}					
Photons	15.1	(2.05 ± 0.27)	8	0.044	(0.19 ± 0.025)
Electrons	68	(2.0 ± 0.3)	this work	0.55	(0.12 ± 0.015)
Electrons (160°)	68	$(2.6 \pm 0.45^{*1})$	5	0.69	(0.15 ± 0.02)
Electrons	93	(2.0 ± 0.3)	this work	0.87	(0.10 ± 0.015)
Electrons	125	(1.5 ± 0.25)	this work	1.10	(0.07 ± 0.012)
3.56-MeV level in Li^6					
Photons	3.56	$(0.92 \pm 0.11^{*2})$	b	0.014	$(1.10 \pm 0.13^{*2})$
Electrons	76	(3.0 ± 0.45)	this work	2.0	(0.38 ± 0.057)
Electrons	106	(1.55 ± 0.23)	this work	2.1	(0.19 ± 0.028)
Electrons	136	(0.90 ± 0.14)	this work	2.0	(0.11 ± 0.017)
11.6-MeV level in Si^{28}					
Photons	11.6	(9.8 ± 2.6)	c	0.016	(1.2 ± 0.32)
Electrons	71.5	(3.5 ± 1.4)	d	0.92	(0.18 ± 0.07)
Electrons	88.4	(3.0 ± 0.75)	this work	1.2	(0.15 ± 0.038)
Electrons	129.4	(1.8 ± 0.45)	this work	1.5	(0.09 ± 0.022)
11-MeV level in Mg^{24}					
Photons	11	(13.1 ± 3.8)	c	0.02	(1.7 ± 0.50)
Electrons	83	(4.4 ± 0.88)	6	1.5	(0.22 ± 0.044)
Electrons	97	(3.5 ± 0.70)	this work	1.6	(0.1 ± 0.034)
Electrons	129	(1.8 ± 0.36)	this work	1.3	(0.081 ± 0.016)

* In units of 10^{-17} MeV-cm² for photons and 10^{-28} cm²/sr for electrons.
 * L. Cohen and R. A. Tobin, Nucl. Phys. 14, 243 (1960).

* A. B. de Nercy, Ann. Phys. (Paris) 6, 1379 (1961).

* R. D. Edge and G. A. Peterson, Phys. Rev. 128, 2750 (1962).




Fig. 2. The experimental values of the quantity $(1/q^2) \times | \langle J_f | T_1^{max}(q) | J_i \rangle |^2$ as a function of the momentum transfer. The theoretical curve was normalized as explained in the text.




FIG. 1. Results obtained for the excitation of the C^{12} 15.1-MeV level in inelastic electron scattering at 180° for different incident electron energies.

METHOD

Betatron; r-chamber

REF. NO.

64 Gr 1 NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.3A	ABY	7-24	C	17-24	EMU-I		4PI

Yield = 5315 ± 30 $\frac{\text{events}}{\text{mole-r}}$ at 24 MeV.Yield = 773 ± 67 $\frac{\text{events}}{\text{mole-r}}$ at 17 MeV.TABLE IV
Results of loading and searching of each plate

Plate No.:	1	2	3	4	5	6	7
Moles of O ¹⁸ per cm ³ of emulsion (dry)	0.0167	0.0355	0.0577	0.0165	0.0238	0.0355	0.0453
Moles of C ¹² per cm ³ of emulsion (dry)	0.0233	0.0373	0.0551	0.023	0.0257	0.0303	0.0466
Number of α events per cm ² (dry) per r	—	—	—	34.3	43.7	70.3	79.3
Number of β events per cm ² (dry) per r	—	—	—	7.0	7.0	7.0	8.0
Number of alpha tracks per cm ² (dry) per r	17.7	20.9	42.1	195	204	205	313
Number of alpha tracks per cm ² (dry) per r	1.3	2.3	3.0	15	15	14	16
Number of alpha tracks per cm ² (dry) per r	124	195	279	535	616	721	838
Number of alpha tracks per cm ² (dry) per r	4	7	9	25	26	25	25




FIG. 6. The variation of the number of three-pronged stars with carbon content for 24-MeV and 17-MeV bremsstrahlung.

REF.

F. Gudden
Phys. Letters 10, 313 (1964)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
Linac				64 Gu 1		JOC	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.E./	FMF	15 (15.1)	D		MAG-D		DST

$$\Gamma_{\gamma 0} = 34.4 \text{ eV } (\pm 10\%)$$

0 0.2-0.5. WIDTH




Fig. 1. Reduced transition probability versus momentum transfer squared. The solid curve has the parameters $B_T(q_T = 0) = 0.875$ and $b = 1.9$ fm, the dashed curve has $B_T(q_T = 0) = 0.79$ and $b = 1.64$ fm. GolB 64: ref. 4, DudT 61: ref. 3, SchS 63: ref. 6.

REF.

S. S. Hanna, R. E. Segel and R. G. Allas
Proc. Paris Conference 1022 (1964)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				64 Ha 3		JDM	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	17-20	D	1-4	NAI-I		DST

$$W(\theta) = A_0 \left[1 + \sum_1^4 a_n P_n(\cos\theta) \right]$$




FIG. 1. — Total yield ($4\pi A_0$) and coefficients in the expansion $1 + \sum_1^4 a_n P_n(\theta)$ for the reaction $B^{11}(p, \gamma) C^{12}$, $1 \text{ MeV} < E_p < 4 \text{ MeV}$.

M. Kihara, K. Baba, K. Miyake, T. Nakamura, T. Yamaki, S. Yasumi
and Y. Yoshimura
J. Phys. Soc. Japan 19, 2236 (1964)

C

12

6

METHOD

REF. NO.

Time-of-flight plus magnetic analysis

64 Ki 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,D	ABY	80-800	C	400-800	TOF-D	45-70	57




Fig. 1. Yield curve of deuterons of the energy of 45 MeV. $\theta_{lab}=57^\circ$. The errors are due to the statistical ones.




Fig. 2. Energy spectrum of deuterons produced by 720 MeV bremsstrahlung. The errors are the same as above.

C	12	6
REF. NO.		NVB
64 Ko 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.XP	SPC	20-32	C	32	SCI-D		90

Target enriched to 55.2% C¹³.

ABX




Fig. 1. Energieverteilung der Photoprotonen aus dem Methantarget (44% C¹², 56% C¹³). Das Histogramm stellt die auf Energieverlust korrigierten Meßwerte dar




Fig. 3. Energieverteilung der Photoprotonen aus C¹³. Das Histogramm wurde aus dem der Fig. 1 gewonnen, indem der Anteil der Protonenemission aus C¹² abgezogen wurde. Nähere Erläuterungen im Text




Fig. 5. Differentieller Wirkungsquerschnitt für Photoprotonenemission aus C¹³. a) Unter 90° zum γ -Strahl unter der Annahme, daß nur Grundzustandsübergänge vorliegen, diese Arbeit (Histogramm —), b) Ergebnisse von Cook⁴ (Kurve).
 Differentieller Wirkungsquerschnitt für Photoprotonenemission aus C¹² nach (1) (Histogramm ---)

$$\frac{\int_{31.5}^{\infty} \sigma(C^{13}) dE\gamma}{\int_{31.5}^{\infty} \sigma(C^{12}) dE\gamma} = 0.55 \pm 25\%, \text{ assuming isotropic angular distribution.}$$

REF. Akira Masaïke
 J. Phys. Soc. Japan 19, 427 (1964)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
64 Ma 4		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.XXX	ABY	150-720	C	150-720	ACT-I		4PI

Table I.

XXX=C 11 FINAL

	Al ²⁷ (γ , π^+) Mg ²⁷	Al ²⁷ \rightarrow Na ²⁶	Al ²⁷ \rightarrow F ¹⁹	C ¹² \rightarrow C ¹¹
Integrated Cross Section (200 MeV \sim 400 MeV)	2.2×10^{-28} cm ² ·MeV	6.2×10^{-28} cm ² ·MeV	2.4×10^{-28} cm ² ·MeV	2.0×10^{-28} cm ² ·MeV




Fig. 7. The yield curve per equivalent quanta for the reaction C¹² \rightarrow C¹¹.

REF.

S. Seki, M. Yamanouchi and M. Miwa
J. Phys. Soc. Japan 19, 1999 (1964)

ELEM. SYM.

C

12

6

METHOD

REF. NO.

64 Se 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	SPC	THR - 24	C	24	EMU-D	2-8	DST


Target thickness 2.5 mg/cm².

Fig. 3. Energy spectrum of photoprotons from carbon.

METHOD

Tandem; $T^3(p,\gamma)He^4$ source; anthracene crystal served as both sample and detector

REF. NO.

64 Sh 4

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.P	ABX	20-23	D	20-23	SCI-D		ΔPI



WIDTHS

TABLE I.

E_γ (MeV)	$\int \sigma dE$ (MeV-mb)*	Γ_γ (eV)
20.57	0.31	19
20.95	0.56	32
21.38	1.17	71

* The integrated cross section from 20.2 to 22.4 MeV is found to be:

$$\sigma_{int} = \int_{20.2}^{22.4} \sigma(E) dE = (16.2 \pm 2.5) \text{ MeV-mb.}$$

Fig. 5. Cross section of the $C^{12}(\gamma, p_0)$ reaction as a function of photon energy.Fig. 6. Comparison of the $C^{12}(\gamma, p_0)$ with the inverse $B^{11}(p, \gamma_0)$ reaction.

REF.

G.G. Taran, A.N. Gorbunov
 Zhur. Eksp. i Teoret. Fiz. 46, 1492-94 (1964)
 Soviet Phys. JETP 19, 1010 (1964)

ELEM. SYM.	A	Z
C	12	6

METHOD			REF. NO.		NVB		
			64 Ta 3		NVB		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	SPC	THR - 100	C	170	CCH-D		4PI
G,NP	SPC	THR - 100	C	170	CCH-D		4PI

$$\int_{170}^{\infty} \sigma(\gamma, p) dE = 122 \pm 5 \text{ MeV-mb}$$


$$\int_{37}^{\infty} \sigma(\gamma, p) dE = 102 \text{ MeV-mb}$$

$$\int_{37}^{\infty} \sigma(\gamma, p + \gamma, n + \gamma, np) dE = 180 \text{ MeV-mb}$$

Yield of various reactions on carbon

Type of reaction	No. of events observed	Yield relative to total yield, %	Type of reaction	No. of events observed	Yield relative to total yield, %
(γ, p)	2207	42	($\gamma, 3\alpha$)	137	2.5
(γ, n)	1541	31	($\gamma, 2\alpha n$)		
(γ, pn)	408	8	($C^{12}(\gamma, nHe^3)2He^4$)	42	0.8
(γ, α) [$C^{12}(\gamma, He^3)Be^9$]	83	1.5	including	301	5.5
($\gamma, \alpha n$)	92	1.8	4-prong stars		
including	542	10	($\gamma, 3pn$)	32	0.6
3-prong stars			($\gamma, 2p2\alpha$)	99	1.8
($\gamma, p\alpha$)	229	4.5	($\gamma, 2\alpha pt$)	141	2.7
($\gamma, p\alpha n$)	52	1	5-prong stars	23	0.5
($\gamma, 2p$)	51	1	6-prong stars	5	0.1
($\gamma, \alpha pn$)	31	0.6			

Total (γ, p) cross section: 1) observed in the present experiment; 2) differential (γ, p) cross section at an angle of 76° from the work of Dodge and Barber^[3], normalized to the maximum cross section obtained by Vanhuyse and Barber^[4]; 3) combined cross section for the reactions (γ, p) and (γ, pn).



Assumed ground state transitions.

REF.

M. Elaine Toms
Nuc. Phys. 50, 561-84 (1964)

ELEM. SYM.	A	Z
C	12	6

METHOD

Betatron



Page 1 of 4

REF. NO.

64 To 1

JCC

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G. 3A	ABX	9-22	C	22	EMU-D		DST

SPECTRUMFig. 3. The combined cross-section for the $^{12}\text{C}(\gamma, 3\alpha)$ reaction as a function of ^{12}C excitation energy.Fig. 4. Measurements of the cross-section for the $^{12}\text{C}(\gamma, 3\alpha)$ reaction as a function of ^{12}C excitation energy.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 5. Distribution of weighted ${}^9\text{Be}$ excitation energy E_x values in 0.3 MeV intervals of E_x for ${}^{12}\text{C}$ excitation energy E_γ regions; less than 12.2 MeV, 12.2-14.2 MeV and 14.3-16.3 MeV.




Fig. 6. Distribution of weighted ${}^9\text{Be}$ excitation energy E_x values in 0.3 MeV intervals of E_x for ${}^{12}\text{C}$ excitation energy E_γ regions; 16.4-17.9 MeV, 18.0-19.5 MeV and 19.6-21.8 MeV.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 7. Cross-sections for the $^{12}\text{C}(\gamma, 3\alpha)$ reaction by mode of breakup: via ^8Be ground state (above), not clearly via either ^8Be ground state or first-excited state (middle) and via ^8Be first-excited state (below).




Fig. 8. Distribution of $I(\theta_1) = N(\theta_1) \text{cosec} \theta_1$ by 18-degree intervals of θ_1 for 77 ground-state events from ^{12}C excitation energy less than 14.3 MeV. The solid curve is E2 interaction, and the broken curve is E1 interaction.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 10. Distribution of alpha energies by 0.2 MeV intervals of alpha energy for first-excited-state events in the C^{13} excitation energy E_1^* range 16.9-18.3 MeV. The solid curve is the predicted distribution for 0.4 E2 and 0.6 M1 interactions.




Fig. 11. Alpha-energy distributions by 0.2 MeV intervals of alpha energy with distributions predicted for photon interactions in ranges of C^{13} excitation energy E_1^* : 12.2-14.2 MeV fitted by 0.25 M1 and 0.75 E1, 14.3-16.3 MeV fitted by 0.8 E1 and 0.2 E2, and 16.4-17.9 MeV fitted by 0.4 E2 and 0.6 M1.




Fig. 12. Alpha-energy distributions by 0.2 MeV intervals of alpha energy with distributions predicted for photon interactions in ranges of C^{13} excitation energy E_1^* : 18.0-19.5 MeV fitted by 0.88 M1 and 0.12 E2 and 19.6-21.8 MeV fitted by all E1.

REF.

G. Walter and A. Coche
 C.R. Acad. Sc. Paris 259, 2817 (1964)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
64 Wa 1		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,3A	ABX	18	D	17	SCI-D		4PI

Based on Schuhl $^{63}\text{Cu}(\gamma, n)\sigma$
 $\sigma = 2.06 \pm 0.4 \times 10^{-22} \text{ cm}^2$

18=17.6 MEV

METHOD

Bremsstrahlung, Penfold-Leiss

[Page 1 of 2]

REF. NO.

65 Ba 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	ABX	THR-52	C	18-52	I BF3		4PI




Fig. 2. Photoneutron cross section on the nucleus ^{12}C . • and ■: data of two independent series of measurements, a and b: mean cross section values, o: the integral cross section $\int [\sigma(\gamma, n) + \sigma(\gamma, np) + 2\sigma(\gamma, 2n)] dE$ (the scale on the right).

NB: In Fig. 1, dots = Ref. 1;
In Fig. 2, dashes = Ref. 2.

References

1. K. Min and W. D. Whitehead, Phys. Rev. 137B (1965) 301.
2. J. Miller, G. Schuhl, G. Tamas and C. Tzara, Phys. Letters 2 (1962) 76.
3. M. I. Thorsen and L. Katz, Proc. Phys. Soc. 77 (1961) 166.
4. F. W. K. Firk, K. H. Lokan and E. M. Bowey, Proc. Intern. Symp. on Direct interactions and nuclear reaction mechanisms, Padua, E. Clementi and C. Villi, eds. (Gordon and Breach, 1962) 204.
5. V. G. Neudachin and V. N. Orlin, Nuclear Phys. 31 (1962) 336.
6. V. Güllert and N. Vinh-Mau, Nuclear Phys. 54 (1964) 321.
7. E. Boeker and C. C. Jonker, Physics Letters 6 (1963) 30.
8. N. W. Reay, N. M. Hintz and L. L. Lee, Nuclear Phys. 44 (1963) 338.




Fig. 1. The $^{12}\text{C}(\gamma, n)^{11}\text{C}$ cross section in the giant resonance range of energy.

METHOD				[Page 2 of 2]		REF. NO.		EGF
Bremsstrahlung, Penfold-Leiss						65 Ba 1		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		

Table 1

Our results			Position of the peaks of the (γ, n) reaction, according to the data of other investigators, in MeV			
E_{max}	ΔE_{max}	$\int_{E_1}^{E_2} \sigma dE$				
(MeV)	(MeV)	(MeV · mb)	[1]	[2]	[3]	[4]
20.9	0.5	2.2 ± 0.3 $E_1 - E_2$ 20.4 - 21.4	(18.8)	(20.6)	20.62 20.90 21.08 21.22 21.58	21.6
22.0	1.0	8.7 ± 1.2 $E_1 - E_2$ 21.4 - 22.9	22.2	22.2	22.02 22.88	22.0
23.5	0.8	10.9 ± 1.7 $E_1 - E_2$ 22.9 - 24.4	23.5	23.3		23.3
25.0	0.7	5.8 ± 2.2 $E_1 - E_2$ 24.4 - 25.4	25.5			25.5

Note: The data in the table are compiled only for the range of 18 - 26 MeV. Peaks shown in brackets are determined with small accuracy.

REF. T. Deforest, Jr., J. D. Walecka, G. Vanpraet, W. C. Barber
 Phys. Lett. 16, 311 (1965)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		65 De 3
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	14- 25	D	65	MAG-D		180




Fig. 1. Cross section for inelastic scattering of 65 MeV electrons at 180° from carbon, plotted as a function of the excitation energy.




Fig. 2. $|(2^{-1})^{1/2} T_2^{+-1} \text{MAG}(m=0)|^2$ for the 20.76 MeV 2^- , $T=1$ state in ^{12}C . As discussed in the text, the theoretical curve has been reduced by a factor of 2. The experimental points are for the state observed at about 19.2 MeV. The circles are obtained from experiments at Darmstadt [11] under the assumption that the single peak observed at 19.46 MeV is caused by the 2^- state. The triangles are previous Stanford results [11], and the square is from the present experiment.

¹¹J. Goldemberg and W.C. Barber, Phys. Rev. 134, B963 (1964).

REF. D. F. Herring, I. C. Nascimento, R. B. Walton, and R. E. Sund
 Phys. Rev. 139, B562 (1965)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		EGF
65 He 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,N	RLY	THR-32	D	14-32	ACT-I		4PI
E+,N	RLY	THR-32	D	14-32	ACT-I		4PI

Ratio of positron to electron induced activity determined.




Fig. 2. The ratio σ^-/σ^+ as a function of energy for a ^{12}C target.




Fig. 4. The ratio σ^-/σ^+ as a function of atomic number at 27-MeV (total) bombarding energy. The straight line is for comparison purposes.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, D	ABY	70-720	D	400-720	MAG-D	40-70	57
G, P	ABY	88-720	D	400-720	MAG-D	70-100	57




Fig. 5. Yield curves of photo-deuterons and protons from carbon. $\theta = 57^\circ$. $E_d = 45$ MeV and $E_p = 72$ MeV. Errors are the statistical standard deviations.




Fig. 7. Excitation function of photo-deuterons of 45 MeV from carbon. $\theta = 57^\circ$. Errors are the statistical standard deviations. Solid curve shows the theoretical results given in the text.




Fig. 6. Energy distributions of deuterons and protons from carbon exposed to 680 MeV bremsstrahlung. $\theta = 57^\circ$. Errors are the statistical standard deviations.




Fig. 8. Energy distribution of deuterons from carbon exposed to photons of 685 MeV. $\theta = 57^\circ$. Errors are the statistical standard deviations. Solid curve shows the theoretical results given in the text.

METHOD

Synchrotron; NBS chamber monitor

REF. NO.

65 M1 1

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XN	ABX	THR-30	C	THR-30	BF3-I		4PI




FIG. 1. Total photoneutron cross section of carbon unfolded in 1-MeV intervals.

TABLE I. Isotopic abundance in the samples and threshold energies.

Isotope	Abundance (%)	(γ,n) (MeV)	(γ,pn) (MeV)	($\gamma,2n$) (MeV)
C ¹²	98.89	18.7	27.4	32.4
C ¹³	1.11	5.0	20.9	23.7
Mg ²⁴	78.60	16.6	24.1	
Mg ²⁵	10.11	7.3	19.0	23.9
Mg ²⁶	11.29	11.1	23.2	18.4




FIG. 2. Total photoneutron cross section of carbon unfolded in 0.5-MeV intervals.

TABLE II. Energy levels observed in carbon (MeV).

Present work (γ,n)	Other experiments	
	(γ,n) ^a	(γ,p) ^b (absorption) ^c
18.8		16.5
		17.6
22.2	19.5	19.1
23.5	22.2	22.5
25.5	23.4	23.5
		23.0
		25.6
		25.6

$\int_{Th}^{29} \sigma dE = 39.2 \text{ MeV}\cdot\text{mb}$

^a Reference 9.
^b Reference 21.
^c Reference 22.

METHOD				REF. NO.			
Emulsions in beam				65 Ro 1		JOC	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,3A	ABX	12-17	C	12-17	EMU-D		4PI




FIG. 1. The yield in events per mole ¹²C per roentgen per MeV interval for the reaction ¹²C(γ, 3α) as a function of maximum photon energy. Open circles, present results; closed circle, from Greenberg *et al.* (1964). The solid curve was used to calculate the cross section in Fig. 2.




FIG. 2. Cross-section curve for the reaction ¹²C(γ, 3α).

METHOD			SOURCE		DETECTOR		ANGLE
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE	
$B^{11}(p,\gamma)C^{12}$	ABX	16-20	D	0-4	NAI-D		DST




FIG. 3. Coefficients obtained in fitting series of Legendre polynomials to the angular distributions for $B^{11}(p,\gamma)C^{12}$.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	SPC	THR-33	C	34	TOF-D	1-14	DST




Fig. 2. Photoneutron spectra from carbon at 55° and 93° to a bremsstrahlung beam produced by 34 MeV electrons penetrating a 0.15 cm thick bismuth disk.

METHOD

REF. NO.

[Page 2 of 2]

65 Ve 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	

TABLE I

Angular distributions for (γ, p) and (γ, n) reactions on ^{12}C and ^{16}O

Mode of excitation	Excited state spin and parity	l	Angular distribution	Relative intensity		
				55°	93°	141°
E1	1^-	0	isotropic	1.00	1.00	1.00
E1	1^-	2	$1 - \frac{2}{3} \sin^2 \theta$	0.81	1.00	0.64
E2.(E1) ^{a)}	$2^+, (1^-)^a$	1	$1 - \frac{1}{2} \sin^2 \theta$	1.32	1.00	1.63
E2	2^+	3	$1 + 2 \sin^2 \theta - \frac{1}{2} \sin^4 \theta$	2.42	1.00	1.40

Target nucleus	Excitation energy (MeV)	Emitted particle	Ref.	Relative intensity		
				55°	93°	141°
^{12}C	22.35	p	Dodge and Barber ²⁰⁾	0.81	1.00	0.45
^{12}C	>22	n	Allum <i>et al.</i> ²¹⁾	0.85	1.00	0.60
^{12}C	25.4	n	this paper	0.80	1.00	
^{14}O	22.3	p	Dodge and Barber ²⁰⁾	0.86	1.00	0.46
^{16}O	22.1	n	this paper	0.81	1.00	0.64
^{16}O	24.2	p	Dodge and Barber ²⁰⁾	0.97	1.00	0.45
^{16}O	24	n	this paper	0.89	1.00	0.68
^{16}O	23	n	this paper	0.81	1.00	0.64
^{16}O	20.8	n	this paper	0.73	1.00	0.63
^{16}O	18-20	n	this paper	0.77	1.00	0.73

^{a)} This angular distribution is also possible for $l = 2$ neutrons or protons from ^{12}C (1^- states) to ^{11}C or ^{11}B ($\frac{1}{2}^-$ ground state).




Fig. 3. The $^{12}\text{C}(\gamma, n_0)^{11}\text{C}$ cross section at 93° , derived on the assumption that all neutrons leave ^{11}C in the ground state. The upper curve is 58 % of the $^{12}\text{C}(\gamma, p_0)^{11}\text{B}$ cross section as calculated from the $^{11}\text{B}(p, \gamma_0)$ cross section of Allas *et al.* ⁹⁾. The vertical lines show the relative strengths of the giant resonance photo-absorption lines from the distorted nucleus calculations of Nilsson *et al.* ¹⁾.

REF.

J.M. Wyckoff, B. Ziegler, H. W. Koch, and R. Uhlig
Phys. Rev. 137, B576-94 (1965)

ELEM. SYM.	A	Z
C	12	6

METHOD

Synchrotron; ion chamber monitor

REF. NO.

65 Wy 1

NVB

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	10 - 35	C	90	SCI-D		4PI

32.1




Fig. 6. Carbon total photonuclear cross section. Points on plots (a) and (b) represent the data of this experiment. The solid line drawn in (a) is the total photonuclear cross section obtained by the Lebedev group using the right-hand ordinate. For ease in comparing the shapes their data has been moved up in energy by 250 keV. The solid line in the lower plot (b) is drawn as the best approximation to the experimental points. The dot-dashed line represents the corrected baseline when the radiative correction is applied to the electronic cross section.




Fig. 7. Carbon total photonuclear cross section integrated over energy. The dashed line is the normalization applied to bring the cross-section values to zero at threshold.

REF.

H. Artus
Z. Physik 189, 355 (1966)

ELEM. SYM.	A	Z
C	12	6

METHOD

60 MeV Linac

REF. NO.

66 Ar 1

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, BE7	ABX	30 - 57	C	30 - 57	ACT-I		4PI

$$\int_{\text{Thr}}^{57} \sigma dE = 6.0 \text{ MeV mb}$$




Fig. 3. Be^7 -Erzeugung aus Kohlenstoff. Die Meßpunkte dieser Arbeit sind durch Kreise (0,12 X_0 Ta-Konverter), Quadrate (0,075 X_0) und Dreiecke (0,012 X_0) dargestellt. Dabei entsprechen die ausgefüllten Symbole Aktivierungen in Targetposition 1, die leeren Zeichen Bestrahlungen in Position 2. Alle Meßwerte sind mit einem zusätzlichen Eichfehler von $\pm 5\%$ behaftet. Die eingezeichnete Kurve für das Ausbeuteverhältnis entspricht dem gestrichelt angegebenen Wirkungsquerschnitt. Die Schwellenenergien E_s für verschiedene Reaktionskanäle werden durch Pfeile angedeutet. Meßwerte anderer Experimentatoren sind die liegenden Kreuze² und das stehende Kreuz¹³. Von der Theorie vorausgesagte Niveaus im C^{12} -Kern werden durch Pfeile gekennzeichnet (bei 34,3 MeV, 35,0 MeV, 36,6 MeV und 41,5 MeV nach ¹⁸; bei 33,7 MeV nach ¹⁹; bei 29,5 MeV und 31,9 MeV nach ²⁰).

REF. H. Arcus, P. Brix, H. G. Clerc, F. Eigenbrod, A. Goldmann,
 F. Gudden, E. Spamer, P. Strehl, M. Stroetzel, O. Titze,
 and K. J. Wetzel
 Proc. Gatlinburg Conference, 314 (1966)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
66 Ar 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.E./	LFT	4	D		MAG-D		
		(4.43)					

$\lambda = 4.43$ MEV

TABLE 1

Summary of Experimental Results^a

Nuclide	E _x (MeV)	Type	Γ^b (eV)	Γ^c/Γ_w	R _{ex} (F)
⁶ Li	2.18	E2	(3.9 ± 0.5) × 10 ⁻⁴	14.4	3.77 ± 0.48
	3.56	M1	3.9 ± 0.4	9.4	2.96 ± 0.11
⁷ Li	11.23 ± 0.05	(M1) or	(1.3 ± 0.4)/g ^b	0.043/g	—
		(M2)	(0.026 ± 0.008)/g	2.6/g	—
⁹ Be	15.97 ± 0.03	M1	(3.7 ± 0.8) g	0.043/g	—
¹¹ B	4.46	E2 and	0.0173 ± 0.0021	3.2	3.44 ± 0.50
		M1	0.64 ± 0.08	0.34	2.50 ± 0.35
¹² C	5.04	M1	1.84 ± 0.14	0.69	2.60 ± 0.11
	4.43	E2	0.0122 ± 0.0008	5.30	3.14 ± 0.30
¹⁶ O	6.92	E2	0.100 ± 0.015	3.28	3.82 ± 0.46
	11.52	E2	0.52 ± 0.13	1.31	—
²⁴ Mg	9.85 ± 0.04	M1	7.95 ± 1.2	0.38	3.50 ± 0.49
	3.97 ± 0.03				
	10.35 ± 0.03	E2	0.24 ± 0.05	0.58	5.05 ± 0.50
	10.70 ± 0.03	M1	22.2 ± 2.4	0.86	3.50 ± 0.36
²⁸ Si	10.93 ± 0.04	E2	0.26 ± 0.11	0.50	—
	4.97 ± 0.02	C0	(2.0 ± 0.5) × 10 ^{-5c}	—	6.30 ± 1.20
⁴⁰ Ca	6.89 ± 0.05	E2	0.29 ± 0.04	2.65	4.60 ± 0.50

^a The Born approximation has been used except for ¹⁶O and ⁴⁰Ca.

^b g = (2I_i + 1)/(2I_f + 1).

^c Γ_w = equivalent to ME = (8.87 ± 1.00) F².

REF.

E. B. Bazhanov, A. P. Komar, A. V. Kulikov and V. I. Ogurtsov
 J. Nucl. Phys. (USSR) **3**, 711 (1966)
 Sov. J. Nucl. Phys. **3**, 522 (1966)

ELEM. SYM.

A

Z

C

12

6

METHOD

Synchrotron

REF. NO.

66 Ba 4

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	THR - 52	C	18 - 52	BF3-I		4PI




Fig. 2. Cross section for photoneutron reactions in the nucleus C^{12} : solid circles—first series of measurements, hollow squares—second series of measurements (left-hand ordinate scale), lines a and b—average values of cross sections, hollow circles—integrated cross section $\int [\sigma(\gamma, n) + \sigma(\gamma, np) + 2\sigma(\gamma, 2n) + \dots] dE_\gamma$ (right-hand scale). The statistical errors are shown. $\int_{18.7}^{52} \sigma dE_\gamma = 65.0 \pm 0.5$ MeV-mb.




Fig. 4. Cross section for the reaction $C^{12}(\gamma, n)C^{11}$ in the giant-resonance region. The statistical errors are shown.




Fig. 5. Comparison of the results of investigation of the $C^{12}(\gamma, n)C^{11}$ reaction. Histogram—our data; smooth curve—data of Ref. 6; experimental points—data of Ref. 9; black columns—results of Ref. 8. In the lower part of the figure are shown the positions of peaks and their half-widths according to the results of a study of neutron spectra.⁽²⁾

METHOD		Linac; S.E.M. Monitor; NBS Ionization Chamber		[Page 1 of 2]	REF. NO.	66 Bi 1	JDM
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XN	ABX	20 - 200	C	20 - 200	BF3-I	0 - 50	4PI

TABLE I. - In this Table are given the cross-sections measured in millibarn averaged over the energy ranges indicated in column 1. The contributions from the internal (I) and external (E) counters add together to give the total (T) cross-section.

		¹² C	²⁷ Al	S
Thres. ÷ 40 MeV	T	3.66	10.15	7.15
	E	0.38	0.80	0.70
	I	3.28	9.35	6.45
(40 ÷ 80) MeV	T	2.52	7.80	6.78
	E	0.94	1.35	1.48
	I	1.58	6.45	5.30
(80 ÷ 120) MeV	T	2.55	7.70	6.12
	E	1.13	1.87	2.37
	I	1.42	5.83	3.73
(120 ÷ 160) MeV	T	2.80	9.18	6.58
	E	1.36	3.00	3.66
	I	1.44	6.18	2.92
(160 ÷ 200) MeV	T	3.18	13.40	12
	E	1.48	5.25	3.60
	I	1.70	8.15	8.6




Fig. 3. - Carbon yield.




Fig. 9. - Total neutron integral cross-sections.




Fig. 10. - Total neutron differential cross-sections.

ELEM. SYM.	A	Z
C	12	6

METHOD
 Linac; S.E.M. Monitor; NBS Ionization Chamber

[Page 2 of 2]

REF. NO.
 66 Bi 1

JDM

TABLE II. - In this table the data from Table I are treated, as discussed in the text to give cross-sections corresponding to photon-neutrons in the approximate energy ranges (0-15) MeV and above 15 MeV. Thus the result a) corresponds to the relation $a = I - 0.4E$, and b) corresponds to $b = 1.4 \cdot E$.

	¹² C	²⁷ Al	S
Thres. ÷ 40 MeV	a) 3.13	9.03	6.17
	b) 0.53	1.12	0.98
(40 ÷ 80) MeV	a) 1.20	5.91	4.71
	b) 1.32	1.89	2.07
(80 ÷ 120) MeV	a) 0.97	5.08	3.35
	b) 1.58	2.62	3.35
(120 ÷ 160) MeV	a) 0.90	4.98	1.46
	b) 1.90	4.20	5.12
(160 ÷ 200) MeV	a) 1.11	6.05	6.95
	b) 2.07	7.35	5.05

a) = $I - 0.4E$, I = int. counter.
 b) = $E \cdot 1.4$, E = ext. counter.

DETECTOR		ANGLE
TYPE	RANGE	

TABLE IV. - The total absorption cross-section for γ -rays is calculated by dividing the total neutron production cross-section by the neutron multiplicity. The result is given in the first column under each element and, for comparison, in the second column the value of b is inserted from Table II.

	¹² C		²⁷ Al		S	
(40 ÷ 80) MeV	2.2	1.3	6.0	1.9	5.2	2.1
(80 ÷ 120) MeV	1.8	1.6	4.7	2.6	3.7	3.4
(120 ÷ 160) MeV	1.8	1.9	4.6	4.2	3.4	5.1
(160 ÷ 200) MeV	1.7	2.1	5.7	7.4	5.1	5.1

TABLE V.

E	$\sigma(\text{Al})/\sigma(\text{C})$	NZ/A
40	3.1	2.25
70	2.75	
100	2.52	
130	2.64	
160	3.15	

TABLE VI.

	$\sigma_{\text{intexp}}^{(14)}$ 0-35	σ_{intexp} 35-140	σ_{intexp}	$\sigma_{\text{int}}^{(18)}$	$\sigma_{\text{int}}(\text{B.L.})$
1	2	3	4	5	6
C	144	220	364 ± 40	373	252
Al	344	590	934 ± 100	840	567
S	466	382	848 ± 90	990	670

METHOD
70 MeV synchrotron

REF. NO.

66 Co 2

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	THR - 65	C	THR - 70	ACT-I		4PI

Cross section integrated to 65 MeV is 77 ± 6 MeV mb.

111




FIG. 1. Comparison of the present results for the $C^{12}(\gamma, n)C^{11}$ cross section with a recent monochromatic- γ -ray result. Error bars have the meaning given in the text in Sec. III. CLSR refers to the least-structure computer routine.




FIG. 4. Comparison of present work with neutron energy spectrum of Firk *et al.* The enhancement of the neutron spectrum at 21.7 MeV is probably due to an excited-state transition.

REF.

H. Crannell
Phys. Rev. 148, B1107 (1966)

ELEM. SYM. A

Z

C

12

6

METHOD

Linac

[Page 1 of 2]

REF. NO.

66 Cr 1

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	4, 10	D	600-800	MAG-D	650-	DST

$$q^2; 2.79 - 11.45 F^{-2}$$

$$\text{RMS } \langle r^2 \rangle = 2.40 \pm 0.02 F$$




Fig. 7. $F^2(q^2)$ as a function of q^2 for the 9.6-MeV level in C^{12} .




Fig. 6. $F^2(q^2)$ as a function of q^2 for the 4.43-MeV level in C^{12} . A diffraction feature can be observed at a q^2 of approximately $7 F^{-2}$.

REF.

H. Crannell
Phys. Rev. 148, B1107 (1966)ELEM. SYM. A Z
C 12 6

METHOD		REF. NO.		JDM	
Linac		66 Cr 1		JDM	
REACTION		SOURCE		DETECTOR	
RESULT		TYPE RANGE		TYPE RANGE	
EXCITATION ENERGY		ANGLE			

[Page 2 of 2]

TABLE II. Elastic and inelastic cross sections for C¹².

E_n (MeV)	θ (degrees)	Differential cross sections in cm ² /sr*		
		Elastic scattering	4.43-MeV level	9.64-MeV level
600	32.0	$(6.35 \pm 0.25) \times 10^{-32}$	$(4.54 \pm 0.05) \times 10^{-31}$	$(2.80 \pm 0.29) \times 10^{-31}$
	33.0	$(2.35 \pm 0.15) \times 10^{-32}$	$(3.58 \pm 0.05) \times 10^{-31}$	$(2.09 \pm 0.24) \times 10^{-31}$
	33.0	$(1.74 \pm 0.20) \times 10^{-32}$	$(3.39 \pm 0.03) \times 10^{-31}$	$(2.37 \pm 0.25) \times 10^{-31}$
	34.0	$(5.22 \pm 0.64) \times 10^{-33}$	$(2.50 \pm 0.02) \times 10^{-31}$	$(1.48 \pm 0.15) \times 10^{-31}$
	35.0	$(2.4 \pm 1.0) \times 10^{-33}$	$(1.72 \pm 0.01) \times 10^{-31}$	$(1.14 \pm 0.11) \times 10^{-31}$
	35.3	$(2.40 \pm 0.29) \times 10^{-33}$	$(1.82 \pm 0.01) \times 10^{-31}$	
	36.0	$(4.51 \pm 0.24) \times 10^{-33}$	$(1.30 \pm 0.01) \times 10^{-31}$	$(9.44 \pm 0.95) \times 10^{-32}$
	37.0	$(7.52 \pm 0.66) \times 10^{-33}$	$(9.15 \pm 0.11) \times 10^{-32}$	$(6.75 \pm 0.68) \times 10^{-32}$
	38.0	$(8.14 \pm 0.54) \times 10^{-33}$	$(5.98 \pm 0.09) \times 10^{-32}$	$(4.49 \pm 0.45) \times 10^{-32}$
	40.0	$(9.76 \pm 0.91) \times 10^{-33}$	$(4.43 \pm 0.11) \times 10^{-32}$	$(2.66 \pm 0.33) \times 10^{-32}$
	40.0	$(7.56 \pm 0.24) \times 10^{-33}$	$(4.48 \pm 0.06) \times 10^{-32}$	$(4.39 \pm 0.44) \times 10^{-32}$
	40.0	$(7.16 \pm 0.57) \times 10^{-33}$	$(3.55 \pm 0.09) \times 10^{-32}$	$(3.36 \pm 0.35) \times 10^{-32}$
	41.5	$(7.00 \pm 0.26) \times 10^{-33}$	$(1.68 \pm 0.05) \times 10^{-32}$	$(1.66 \pm 0.24) \times 10^{-32}$
	45.0	$(5.65 \pm 0.20) \times 10^{-33}$	$(5.01 \pm 0.28) \times 10^{-32}$	$(5.69 \pm 0.59) \times 10^{-32}$
	47.5	$(5.35 \pm 0.11) \times 10^{-33}$	$(1.68 \pm 0.14) \times 10^{-32}$	$(2.15 \pm 0.72) \times 10^{-32}$
	48.0	$(2.53 \pm 0.10) \times 10^{-33}$	$(1.06 \pm 0.12) \times 10^{-32}$	$(2.53 \pm 0.27) \times 10^{-32}$
	50.0	$(1.46 \pm 0.06) \times 10^{-33}$	$(4.2 \pm 1.3) \times 10^{-34}$	$(8.4 \pm 2.1) \times 10^{-34}$
	50.0	$(1.50 \pm 0.10) \times 10^{-33}$	$(2.03 \pm 0.98) \times 10^{-34}$	$(7.9 \pm 1.5) \times 10^{-34}$
	50.0	$(1.43 \pm 0.03) \times 10^{-33}$	$(3.99 \pm 0.40) \times 10^{-34}$	$(1.02 \pm 0.10) \times 10^{-34}$
	52.0	$(9.31 \pm 0.43) \times 10^{-33}$	$(1.16 \pm 0.35) \times 10^{-34}$	
55.0	$(4.38 \pm 0.26) \times 10^{-33}$	$(4.5 \pm 2.6) \times 10^{-34}$		
60.0	$(7.6 \pm 3.8) \times 10^{-33}$			
800	32.0	$(1.24 \pm 0.08) \times 10^{-32}$	$(2.12 \pm 0.11) \times 10^{-33}$	$(1.86 \pm 0.19) \times 10^{-33}$
	33.0	$(3.6 \pm 1.8) \times 10^{-33}$	$(1.14 \pm 0.29) \times 10^{-33}$	
	35.0	$(5.52 \pm 0.20) \times 10^{-33}$	$(3.91 \pm 0.20) \times 10^{-33}$	$(3.9 \pm 3.9) \times 10^{-33}$
	37.5	$(2.66 \pm 0.22) \times 10^{-33}$	$(8.3 \pm 1.7) \times 10^{-34}$	$(1.1 \pm 1.1) \times 10^{-33}$
	40.0	$(9.74 \pm 0.48) \times 10^{-34}$	$(2.1 \pm 1.0) \times 10^{-34}$	$(3.8 \pm 2.1) \times 10^{-34}$
	40.0	$(9.51 \pm 0.62) \times 10^{-34}$	$(1.99 \pm 0.99) \times 10^{-34}$	$(5.6 \pm 1.7) \times 10^{-34}$
	40.0	$(1.02 \pm 0.03) \times 10^{-33}$	$(2.3 \pm 2.8) \times 10^{-34}$	$(3.6 \pm 8.6) \times 10^{-34}$
	41.0	$(4.54 \pm 0.12) \times 10^{-34}$	$(1.12 \pm 0.61) \times 10^{-34}$	$(2.5 \pm 1.2) \times 10^{-34}$
	42.5	$(4.48 \pm 0.27) \times 10^{-34}$	$(1.65 \pm 0.83) \times 10^{-34}$	$(4.0 \pm 1.3) \times 10^{-34}$
	44.0	$(1.56 \pm 0.32) \times 10^{-34}$	$(6.9 \pm 3.4) \times 10^{-35}$	$(1.0 \pm 1.0) \times 10^{-34}$
	45.0	$(9.9 \pm 4.9) \times 10^{-35}$	$(9.9 \pm 4.9) \times 10^{-35}$	$(9.9 \pm 4.9) \times 10^{-35}$
	45.0	$(1.16 \pm 0.16) \times 10^{-34}$	$(1.17 \pm 0.32) \times 10^{-34}$	$(1.17 \pm 0.59) \times 10^{-34}$
	45.0	$(1.27 \pm 0.17) \times 10^{-34}$	$(8.7 \pm 2.0) \times 10^{-35}$	$(1.52 \pm 0.68) \times 10^{-34}$
	45.0	$(8.0 \pm 1.6) \times 10^{-35}$	$(8.0 \pm 1.3) \times 10^{-35}$	$(4.0 \pm 2.7) \times 10^{-35}$
	46.0	$(7.4 \pm 1.3) \times 10^{-35}$	$(7.4 \pm 1.8) \times 10^{-35}$	$(6.4 \pm 5.2) \times 10^{-35}$
	47.0	$(2.5 \pm 1.0) \times 10^{-35}$	$(4.5 \pm 1.4) \times 10^{-35}$	$(3.9 \pm 1.2) \times 10^{-35}$
48.0	$(1.02 \pm 0.51) \times 10^{-35}$	$(3.58 \pm 0.95) \times 10^{-35}$	$(1.8 \pm 1.3) \times 10^{-35}$	
48.0	$(1.83 \pm 0.54) \times 10^{-35}$	$(4.30 \pm 0.40) \times 10^{-35}$	$(4.0 \pm 2.7) \times 10^{-35}$	
49.0	$(1.0 \pm 1.0) \times 10^{-35}$	$(3.2 \pm 1.7) \times 10^{-35}$		
50.0	$(< 1.2) \times 10^{-35}$	$(2.7 \pm 1.3) \times 10^{-35}$		

* Errors given are those due to counting statistics only. All cross sections were measured with an angular acceptance in the horizontal plane ($\Delta\theta$) of 1.93°.

REF.

F. W. K. Firk
Proc. Gatlinburg Conference 352 (1966)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

66 Fi 2

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XN	SPC	THR-65	C	65	TOF-D	5-40	90




Fig. 1. Observed photoneutron time-of-flight spectra of C, O, Mg, Si, Ca, Co, V, and Ta.

[over]




Fig. 2. The 90° relative yields of photoneutrons from ^{12}C and ^{16}O compared with the total (γ, n) yields.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		JDM
66 Fo 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, N	ABX	18 - 70	C	18 - 70	ACT-I		4PI

$$\int_{\text{Thr}}^{70} \sigma(\gamma, n) = 64 \pm 5 \text{ MeV-mb}$$

$$\int_{\text{Thr}}^{70} \sigma(\gamma, n) = 88 \pm 7 \text{ MeV-mb}$$

$$\int_{35}^{70} \sigma(\gamma, n) = 30 \pm 5 \text{ MeV-mb}$$

Table compares with other measurements.




Fig. 1. - $^{12}\text{C}(\gamma, n)^{12}\text{C}$ cross-section: bin width $\Delta = 2 \text{ MeV}$.

REF. S.C. Fultz, J.T. Caldwell, B.L. Berman, R.L. Bramblett,
and R.R. Harvey
Phys. Rev. 143, B790 (1966)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

Nearly monochromatic x-rays from positron annihilation

66 Fu 1

jdm

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	18- 37	D	18- 37	BF3-I		4PI

^{12}C emits neutrons from mostly ground-state transitions.

112+

$$\int_0^{38} \sigma dE = 46 \pm 4 \text{ MeV mb}$$

TABLE I. Energy levels observed in C^{12} .

$E(\text{MeV})$	$\sigma_0(\text{mb})$	$\Gamma(\text{MeV})$	$\int_0^{\infty} \sigma dE$ MeV mb*	Percent dipole strength ^b
22.0	4.50	1.5	10.6	19
23.2	5.75	2.0	18.1	33
24.0				
25.5	2.50	2.0	7.9	14
27.1	1.50	1.5	3.5	6
28.3	1.50	1.5	3.5	6
30.5	1.95	2.0	6.1	11
35.2	~1.0	~3.5	~5.5	~11

* The integrated cross sections presented here are the areas under Lorentz lines which have been fitted to the resonances, (i.e., $(\pi/2)\sigma_0\Gamma$).
^b The percentage of dipole strengths represented here are the fractional integrated cross sections for the resonances listed.




FIG. 4. Top data points (A) are cross sections for photoneutrons emitted in ground-state transitions. The dashed curve represents the data of Verbinsky *et al.* (Ref. 16). The lower data points (B) are cross sections for neutrons emitted in transitions to the first excited state of C^{12} , while data points (C) are cross sections for neutrons emitted in transitions to the third and higher excited states.




FIG. 1. Photoneutron-cross-section data for carbon [$\sigma(\gamma,n) + \sigma(\gamma,np)$] obtained by the use of monochromatic photons, and measurements of the total neutron yield.

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				66 Ke 1		HMG	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$^{12}\text{C}(p, p')$	ABX	21-22	D	21-22	EMU-D		DST
				(21.3), (21.6)			




FIG. 3. Numbers of observed photoprotons plotted against $\cos^2 \phi$ for $E_\gamma = 21.6$ MeV. The distribution to be expected for pure $E1$ is shown as a histogram.




FIG. 4. Photoproton cross section as a function of photon energy. Solid line derived by detailed balance from the $^{11}\text{B}(p, \gamma_0)$ work of Allas *et al.* (Ref. 2). Dashed curve from the $^{12}\text{C}(e, p'e')$ results of Dodge and Barber (Ref. 10) assuming ground state transitions and $1 + 1.5 \sin^2 \theta$ angular distribution. Dot-dashed curve shows the $^{12}\text{C}(\gamma, p_0)$ results of Shin and Stephens (Ref. 3). The crosses represent the present results.

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	
Monochromatic x-rays from T(p, γ)He reaction	66 Lo 1	JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	21 - 27	D	21 - 27	ACT-I		4PI




FIG. 3. $^{12}\text{C}(\gamma,n)^{12}\text{C}$ activation cross-section curve as a function of photon energy.

- ¹⁰ K. Min and W. D. Whitehead, Phys. Rev. 137, B301 (1965).
¹¹ B. C. Cook (private communication).
¹² S. C. Fultz (private communication).
¹³ B. M. Spicer, Nuovo Cimento 2, 243 (1964).




FIG. 4. Comparison of time-of-flight neutron-energy curve (dashed line) of Firk and Bowey with $^{12}\text{C}(\gamma,n)^{12}\text{C}$ activation curve of present experiment (solid line).

REF.

J. M. Maison, M. Langevin and J. M. Loiseaux
 J. Physique, Colloq. 1, 36 (1966)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

66 Ma 2

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,NG/	RLX	21-31	C	21-31	NAI-D		
G,PG/	RLX	21-31	C	21-31	NAI-D		




FIG. 1. — Variation de la section efficace relative en fonction de l'énergie d'excitation pour l'émission de neutrons et protons vers les états $5/2^-$ (4,32 et 4,46 MeV) et $3/2^-$ (4,81 et 5,01 MeV) de ^{11}C et ^{11}B . Les + et les o correspondent à deux séries de mesures indépendantes.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.N	ABX	18-26	D	18-26	BF3-I		4PI

473

TABLEAU V

σ_{int} à 26 MeV en MeV.mb

	¹⁶ O	⁴⁰ Ca	¹² C	Mg
γ, n	$41,5 \pm 4$	73 ± 7	$29,4 \pm 3$	58 ± 6




FIG. 3. — $\sigma(\gamma, n)$ dans ¹²C. Points expérimentaux d'avril 1962. La courbe tient compte de la largeur finie de la raie de photons (400 keV).

V. Parikh
Nucl. Phys. 77, 157 (1966)

C

12

6

METHOD

REF. NO.

Linac, S.E.M. Monitor, Faraday Cup

66 Pa 2

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.P	SPC	THR - 55	C	55	SCI-D	34	DST




Fig. 1. Angular distribution of 34.2 MeV photoprotons from ^{12}C .

REF. G. Proca, D. Isabelle and J. Goldemberg
 J. Physique Colloq. 1, 105 (1966)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	66 Pr 1

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E, E/$	NOX	14-21	D	100-200	MAG-D		180

Level structure observed

Level	E_x	$B_{M_1}(q=0)$	$B_{E_2}(q=0)$
$1^+, T=1$	15.11	1, 1 F^+	
$2^-, T=1$	16.1 18.8 20.3		$(0.95 \pm 0.24) F^+$
$2^-, T=1$	16.0 19.4		
$1^-, T=1$	17.2 21 22.5 25.5		

REF. N. G. Afanas'ev, V. A. Gol'dshtein, S. V. Dementii, V. G. Vlasenko,
I. M. Arkatov, and Yu. I. Titov
J. Nucl. Phys. (USSR) **5**, 1134 (1967)
Sov. J. Nucl. Phys. **5**, 808 (1967)


ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	HMG
67 Af 1	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	RLX	0-100	D	346	MAG-D		DST

$d^2\sigma/dE d\Omega$, rel. un.



Spectrum of inelastic scattering of 846-MeV electrons by C^{12} (full circles) and elastic scattering by hydrogen (crosses) at different angles.

θ , deg	$\sigma_C/\sigma_p, \text{MeV}^{-1}$	R	α , MeV/c
23	$5.6 \cdot 10^{-3} \pm 11\%$	2.4 ± 0.3	216 ± 24
35	$6.1 \cdot 10^{-3} \pm 13\%$	2.8 ± 0.4	188 ± 24
45	$5.8 \cdot 10^{-3} \pm 21\%$	3.0 ± 0.6	176 ± 37

" σ_C/σ_p is the measured cross section ratio for inelastic scattering of 346 MeV electrons by C^{12} and elastic scattering by hydrogen"

REF. U. Amaldi Jr., G. Campos Venuti, G. Cortellessa, E. De Sanctis,
S. Frullani, R. Lombard and P. Salvadori
Phys. Letters 25B, 24 (1967)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		EGF
67 Am 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	RLX	130, 160	D	635	MAG-D		51




Fig. 1. Momentum distribution of the 1s protons. The dashed and full lines represent the harmonic-oscillator and square-well fits, respectively. In the inset a kinematical diagram of the reaction is shown.




Fig. 2. Momentum distribution of the 1p protons. The dashed and full lines represent the harmonic-oscillator and square-well fits, respectively.

REF. Yu. P. Antuf'ev, I. I. Miroshnichenko, V. I. Noga, and P. V. Sorokin
 Yad. Fiz. 6, 431 (1967)
 Sov. J. Nucl. Phys. 6, 312 (1968)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	HMG
	67 An 1	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	RLX	THR-999	C	THR-999	ACT-I		4PI

999 = 1.6 GEV




Fig. 1. Cross section σ_0 of the reaction $C^{12} \rightarrow C^{11}$ per equivalent γ quantum as a function of the maximum bremsstrahlung energy: \circ - from [2], \bullet - our data.

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		67 Be 3
		EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	4.43	D	113-390	MAG-D	100-400	DST




Fig. 1. Electro-excitation of the first excited state of ^{12}C at $q = 1.007 \text{ fm}^{-1}$. The straight line is a least-squares fit to the points.

Table 1
 Experimental cross section for the 4.43 MeV level of ^{12}C at $q = 1.007 \text{ fm}^{-1}$.

E (MeV)	θ	$d\sigma/d\Omega$ (10^{-32} cm^2)
389.85	30°	410 ± 26
295.05	40°	211 ± 10
235.40	50°	117 ± 7
202.00	60°	81.3 ± 6.5
176.25	70°	56.4 ± 3.6
157.35	80°	41.0 ± 2.2
144.50	90°	27.4 ± 1.7
132.50	100°	18.8 ± 1.1
124.20	110°	13.2 ± 0.8
117.70	120°	9.50 ± 0.59
112.60	130°	6.14 ± 0.36

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
[Page 1 of 2]		67 Cr 1
		JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	0-10	D	100-200	MAG-D		DST

$$q^2 = 0.12 \text{ fm}^{-2} \rightarrow 0.48 \text{ fm}^{-2}$$

TABLE 1
Measured transition widths for excited states in ^{12}C

Energy (MeV)	λ	Γ_{el} (in eV)		
		Present determinations	Ref. 1)	Previous determinations
4.43	2	$(10.6 \pm 1.1) \times 10^{-3}$	$(11.2 \pm 1.2) \times 10^{-3}$	$(10.5 \pm 2.0) \times 10^{-3}$ a)
7.66	0	$(6.2 \pm 0.6) \times 10^{-4}$	$(6.5 \pm 0.7) \times 10^{-4}$	$(5.5 \pm 3) \times 10^{-4}$ b)
9.4	3	$(3.1 \pm 0.4) \times 10^{-4}$	$(3.6 \pm 0.4) \times 10^{-4}$	

a) Ref. 6).

b) Ref. 7).




Fig. 1. Transition plot for the excitation of the 7.66 MeV level in ^{12}C . The ratio R_λ defined in eq. (1) is plotted for $\lambda = 0$ and $\lambda = 1$ using data from ref. 1). The large extrapolation necessary in the previous determinations of transition width is shown, and the fact that only the correct choice of multipolarity ($\lambda = 0$) will yield a straight line.

METHOD

[Page 2 of 2]

REF. NO. 67 Cr 1 JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	






Fig. 3. Transition plots for the first three excited states in ¹²C. The ratio R_λ defined in eq. (1) is plotted for the appropriate value of λ . Both the data from ref. 1) and this experiment are shown.

REF.

H. Crannell and H. A. Dahl
Phys. Rev. 155, 1062 (1967)

ELEM. SYM.

A

Z

C

12

6

METHOD

Linac

REF. NO.

67 Cr 2

JDM

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	19	D	400-800	MAG-D		DST





FIG. 1. Spectra of electrons scattered from C^{12} . The ordinate gives counts arbitrarily normalized. The abscissas give the scattered momentum as determined by a rotating-coil monitor. These readings are approximately in units of MeV/c, the accuracy depending on adjustment of the rotating-coil system. Corrections that have been applied to the data points are discussed in the text. Excitation energies at which peaks might be expected to appear are labeled. In each of these spectra the level at 19.5 MeV is clearly visible. No levels at higher energy are observed.

TABLE I. Differential cross section in cm^2/sr for the 19.5-MeV level in C^{12} .

E_0 (MeV)	θ (deg)	Δ (MeV/c)	Cross section	
			Method I	Method II
400.0	40.0	267	1.3×10^{-21}	...
434.0	40.0	290	9.9×10^{-22}	...
434.0	40.0	290	9.2×10^{-22}	...
490.0	40.0	323	7.7×10^{-22}	...
550.0	40.0	369	4.0×10^{-22}	1.7×10^{-22}
580.5	40.0	390	3.0×10^{-22}	1.0×10^{-22}
663.0	40.0	447	2.2×10^{-22}	8.6×10^{-23}
726.1	40.0	490	7.7×10^{-23}	3.0×10^{-23}
782.0	40.0	528	5.1×10^{-23}	2.1×10^{-23}
820.0	40.0	554	3.1×10^{-23}	6.3×10^{-24}
800.0	41.0	553	2.3×10^{-23}	...
800.0	44.0	592	1.0×10^{-23}	...
800.0	45.0	605	6.5×10^{-24}	...
800.0	47.0	630	3.2×10^{-24}	...
800.0	48.0	643	2.1×10^{-24}	...
800.0	48.0	643	1.8×10^{-24}	...




FIG. 2. Form factors for the 19.5-MeV level in C^{12} . Experimental determinations of the form factors by each of two methods, discussed in the text, are shown. The theoretical form factors for the $T=1$, 1^- and 2^- levels, and the sum of both contributions are also plotted.

METHOD				REF. NO.		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR	
			TYPE	RANGE	TYPE	RANGE
G, BE7	ABY	THR-999	C	300-999	ACT-I	4PI
G, N	ABY	18-999	C	300-999	ACT-I	4PI

67 Di 2 EGF

Production of C^{11} and Be^7 from C^{12} by High Energy Photons

999 = 1 GEV



V. di Napoli, F. Dobici, F. Salvetti and O. Firina

Istituto di Chimica Generale ed Inorganica dell'Università - Roma - Italy

H. G. de Carvalho - Centro Brasileiro de Pesquisas Físicas - Rio de Janeiro - Brasil

The present paper is part of a systematic work to study in detail the inelastic interactions of high energy photons with complex nuclei,¹ by means of bremsstrahlung beams with peak energies in the range 300 MeV up to 1000 MeV, from the Electron Synchrotron of Frascati. Fig. I and II show the cross-sections per equivalent quantum found for the C^{11} and Be^7 production, using the activation method. Within our experimental errors the cross-sections σ_k obtained by means of the photon difference method seem to be constant in the energy-range and equal to 1.2 ± 0.2 mb for C^{11} and 0.08 ± 0.04 for Be^7 production. To verify the accuracy of the σ_k measurements, σ_0 has been computed at two energies i.e. 300 MeV and 1000 MeV using the best σ_k data available at present in the literature,² by means of the numerical integral: $\sigma_0 = \int \sigma_k n(k, E) dk$, where $n(k, E)$ is the number of photons in the energy interval (k, dk) of the bremsstrahlung beam, normalised to one equivalent quantum. For comparison³ the results are represented in Fig. I and II. The $C^{12}(\gamma, n)C^{11}$ cross-section, in the energy range, is only 10% lower than the predicted value from the photon-meson model i.e.

$\sigma(k) = (A-2)P(k)[(\sigma_{\pi^+}(k) + \sigma_{\pi^-}(k))]$; $P(k)$ is the probability of simultaneous escape of the produced meson and the recoil nucleon without meson absorption or particle scattering; the σ 's are the photomeson cross-section production from free neutrons corrected for nucleon motion in the nucleus. For medium and heavy nuclei, however, we found quite a bad agreement with the predicted (γ, n) cross-sections from this equation.¹ The yields ratio C^{11}/Be^7 is larger than the one from high energy proton induced reactions in C^{12} , because photopions and recoil nucleons originate uniformly within the nucleus, rather than impinge from the outside and hence have only 1/2 the average path length, increasing by 4 the transparency.



References: 1) V. di Napoli et al.: Nuovo Cimento 48 (1967) 1; and references.

- 2) S.G. Fujita et al.: Phys. Rev. 143 (1966) 790; H. Artus, Z. Naturf. 20a (1965) 320.
- 3) A. Mosaiker: J. Phys. Soc. Japan 19 (1964) 427.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		eg ⁱ
67 Di 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.N	ABX	300-999	C	300-999	ACT-I		4PI
G.BE7	ABX	300-999	C	300-999	ACT-I		4PI

999=1BEV




FIG. 4

[over]





Fig. 2

○ MASAIKE
 ○ PRESENT WORK

ELEM. SYM.	A	Z
C	12	6
REF. NO.		HMG
67 Fe 1		

METHOD

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	25-30	D	10-15	NAI-D		90

FIG. 11. Yield curves for $B^{11}(p,\gamma)C^{12}$ and $B^{11}(p,\gamma_1)C^{12}$.TABLE II. Comparison of C^{13} excitation energies (in MeV) at which fine-structure peaks have been observed.

Present experiment (p,γ)	Gove <i>et al.</i> ^a (p,γ)	Becker and Fox ^b (p,γ)	Allas <i>et al.</i> ^c (p,γ)	Dodge and Barber ^d (γ,ρ)
	22.25 (γ_1)	22.1 (γ_1)	22.1 (γ_1)	...
	22.5 (γ_0)	22.5 (γ_0)	22.6 (γ_0)	22.5
	23.6 (γ_1)	23.6 (γ_1)	23.7 (γ_1)	23.2, 23.9
25.5 (γ_0,γ_1)	25.5 (γ_0,γ_1)	25.5 (γ_0,γ_1)	25.5 γ_0	25.6
			25.6 γ_1	
26.9 (γ_1)		26.9 (γ_1)	26.8 (γ_1)	26.6
27.45 (γ_0)		27.14
28.0 (γ_0,γ_1)		...	28.0 (γ_1)	27.9
		...	28.1 (γ_0)	
28.45 (γ_1)	
28.9 (γ_0)				28.9

^a Reference 5.
^b Reference 13.

^c Reference 14.
^d Reference 16.

^a H. E. Gove, A. E. Litherland, and R. Batchelor, Nucl. Phys. **26**, 480 (1961).

^b J. A. Becker and J. D. Fox, Nucl. Phys. **42**, 669 (1963).

^c R. G. Allas, S. S. Hanna, L. Meyer-Schützmeister, and R. E. Segel, Nucl. Phys. **58**, 122 (1964).

^d W. R. Dodge and W. C. Barber, Phys. Rev. **127**, 1746 (1962).

TABLE III. Absolute cross sections for $B^{11}(p,\gamma)C^{12}$ and $B^{11}(p,\gamma_1)C^{12}$, calculated at peaks of yield curve, assuming isotropic angular distributions.

Proton energy (MeV)	Cross sections (μb)	
	$\sigma(\gamma_0)$	$\sigma(\gamma_1)$
11.76	23.2 ± 6.5	44.7 ± 10.5
12.55	20.5 ± 5.7	35.7 ± 8.9
13.09	18.7 ± 4.4	37.6 ± 9.4
13.98	10.8 ± 2.9	25.1 ± 6.2
14.19	15.6 ± 5.0	22.7 ± 8.8

REF. S. Ferroni, V. Gracco, D. B. Isabelle, R. Malvano and M. Sanzone
Nucl. Phys. A98, 409 (1967)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		JOC
67 Fe 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	ABX	100-150	C	150	BF3-I		4PI




Fig. 3. Yields of the (γ , Tn) reaction for the three nuclei as a function of energy.




Fig. 6. The absorption cross sections for the three nuclei after all corrections have been applied.

TABLE 2
Values of the parameters for the best fit to the bremsstrahlung weighted cross sections

$$\int_{E_0}^{E_\gamma} \frac{\sigma(E)}{E} dE = (a_0 + a_1 E_\gamma) E_\gamma$$

	¹² C	¹⁶ O	⁴⁰ Ca
a_0 (10^{28})	7.03 \pm 0.05	7.9 \pm 0.5	24.6 \pm 0.8
a_1 (10^{28})	-0.0173 \pm 0.0004	-0.0072 \pm 0.0036	-0.09 \pm 0.006

REF. H. M. Gerstenberg and E. G. Fuller
NBS Tech. Note 416, June 1967

ELEM. SYM.	A	Z
C	12	6
REF. NO.		HMG
67 Ge 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABY	THR-27	C	22,27	BF3-I		4PI

Table 7. Comparison of neutron yields. Yields are given in units of (neutron cm²/MeV nucleus) × 10⁻²⁸. The estimated uncertainties in Y and Y_c are of the order of 6% and 10%, respectively.

Element	E ₀	Y(E ₀)	UCRL	Saclay	Va.	NBS(Old)	UCRL	Saclay	Va.	NBS(Old)	Ref.
							Exp	Exp	Exp	Exp	
			Y _c				Y _c /Y				
Pb	27	103	86				0.83				26,30
	22	111	92	116			0.83	1.05			
Au	27	89	97			115	1.09			1.25	24,30, 38
	22	92	98	88			1.07	0.96			
Ta	27	81	32	77		113	1.01	0.95		1.33	27,30, 38
	22	85	79	80			0.93	0.94			
Ho	27	67	75			103	1.12			1.49	27,31, 39
	22	69	77	32			1.12	1.19			
Ag	27	36									
	22	34.8									
Cu	27	14.4	13.2				0.92				28,30
	22	12.6	11.5	12.4			0.91	0.98			
Co	27	12.7	12.1		13.5		0.95		1.27		29,34
	22	10.6	9.9				0.94				
Ca	27	1.69		1.13	1.01			0.67	0.60		32,35
P	27	2.35			1.76				0.75		36
Al	27	1.92	1.62		1.38		0.84		0.72		25,37
O ¹⁶	27	0.54	0.42	0.48	0.42		0.78	0.89			16,32, 37
C	27	0.50	0.35	0.33	0.46		0.70	0.66			25,32, 33




Fig. 31. Absolute neutron yield as a function of atomic number. The neutron yield from calcium ($Z = 20$) is particularly low in comparison with the other elements because its (γ, n) threshold is high compared to the mean energy of the giant resonance.

REF. S. A. E. Johansson and L. Nilsson
Arkiv Fysik 35, 403 (1967)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	67 Jo 1	egf
----------	---------	-----

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	19-20	D	19-20	ACT-I		4PI

${}^7\text{Li}(p,\gamma){}^4\text{Be}$ radiation




Fig. 1. The cross section of the ${}^{12}\text{C}(\gamma,n){}^{11}\text{C}$ reaction. The two curves show two possible types of cross-section curves. The horizontal bars indicate the energy resolution. The arrows show the corrected position of the breaks in the activation curve of [12]. The height of the arrows is proportional to the integrated cross sections.

REF.

K. Krämer, H. v. Buttlar, A. Goldmann, and B. Huber
Z. Physik 207, 1 (1967)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

67 Kr 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.T	RLX	THR-55	C	30-55	ACT-I		4PI




Fig. 3. Yields of photoneutrons from ^{12}C and ^{16}O . Error bars are of the same order of magnitude as the diameter of the circles. The lowest thresholds are indicated by arrows. Curves have been drawn arbitrarily through the data points. The dashed line shows σ_y for the reaction $^{12}\text{C}(\gamma, pt)^3\text{Be}$ calculated from the cross section measured by МАЙКОВ¹⁸

¹⁸Maikov, V.N.: Soviet Phys. JETP 7, 973 (1958).
[Zhur. Eksptl. i Teoret. Fiz. 34, 1406 (1958)].

Table 2. Measured yields relative to that of $^{12}\text{C}(\gamma, n)^{11}\text{C}$, and σ_y for the process $^{12}\text{C}(\gamma, t)$

E_m MeV	$Y[^{12}\text{C}(\gamma, t)] Y_M$ 10^{-2}	$\sigma_y[^{12}\text{C}(\gamma, t)]$ μbarn
35	1.15 ± 0.06	20.3 ± 1.1
40	2.52 ± 0.08	50.4 ± 1.6
45	3.75 ± 0.11	30.9 ± 2.4
50	4.94 ± 0.11	112.4 ± 2.5

REF. H. W. Kuehne, P. Axel, and D. C. Sutton
 Phys. Rev. 163, 1278 (1967)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	HMG
	67 Ku 2	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.G	LFT	15	D	15	NAI-D	15	135

Photons, defined in energy to about 1% with the aid of a bremsstrahlung monochromator, were scattered by isolated energy levels in C, Mg, and Si. Parameters for the six observed levels are:

Isotope	Energy (MeV)	Γ_0/Γ (eV)	Γ_0/Γ	$B(M1)/(ch/2M\omega)^2$
C ¹³	15.11	36	1	0.93
Mg ²⁴	10.66±0.02	14	0.8	1.21
Si ²⁸	11.42±0.02	23	1	1.33
Si ²⁸	12.33±0.03			
Mg ²⁴	9.92±0.03	3.0	0.5	0.49
Mg ²⁴	10.07±0.05	4.2		≥0.36

The 15.11-MeV level in C¹³, the 9.92- and 10.66-MeV levels in Mg²⁴, and the 11.42-MeV level in Si²⁸ are $T=1$, $T_z=0$ analogs of low-lying $1+$ states in the neighboring odd-odd nuclei. These levels exhaust most of the magnetic dipole transition strength of the respective nuclei, and therefore give information about the expectation value of $1s$ in the ground state.

REF. J. M. Loiseaux, J. M. Maison, and M. Langevin
 J. de Physique 28, 11 (1967)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.		JOC	
				67 Lo 1			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G/	ABX	18-34	C	34	NAI-D		DST




FIG. 4.

- I : ¹²C Section efficace différentielle de diffusion ($E_m = 34$ MeV).
- II : ¹²C Section efficace prévue par la relation de dispersion et multipliée par 0,55.
- III : ¹⁴N Section efficace différentielle de diffusion à 140° ($E_m = 27$ MeV), ($\circ E_m = 32$ MeV).

REF.

G. A. Peterson
Phys. Letters 25B, 549 (1967)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		EGF
67 Pe 1		

REACTION	RESULT	EXCITAT ENERG	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.E./	LFT	15	D	40-65	D	20-65	DST

$$\Gamma_Y(C^{12}) = 36 \pm 3 \text{ eV}$$




Fig.1. Ratio of differential cross sections for the excitation of the 15.11 MeV state in ¹³C to that of the 15.11 MeV state in ¹²C versus momentum transferred to the nucleus in units of fm⁻¹.

REF.

I. L. Smith, J. Garvey, J. G. Rutherglen, and G. R. Brookes
Nucl. Phys. B1, 483 (1967)

ELEM. SYM. A Z

C 12 6

METHOD

REF. NO.

67 Sm 1

JOC

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,NP	ABX	150-250	C	250	TOF-D		DST




Fig. 4. Vertical angular distribution of coincident protons from carbon (the curve is explained in the text).

METHOD

REF. NO.

[Page 1 of 2]

67 Ta 1

HMG

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N G,3PN	ABX	THR-170	C	170	CCH-I		DST
G,NA G,PNA							
G,NHE3 G,2P2N2A							
G,P G,PA							
G,2P G,PT2A							
G,PN G,HE3							
G,2PN G,3A							

Table I

Yields of photonuclear reaction of carbon

Reaction	Reaction thresh- old, MeV	Number of events	Relative yield, %	Absolute yield, mb	Integrated cross section, MeV-mb*
(γ, p)	15.95	2213	42.6	4.41	116±4
(γ, n)	18.72	1541**	29.6	3.12	(82±7)***
(γ, pn)	27.41	402	7.8	0.80	42±5
(γ, He^3)	26.28	45	0.9	0.09	3.0±0.5
($\gamma, n\alpha$)	25.26	130	2.5	0.27	8.6±1.4
($\gamma, p\alpha$)	24.52	159	3.1	0.33	10.2±1.1
(γ, pna)	31.87	114	2.4	0.25	22.5±4.5
($\gamma, 2p$)	27.18	45	0.9	0.09	5.7±1.0
($\gamma, 2pn$)	34.0	35	0.7	0.07	4.2±1.1
($\gamma, 3\alpha$)	7.28	137**	2.5	0.28	(3.8±1.1)***
(γ, nHe^4)	27.85	42**	0.8	0.09	5.6±1.3
($\gamma, p2\alpha$)	27.08	70	1.4	0.14	9.4±1.2
($\gamma, 2p2n2\alpha$)	35.5	170**			
($\gamma, 3pn$)	60	32**	3.8	0.40	~29±6***
5-Prong stars	~55	28	0.5	0.05	(5±2)***
6-Prong stars	~85	5	0.1	0.01	(2±2)***
Total:		5207		10.7±0.2	345±15

* Only the statistical errors are included

** Visual identification

*** Determined from the reaction yield.

Table II

Parameters of the peaks in the cross section of the (γ, p) reaction

E_γ , MeV	σ_{max} , mb	Γ , MeV	$\int_0^\infty \sigma dE$, MeV-mb	$f_{0\%}$, %
17.5 ± 0.1	3.6 ± 2.0	0.4 ± 0.3	2.3 ± 2.0	2.3
19.3 ± 0.1	4.2 ± 1.8	0.46 ± 0.34	3.5 ± 2.5	3.5
20.4 ± 0.1	7.3 ± 1.6	0.92 ± 0.38	11.3 ± 4.3	11.3
(21.38 ± 0.05)	8.0 ± 2.7	0.15 ± 0.20	1.7 ± 2.5	~1.7
22.1 ± 0.1	12.0 ± 1.7	1.00 ± 0.23	19.0 ± 6.3	19
23.6 ± 0.1	11.2 ± 2.0	0.93 ± 0.23	18.0 ± 4.7	18
(24.75 ± 0.05)	5.2 ± 1.6	1.6 ± 0.77	11.2 ± 5.5	11.3
(25.8 ± 0.1)	5.8 ± 1.1	2.3 ± 0.87	17.5 ± 7	17.5
28.0 ± 0.2	3.2 ± 1.6	0.9 ± 0.4	4.3 ± 2.8	4.3
(31.1 ± 0.8)	3.5 ± 0.8	2.3 ± 0.68	10.4 ± 3.5	10.4

METHOD

[Page 2 of 2]

REF. NO.

67 Ta 1

HMG

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 1. Cross section of the (γ, p) reaction of carbon.

"The cross sections of the reaction (γ, p) were calculated assuming that the daughter nucleus B^{11} was left in its ground state."




Fig. 2. Proton angular distributions of the (γ, p) reaction of carbon for different intervals of the energy E_γ (MeV).




Fig. 3. Dependence of the coefficients of the angular distribution on the energy E_γ (MeV).

C	12	6
REF. NO.		HMG
68 Be 6		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	18-20	D	63-128	MAG-D		DST
				(63.7, 94.9, 127.2)			(125.155)

19.4 MEV LEVEL




Fig. 3. - Present and previous measurements in the low-momentum-transfer region. The data of Goldemberg and Barber were obtained at scattering angles of 152° and 180°; those of Vanpraet at 180°. The theoretical curve is the 2^- calculation by Hill.
 ▲ Stanford, GOLDEMBERG and BARBER;
 ■ Stanford, VANPRAET; ○ Saskatchewan;
 — Hill; × Darmstadt, GOLDEMBERG and BARBER.




Fig. 2 - The angular dependence of the square of the total form factor $|F_T|^2$. Since the line passes through the origin any longitudinal contribution is small.




Fig. 4. - The present results in the form $(Z^2/4\pi)(\frac{1}{2} - \text{tg}^2 20^\circ) F_T^2(q)$ and the results of Crannell *et al.* The curves showing minima at ~ 200 MeV/c represent the particle-hole predictions for a 1^- assignment.
 — $\langle 1^- | M_1(q) | 0^+ \rangle|^2$; --- $\langle 1^- | T_1(q) | 0^+ \rangle|^2$;
 • Experimental I; ▲ Experimental II;
 × Saskatchewan data.

REF. V. F. Borznkovskii, A. S. Cherkasov, N. G. Afanas'ev, I. A. Grisaev,
 and I. I. Zalyubovskii
 Yad. Fiz. 7, 261 (1968)
 Sov. J. Nucl. Phys. 7, 181 (1968)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
68 Bo 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.XXX	RLY	THR-110	D	110	MAG-D		76

XXX=MASS SPECTRUM





FIG. 2. Spectrum of pulses in AI-100 analyzer at $\theta = 76^\circ$. Target - polyethylene (6 mg/cm^2). Electron energy 110 MeV. Field of magnetic spectrometer corresponds to proton energy 5.3 MeV. Bias voltage on counter 60 V.

REF. S. V. Dementii, N. G. Afanas'ev, I.M. Arkatov, V. G. Vlasenko,
 V. A. Gol'dshtein and E. L. Kuplennikov
 Yad. Fiz. 8, 429 (1968)
 Sov. J. Nucl. Phys. 8, 243 (1969)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
68 De 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	55-202	D	580-968	MAG-D	450-968	DST
				(580, 805, 968)			(16°-60°)



q_{μ}^2, F^{-2}	ω, MeV	E, MeV	θ, deg	$\sigma_C \text{ max. cm}^2/\text{MeV-}\pi$	$\sigma_C \text{ max. } \sigma_M, \text{ MeV}^{-1}$	$W_1 \cdot 10^{-4} \text{ MeV}^{-1}$	$W_2 \cdot 10^{-4} \text{ MeV}^{-1}$
1.9	55	580 28		$2.19 \pm 0.12 \cdot 10^{-28}$	$1.47 \cdot 10^{-3} \pm 5.5\%$	46 ± 6	8.9 ± 0.3
		805 20		$3.55 \pm 0.14 \cdot 10^{-28}$	$1.16 \cdot 10^{-3} \pm 4.1\%$		
		968 16		$5.06 \pm 0.33 \cdot 10^{-28}$	$1.08 \cdot 10^{-3} \pm 6.3\%$		
2.8	75	580 35.5		$5.45 \pm 0.24 \cdot 10^{-28}$	$0.934 \cdot 10^{-3} \pm 6.2\%$	1.4 ± 2.2	9.1 ± 0.2
		805 25		$1.18 \pm 0.08 \cdot 10^{-28}$	$0.945 \cdot 10^{-3} \pm 6.5\%$		
		968 20.5		$1.78 \pm 0.07 \cdot 10^{-28}$	$0.926 \cdot 10^{-3} \pm 3.9\%$		
3.9	97	580 43		$1.96 \pm 0.14 \cdot 10^{-28}$	$7.48 \cdot 10^{-4} \pm 7\%$	0.01 ± 3.0	7.4 ± 0.4
		805 30		$5.02 \pm 0.47 \cdot 10^{-28}$	$8.40 \cdot 10^{-4} \pm 9.4\%$		
		968 24.5		$7.05 \pm 0.17 \cdot 10^{-28}$	$7.50 \cdot 10^{-4} \pm 2.4\%$		
6.3	149	580 59.5		$4.17 \pm 0.39 \cdot 10^{-28}$	$8.06 \cdot 10^{-4} \pm 9.4\%$	1.86 ± 1.69	4.6 ± 0.5
		805 40		$8.76 \pm 0.89 \cdot 10^{-28}$	$4.71 \cdot 10^{-4} \pm 7.8\%$		
		968 32		$1.59 \pm 0.08 \cdot 10^{-28}$	$5.11 \cdot 10^{-4} \pm 4.9\%$		
7.7	177	805 45		$4.95 \pm 0.35 \cdot 10^{-28}$	$4.32 \cdot 10^{-4} \pm 7\%$	8.61 ± 1.67	1.7 ± 0.8
		968 38		$6.45 \pm 0.29 \cdot 10^{-28}$	$3.23 \cdot 10^{-4} \pm 4.3\%$		
8.8	202	805 50		$2.61 \pm 0.15 \cdot 10^{-28}$	$3.52 \cdot 10^{-4} \pm 5.7\%$	2.2 ± 2.2	2.5 ± 1.1
		968 39.5		$4.22 \pm 0.27 \cdot 10^{-28}$	$3.18 \cdot 10^{-4} \pm 1.3\%$		

Inelastic form factors for carbon: a - $W_1(q_{\mu}^2, \omega)$, theoretical values calculated from Eq. (7); b - $W_2(q_{\mu}^2, \omega)$, theoretical values calculated from Eq. (8). In the $q_{\mu}^2 - \omega$ plane the straight line 1 corresponds to elastic scattering of electrons by carbon; region 2 is the region of excitation of discrete levels; line 3 is the threshold for electrodisintegration of the nucleus (see Eqs. (2) and (3)). The experimental points lie in a plane perpendicular to the $q_{\mu}^2 - \omega$ plane and passing through the "line of maxima" ϵ_M .

REF.

T. E. Drake, E. L. Tomusiak and H. S. Caplan
Nucl. Phys. A118, 138 (1968)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

68 Dr 1

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	19	D	140	MAG-D		DST

19=19.4


19.4 MeV Level in ^{12}C $J^\pi = 2^-, T=1$ 

Fig. 6. The momentum-transfer dependence of the square of the transverse matrix element for the 19.4 MeV level in ^{12}C . Theoretical curve I was obtained using TDA with harmonic-oscillator single-particle wave functions chosen to give the experimental rms radius of the ground state charge distribution. Curve II was obtained using RPA with Saxon-Woods single-particle wave functions chosen with the $\text{OP}_{\frac{1}{2}}$ radial function more confined to the origin. A Boeker-Brink potential (see appendix 1) was used for both curves.




Fig. 7. Plot of exhibits $|F_z(q, \theta)|^2 = |F_{\text{Cou}}(q)|^2 - (\frac{1}{2} + \frac{1}{2} \text{tg}^2 \theta) |F_{\text{Trans}}(q)|^2$ for the 19.4 MeV level of ^{12}C at a scattering angle $\theta = 40^\circ$. The experimental data of Crannell *et al.* ³⁾ are displayed as well as

our own.

FORM NBS-418
(REV. 7-14-64)
USCOMM-DC 28010-P64

214

METHOD

REF. NO.

[Page 1 of 2]

68 Fr 2

HMG

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	RLX	THR-32	C	30,32	TEL-D		DST
		(32.1)		(30.5)			
				(32.1)			




FIG. 8. Comparison of the total cross section shown in Fig. 7 to the cross section obtained from the inverse $B^{11}(p, \gamma)$ reaction to show nonground-state contamination. Curves are normalized together between 25.6 and 28.6 MeV.




FIG. 10. Angular-distribution fits at three particular photon energies from data obtained with 32.1-MeV bremsstrahlung. Data have been grouped into 200-keV intervals to emphasize smallness of systematic errors.




FIG. 11. The asymmetry parameter A_1 for data grouped into 50-keV intervals and $E_{\beta} = 32.1$ MeV. The solid curve is a smoothed fit to these experimental points. Errors shown are statistical only. Because of their similarity in energy dependence and poorer statistics, values of A_1 for $E_{\beta} = 30.5$ MeV are given in 200-keV intervals only. The latter set of points has been shifted downward 0.025, in view of the similarity and the bias limits shown in Table I, to remove an absolute discrepancy of 0.025.

METHOD					REF. NO.		HMG	
					68 Fr 2			
					[Page 2 of 2]			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		




FIG. 12. The asymmetry parameter A_3 for data grouped into 100-keV intervals and $E_0=32.1$ MeV. The values of A_3 for $E_0=30.5$ MeV are given in 200-keV intervals only.




FIG. 14. The anisotropy parameter A_2 for $E_0=32.1$ MeV.




FIG. 13. The anisotropy parameter A_2 for $E_0=30.5$ MeV.




FIG. 15. The quadrupole interaction parameter A_4 plotted in the manner of A_2 except that an average difference of 0.05 between sets of points was not arbitrarily removed. Because of the controls described in Sec. II, the $E_0=32.1$ -MeV results are more reliable, indicating $A_4 \approx 0$ at all energies.

REF. D. E. Frederick
Nucl. Phys. A119, 347 (1968)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
68 Fr 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	RLX	20-38	C	38	TEL-D	4-22	90
				(37.7)			

38 = 37.7 MEV

No structure 32-37 MeV with resolute ≤ 500 keV.




Fig. 1. Photoproton spectrum at 90° obtained with bremsstrahlung of end-point energy $E_\beta = 37.7$ MeV under the assumption of ground state transitions only.

ELEM. SYM.	A	Z
C	12	6

METHOD					REF. NO.	ANGLE	
					68 Ka 1	HMG	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	50-85	C	55-85	TOF-D	10-85	67 (67.5)

NEUT ENGY SPEC




FIG. 6. Observed neutron spectra due to 55-85-MeV difference photon spectra. The effective cross sections have been divided by NZ/A .

TABLE I. Comparison of present cross-section values in mb for production of high-energy photoneutrons by 55-85-MeV photons with measured cross sections $\sigma(\gamma, Tn)$, also in mb, for total photoneutron production. The present cross-section values are uncertain by 8 to 10% because of counting statistics and normalization errors; in addition all values depend on an absolute normalization in terms of the deuteron photodisintegration cross section, which is known to about 10% at these energies.

Target	$4\pi(d\sigma/d\omega)_{\gamma}^a$ ($E_n > 10$ MeV) [Present experiment]	$\sigma(\gamma, Tn)$ Jones and Terwilliger ^a	Costa <i>et al.</i> ^b	Other results
Li	0.75		1.0	
Be	1.0	2.7	2.3	2.3 ^c
B	1.0		1.4	
C	1.5	1.3	1.4	2.4 ^d
O	1.3		1.6	
Al	2.8	5.5	4.6	8 ^d
S	2.1		4.4	6.5 ^d
Fe	4.2	16	12	
Cu	4.3	20	19	
Zn	4.4		15	
In	7.4			
Sn	7.0			
Ta	10.7	95		
Tl	10.7			
Pb	8.3	100		
Bi	13			
U	16	65		

^a Average cross sections between 55 and 85 MeV, as read from Figs. 4 and 5 of Ref. 4.
^b $\int_0^{\infty} \sigma_{\text{pd}} dE - \int_0^{\infty} \sigma_{\text{pd}} E/50$, as taken from Fig. 4 of Ref. 5 and Table I of Ref. 6.
^c S. Costa, L. Pasqualini, G. Piragino, and L. Roasin, Nuovo Cimento 42, 306 (1966).
^d G. Bishop, S. Costa, S. Ferroni, R. Malvano, and G. Ricco, Nuovo Cimento 42, 148 (1966).




FIG. 7. Effective cross sections for production of fast neutrons with energies greater than 10 MeV (solid circles) and 30 MeV (open circles) by the 55-85-MeV photon difference spectrum. The dashed curves are modified quasideuteron model predictions as discussed in the text.

REF. J. L. Matthews, W. Bertozzi, S. Kowalski, C. P. Sargent and W. Turchinetz
Nucl. Phys. A112, 654 (1968)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		EGF
68 Ma 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.P	ABX	100	C	95-102	TFL-D	40-95	DST





Fig. 7. Photoproton spectra from ^{12}C with 98.5 MeV bremsstrahlung. The points labeled MIT are the result of the present experiment. Those labeled Oxford were derived from the results given in ref. ¹³⁾ (see text). The MIT data have been normalized to the Oxford measurement at $E_p = 55$ MeV, $\theta_p = 72^\circ$ in order to obtain the cross-section scale for the present experiment.

Fig. 8. Comparison of the ^{12}C and ^6Li photoproton spectra at $\theta_p = 92^\circ$, with 98.5 MeV bremsstrahlung. The arrows indicate the maximum proton energies allowed by the kinematics in each reaction.



REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.P	RLX	50-80	C	50-80	TEL-I	35-60	45

(E_p is the photon energy at which protons of energy E_p appear.)




FIG. 1. Yield (a) and cross section (b) for photoproduction of protons in the energy interval $E_p = 46.3-48.5$ MeV.




FIG. 2. The energy of the observed resonances versus the energy of the detected proton. The straight lines represent the kinematics as described in the text.

REF. G. A. Proca and D. B. Isabelle
 Nucl. Phys. A109, 177 (1968)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		68 Pr 1
		EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	14-21	D	100-200	D	30-200	180

Data given on

15.11 MeV	π 1+	\bar{T} 1
16.11	2+	1
16.57	2-	1
18.85	2+	1
19.42	2-	1
20.27	(2+)	1

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
68 Ra 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.XN	SPC	THR-103	C	103	TOF-D	3-90	DST




Fig. 10. High-energy spectra.

[over]




Fig. 8. Angular distribution coefficients for giant resonance region (curves from Boeker). The solid lines under the energy spectrum are from Kamimura *et al.*¹⁷; the dotted lines are from Drechsel *et al.*¹⁸. The smooth curves for A_j/A_0 are from Boeker²⁴.

METHOD						REF. NO.		EGF
						68 Ri 1		
[Page 1 of 2]								
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
E, E/	FMF	26-35	D	60-100	MAG-D	40-80	DST	




Fig. 2. Inelastic electron scattering cross section in ^{12}C at $E_i = 102.5 \text{ MeV}$ and $\theta = 60^\circ$. The full line is the quasi-elastic cross section calculated as described in the text.




Fig. 3. Inelastic electron scattering cross section in ^{12}C at $E_i = 60 \text{ MeV}$ and $\theta = 155^\circ$. The full line is the quasi-elastic cross section calculated as described in the text.

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		[Page 2 of 2] 68 Ri 1
		EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	

TABLE I
Experimental conditions of runs performed

E_i (MeV)	θ (deg)	q (fm ⁻¹)
102.5	60	0.47
60	155	0.47
71	155	0.58
82	155	0.70
100	155	0.87




Fig. 6. Inelastic scattering cross section in ¹²C at $E_i = 100$ MeV and $\theta = 155^\circ$. The full line is the quasi-elastic cross section calculated as described in the text.




Fig. 8. Form factors per MeV at excitation energy E_x and $\theta = 155^\circ$ as a function of the momentum transfer.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
(G.PN) (G.PT)	ABX	THR-170	C	170	CCH-1		DST
(G.PNA) (G.NHE3)							
(G.PA)							
(G.NA)							

Table I. Parameters of the angular distributions in the reactions (1), (3), (4), (5).

Reaction	Energy, MeV	Particle type	a_0	a_2	a_4
(1)	27-40	p	1	-0.33 ± 0.17	-0.08 ± 0.22
		n	1	-0.05 ± 0.21	-0.08 ± 0.27
	40-50	p	1	-0.07 ± 0.24	-0.14 ± 0.31
		n	1	-0.09 ± 0.30	$+0.12 \pm 0.35$
60-170	p	1	-0.10 ± 0.32	$+0.36 \pm 0.36$	
	n	1	-0.42 ± 0.44	$+0.32 \pm 0.47$	
(3)	24-40	p	1	$+0.27 \pm 0.18$	$+0.22 \pm 0.22$
		n	1	-0.30 ± 0.16	-0.31 ± 0.20
	40-170	p	1	-1.39 ± 1.25	$+0.43 \pm 1.31$
		n	1	$+0.23 \pm 0.22$	-0.21 ± 0.29
(4)	26-40	n	1	-0.02 ± 0.27	$+0.46 \pm 0.33$
		α	1	-0.18 ± 0.22	$+0.23 \pm 0.34$
	40-170	n	1	-0.29 ± 0.33	-0.04 ± 0.40
		α	1	$+0.97 \pm 0.30$	$+0.55 \pm 0.55$
(5)	27-170	p	1	-0.29 ± 0.17	-0.55 ± 0.24
		t	1	-0.28 ± 0.20	-0.13 ± 0.27




FIG. 1. (a) Cross sections of reaction (1) (full line) and sum of the cross sections of reactions (1) and (2) (dashed line); (b) Cross section of reaction (2); (c) Cross section of reaction (3); (d) Sum of the cross sections of reactions (5) and (6). Cross sections were computed with the assumption that the final nuclei were generated in the ground state.

Table II. Integrated cross sections for the reactions (1)-(6), MeV-mb.

Reaction type	E_p , MeV					
	0-40	40-50	50-100	100-140	140-170	0-170*
(γ , pn)	11.5	10.7	7.4	7.7	5.2	42 \pm 5
(γ , pna)	0.4	7.1	7.7	5.4	1.0	22.6 \pm 4.5
(γ , pa)	4.9	3.5	1.2	0.6	—	10.2 \pm 1.0
(γ , na)	4.1	3.0	1.2	0.3	—	8.6 \pm 1.7
(γ , pt)	1.6	2.9	3.0	1.9	—	9.4 \pm 1.2
(γ , nHe)	1.0	1.9	0.8	1.9	—	5.6 \pm 1.2

*Only the statistical errors are indicated here.




FIG. 10. Angular distributions: a: of nucleons and α particles in the reactions (3) and (4), b: of protons and tritons in reaction (5).




FIG. 9. Angular distributions of nucleons: a: reaction (1), b: reaction (2).

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				68 Wu 1		EGF	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	THR - 40	C	20-40	TOF-D	2-25	90

222+





FIG. 1. The observed $C^{12}(\gamma, n)C^{11}$ differential cross section at 90° compared with the $C^{12}(\gamma, p)B^{11}$ data deduced from detailed balance analyses using the $B^{11}(p, \gamma)C^{12}$ results of Allas *et al.* (Ref. 11) up to 28 MeV and Brassard, Scholz, and Bromley (Ref. 12) above 28 MeV. The ratio of the amplitudes a_0/a_2 is shown for penetration factors corresponding to $l=0$ and $l=2$ particle emission. The cross-hatched region indicates the error due to statistics: The absolute $C^{12}(\gamma, n)C^{11}$ cross section has a systematic uncertainty of 20% which is not included in the above diagrams.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABY	113-999	C	400-999	MAG-D	97,340	30
G,D	ABY	219-999	C	400-999	MAG-D	194	30

999 = 1.4 GEV



Excitation functions: cases a, b - for protons with energies 97 and 340 MeV, respectively, c - for deuterons with energy 194 MeV, at an l.s. angle 30°.

REF.

Yu. P. Antuf'ev, V. L. Agranovich, V. B. Ganenko, V. S. Kuzmenko,
I. I. Miroshnichenko, and P. V. Sorokin
Ukr. Fiz. Zh. 14, 248 (1969)

ELEM. SYM.	A	Z
C	12	6

METHOD					REF. NO.		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABY	114-999	C	700,999	TEL-D	97-230	DST
G,D	ABY	116-999	C	700,999	TEL-D	97-205	DST

999 = 1.2 GEV

Summary

The cross-sections of the (γ, p) (γ, d) reactions were investigated. Li^7 , Be^9 , C^{12} , Si^{28} , Cu^{63} , Mo^{98} and Ta^{181} targets were irradiated with the bremsstrahlung of 700 and 1200 MeV maximum energy from the Kharkov PhTI Ac. Sci. UkrSSR linear accelerator. The photo-protons and deuterons were detected by the scintillation telescope at 30° , 60° , and 120° with the beam. Possible mechanisms of the proton and deuteron photoproduction are discussed. The qualitative agreement of A dependence of the cross-sections is observed with a suggestion on the meson mechanism for these reactions.




Рис. 1. Залежність перерізу (γ, p)-реакції від A : $a - \phi = 30^\circ$, $E_p = 97$ Mev; $b - E_p = 205$ Mev; $c - \phi = 60^\circ$, $E_p = 230$ Mev; $d - E_p = 157$ Mev (\times — дані [3]); $e - \phi = 120^\circ$, $\circ - E_p = 120$ Mev, $\Delta - E_p = 157$ Mev, $\square - E_p = 230$ Mev. Абсолютне значення перерізу наведено при енергії протонів $E_p = 120$ Mev. Інші дані нормовані до перерізу для Li^7 при $E_p = 120$ Mev.

Рис. 2. Залежність перерізу (γ, d)-реакції від NZ/A : $a - \phi = 30^\circ$, $E_d = 97$ Mev; $b - \phi = 30^\circ$, $E_d = 205$ Mev; $c - \phi = 60^\circ$, $E_d = 97$ Mev, $\circ - \phi = 120^\circ$, $E_d = 97$ Mev (перерізи наведені в одиницях 10^{-22} см²/стер·Мев·Q).

REF.

Yu. P. Antufyev, V. L. Agranovich, V. B. Ganenko, V. S. Kuzmenko,
I. I. Miroshnichenko, P. V. Sorokin
Ukr. Fiz. 14, 496 (1969)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
1)	E, P	ABX	THR-999	C 400-999	MAG-D		DST
2)	E, D	ABX	THR-999	C 400-999	MAG-D		DST
3)	E, T	ABX	THR-999	C 400-999	MAG-D		DST




Рис. 1. Те саме, що на рис. 2 при $E_0 = 1200$ Mev.




Рис. 2. Енергетична залежність перерізів реакцій (γ, p) , (γ, d) , (γ, t) на ядрі C^{12} при максимальній енергії фотонів $E_0 = 700$ Mev (\circ — протони, Δ — дейтрони, \square — тритони).

Рис. 6. Порівняння експериментальних спектрів дейтронів з розрахунковими [11] для C^{12} , $E_0 = 700$ Mev: \circ — протони, $\theta = 60^\circ$, \square — протони, $\theta = 120^\circ$; \circ — дейтрони, $\theta = 60^\circ$, \square — дейтрони, $\theta = 120^\circ$.

- 1) 999 = 1 GEV, REL P/D
- 2) 1.3 GEV, REL P/D, D/T
- 3) 999 = 1 GEV, REL D/T

The differential cross-sections of (γ, p) , (γ, d) and (γ, t) reactions on carbon for 700 and 1200 MeV maximum energies of photons and energy distributions of the secondary particles were measured at 30, 60 and 120° of particle emission angles. Excitation function for protons with the energy of 97 MeV is given for the maximum incident photon energy from 400 to 1300 MeV. Deuteron to proton and triton to deuteron yield ratios for various nuclei are also shown.

Experiment was carried out at the Kharkov linear accelerator. The particles were detected by scintillation counters after a magnetic spectrometer.

Possible mechanisms of the high energy photon-nuclei interactions are discussed (6 Figs).

Таблиця 2

	θ°	$\left[\frac{d^2\sigma_d(2p)}{d\Omega dP} / \frac{d^2\sigma_p(2p)}{d\Omega dP} \right] \cdot 10^2$	$\left[\frac{d^2\sigma_t(3p)}{d\Omega dP} / \frac{d^2\sigma_d(2p)}{d\Omega dP} \right] \cdot 10^2$
Li ⁷	60	1.94 ± 0.19	2.1 ± 0.3
	120	0.8 ± 0.08	0.9 ± 0.13
C ¹²	60	2.2 ± 0.2	1.8 ± 0.3
	120	—	—
Ta ¹⁸¹	60	3.1 ± 0.3	2.8 ± 0.4
	120	1.89 ± 0.2	1.85 ± 0.28

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XN	ABX	19-25	C	19-25	BF3-I		4PI

314




Fig. 1. a) Our experimental results for the $C^{12}(\gamma, n)C^{11}$ cross section; b) theoretical results of Drechsel *et al.*; c-f) theoretical results of Kamimura *et al.* in the case of the forces I, II, III and IV, respectively

METHOD	REF. NO.
	69 Be 2
	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.MU-T	ABX	10-30	C	35	MGC-D	10-30	4PI




Fig. 1. The nuclear part of the photo-absorption cross section for ^{12}C compared to $\sigma(\gamma, p_0)$ (obtained by detail balance from the inverse reaction) and $\sigma(\gamma, n)$ from refs. ¹³⁻¹⁵). The estimated error of the position of the zero line is ± 0.7 mb.

TABLE 1
Weighted energy-integrated total photonuclear cross sections

	σ_{int} (mb · MeV)	$\frac{\sigma_{\text{int}}}{60(NZ/A)}$ (mb · MeV)	σ_{-1} (mb)	$\frac{\sigma_{-1}}{0.30A^{1/3}}$ mb	σ_{-1} (mb · MeV ⁻¹)	$\frac{\sigma_{-1}}{3.5A^{1/3}}$ ($\mu\text{b} \cdot \text{MeV}^{-1}$)
^{12}C	133 ± 13	0.74 ± 0.07	5.4 ± 0.6	0.65 ± 0.07	0.23 ± 0.03	1.04 ± 0.15
^{14}N	195 ± 37	0.93 ± 0.08	8.4 ± 1.7	0.83 ± 0.20	0.36 ± 0.08	1.28 ± 0.29
^{16}O	171 ± 17	0.71 ± 0.07	7.2 ± 0.8	0.60 ± 0.07	0.31 ± 0.04	0.87 ± 0.11
^{19}F	271 ± 50	0.94 ± 0.17	14.1 ± 2.7	0.82 ± 0.19	0.74 ± 0.17	1.60 ± 0.35
^{28}Si	360 ± 30	0.36 ± 0.07	17.5 ± 1.7	0.68 ± 0.07	0.83 ± 0.10	0.93 ± 0.11
^{40}Ca	580 ± 60	0.96 ± 0.10	29 ± 3	0.71 ± 0.08	1.5 ± 0.2	0.92 ± 0.12

The interval of integration is 10-30 MeV.

¹³R.G. Allas, S.S. Hanna, L. Meyer-Schützmeister and R. Segel, Nucl.Phys. 53 (1964) 122.

¹⁴G. Kernel and W.M. Mason, Nucl.Phys. A123 (1969) 205.

¹⁵S.C. Fultz, J.T. Caldwell, B.L. Berman, R.L. Bramblett and R.R. Harvey, Phys.Rev. 133 (1966) 790.

C	12	6
REF. NO.		egf
69 Be 5		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.E./	FMF	20-180	D	200-300	MAG-D	180-300	60

DIFFUSION QUASI-ELASTIQUE D'ELECTRONS DE HAUTE ENERGIE
 SUR LE CARBONE


J. BERTHOT, P.Y. BERTIN, J.P. GIRARDEAU-MONTAUT et D.B. ISABELLE, Laboratoire de Physique
 Nucléaire, F.63 - CLERMONT-FERRAND.

La diffusion quasi-élastique des électrons sur les noyaux permet d'obtenir directement des informations sur les fonctions de corrélation des nucléons dans les noyaux (1). La méthode d'analyse la plus simple, à savoir la mesure de règles de somme, s'est révélée peu fructueuse, car ces quantités sont dominées par les effets de corrélation à longue portée (Pauli). Toutefois, l'étude de la forme du spectre en énergie des électrons diffusés devrait nous permettre d'obtenir des informations plus détaillées.

C'est pourquoi nous avons effectué au près de l'Accélérateur Linéaire d'Orsay, une série de mesures à transfert de quantité de mouvement q et à angle de diffusion θ constants, ce qui implique une variation de l'énergie incidente en fonction de l'énergie d'excitation. Nous avons choisi $\theta = 60^\circ$ et $q = 190, 250$ et 300 MeV/c.

Les corrections radiatives (1) qu'il est nécessaire d'appliquer à toute expérience de diffusion d'électrons ont été effectuées grâce à un programme de déconvolution. Aucun modèle nucléaire n'est introduit dans cette analyse qui n'utilise que les résultats expérimentaux obtenus par des mesures complémentaires effectuées à énergie incidente et angle de diffusion constants.

Nous avons comparé (Figure 1) nos spectres expérimentaux avec ceux calculés théoriquement par De Forest (2). Celui-ci a utilisé un modèle en couche pour décrire les nucléons à l'intérieur du noyau et une fonction d'onde de l'oscillateur harmonique pour prendre compte de la distorsion de l'onde du nucléon sortant. Nous avons constaté, dans tous les cas mesurés, un bon accord entre l'expérience et la théorie en ce qui concerne le maximum du pic quasi-élastique



tant en position, qu'en valeur absolue de la section efficace.

Dans la partie décroissante du spectre quasi-élastique, nous avons constaté un accord satisfaisant entre les valeurs données par la théorie et celles observées expérimentalement. Cependant, nous devons remarquer que, tandis que le calcul théorique prévoit une décroissance monotone, les spectres expérimentaux présentent dans cette région, au moins deux épaulements. Pour les spectres considérés, la position en énergie de chaque épaulements, déterminée à 5 MeV près, est la même. Si nous choisissons comme zéro la valeur de l'énergie de recul du nucléon libre ($q^2/2M$), nous trouvons que pour le premier épaulements $\omega' = 45$ MeV, tandis que pour le second $\omega' = 75$ MeV.

La présence de ces épaulements pourrait s'expliquer par l'existence d'effets de corrélation à courte portée entre les nucléons dans les noyaux (effet quasi-deuteron par exemple). Toutefois, pour pouvoir réaliser une analyse suffisamment précise, nous devons effectuer des mesures complémentaires, à ces transferts de quantité de mouvement identiques mais pour des angles de diffusion différents. Ce programme est actuellement en cours.

1) D.B. ISABELLE : Rend. Scuola Intern. di Fisica E. Fermi, Corso XXXVIII, p. 392 (1967)
 2) T. DE FOREST : to be published.

METHOD				REF. NO.			
					69 Ca 3	hmg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, P_0	ABX	70-27	D	32	TEL-D	0-13	DST
				(31.5)		(.32-12.3)	

316




Fig. 3.—Differential cross section at 90° for $^{12}\text{C}(\gamma, p)^{11}\text{B}$.




Fig. 4.—Angular distribution coefficients.

REF.

H. G. De Carvalho, V. Di Napoli, D. Margadonna and F. Salvetti
and K. Tesch
Nucl. Phys. A126, 505 (1969)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	69 De 1

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABY	THR-999	C	1-6 (1.0-5.5)	ACT-I		4PI

Yield per equivalent quantum.

999 = 5.5 GEV




Fig. 1. Cross sections per equivalent quantum for (γ, n) reactions as a function of the maximum bremsstrahlung energy. A straight line is adjusted to the experimental points by means of a least-squares fit. The indicated errors are due to the γ -ray spectroscopy. The dashed line gives the result of simple estimates.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G _n N	NOX	19-260	C	260	ACT-I		DST

C11 RECOIL DS




FIG. 1. Angular distributions of the C^{11} recoil nuclei for different values of the maximum energy. The yield of the recoil nuclei in relative units is plotted against the angle of the recoil in the laboratory system. a) Energy of the recoil nuclei higher than 2.27 MeV; the solid and dashed lines are calculated with the help of the expression (1) for a "thick" and a "thin" target and for an excitation energy $E_{1exc} = 30$ MeV of the final nucleus C^{11} . b) Energy of the recoil nuclei higher than 1.37 MeV ("thick" target). The solid line was calculated using expression (1). In the energy range $63 \text{ MeV} < E_\gamma < 75 \text{ MeV}$ the excitation energies of the C^{11} nuclei of $E_{1exc} = 25, 28, 30$ MeV correspond to the curves I, II, and III, respectively. In the energy range of the γ rays from 75 MeV to 260 MeV the excitation energy is $E_{1exc} = 30$ MeV. c) Energy of the recoil nuclei higher than 0.4 MeV ("thick" target). The calculations were performed for the following values of the excitation energy of C^{11} and energy ranges of the γ rays: for E_γ from E_{7thr} to 63 MeV, $E_{1exc} = 20, 22, 25$ MeV (curves I, II, and III); for E_γ from 63 to 75 MeV, $E_{1exc} = 28$ MeV; and for E_γ from 75 MeV to 260 MeV, $E_{1exc} = 30$ MeV. d) Energy of the recoil nuclei higher than 0.12 MeV ("thick" target). It was assumed in the calculations that the excitation energy of C^{11} in the energy range of E_γ from E_{7thr} to 50 MeV is $E_{1exc} = 18, 20, 22$ MeV (curves I, II, III); for E_γ from 50 MeV to 63 MeV, $E_{1exc} = 22$ MeV; for E_γ from 63 MeV to 75 MeV, $E_{1exc} = 28$ MeV; and for E_γ from 75 MeV to 260 MeV, $E_{1exc} = 30$ MeV. e) Energy of the recoil nuclei higher than 0.01 MeV ("thick" target). The theoretical curves were calculated with the following assumptions made about the excitation energy of C^{11} : for E_γ from E_{7thr} to 38 MeV, $E_{1exc} = 5$ and 10 MeV (curves I and II); for E_γ from 38 MeV to 50 MeV, $E_{1exc} = 20$ MeV; for E_γ from 50 MeV to 63 MeV, $E_{1exc} = 22$ MeV; for E_γ from 63 MeV to 75 MeV, $E_{1exc} = 28$ MeV; for E_γ from 75 MeV to 260 MeV, $E_{1exc} = 30$ MeV. f) Energy of the recoil nuclei higher than 0.01 MeV. The experimental data for the "thick" target are shown. The calculation were carried out using expression (1) for a "thick" target (solid line) and "thin" target (dashed line). In both cases the excitation energy of C^{11} in the range from E_{7thr} to 38 MeV is $E_{1exc} = 5$ MeV, in the other ranges it is the same as for Fig. 1e.




FIG. 2. The excitation energy E_{exc} of C^{11} vs. γ -ray energy.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	300	D	580-968	MAG-D		DST




FIG. 3. Spectra of electrons with energy 580 (a), 805 (b), and 968 MeV (c) scattered by carbon. On the first two plots of Fig. 3a, the "elastic" peaks and the corresponding experimental points are shown in a reduced scale; in these cases, the corresponding "radiative tails" in the normal scale are shown to the left of the "elastic" peaks. Curves 1, 2, 2a, and 2b are explained in the text.




FIG. 3. (cont'd)




FIG. 2. Typical experimental yield curves measured at 580 MeV and 43°. O - results of measurements on carbon, X - results of difference experiment ($CH_2 - C^{12}$), Δ - background due to positively charged particles.

REF. I. S. Gulkarov, N. G. Afanas'ev, A. A. Khomich, V. D. Afanas'ev,
 V. M. Khvastunov, G. A. Savitskii, and N. G. Shevchenko
 Yad. Fiz. 9, 1138 (1969)
 Sov. J. Nucl. Phys. 9, 666 (1969)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
69 Gu 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	0-35	D	115,200	MAG-D	80-200	DST




FIG. 1. Spectra of inelastically scattered electrons with excitation of giant resonance in C^{12} , measured at initial electron energy 200 MeV and at different momenta transferred to the nucleus: a- $\theta = 20^\circ$, $q = 68.8$ MeV/c; b- $\theta = 30^\circ$, $q = 99.7$ MeV/c; c- $\theta = 40^\circ$, $q = 130.3$ MeV/c; d- $\theta = 50^\circ$, $q = 160.4$ MeV/c; e- $\theta = 60^\circ$, $q = 189.1$ MeV/c; f- $\theta = 70^\circ$, $q = 218.9$ MeV/c. The abscissas represent the excitation energy ϵ throughout.




FIG. 2. Spectra of inelastically scattered electrons with excitation of giant resonance in C^{12} , measured at initial electron energy 115 MeV, i.e., with a better energy resolution: a- $\theta = 35^\circ$, $q = 66.3$ MeV/c; b- $\theta = 75^\circ$, $q = 129.5$ MeV/c. The abscissas represent the excitation energy ϵ . The reason for the energy shift between the maxima of the two spectra has not been established.




FIG. 3. Dependence of the form factor of the level with excitation energy 16.1 MeV of the C^{12} nucleus ($\lambda = 2$) on the momentum transfer. Points: \circ - total form factor; \triangle - contribution from the level with excitation energy 15.1 MeV subtracted; \times - data obtained at Orsay [3,5] for the 16.1 MeV level. Explanations of the theoretical curves 1 and 2 are given in the text.




FIG. 4. Dependence of the quantity $Z^2 F_\lambda^2 [4\pi(\Delta^2/2q^2 + \tan^2(\theta/2))]^{-1}$ of the giant resonance on the momentum transfer. Points: \circ - our data, Δ - photon point (integration limits from 20 to 26 MeV) [6]; \triangle - photon point (integration limits from 12 to 35 MeV) [13]; \times - photon point obtained in the study of the partial reaction [14]; upper limit of integration 35 MeV; \square - data obtained in 180° scattering of the electrons; \bullet - value predicted from the sum rules. The curves are explained in the text.




FIG. 5. Coulomb form factor of the giant resonance vs. the momentum transfer. Curves 1 and 2 - calculation by Goldhaber-Teller and Brown models. On the upper right is shown the dependence of the excitation cross section of the giant resonance on the scattering angle.

METHOD				[Page 1 of 2]		REF. NO.	69 Ke 1	egf
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
P.G	ABX	28-37	D	13-21	NAI-D	20-38	DST	

2277




Fig. 7. The total cross section A_0 and coefficients $A_1^{c.m.}/A_0$ obtained from fitting the 24 cm x 30 cm NaI(Tl) angular distribution data to the series $W(\theta) = \sum_{l=0}^3 A_l P_l(\cos \theta)$ and transforming to the centre-of-mass reference frame.

METHOD				REF. NO.		69 Ke 1		egf	
[Page 2 of 2]									
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE		
			TYPE	RANGE	TYPE	RANGE			




Fig. 8. The total cross section A_0 and coefficients $A_1^{c.m.}/A_0$ and $A_2^{c.m.}/A_0$ calculated from the gamma-ray yields at 45° , 90° and 135° .

over




Fig. 9. Total cross sections for the $^{11}\text{B}(p, \gamma)^{12}\text{C}$ and $^{11}\text{B}(p, \gamma)^{11}\text{C}_{4,43}^c$ reactions up to $E_p = 21$ MeV.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G ₁ P	ABX	36-80	C	50-80	TEL-D	20-70	45

TABLE 1

Parameters of the resonant cross sections for $J E_p = 2.3$ MeV

E_p (MeV)	E_R (MeV)	$J E_R$ (MeV)	$d\sigma_R/d\Omega$ ($\mu\text{b} \cdot \text{sr}^{-1}$)
31.3	49.5	6	152 ± 9
33.6	50.5	7	134 ± 9
35.9	54.5	6	124 ± 11
38.2	58.0	7	113 ± 9
40.5	60.0	6	99 ± 8
42.8	62.0	6	76 ± 7
45.1	65.0	6	74 ± 7
47.4	67.5	7	62 ± 6
49.7	69.5	6	57 ± 6
52.0	72.5	7	46 ± 6
54.3	75.5	6	44 ± 7
56.6	77.0		
all ± 1.2	all ± 1.5	all ± 1.5	




Fig. 4. The proton spectrum derived from the data of fig. 2.




Fig. 6. The energy of the observed resonances versus the energy of the detected proton. The straight lines represent the kinematical relation, as described in the text.

[over]




Fig. 5. Proton yields and cross sections at 45° , relative to proton energy intervals $\Delta E_p = 2.3$ MeV. The solid lines in the yields are a smoothing by eye of the experimental values. The cross sections have been derived from the smooth curves using the Penfold and Leiss²⁰ unfolding procedure. The errors shown are statistical.




Fig. 7. Comparison between the integrated cross section below the resonances, normalized to unitary proton energy interval, and the theoretical value for the 1p protons using the momentum distributions of fig. 3. Also shown are the ground state cross sections measured from Penner and Leiss¹³) and a point from Whitehead *et al.*¹²).

¹²C. Whitehead, W. R. McMurray, M. J. Aitken, N. Middlemas and C. H. Collie, Phys. Rev. 110 (1953) 1130

¹³S. Penner and J. E. Leiss, Phys. Rev. 114 (1959) 1101

²⁰A. S. Penfold and J. E. Leiss, Phys. Rev. 117 (1959) 1332

METHOD				REF. NO.			
				69 0w 2		egf	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,NA	ABX	26-31	C	26-31	ACT-I		4PI

276




Fig. 1. Cross section to 31 MeV of the $^{12}\text{C}(\gamma, n\alpha)^7\text{Be}$ reaction.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XXX	ABX	16-60	C	170	CCH-D		4PI
G,P	3.7	16-60	C	170	CCH-D		DST

XXX= CHARGED PARTICLES

217

Table I. Coefficients of angular distributions of nucleons in certain partial channels of photodisintegration of carbon

Type of reaction	a ₀	a ₂	a ₄	a ₆
(γ, p)	1	+0.2 ± 0.2	-0.7 ± 0.2	-0.1 ± 0.3
(γ, pn)	1	-0.0 ± 0.2	-0.3 ± 0.3	-0.0 ± 0.4
(γ, pα)	1	+0.3 ± 0.3	-0.0 ± 0.3	+0.1 ± 0.4
(γ, pα) + (γ, p)				

Table II. Integral cross sections of partial reactions of the photodisintegration of carbon in the energy region 32-37 MeV (only statistical errors are indicated)

Type of reaction	Integral cross section		Type of reaction	Integral cross section	
	MeV·mb	%		MeV·mb	%
(γ, p)	0.0 ± 1.1	33 ± 6	(γ, pn) + (γ, n He*)	1.5 ± 0.5	5.7 ± 1.7
(γ, n)	6.5 ± 1.0	23 ± 5	(γ, 2p)	0.6	2.2
(γ, pα)	5.5 ± 1.0	20.0 ± 4.5	(γ, He ³)	0.58	3
(γ, pα) + (γ, n)	3.3 ± 0.5	12 ± 2.5			

*Estimated from the cross section of the reaction (γ, p).




FIG. 1. Summary effective cross section of the reactions (γ, p), (γ, pn), (γ, pn), (γ, 2pn), (γ, 2p), (γ, pα), (γ, nα), (γ, pα), (γ, nHe³), and (γ, He³). The dashed lines shows the cross section of the reaction (γ, p).




FIG. 3. Angular distribution of photoprotons in different partial channels in the photon energy region 32-37 MeV.




FIG. 2. Spectrum of photoprotons from carbon in the photon energy region E_γ = 32-37 MeV.

Figure 2 shows the spectrum of the photoprotons emitted by carbon after absorbing photons with energy 32-37 MeV (obtained from the data on the reactions (γ, p), (γ, pn), (γ, pα), (γ, pt), and (γ, 2p) under the assumption that the final nuclei are produced in the ground state). The spectrum consists of two components—soft (maximum at E_p = 4 MeV) and hard (E_p = 13-19 MeV). The main contribution to the soft component is made by protons from the reaction (γ, pn), while the hard component is represented exclusively by protons from the reaction (γ, p).

REF. Yu. I. Titov, E.V. Stepula, R.V. Akhmerov, S.A. Byvalin,
 N.F. Severin, E.M. Smelov, L.D. Yaroshevskii
 Yad. Fiz. 2, 1326 (1969)
 Sov. J. Nucl. Phys. 2, 772 (1969)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				69 Ti 2		hmg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E ₁ E ₂	ABX	0-700	D	999	MAG-D		25

Second peak interpreted as pion production.

999 = 1.126 GEV




FIG. 1. Cross section for inelastic scattering by C¹² at an angle of 25°C for an initial electron energy of 1126 MeV. Solid curve—quasielastic scattering cross section.

REF.

Y. Torizuka, M. Oyamada, K. Nakahara, K. Sugiyama, Y. Kojima,
T. Terasawa, K. Itoh, A. Yamaguchi, M. Kimura
Phys. Rev. Letters 22, 544 (1969)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

69 To 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	11	D	183, 250	MAG-D	150-250	DST

11=10.3 MEV




FIG. 1. Energy spectra of scattered electrons.
(a) Carbon, $E_0 = 250$ MeV, $\theta = 85^\circ$. The dashed lines are assumed to reproduce the 4.43-MeV peak shape for the 9.63-MeV peak. (b) Beryllium oxide, $E_0 = 183$ MeV, $\theta = 75^\circ$. The crosses indicate the energy spectrum obtained for beryllium target. The 7-MeV peak was decomposed into two components by using shapes of other peaks.

[over]




FIG. 2. Experimental and theoretical form factors squared. (a) The 10.5-MeV ($1^-; T=0$) level of ^{12}C and (b) the 7.12-MeV ($1^-; T=0$) level of ^{16}O . The dashed and solid curves in (b) are theoretical Coulomb and sum of Coulomb and transverse form factors for $E_0 = 183$ MeV, respectively, which have been calculated by Seaborn and Eisenberg (Ref. 2) in the 1p-1h model. They are multiplied by 3 in the figure. The dotted curves are $E1 \Delta T = 1$ form factors calculated using random-phase-approximation wave functions of Gillet and N. Vinh Mau [Nucl. Phys. 54, 321 (1964)].

METHOD

REF. NO.

69 To 4

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E. E./	ABX	18-39	D	177,250	MAG-D	138-232	DST




FIG. 1. The spectra (A), (B), (C), and (D) correspond to the cross sections at 35° (250 MeV), 60° (250 MeV), 80° (250 MeV), and 135° (177.6 MeV), respectively. The solid curve in (A) is the $^{12}\text{C}(\gamma, n)^{11}\text{C}$ cross section reported by Cook *et al.* Phys. Rev. 143, 724 (1966). Our calculated cross sections for the 1^- states, and those calculated by Kelly *et al.* (Ref. 9) for the 2^- states, using the wave functions of the KIA model (Ref. 11) are shown in the spectra with solid lines for the 1^- states and dotted lines for the 2^- states.

The unfolded spectra of the scattered electrons with initial electron energy of 250 MeV at 35° (A), 60° (B), and 80° (C) are displayed in Fig. 1. The scales on the right-hand side indicate the experimental cross section. The spectrum (D) corresponding to the incident energy of 177 MeV at an angle of 135° was not unfolded. The

²F. J. Kelly and H. Überall, Phys. Rev. 175, 1235 (1968).

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E_e, E_e'	ABX	13-22	D	50-70	MAG-D	23-70	180

Table. Data from $^{12}\text{C}(e, e')$ at 180°

15.1, 19.2 MEV

Levels (MeV) (± 0.25 MeV)	q (MeV/c)	Inelastic cross section (10^{-32} cm 2 .sr)	$ \langle \ T(q)\ \rangle ^2 \times 10^{-3}$
15.1	85	2.00	(0.78 ± 0.08)
	115	1.75	(1.14 ± 0.12)
	125	1.62	(1.23 ± 0.12)
19.2	91	1.41	(0.56 ± 0.09)
	111	2.40	(1.57 ± 0.16)
	121	2.38	(1.30 ± 0.13)



Fig. 1. Cross section for inelastic scattering of 50 and 70 MeV electrons at 180° from carbon, plotted as a function of the excitation energy

Fig. 2




Fig. 3

Fig. 2. The experimental values of the reduced transition probability $B_{T1}(1^- \rightarrow 0^-)$ for ^{12}C shown as a function of the square of the momentum transfer. The theoretical curves are obtained for two different values of the harmonic oscillator parameter b . They are taken from Ref. ²⁰ after requiring to go through the photon point corresponding with a value of $B_{T1}(0) = 0.93$ (Ref. ²¹)

Fig. 3. The quantity $|\langle 2^- T_2 V(q) 0^- \rangle|^2$ for the 20.76 MeV 2^- , $T=1$ state in ^{12}C and the experimental results for the peak strength observed at 19.2 MeV. The dashed curve represents 50% of the calculated strength in the particle-hole model

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
70 Ah 1		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	15-80	C	108	NAI-D	5-75	135




Fig. 1. Pulse height distribution for ^{12}C and ^{16}O . The simultaneously measured neutron background distribution is subtracted. The 15.11 MeV line in the oxygen distribution is the deexcitation of the 15.11 MeV level in ^{12}C , populated by $^{16}\text{O}(\gamma, \alpha)^{12}\text{C}$ [7]. Lines below 10 MeV are γ -transitions in the daughter nuclei following (γ, n) or (γ, p) reactions. The heavy line is calculated with the assumptions given in the text and with resonance parameters adjusted to give a good fit to the measured distribution.




Fig. 2. The full line is the total absorption cross section σ_{tot} calculated from the imaginary part of the scattering amplitude. The points are taken from total absorption measurements [8]. Since particle threshold effects are not built into the scattering amplitude, agreement between the calculated and measured cross section cannot be expected in the threshold regions.

Table 1
 Resonance energies (MeV) for the elements investigated.

Carbon	22.7 ± 0.3	25.9 ± 0.4	29.5 ± 0.6	45 ± 3
Oxygen	22.7 ± 0.3	25.2 ± 0.3	31.3 ± 0.6	50 ± 3
Magnesium	16.4 ± 0.3	19.1 ± 0.3	24.7 ± 0.5	34 ± 2
Silicon	17.8 ± 0.3	19.3 ± 0.3	21.2 ± 0.4	28.5 ± 1.0
Calcium	19.5 ± 0.3	24.4 ± 0.5	35 ± 2	
Silver	15.3 ± 0.3			

REF. P. Antony-Spies, P. P. Delsanto, E. Spamer, A. Goldmann and
O. Titze
Phys. Letters 31B, 632 (1970)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.				
REACTION		RESULT	EXCITATION ENERGY	SOURCE		DETECTOR	ANGLE	
				TYPE	RANGE	TYPE	RANGE	
E _i E/		ABX	16-30	D	55	MAG-D	25-40	141




Fig. 2. Differential cross section for electroexcitation of ¹²C plotted against excitation energy. (a) Contributions of the various multipoles to the cross section. (b) Total theoretical cross section (negative parity states) folded with an experimental resolution of 0.2 MeV. (c) Experimental cross section.

REF. Yu. P. Antuf'ev, V. L. Agranovich, V. G. Ganenko, V. S. Kuz'menko,
I. I. Miroshnichenko, and P. V. Sorokin
Yad. Fiz. 12, 1143 (1970); Sov. J. Nucl. Phys. 12, 627 (1971)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
70 An 5		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, P	RLY	96-999	C	620,999	TEL-D	80-265	DST

999 = 1140 MEV




FIG. 1. Angular distributions of protons with energies of 80, 160, and 285 MeV produced from C^{12} , Al^{27} , Ni^{59} , and Ta^{181} nuclei by photons with maximum energy 1140 MeV. Only the statistical errors are shown.

FIG. 2. Angular distributions of 80-MeV protons (normalized at $\theta = 40^\circ$) for C^{12} (O) and Ta^{181} (X), $E_{\gamma \max} = 1140$ MeV.





FIG. 3




FIG. 4

FIG. 3. Total cross section for proton production per nucleon. $E_{\gamma \max} = 1140$ MeV. Dashed curve—theory from ref. 11.

FIG. 4. Total cross section for proton production from C^{12} as a function of $E_{\gamma \max}$. Dashed curve—total cross section of photomesonic reactions, solid curve—sum of total cross sections for photomesonic reactions and the deuteron photodisintegration reaction.

¹¹ K. S. Kölbig and B. Margolis, Nucl. Phys. B6, 85 (1968).

REF. S.V. Dementii, N.G. Afanas'ev, I.M. Arkatov, V.G. Vlasenko,
 V.A. Gol'dshtein, and E.L. Kuplennikov
 Yad. Fiz. 11, 19 (1970)
 Sov. J. Nucl. Phys. 11, 10 (1970)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
70 De 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E ₁ E ₂	ABX	400-999 (400-1115)	D	690-999 (690-1115)	MAG-D		DST

690=1115 MEV




FIG. 1. Cross section for scattering of 690-MeV electrons by C¹² nuclei as a function of the energy E' for various scattering angles. Curves 1, 2, 3 have been calculated respectively from Eqs. (1), (15), and (14).




FIG. 2. Cross sections for scattering of 970-MeV electrons by C¹² nuclei as a function of the energy E' for various scattering angles. Curves 1-3 have been calculated as in Fig. 1.




FIG. 3. Cross sections for scattering of 1115-MeV electrons by C^{12} nuclei as a function of the energy E' for various scattering angles. Curves 1-3 have been calculated as in Fig. 1.

Gunnar Hylten
Nucl. Phys. A158, 225 (1970)

C | 12 | 6

METHOD

REF. NO.

70 Hy 1

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, N	ABX	200-999	C	100-999	ACT-I		4PI

999 = 1050 MEV




Fig. 1. The yield in mb per equivalent quantum as a function of the bremsstrahlung end-point energy. The lower solid curve is obtained from relation (1) with the cross section of Barber *et al.*² up to 150 MeV and then $\sigma \equiv 0$. The upper solid curve corresponds to the integrated cross-section curve in fig. 2.

²W. C. Barber, W. D. George and D. D. Reagan, Phys. Rev. 98 (1955) 73.




Fig. 2. The integrated cross section and cross section above 150 MeV as a function of the photo energy E . The cross section is the derivative of the integrated cross-section curve.

REF. K. Kayser, W. Collin, P. Filss, S. Guldbakke, G. Nolte,
H. Reich, J. O. Trier, W. Witschel
Z. Physik 239, 447 (1970)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	egf
	70 Ka 2	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, N	ABX	18-100	C	20-140	ACT-I		4PI
G, 2N	ABX	30-120	C	35-130	ACT-I		4PI

Investigation of the Photo Nuclear Reactions $^{12}\text{C}(\gamma, n)$, $^{12}\text{C}(\gamma, 2n)$, $^{39}\text{K}(\gamma, n)$ and $^{40}\text{Ca}(\gamma, np)$ up to the Meson Threshold

In order to obtain data on the photon absorption process between the giant resonance and the meson threshold the cross sections of the reactions $^{12}\text{C}(\gamma, n)^{11}\text{C}$, $^{12}\text{C}(\gamma, 2n)^{10}\text{C}$, $^{39}\text{K}(\gamma, n)^{38}\text{K}$, and $^{40}\text{Ca}(\gamma, np)^{38}\text{K}$ have been determined by the analysis of yield curves at the 140 MeV electron synchrotron of the PTB.

Though the (γ, n) cross sections grow small with increasing photon energy they are different from zero up to energies of 60 MeV and above. The cross section of the reaction $^{12}\text{C}(\gamma, 2n)$ is extremely small; its highest value amounts to 0.15% of the highest value of the $^{12}\text{C}(\gamma, n)$ reaction. The measured $^{40}\text{Ca}(\gamma, np)$ cross section is of the order predicted by the naive quasi-deuteron model. The integrated cross sections of the above reactions up to 140 MeV are 85 ± 7 , 0.90 ± 0.10 , 139 ± 16 , and 76 ± 7 MeV mb respectively.




Fig. 4. Wirkungsquerschnitte der Reaktionen $^{12}\text{C}(\gamma, n)^{11}\text{C}$ (linke Skala) und $^{12}\text{C}(\gamma, 2n)^{10}\text{C}$ (rechte Skala). Die Fehlerbalken entsprechen dem nach dem Fehlerfortpflanzungsgesetz aus der Standardabweichung der Ausbeutewerte berechneten Fehler

METHOD

REF. NO.

70 Li 1 hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	22-37	D	52-102	MAG-D		DST


RED. MATRIX ELEMENTS

FIG. 1. Reduced matrix elements squared for the 22- to 26-MeV excitation region versus q : (a) transverse and (b) longitudinal. Shown are experimental results and the theoretical results of deForest (quasi-elastic model using oscillator-model (dashed line), and plane-wave (solid line) final-state nucleon wave functions), Lewis and Walecka (single particle-hole model (dash-dotted line)), and Überall (generalized Goldhaber-Teller model (dotted line)).




FIG. 2. Experimental results for the transverse reduced matrix elements squared, together with calculated results of deForest for the indicated energy intervals (oscillator-model (dashed line) and plane-wave (solid line) final-state nucleon wave functions.)




FIG. 3. Same as Fig. 2 except for longitudinal reduced matrix elements squared [oscillator-model (dashed line) and plane-wave (solid line) final-state nucleon wave functions].

REF.

J. W. Lightbody Jr.
 Phys. Letters 33B, 129 (1970)

ELEM. SYM. A Z

C 12 6

METHOD

REF. NO.

70 Li 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABI	17-37	D	52-102	MAG-D		DST

SEE 70LI1




Fig. 1. $A(q)/Z$ versus q , calculated for an oscillator parameter $b = 1.64$ fm. Squares are data integrated to 37 MeV excitation. Circles are data corrected for missed excitation ($\omega > 37$ MeV). The Berthot et al. data point (see text) is the triangle.




Fig. 2. $B(q)/4\pi$ versus q , calculated for an oscillator parameter $b = 1.64$ fm (solid line). Also shown is $B(q)/4\pi$ before removing the contribution of spurious states to the sum (dashed line). Squares are data integrated to 37 MeV excitation, circles are data corrected for excitations $\omega > 37$ MeV. For comparison we show the Czyz et al. $E1, T = 1$ sum rule (dash-dot line), and the photoabsorption data of Wyckoff et al. (triangle).

REF. H. A. Medicus, E. M. Bowey, D. B. Gayther, B. H. Patrick,
and E. J. Winhold
Nucl. Phys. A156, 257 (1970)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		eg
70 Me 4		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PG/	ABY	18-42	C	24-42	SCD-D	1-3	112
G,NG/	ABY	21-42	C	24-42	SCD-D	1-3	112

TABLE 2

The fractions of protons and neutrons emitted to the ground and excited states of ^{11}B and ^{11}C for each of the four bremsstrahlung end-point energies

Energy of level (MeV)		J^π	Percentage of transitions into the level								Error in percentage of transitions (%)
^{11}B	^{11}C		$E_0 = 24.5$ MeV		$E_0 = 27$ MeV		$E_0 = 33$ MeV		$E_0 = 42$ MeV		
^{11}B	^{11}C		^{11}B	^{11}C	^{11}B	^{11}C	^{11}B	^{11}C	^{11}B	^{11}C	
8.57	8.11	3^-					0.1		0.4	± 30	
8.00	7.51	2^+					0.8		0.5	± 30	
7.30	6.91	2^+			0.5	0.6	0.4	0.4	0.5	± 30	
6.79	6.34	2^+			0.3	0.6	0.7	0.9	0.5	± 30	
6.74	6.48	2^-			0.4	0.5	0.8	0.4	0.7	± 30	
5.02	4.79	3^-	1.5		1.6	1.8	2.0	2.8	2.5	± 15	
4.44	4.30	2^-	0.7		0.8	0.5	1.4	1.6	1.8	± 15	
2.12	2.00	1^-	6.0	6.2	4.0	3.4	5.4	5.6	4.5	± 10	
0	0	2^-	91.8	93.8	92.4	92.6	88.4	88.3	88.6	88.3	

The final column gives the percentage error for each transition, this being independent of bremsstrahlung energy.

TABLE 3

Integrated cross sections, from threshold to the four bremsstrahlung end-point energies E_0 , for transitions to each of the various levels in ^{11}B and ^{11}C deduced from the yield ratios given in table 2

Energy of level (MeV)	$\int_0^{E_0} \left(\frac{d\sigma}{d\Omega}\right) dE$ (MeV · mb/sr) for $^{12}\text{C}(\gamma, p)^{11}\text{B}$				$\int_0^{E_0} \left(\frac{d\sigma}{d\Omega}\right) dE$ (MeV · mb/sr) for $^{12}\text{C}(\gamma, n)^{11}\text{C}$			
	$E_0 = 24.5$	$E_0 = 27$	$E_0 = 33$	$E_0 = 42$	$E_0 = 24.5$	$E_0 = 27$	$E_0 = 33$	$E_0 = 42$
8.57	0	0	0.03	0.08	8.11	0	0	0
8.00	0	0.02	0.1	0.1	7.51	0	0	0
7.30	0	0.06	0.06	0.06	6.91	0	0.03	0.03
6.79	0	0.05	0.06	0.06	6.34	0	0.04	0.04
6.74	0	0.05	0.08	0.08	6.48	0	0.03	0.03
5.02	0.1	0.1	0.2	0.3	4.79	0	0.1	0.1(5)
4.44	0.05	0.05	0.2	0.2(5)	4.30	0	0.03	0.1
2.12	0.2(5)	0.3	0.5	0.5	2.60	0.1(5)	0.1(5)	0.2(5)
Total excited states	0.4	0.6(5)	1.2(5)	1.4(5)	Total excited states	0.1(5)	0.4	0.6
Assumed ground state	5.0	6.2	7.8	9.0	Assumed ground state	2.3	3.1	4.0

REF.

A. Murakami
 J. Phys. Soc. Japan 28, 1 (1970)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	egf
70 Mu 1	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,A	ABX	12-32	C	32	EMU-D		4PI

3.3




Fig. 4. The cross section obtained for the reaction $^{12}\text{C}(\gamma, 3\alpha)$ compared with those by other authors.
 — results of the present work
 - - - results of Maikov²³⁾
 - · - results of Goward and Wilkins²⁴⁾

ELEM. SYM.	A	Z
C	12	6

METHOD						REF. NO.	egf
						70 St 2	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E _e E/	ABX	7	D	31-59	MAG-D		DST
		(7.65)					

7 = 7.65 0+

Tabelle 3. Experimentelle Parameter und Meßwerte. E₀, Θ Primärenergie, Streuwinkel im Laborsystem. q unelastischer Impulsübertrag. σ/σ_E gemessenes Verhältnis von unelastischem zu elastischem differentiellen Wirkungsquerschnitt; in Klammer ist der statistische Fehler in % angegeben. dσ/dΩ unelastischer differentieller Wirkungsquerschnitt; wegen der Fehlerangaben vgl. Text. Die Meßwerte für Si und S sind als Ergebnisse für ^{nat}Si und ^{nat}S aufzufassen (vgl. Text)

	E ₀ (MeV)	Θ (°)	q ² (fm ⁻²)	σ/σ _E (10 ⁻⁴)	dσ/dΩ (10 ⁻³³ cm ² ster.)
¹² C 7,65 MeV	59,58	117,04	0,231	14,90 (0,6)	24,69 ± 1,60
	56,94	129,02	0,231	13,79 (1,0)	13,68 ± 0,96
	54,12	141,11	0,229	14,98 (0,7)	8,15 ± 0,55
	52,75	153,15	0,231	15,24 (1,0)	6,62 ± 0,47
	51,90	165,05	0,231	15,78 (1,8)	1,16 ± 0,09
	51,18	104,98	0,145	5,80 (0,7)	28,80 ± 1,69
	47,90	117,04	0,145	5,51 (0,7)	17,19 ± 1,01
	45,48	129,02	0,145	5,97 (1,2)	11,14 ± 0,71
	43,57	141,11	0,143	5,95 (1,1)	6,09 ± 0,38
	42,54	153,15	0,145	5,82 (1,4)	2,67 ± 0,18
	37,51	104,98	0,074	1,41 (1,4)	15,38 ± 1,07
	35,08	117,04	0,073	1,33 (1,6)	9,14 ± 0,57
	33,39	129,02	0,073	1,44 (1,7)	5,89 ± 0,37
	32,04	141,11	0,073	1,39 (2,0)	3,11 ± 0,21
31,36	153,15	0,073	1,39 (3,2)	1,39 ± 0,11	
²⁴ Mg 6,44 MeV	59,01	116,94	0,232	5,53 (2,2)	28,28 ± 2,24
	55,81	129,03	0,231	5,41 (1,5)	16,65 ± 1,20
	53,64	140,95	0,231	5,65 (1,6)	9,51 ± 0,70
	52,14	153,00	0,231	5,17 (2,1)	3,95 ± 0,32
	36,98	104,96	0,074	0,59 (2,6)	25,08 ± 1,78
	34,59	116,94	0,074	0,56 (3,2)	15,09 ± 1,16
	32,87	129,03	0,074	0,58 (2,8)	9,37 ± 0,68
	31,72	140,95	0,074	0,58 (3,1)	5,10 ± 0,39
	30,86	153,00	0,074	0,68 (5,5)	2,69 ± 0,27
	Si 4,98 MeV	58,38	116,94	0,232	5,61 (1,9)
55,12		129,03	0,231	5,27 (0,9)	21,98 ± 1,34
52,87		141,11	0,231	5,58 (1,3)	12,72 ± 0,81
51,28		153,00	0,230	5,69 (1,9)	6,00 ± 0,42
49,93		104,96	0,145	2,05 (1,6)	49,60 ± 3,13
49,97		104,96	0,146	2,10 (1,4)	50,68 ± 3,09
46,77		116,94	0,146	2,23 (2,2)	33,90 ± 2,31
44,28		129,03	0,146	2,14 (2,0)	19,48 ± 1,29
42,44		141,11	0,146	2,20 (1,7)	10,98 ± 0,70
41,23		153,00	0,145	2,45 (2,1)	5,64 ± 0,38
41,37		153,00	0,146	2,48 (2,5)	5,67 ± 0,40
36,28		104,96	0,074	0,53 (4,5)	32,51 ± 2,90
36,26		104,96	0,074	0,56 (4,0)	34,35 ± 2,89
33,98		116,94	0,074	0,55 (4,3)	21,28 ± 1,83
32,17		129,03	0,074	0,52 (4,0)	11,98 ± 1,00
30,87		141,11	0,073	0,59 (5,3)	7,54 ± 0,73
29,96	153,00	0,073	0,66 (5,7)	3,89 ± 0,39	
30,09	153,00	0,074	0,60 (5,0)	3,52 ± 0,33	
Si 6,69 MeV	58,38	116,94	0,225	0,48 (25)	3,30 ± 1,00
S 3,78 MeV	59,19	104,96	0,210	0,32 (37)	5,17 ± 3,36
	58,80	104,96	0,209	0,27 (45)	4,40 ± 3,30
⁴⁰ Ca 3,35 MeV	53,98	140,95	0,250	0,92 (9)	3,03 ± 1,21

over




Fig. 2. Meßergebnisse für den 7,65 MeV-Übergang in ^{12}C als Funktion von q^2 . Meßpunkte und eingezeichnete Kurve gelten für eine Auswertung mit Faktoren f_c und x_1, x_2 -Werten nach Modell I (Fig. 1). Man beachte den unterdrückten Nullpunkt der Ordinatenkala

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	15-30	D	250	MAG-D		DST

The longitudinal and transverse form-factor spectra for electroexcitation of the ^{12}C giant resonance were determined separately. Results are presented for an excitation energy range of from 15 to 30 MeV and for a momentum-transfer range of $0.84\text{--}1.56\text{ F}^{-1}$. We have found levels with the following excitation energies (MeV), spins, and parities: 18.6 (3^-), 19.6 (4^-), 20.0 (2^+), 20.6 (3^+), 21.6 (3^-), 22.0 (1^-), 22.7 (1^-), and 23.8 (1^-). We have also found new evidence of the spin-isospin mode for the 22.7-MeV (1^-) excitation.




FIG. 1. The longitudinal form factors $|W_L(q, \omega)|^2$ for electroexcitation in ^{12}C in the excitation energy range from 15 to 30 MeV corresponding to the momentum transfers of $0.84, 1.04, 1.22,$ and 1.56 F^{-1} . The widths of elastic peaks for these spectra are about 650 keV. The form-factor area for the peaks other than the giant dipole resonance around 22 MeV were determined by a fitting procedure using the shape of the elastic peak. The arrows show the positions of the form-factor peaks. The levels for which spins and parities are assigned tentatively from the present study are shown in the parentheses.




FIG. 2. The same as for Fig. 1 except that the transverse form factors $|W_T(q, \omega)|^2$ are shown.

REF.

S.G. Tonapetyan, O.G. Konovalov, A.I. Derebchinskii, A.A. Zybalov,
 V.M. Khvorostyan, N.V. Goncharov & V.A. Gol'dshtein
 ZhETF Pis. Red. 11, 165 (1970)
 JETP Letters 11, 101 (1970)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
70 To 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
SG,P	NOX	680-840	C	700-930	MAG-D		41

P-POLARIZATION

The table lists the preliminary results for the nuclei ${}^7_3\text{Li}$ and ${}^{12}_6\text{C}$.

Nucleus	E_{mes} MeV	E_{eff} MeV	ΔE_{p} MeV	P MeV/c	ΔP MeV/c	Polarization	Kinematic region
Li^7	700	-	-	755	23	-0.15 ± 0.19	I
	810			790	25	-0.15 ± 0.22	
C^{12}	700			755	23	$+0.05 \pm 0.31$	
	810	-	-	790	24	-0.05 ± 0.20	
	930			840	25	0.08 ± 0.43	
Li^7	700	650	33	618	18	-0.76 ± 0.20	
	810	715	38	655	20	-0.31 ± 0.19	
C^{12}	700	650	33	618	18	-0.71 ± 0.21	
	810	715	38	655	20	-0.35 ± 0.26	
	930	840	43	754	23	$+0.48 \pm 0.35$	

Region I - no real mesons possible.

" II - real mesons possible.

REF. A. V. Vysotskaya and N. G. Afanas'ev
 Yad. Fiz. 11, 942 (1970)
 Sov. J. Nucl. Phys. 11, 523 (1970)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	70 Vy 1

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, P	ABX	31-200	D	100-200	MAG-D	15-60	DST




FIG. 2. Proton spectra measured at different angles for electrons with $E_1 = 200$ MeV.




FIG. 3

FIG. 4

FIG. 3. Angular distributions for different proton energies ($E_1 = 200$ MeV).

FIG. 4. Excitation functions for different proton energies, measured for 100-, 150-, and 200-MeV electrons at the proton emission angle $\theta_p = 40^\circ$.

REF.

C. P. Wu, F. W. K. Firk, and B. L. Berman
Nucl. Instr. & Methods 79, 346 (1970)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	egf
70 Wu 3	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,MUT	ABX	20- 30	C	50	RSP-D		4PI




Fig. 1. A comparison between the present observed total photon absorption cross section of carbon and the results of Burgov et al. ³⁾.

- 3) N.A. Burgov, G.V. Danilyan, B.S. Dolbilkin, L.E. Lazareva and F.A. Nikolaev, Zh. Eksperim. i. Teor. Fiz 45, 1693 (1963).

REF. J. Ahrens, H. Borchert, H. B. Eppler, H. Gimm, H. Gundrum, P. Riehn,
 G. Sita Ram, A. Zieger, and B. Ziegler
 Elba-71, Tagungsbericht Elektronen Beschleuniger Arbeits Gruppen
 (Sept. 1971) Justus Liebig-Universität Giessen. p.359

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	71 Ah 1 hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	THR-150	C	10-150	MGC-D		4PI




fig. 4

REF.

Yu. P. Antuf'ev, V. L. Agranovich, V. B. Ganenko, V. S. Kuz'menko,
I. I. Miroshnichenko, and P. V. Sorokin
Yad. Fiz. 13, 473 (1971); Sov. J. Nucl. Phys. 13, 265 (1971)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
71 An 1		

METHOD

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	SPC	49-999	C	700,999	TEL-D	25-400	DST
G,D	SPC	55-999	C	700,999	TEL-D	25-400	DST

999=1.2 GEV, REL D/P




FIG. 1. Energy spectra of secondary particles. Points: ○ - protons, $\theta = 30^\circ$; ● - deuterons, $\theta = 30^\circ$; Δ - protons, $\theta = 120^\circ$; ▲ - deuterons, $\theta = 120^\circ$. Solid curves - calculation with Eq. (1); dashed curves - calculation with the quasideuteron model, Eq. (2); dotted curve - calculation with the photomeson model.

Yield of protons 30-400 MeV, deuterons 30-200 MeV.




FIG. 2. Ratios of cross sections for photoproduction of deuterons and protons σ_d/σ_p as a function of their kinetic energy. Points: ○ $\theta = 30^\circ$, Δ $\theta = 120^\circ$. Solid curve - calculation on the assumption of the photomeson model for production of deuterons and protons.

Table I. Values of the parameter τ , MeV

Target	$E_0 = 700$ MeV										$E_0 = 1200$ MeV					
	Protons					Deuterons					Protons			Deuterons		
	40°	50°	90°	100°	120°	40°	50°	90°	100°	120°	30°	50°	120°	30°	50°	120°
Li	46	42	34	30	27	28	24	22	21	20	45	28		27	24	
Be	48	43	36	30	27	28	26	24	22	19	45	28		27	24	
C	50	44	38	30	28	34	33	29	23	19	60	48	37	34	22	
Si		43			28		27			22		46	35	28	25	
Cu											45	29	27	24		
Ta					28					21	45	34	27	24		
Pb										51	29	36	22			




FIG. 3. Ratio of deuteron and proton yields as a function of momentum at various angles. $E_0 = 700$ MeV; points: ● $\theta = 40^\circ$, Δ $\theta = 60^\circ$, □ $\theta = 80^\circ$, ▽ $\theta = 100^\circ$, ○ $\theta = 120^\circ$.




FIG. 4. The ratios N_d/N_p as a function of target-nucleus mass number A at an angle $\theta = 60^\circ$ for $E_0 = 1200$ MeV. Solid curve - $A^{0.13}$.

The measured secondary-particle spectra for kinetic energies $T > 80$ MeV are well described by the expression

$$d^2\sigma / d\Omega dTQ = \text{const } T \exp(-T/\tau), \quad (1)$$

which is identical to the formula for the evaporation process.⁽¹⁾ In Table I we have given the values of the parameter τ for the nuclei studied, at various angles. The accuracy in determination of τ is about 10%.

REF. Yu. P. Antuf ev, -V.L. Agranovich, V.B. Ganenko, V.S. Kuz menko,
I.I. Miroshnichenko, and P.V. Sorokin
Yad. Fiz. 14, 898 (1971)
Sov. J. Nucl. Phys. 14, 502 (1972)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
71 An 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XD	ABX	115-999	C	620,999	MAG-D		DST

999 = 1.14 GEV




FIG. 1. Angular distributions of deuterons in (γ , d) reactions in nuclei for $E_0 = 620$ MeV (a-c) and $E_0 = 1140$ MeV (d-k). The statistical errors are shown. a-g- angular distributions of deuterons with energies of 90 MeV, h-k-with energy 160 MeV.




FIG. 2. Angular distributions of 160-MeV deuterons normalized at $\theta = 40^\circ$ for C^{12} (Δ) and Ta^{181} (\circ) for $E_0 = 1140$ MeV.




FIG. 3. Angular distributions of 90-MeV deuterons normalized at $\theta = 40^\circ$ for $E_0 = 1140$ MeV (\circ) and $E_0 = 620$ MeV (Δ).

REF.

J. C. Bergstrom, H. Crannell, E. J. Kline, J. T. O'Brien,
 J. W. Lightbody, Jr., and S. P. Fivozinsky
 Phys. Rev. C4, 1514 (1971)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	71 Be 2

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	3-44	D	77-106	MAG-D		DST

Figures also given for: 77 MeV, 75°
 55.4 MeV, 145.7°
 81 MeV, 145.7°

964




FIG. 11. The ratio of the ^{13}C to ^{12}C form factors (squared) in the 10-29-MeV region (circles) and the 17-29-MeV region (boxes). The form factors were obtained by integrating the differential form factors over the indicated energy regions. The errors reflect counting statistics and the elastic peak tail uncertainty (see text). The tail uncertainty tends to cancel in the formation of the ratios.




Fig. 10. The form factor (squared) for ^{13}C . The differential form factors have been integrated over the region 10-17, 17-21.7, 21.7-26.7, and 26.7-31 MeV. Results are presented for 75 and 145.7° scattering angles. The errors reflect both counting statistics and the uncertainty in the elastic peak tail (see text).





FIG. 7. Similar to Fig. 2, but spectrum is for ^{12}C and incident energy is 106 MeV.

METHOD				REF. NO.			
				71 Bu 2		egf	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	ABX	5-85	D	500	MAG-D		51

TABLE I.

PROT P=25,100 MEV/C

		$p = 25 \text{ MeV}/c$		$p = 100 \text{ MeV}/c$			
		σ_{exp}	σ_{th}	$\sigma_{\text{exp}}/\sigma_{\text{th}}$	σ_{exp}	σ_{th}	$\sigma_{\text{exp}}/\sigma_{\text{th}}$
^{12}C	1p	0.25	0.65	0.35	0.83	3.15	0.38
	1s	0.38	1.28	0.30	0.31	0.87	0.38
^{40}Ca	1d _{1/2}		0.03		0.29	2.62	0.11
	2s _{1/2}	1.41	7.28	0.19		0.86	
	1d _{3/2}	(0.28)	0.05		0.76	3.94	0.19
	1p	(0.65)			(0.93)		
	1s	(0.63)			(0.67)		

Cross-sections are given in units $10^{-28} \text{ (cm}^2\text{MeV/}\mu\text{r}^2\text{)}$.Fig. 1. - $^{12}\text{C}(\alpha, p)$, $\epsilon = 500 \text{ MeV}$, $T' = 36 \text{ MeV}$: a) $p = 100 \text{ MeV}/c$, b) $p = 35 \text{ MeV}/c$.

ELEM. SYM.	A	Z
C	12	6
REF. NO.	71 Co 2	
	egf	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XN	ABI	36-64	C	10-64	BF3-I		4PI

FAST N YIELD




Fig. 2.




Fig. 3.

Fig. 2. - Experimental photoneutron cross-sections integrated over photon energy between 36 and 64 MeV and divided by NZ/A are plotted as a function of the mass number. Black dots are total cross-sections not corrected for neutron multiplicity; open circles represent fast neutron cross-sections (see text). The dashed lines are drawn only to guide the eye.

Fig. 3. - The ratio between «fast» and total photoneutron integrated cross-sections as a function of the mass number A . The solid line represents a fit of the ratios calculated for some nuclei by taking into account the theoretical neutron energy spectra given by GABRIEL and ALSMILLER (*) and the efficiencies of our detector (see Fig. 1).

REF. V. Di Napoli, D. Margadonna, F. Salvetti, H.G. De Carvalho,
J.B. Martins
Nucl. Inst. Meth. 93, 77 (1971)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

71 Di 5

hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, BE7	ABY	26-999	C	300-999	ACT-I		4PI

The use of the $^{19}\text{F}(\gamma, n)^{18}\text{F}$, $^{27}\text{Al}(\gamma, x)^{24}\text{Na}$, $^{197}\text{Au}(\gamma, n)^{196}\text{Au}$, and $^{12}\text{C}(\gamma, x)^7\text{Be}$ reactions as absolute monitors for high-energy, high-intensity bremsstrahlung beams is discussed. The cross sections per equivalent quantum and the absolute cross sections, in the energy range 300-1000 MeV, are reported for these reactions. $^{19}\text{F}(\gamma, n)^{18}\text{F}$, $^{27}\text{Al}(\gamma, x)^{24}\text{Na}$, and $^{12}\text{C}(\gamma, x)^7\text{Be}$ reactions are proposed as very simple and suitable systems for monitoring purposes.

999 = 1 GEV




Fig. 4. $^{12}\text{C} \rightarrow ^7\text{Be}$ reaction cross sections per e.q. Circles: present work; triangles: Frascati²⁾; squares: Orsay¹⁾. The straight line is a least-squares fit of all values.

TABLE 2

Energy (MeV)	$\sigma_q(\text{mb})$			
	$^{19}\text{F} \rightarrow ^{18}\text{F}$	$^{27}\text{Al} \rightarrow ^{24}\text{Na}$	$^{197}\text{Au} \rightarrow ^{196}\text{Au}^*$	$^{12}\text{C} \rightarrow ^7\text{Be}^*$
260			270 = 14	
300	5.90 ± 0.20	0.37 ± 0.01	258 = 13	0.37 ± 0.01
320		0.39 ± 0.01		
350	5.90 ± 0.20	0.39 ± 0.01		0.38 ± 0.01
380	5.95 ± 0.20	0.41 ± 0.01		
400				0.42 ± 0.01
420		0.45 ± 0.01		
450	5.95 ± 0.20	0.48 ± 0.01	249 ± 12	0.44 ± 0.02
500	6.65 ± 0.20	0.49 ± 0.01		
550	6.65 ± 0.20	0.50 ± 0.02		0.47 ± 0.02
600	6.80 ± 0.20	0.52 ± 0.02		
650		0.52 ± 0.02	266 = 13	0.47 ± 0.02
700	7.20 ± 0.30	0.56 ± 0.02		
750		0.55 ± 0.02		
850	7.60 ± 0.30	0.60 ± 0.02	246 = 12	0.49 ± 0.03
900		0.59 ± 0.02		
1000	7.10 ± 0.30	0.66 ± 0.02	249 = 12	0.51 ± 0.03

* The values given in the last two columns are, in most cases, an average of two or more measurements (see figs. 3 and 4).

ELEM. SYM.	A	Z
C	12	6
METHOD	REF. NO.	
	71 Eg 1	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, XP	ABY	56-130	D	130	MAG-D		DST

The angular and the energy distributions of the secondary protons in (γp) and (ep) reactions on C^{12} nucleus for 130 MeV excitation energy are studied. The experimental results are compared with the quasideuteron and one particle absorption models of γ -quanta in the nucleus. For both cases the parameters of the proton wave function in the nucleus are estimated.




Рис. 2. Угловое распределение протонов из реакции (γp) и (ep) на C^{12} для энергии возбуждения до 130 Мэв. \circ — экспериментальные точки при энергии: $T_p = 40$ Мэв, \triangle — при $T_p = 63$ Мэв. Сплошные кривые — результаты расчетов по квазидейтонной модели Дедрива [19] для $T_p = 40$ Мэв (верхняя) и $T_p = 63$ Мэв (нижняя). Кривые нормированы по экспериментальным точкам при $\theta_p = 40^\circ$ Мэв. Пунктирные кривые — те же результаты после смещения на 15° в области малых углов (см. в тексте).




Рис. 1. Энергетический спектр протонов (γp) и (ep) на ядре C^{12} для энергии возбуждения до 130 Мэв. \circ — экспериментальные точки для $\theta_p = 30^\circ$, \triangle — для $\theta_p = 60^\circ$. Сплошная кривая — результаты расчетов по квазидейтонной модели Дедрива [19] при $\theta_p = 30^\circ$, пунктирные кривые — при $\theta_p = 60^\circ$. Экспериментальные и теоретические данные произвольно нормированы при энергии протонов 50 Мэв.

Таблица
Сечение образования протонов на ядре C^{12} фотонами и электронами с энергией до 130 Мэв, в мкбарн:

$T_p = 40$ Мэв		$T_p = 63$ Мэв		$\theta_p = 30^\circ$		$\theta_p = 60^\circ$	
$d^2\sigma/d\Omega_p E_p Q$	θ_p°	$d^2\sigma/d\Omega_p E_p Q$	θ_p°	$d^2\sigma/d\Omega_p E_p Q$	$T_p, \text{ Мэв}$	$d^2\sigma/d\Omega_p E_p Q$	$T_p, \text{ Мэв}$
2.02 ± 0.08	25	0.815 ± 0.065	25	2.08 ± 0.05	40	1.65 ± 0.07	40
2.08 ± 0.05	30	0.79 ± 0.06	30	1.25 ± 0.07	50	0.925 ± 0.05	50
1.97 ± 0.05	35	0.78 ± 0.08	35	0.705 ± 0.065	63	0.41 ± 0.055	63
2.16 ± 0.11	40	0.694 ± 0.065	40	0.55 ± 0.04	72	0.24 ± 0.015	72
2.02 ± 0.10	50	0.56 ± 0.08	50	0.23 ± 0.02	82	0.11 ± 0.04	82
1.65 ± 0.07	50	0.41 ± 0.055	60	0.21 ± 0.01	92	0.06 ± 0.01	92
1.26 ± 0.06	70	0.300 ± 0.04	70	0.144 ± 0.01	100	—	100
1.01 ± 0.06	80	—	80	—	—	—	—

METHOD

REF. NO.
71 Eg 2
egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, XP	ABY	56-250	C	100-250	MAG-D		30

EXCITATION FUNCTION OF (γp) AND (ep) REACTIONS ON C^{12} AT PHOTON AND ELECTRON ENERGIES UP TO 100+250 MeV

K. M. EGIAN, G. L. BOCEK, V. M. KULIBABA

The results of excitation function measurements are given for protons, produced by photons and electrons with energies of 100+250 Mev.

In the kinematically allowable region for π -photoproduction on nucleon nucleons ($E_p \approx 40$ MeV, $\nu_p = 30^\circ$), the slope of excitation function curve could be explained assuming that protons are produced in two processes: γ -quanta absorption by inter-nuclear quasideuterons with subsequent disintegration (the main process) and the photoproduction of π -mesons. In the kinematically forbidden region, experimental data are in good agreement with calculations by the quasideuteron model.




Рис. 2. Функция возбуждения реакций (γp) и (ep) на углероде при энергии протонов 83 Mev под углом 30° в л. с. Кривая рассчитана по квазидейтонной модели и нормирована при $(E_\gamma)_{max} = 200$ Mev.




Рис. 1. Функция возбуждения реакций (γp) и (ep) на углероде при энергии протонов на 40 Mev под углом 30° в л. с. Сплошная кривая рассчитана по квазидейтонной, пунктирная — по одночастичной, штрих пунктирная — по фотомезонной модели поглощения γ -квантов. Порядок две кривые нормированы при $(E_\gamma)_{max} = 100$ Mev. Третья кривая нормирована при $(E_\gamma)_{max} = 258$ Mev по разности значений экспериментального сечения и нормированного сечения, рассчитанного по квазидейтонной модели.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
SG,P	NOX	16-800	C	650-800	MAG-D		UKN

POLARIZED PROTONS

E_{max} MeV	E_{eff} MeV	$\pm \Delta E$ MeV	Momen- tum p. MeV/c	MeV/c	Polariza- tion P
650	549	27	539	19	-0.25 ± 0.18
700	598	29	578	21	-0.30 ± 0.16
750	648	33	616	22	-0.47 ± 0.17
800	715	38	667	23.5	-0.45 ± 0.13



Proton polarization from C^{12} as a function of γ -ray energy corresponding to the kinematics of single-pion photoproduction from free nucleons. Points: \circ —experimental data obtained in the present work; \square —polarization values calculated with Eq. (3) with the value of P_{π^-} from Benevntano et al. [8]; Δ —Theoretical polarization value for the value of P_{π^-} from Kenemuth and Stein. [7]

⁷ J.R. Kenemuth & P.C. Stein,
 Phys. Rev. 129, 2259 (1963).

⁸ M. Benevntano et al., Proc. XV
 Intern. Conf. on High Energy Physics,
 Kiev, 1970.

REF.

N.V. Goncharov, A.I. Derebchinskii, O.P. Konovalov, S.G. Tonapetyan,
and V.M. Khvorostyan
Yad. Fiz. 14, 31 (1971)
Sov. J. Nucl. Phys. 14, 18 (1972)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

71 Go 2

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, PI+	RLY	150-500	C	500	CCH-D		DST

PI+/PI+ YIELD RATIO

Measurements are reported of the relative yield of π^+ mesons and the π^+/π^- yield ratio for mesons with energy 40 ± 10 MeV emitted in the angular range $\theta_{\text{lab}} = 50-160^\circ$ in photon-induced reactions with $E_{\text{max}} = 500$ MeV with light and medium nuclei. The charged π -meson detector was a 34-cm Freon bubble chamber with a tube for the beam. The π^+/π^- yield ratio for He^4 , Li^7 , C^{12} , Si^{28} , S^{32} , Ca^{40} , and Nb^{93} was found to be respectively 0.94 ± 0.14 , 2.15 ± 0.31 , 1.22 ± 0.21 , 1.25 ± 0.15 , 1.0 ± 0.13 , 1.11 ± 0.13 , and 1.53 ± 0.25 . It was established that the π^+ -meson yield follows a $ZA^{-1/3}$ law.





FIG. 2. Relative yield of π^+ mesons per nucleus as a function of mass number A.

FIG. 3. π^+/π^- yield ratio as a function of mass number A.



REF.


B.I. Goryachev, B.S. Ishkhanov, and V.G. Shevchenko
 Proceedings of the Second Symposium on the Problems
 of Nuclear Physics, Novosibirsk, USSR, June 1970
 (Kolybasov, V.M., Ed., Izdatel'stvo Nauka, Moscow 1971),
 pp.362-78

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	
71 Go 3	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
C, γ	ABX	19- 24	C	19- 24	MOD-I		4PI



Фиг. 1. Сечение реакции $C^{12}(\gamma, n)$.

Т а б л и ц а 1

Энергии резонансов (Мэв)

C^{12}	O^{16}	Ca^{40}
20,0	18,9	16,9
20,4	17,3	17,2
20,7	17,9	17,8
21,1	18,4	18,2
21,4	18,9	18,5
21,9	19,3	18,8
22,4	19,5	19,0
22,9	20,3	19,3
23,3	20,9	19,6
23,7	21,3	19,9
	21,8	20,4
	22,2	21,0
	22,5	21,2
	22,9	21,5
	23,3	22,0
	23,8	22,4
	24,2	22,6
	24,6	23,3
		24,2
		24,6
		25,5

REF.

I.A. Grishaev, A.N. Krinitsyn, N.I. Lapin, V.I. Nikiforov,
G.D. Pugachev, and B.I. Shramenko
Yad. Fiz. 14, 35 (1971)
Sov. J. Nucl. Phys. 14, 20 (1972)

ELEM. SYM. A Z

C 12 6

METHOD

REF. NO.

71 Gr 2

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABY	150-560		560	EMU-D		DST
G,PI-	ABY	150-560		560	EMU-D		DST

PI-/PI+ YIELD RATIO

Cross section for photoproduction of π^- and π^+ mesons for
 $E_\gamma = 560$ MeV

Nucleus	10^{22} cm ² /sr-MeV-quant						
	$\theta = 60^\circ$			$\theta = 120^\circ$			
	Data of ref. 1, T = 33 MeV	Our data, T = 40 MeV		Our data $\pm \sigma$			
		π^+	π^-	T = 40 MeV		T = 60 MeV	
	π^+	π^-	π^+	π^-	π^+	π^-	
C	21.4 ± 0.5	20.6 ± 1.5	26.2 ± 2	27.5 ± 2.1	36.8 ± 2.8	21.6 ± 2.1	26.4 ± 2.6
Al	42.4 ± 1.0	38.5 ± 3	47.3 ± 4	57 ± 4	76 ± 5.4	40.2 ± 3.5	52 ± 4.7
Cu	78.8 ± 1.8	71.6 ± 3.6	96 ± 8.8	109 ± 7.5	152 ± 10.8	115 ± 8.6	135 ± 10.7
Pb				208 ± 18.5	369 ± 27	170 ± 19	260 ± 26.5




FIG. 1. Total yield of charged mesons as a function of atomic weight. The solid straight line is the experimental dependence, and the dashed straight line is the $A^{2/3}$ law. a- $\theta = 60^\circ$, T = 40 MeV; b- $\theta = 120^\circ$. Points: O-T = 40 MeV, Δ -T = 65 MeV.




FIG. 3. π^-/π^+ yield ratio as a function of N/Z. a- $\theta = 60^\circ$, T = 40 MeV; b- $\theta = 120^\circ$, T = 40 MeV; c- $\theta = 120^\circ$, T = 65 MeV

REF. B. S. Ishkhanov, I. M. Kapitonov, I. M. Piskarev, V. G. Shevchenko
 Yad. Fiz. 14, 253 (1971)
 Sov. J. Nucl. Phys. 14, 142 (1972)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
71 Is 3		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	19-25	C	19-25	BF3-I		4PI

412




FIG. 1. Cross section for the reaction $C^{12}(\gamma, n)C^{11}$.

Comparison of the location of resonances (in MeV) obtained in the present work and the locations of the more reliable resonances observed in other work

Data of present work	Analyzed data of other groups	Data of Cook et al. [1]	Data of Furk et al. [1]	Data of present work	Analyzed data of other groups	Data of Cook et al. [1]	Data of Furk et al. [1]
19.2	17.5-19.0			21.40	21.3-21.4	(21.30)	
19.7	19.3	(19.20)		(21.70)	21.7	(21.65)	21.7
20.1	19.8-19.9	19.80		21.95	22.0	22.15	22.0
20.45	20.1-20.2	20.10		22.5		22.75	
20.75	20.5-20.6	20.33		23.0			23.1
21.05	20.8-21.0	(20.60)		23.3		23.60	
		(20.90)		23.7			(23.7)

REF.

H. G. Miller, W. Buss and J. A. Rawlins
 Nucl. Phys. A163, 637 (1971)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

71 Mi 1

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	70-90	C	64-128	TOF-D	50-90	DST




Fig. 3. Reduced neutron yield and cross section, $E_a \approx 61$ MeV




Fig. 2. Reduced neutron yield for $E_a \approx 49$ MeV and (γ, n) cross section from ^{12}C as obtained by the method of least structure.




Fig. 4. Reduced neutron yield and cross section, $E_n \approx 69$ MeV.

2 K. G. Dedrick, Phys. Rev. 100 (1955) 58.

4 V. I. Mamasakhlisov and R. I. Dzhibuti, JETP (Sov. Phys.) 14 (1962) 1066.




Fig. 7. Comparison of the $^{12}\text{C}(\alpha, n)^{11}\text{C}$ cross section at $\theta = 90^\circ$ with the predictions of quasi-alpha ⁴⁾ and quasi-deuteron ²⁾ models.




Fig. 6. Comparison of the neutron angular distribution with theoretical predictions of quasi-alpha ⁴⁾ and quasi-deuteron ²⁾ models.

E. J. Moniz, I. Sick, R. R. Whitney, J. R. Ficenec, R. G. Kephart
and W. P. Trower
Phys. Rev. Letters 26, 445 (1971)

C 12 6

METHOD

REF. NO.

71 Mo 3

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	0-240	D	500	MAG-D		60




Fig. 1. Cross sections $d^2\sigma/d\Omega d\epsilon$ versus electron energy loss $\omega = \epsilon_1 - \epsilon_2$ for inelastic scattering of 500-MeV electrons at 60° from (a) carbon, (b) nickel, and (c) lead. Solid lines are the results of the Fermi-gas calculation with the nuclear parameters indicated on the figures.

Table I. Nuclear Fermi momentum k_F and average nucleon interaction energy $\bar{\epsilon}$ determined by least-squares fit of theory to quasielastic peak.

Nucleus	k_F (MeV/c) ^a	$\bar{\epsilon}$ (MeV) ^b
${}^3\text{Li}^6$	160	17
${}^6\text{C}^{12}$	221	25
${}^{12}\text{Mg}^{24}$	235	32
${}^{20}\text{Ca}^{40}$	251	29
${}^{28}\text{Ni}^{58.7}$	260	36
${}^{39}\text{Y}^{89}$	254	30
${}^{50}\text{Sn}^{118.7}$	260	42
${}^{73}\text{Ta}^{181}$	265	42
${}^{82}\text{Pb}^{208}$	265	44

^aThe fitting uncertainty in these numbers is approximately ± 5 MeV/c.

^bThe fitting uncertainty in these numbers is approximately ± 3 MeV. Simple estimates for $\bar{\epsilon}$ give numbers in reasonable agreement with those in the table.

REF. A. Nakada, Y. Torizuka, and Y. Horikawa
Erratum: Phys. Rev. Letters 27, 1102 (1971)

ELEM. SYM.	A	Z
C	12	6

See: Phys. Rev. Letters 27, 745 (1971)

METHOD	PL. NO.	hmg
	71 Na 1	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	9-17	D	250	MAG-D		DST

ERRATUM 71 NA1

Erratum for Reference 71 Na 1.

See: Erratum - Phys. Rev. Letters 27, 1102 (1971)

METHOD					REF. NO.		
					71 Na 1		hmg
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	4- 17	D	250	MAG-D		DST

LEVELS 14.1, 4.43




FIG. 1. The inelastic electron scattering spectrum for excitation energies of 9-17 MeV. The 14.1-MeV peak is seen superimposed on the inelastic continuum and the background was estimated as indicated with the dashed line.




FIG. 2. The experimental form factors for elastic, 4.43-MeV (2⁺), and 14.1-MeV (4⁺) states in ¹²C. The form factors calculated from the Nilsson orbits with suitable parameters are shown along with the experimental data.

TABLE I. The values of ¹²C parameters as calculated from the orbits which reproduce the experimental data.

$\langle r^2 \rangle^{1/2}$ (F)	2.42
$\langle r^4 \rangle^{1/4}$ (F)	2.68
Q_0^a (eF ²)	-20.1
Q_{40}^b (eF ⁴)	+21.5

^a Q_0 is defined by $Q_0 = (16\pi/5)^{1/2} \langle \chi | \sum r^2 Y_{20} | \chi \rangle$.

^b $Q_{40} = \langle \chi | \sum r^4 Y_{40} | \chi \rangle$.

V.I. Noga, Yu. N. Ranyuk, P.V. Sorokin, V.A. Tkachenko
Ukr. Fiz. Zh. 16, 1850 (1971)

C 12 6

METHOD				REF. NO.			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,Be7	ABY	THR-999	C	80-999	ACT-I		4PI

999=1.2 GEV




Рис. 3. Энергетическая зависимость сечений и выходов реакций: \circ — экспериментальные зависимости выхода от E_{max} ; сплошная кривая — σ_x — функция возбуждения, вычисленная по способу наименьших квадратов по экспериментальным данным; пунктирная — σ_q — выход, соответствующий функции возбуждения.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABY	18-68	C	10-68	ACT-I		4PI

Nippon Kagaku Zasshi. 92, 164~168(1971)

The Yields of Radioactivities Induced by (γ, n) Reactions with Bremsstrahlung up to 68 MeV

by Tatsuya SAITO


The (γ, n) yields of 12 target nuclides have been measured at 10, 13, 16, 30, 45 and 68 MeV bremsstrahlung by observing the induced activities.

The energy dependence of the yields has been investigated extensively in the same way as in the previous work at 20 MeV bremsstrahlung.

In the case of heavy nuclides, the yields rise greatly as a function of maximum bombarding energy up to 20 MeV, and rise gradually from 20 MeV up to 68 MeV. However, in the case of light nuclides, the yields rise greatly up to 30 MeV, because the neutron separation energies of light ones are larger than those of heavy ones, and the bremsstrahlung spectrum covers the giant resonance and so the yields rise gradually from 30 MeV up to 68 MeV.


The yields have approximately been estimated from the parameter of the giant resonance, that is the peak cross section and the half width, in order to compare with the experimental data. As a result, the experimental data of light nuclides and heavy ones are nearly in agreement with the estimated data of Nathans et al., Johns et al. and Montalbetti et al., but those of medium weight ones are relatively lower values.

Department of Chemistry, Faculty of Science, Tohoku University;
Katahira-cho, Sendai-shi, Japan



射線照射最大エネルギー

I: $^{197}\text{Au}(\gamma, n)^{196}\text{Au}$, II: $^{127}\text{I}(\gamma, n)^{126}\text{I}$
 III: $^{55}\text{Mn}(\gamma, n)^{54}\text{Mn}$, IV: $^{23}\text{Na}(\gamma, n)^{22}\text{Na}$
 V: $^{16}\text{O}(\gamma, n)^{15}\text{O}$, VI: $^{12}\text{C}(\gamma, n)^{11}\text{C}$

図3 (γ, n) 反応の収率

原子番号

●: 実験値, ⊕: Johnsら,
 ○: Nathansら, ○: Montalbettiら

図4 (γ, n) 反応の収率の比較

REF.

Y. M. Shin, C. F. Wong, H. S. Caplan
Nucl. Phys. A166, 162 (1971)

ELEM. SYM.	A	Z
C	12	6

METHOD			REF. NO.				
			71 Sh 1				egf
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,P	SPC	16-29	D	43	MAG-D	3-13	DST





Fig. 1. Proton energy spectra from the reactions $^{12}\text{C}(e, e'p)^{11}\text{B}$ and $^{13}\text{C}(e, e'p)^{12}\text{B}$ with 43 MeV electrons.




Fig. 2. Differential cross sections of the $^{12}\text{C}(\gamma, p)^{11}\text{B}$ and $^{13}\text{C}(\gamma, p)^{12}\text{B}$ reactions at 90° with $E = 43$ MeV. The solid line is from the results of Allas *et al.*




Fig. 3. Energy dependence of coefficients in the angular distribution of the reaction $^{12}\text{C}(e, e'p)^{11}\text{B}$. The broken line is the result of (p, γ) experiment by Allas *et al.*

REF.

H. J. Von Eyss, H. Schier, and B. Schoch
 Elba-71, Tagungsbericht Elektronen Beschleuniger Arbeits Gruppen
 (Sept. 1971) Justus Liebig-Universität Giessen. p.391

ELEM. SYM. A Z

C

12

6

METHOD


REF. NO.

71 Vo 1

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,N	ABX	THR-266	C	150-266	TOF-D		90

See over for figure.



Figur 3: Differentieller Neutronenproduktions-Wirkungsquerschnitt als Funktion der Massenzahl A für einige Neutronenenergien

METHOD

REF. NO. 71 Ya 2 hmg

[Page 1 of 3]

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	15-100	D	88-250	MAG-D		DST

J-PI, FMF, 10 LEVEL.

The longitudinal and transverse form factors were determined separately as functions of the excitation energy from 15 to 40 MeV in the momentum-transfer range $0.75-1.56 \text{ F}^{-1}$. For some of the form-factor spectra the measurements were extended up to an excitation energy of 100 MeV and the results were compared with the quasielastic model presented by de Forest. The q dependence of the experimental form factors for levels were compared with the predictions of the particle-hole model and with the intermediate-coupling model. We have then found the levels with the following excitation energies (MeV), spins, and parities: 19.1 (1^-), 18.6 (3^-), 19.6 (4^-), 20.0 (2^-), 20.6 (3^-), 21.6 (3^-), 22.0 (1^-), 22.7 (1^-), and 23.3 (1^-). We have also found new evidence of the possibility of the spin-isospin mode for the 22.7-MeV (1^-) excitation.

TABLE I. The theoretical and observed energy levels (MeV) for $J^\pi = 1^-, 2^-, 3^-$, and 4^- , $T=1$ in ^{12}C .

J^π	Donnelly (Ref. 28)	Main configuration	Present work
1^-	19.52	$(2s_{1/2}, 1p_{3/2}^{-1})$	18.1
	23.09	$(1d_{3/2}, 1p_{1/2}^{-1})$	22.0, 23.8, 25.5
	24.89	$(1d_{5/2}, 1p_{3/2}^{-1})$	22.7, 25.5
	35.55	$(1p_{1/2}, 1s_{1/2}^{-1})$	
2^-	18.80	$(2s_{1/2}, 1p_{3/2}^{-1})$	16.6
	20.60	$(1d_{3/2}, 1p_{3/2}^{-1})$	19.3
	23.83	$(1d_{5/2}, 1p_{3/2}^{-1})$	
3^-	19.24	$(1d_{3/2}, 1p_{3/2}^{-1})$	18.6
	25.09	$(1d_{5/2}, 1p_{3/2}^{-1})$	25.5
4^-	20.17	$(1d_{3/2}, 1p_{3/2}^{-1})$	19.6




FIG. 7. The transverse form factor for the 15.1-MeV level. Previous data (Refs. 3-7, 14, 15) are also plotted. The curves were calculated using the intermediate-coupling model (Boyarukina) with the different oscillator-length parameters b .




FIG. 2. Spectra of scattered electrons at 35, 40, 50, 60, and 80° from ^{12}C for the incident energy of 250 MeV. The cross section is shown in the unit of $10^{-32} \text{ cm}^2/\text{sr MeV}$. The arrows indicate the positions of the peaks. The values of momentum transfer for each spectrum were calculated assuming an excitation energy of 25 MeV.

³G.A. Proca and D.B. Isabelle, Nucl. Phys. A109, 177 (1968).

⁴G.R. Bishop and A. Bottino, Phys. Letters 10, 308 (1964).

⁵B. Dudelzak and R.E. Taylor, J. Phys. Radium 22, 544 (1961).

⁶W.C. Barber, F. Berthold, G. Fricke, and F.E. Gudden, Phys. Rev. 120, 2081 (1961).

⁷F. Gudden, Phys. Letters 10, 313 (1964).

¹⁴J. Goldemberg, W.C. Barber, F.H. Lewis, Jr., and J.D. Walecka, Phys. Rev. 134, B1022 (1964).

¹⁵T.W. Donnelly, J.D. Walecka, I. Sick, and E.B. Hughes, Phys. Rev. Letters 21, 1196 (1968).

²⁰N. Bezić, D. Brajinik, D. Jamnik, and G. Kernel, Nucl. Phys. A128, 426 (1969).

METHOD

[Page 3 of 3]

REF. NO.

71 Ya 2

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




FIG. 8. The longitudinal and transverse form factors for the 16.1-MeV level. The curves correspond to the particle-hole model (Gillet and Vinh Mau) with $b = 1.64 F$ (dashed), the intermediate-coupling model (Boyarkina) with $b = 1.64 F$ (solid), and with $b = 1.9 F$ (dotted).




FIG. 15. The longitudinal form factors for the 22.0- and 23.8-MeV levels and the sum of both are plotted against q . The form factors at the photon point were estimated from the data of Bezić *et al.* (Ref. 20). Also shown are the theoretical form factors for the dipole isospin state with a dominant $(1d_{3/2}, 1p_{3/2}^{-1})$ configuration.

TABLE II. The observed and theoretical energy levels (MeV) for $J^\pi = 1^+, 2^+$, and 3^+ , $T = 1$ in ^{12}C .

J^π	Present work	Boyarkina (Ref. 29)	Gillet and Vinh Mau (Ref. 25)
1^+	15.1	15.5	16.6
2^+	16.1	15.6	16.3
	20.0	23.9	
3^+	20.6	21.0	
		24.5	



REF. J. Ahrens, H. Borchert, H. B. Eppler, H. Gimm, H. Gundrum,
 P. Riehn, G. Sita Ram, A. Zieger, M. Kroning, B. Ziegler
 Proc. International Conference on Nuclear Structure Studies
 Using Electron Scattering and Photoreaction, Sendai, Japan
 p. 213 (1972).

ELEM. SYM.	A	Z
C	12	6
METHOD	REF. NO.	
	72 Ah 7	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MUI	ABX	16-140	C	140	MGC-D		4PI

593

Fig. 8. Total nuclear cross-sections for C.



$$\Sigma_c = 50 \frac{NZ}{A}, \quad \Sigma_1(k) = \int_0^k \sigma(k') dk', \quad \Sigma_2(k) = \int_0^k \sigma(k') \frac{dk'}{k'}$$

• NBS $k=35$ MeV, ○ $k=140$ MeV

Fig. 11 Integrated cross-sections

METHOD			SOURCE		DETECTOR		REF. NO.	ANGLE
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE	72 An 3	hmg
G,T	SPC	107-999	C	700,999	MAG-D			DST
G,D	RLY	105-999	C	700,999	MAG-D			DST

999=1.2 GEV

FIG. 1. Triton spectra from C^{12} for $E_0 = 700$ MeV. Solid curve—theoretical yield of tritons from reaction (6) at 70° . Points: \circ —emission angle 40° ; \square — 60° ; \circ — 80° ; Δ — 120° .





FIG. 2. Triton spectra from C^{12} for $E_0 = 1200$ MeV at 120° . Solid curve—calculation with Eq. (7). Points: \square —proton spectra from C^{12} at 120° , \circ —triton spectra. The dashed curve is drawn through the experimental points.




FIG. 6

FIG. 6. Ratio of triton and deuteron yields N_T/N_D for the nucleus C^{12} as a function of secondary-particle momentum for $E_0 = 700$ MeV. Points: \circ — $\theta_T = 40^\circ$; \square — 60° ; Δ — 120° ; \blacktriangle and \blacksquare — N_T/N_D measured in a proton beam at angles of 120 and 60° . [17]




FIG. 4

FIG. 4. Ratio of triton and deuteron photoproduction cross sections $d\sigma_T/d\sigma_D$ as a function of energy for C^{12} for $E_0 = 700$ MeV. Points: \circ — $\theta_T = 40^\circ$; \square — 60° ; Δ — 120° ; point \bullet — $d\sigma_T/d\sigma_D$ measured in a proton beam [16] ($\theta = 40^\circ$, averaged over the secondary-particle energy range 80 – 150 MeV).




FIG. 5

FIG. 5. Triton and deuteron photoproduction cross-section ratio as a function of energy in C^{12} for $E_0 = 1200$ MeV. Points: \square — $\theta_T = 60^\circ$; Δ — 120° .

G. Anderson, I. Blomqvist, B. Forkman, G. G. Jonsson, A. Jarund,
 REF. I. Kroon, K. Lindgren and B. Schroder
 Nucl. Phys. A197, 44 (1972)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		72 An 8
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	1 * 7	C	1 * 7	ACT-I		4PI

¹⁰G. Hylten, Nucl. Phys. A158, 225 (1970).

²⁵W. C. Barber, W. D. George, D. D. Reagan,
 Phys. Rev. 98, 73 (1955).

* ENERGIES GEV




Fig. 15. Yield and cross section for the $^{12}\text{C}(\gamma, n)^{11}\text{C}$ reaction: \circ this work and \bullet some of the points from ref. ¹⁰. The yield curve drawn from 50 MeV is obtained from the cross section by Barber *et al.* ²⁵. The solid curve shows the cross section deduced from all points and the dashed curve gives the result obtained by Hyltén ¹⁰.

(over)

TABLE 6
 Partial reaction cross sections relative to the total photoabsorption cross section, $\sigma/\sigma_{\text{tot}}$ (in %)

Reaction	0.3 GeV	1.0 GeV	5.0 GeV
$^{12}\text{C}(\gamma, n)$	10 ± 5	6.8 ± 3.4	0 ± 5
$^{27}\text{Al}(\gamma, \pi^+)$	0.2 ± 0.1	0.1 ± 0.1	0.04 ± 0.02
$(\gamma, 2pn)$	1.7 ± 0.9	1.5 ± 0.8	1.7 ± 0.6
$^{127}\text{I}(\gamma, 3n)$	1.6 ± 0.8	0 ± 3	5.5 ± 2.3
$(\gamma, 6n)$	0.9 ± 0.5	1.2 ± 0.6	1.5 ± 0.8
$(\gamma, 7n)$	0.9 ± 0.5	0.5 ± 0.3	0.1 ± 0.1
$(\gamma, 8n)$	0.7 ± 0.4	0.6 ± 0.3	0.4 ± 0.2
$(\gamma, 9n)$	0.3 ± 0.2	0.2 ± 0.1	0.5 ± 0.3
$\Sigma(\gamma, xn)$	4.4 ± 2.2	2.6 ± 1.3	7.8 ± 3.9
$(\gamma, p5n)$	1.7 ± 0.9	1.7 ± 0.9	0.9 ± 0.5
$(\gamma, p7n)$	1.9 ± 1.0	2.1 ± 1.1	0.9 ± 0.5
$(\gamma, p9n)$	1.6 ± 0.8	1.6 ± 0.8	1.3 ± 0.7
$\Sigma(\gamma, pxn)$	5.2 ± 2.6	5.4 ± 2.7	3.1 ± 1.6
$(\gamma, 2p3n)$	0.6 ± 0.3	0.3 ± 0.2	0.3 ± 0.2
$(\gamma, 2p5n)$	0.5 ± 0.3	0.6 ± 0.3	0.8 ± 0.4
$(\gamma, 2p7n)$	1.3 ± 0.7	1.0 ± 0.5	0.4 ± 0.2
$(\gamma, 2p9n)$	1.1 ± 0.6	1.7 ± 0.9	2.4 ± 1.2
$(\gamma, 2p10n)$	2.7 ± 1.4	1.9 ± 1.0	5.0 ± 2.5
$\Sigma(\gamma, 2pxn)$	6.3 ± 3.2	5.6 ± 2.8	8.8 ± 4.4
$(\gamma, 4p12n)$	1.5 ± 0.8	2.4 ± 1.2	2.0 ± 1.0
$(\gamma, 4p14n)$	0.8 ± 0.4	1.9 ± 1.0	2.5 ± 1.3
$(\gamma, 4p15n)$	0.4 ± 0.2	0.6 ± 0.3	1.5 ± 0.8
$\Sigma(\gamma, 4pxn)$	2.7 ± 1.4	4.9 ± 2.5	6.0 ± 3.0
$(\gamma, 6p17n)$		0.5 ± 0.3	1.5 ± 0.8
$(\gamma, 6p18n)$	0.1 ± 0.1	0.2 ± 0.1	0.1 ± 0.1
$\Sigma(\gamma, 6pxn)$	0.1 ± 0.1	0.7 ± 0.4	1.6 ± 0.8
$(\gamma, 8p18n)$		0.8 ± 0.4	2.5 ± 1.3
$\Sigma(\gamma, ypxn)$	19 ± 10	20 ± 10	30 ± 15
$^{197}\text{Au}(\gamma, 3n)$	0.9 ± 0.5	2.2 ± 1.1	1.5 ± 0.8
$(\gamma, 7n)$	0.8 ± 0.4	1.3 ± 0.7	2.4 ± 1.2
$(\gamma, 9n)$	0.8 ± 0.4	0.6 ± 0.3	0.3 ± 0.2
$(\gamma, 11n)$	0.4 ± 0.2	0.9 ± 0.5	0.5 ± 0.3
$\Sigma(\gamma, xn)$	2.9 ± 1.5	5.1 ± 2.6	4.6 ± 2.3
$\text{Au}(\gamma, f)$	1.5 ± 0.4	2.7 ± 0.7	6.9 ± 2.0
$\text{Pb}(\gamma, f)$	3.1 ± 0.8	6.9 ± 1.7	17 ± 5

REF.

J. Berthot and D. B. Isabelle
 Lettere al Nuovo Cimento 5, 155 (1972)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

72 Be 10

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	SPC	48- 88	D	213	MAG-D		60

Bumps are at excitation energies of 48, 68, 84 MeV.

Paper also analyses other data for similar bumps (Orsay & Karkov).






Fig. 1. - Quasi-elastic scattering spectrum of 213.4 MeV electrons on ^{12}C corrected for radiative effects. The smooth curve was obtained with De Forest theory. The arrows labelled 0, 1, 2, 3 indicate, respectively, the energy position of the quasi-elastic peak maximum and of the three bumps. $\theta = 60^\circ$.

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				72 Be 12		hvm	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	30-140	D	200-350	MAG-D		DST



Références

- (1) J. BERTHOT and D.B. ISABELLE, Jour. Phys. Appl., to be published, 1972
- (2) K.V. MAC VOY and L. VAN HOVE, Phys. Rev., 1962, 125, 1034
- (3) S.D. DRELL and C.L. SCHWARTZ, Phys. Rev., 1958, 112, 568
- (4) T. DE FOREST, Nucl. Phys., 1969, A132, 305
- (5) T.W. DONNELLY, Nucl. Phys., 1970, A150, 393

METHOD					REF. NO.		
					72 B1 6		
					hmg		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
HE,G	ABX	26-32	D	1-6	NAI-D		DST

Angular distribution at 3.5 MeV was found to be:

HE=HE3,G-WIDTH,J-PI

$$W_0(\theta) \approx 1 - (.78 \pm .19) P_2(\cos\theta)$$




FIG. 4. 90° excitation curves for radiative capture of ³He to the first three states of ¹²C. The smooth curves drawn through the γ_0 and γ_2 data represent a single resonance in ¹²C formed by *s* and *d* waves (see text). The curve for γ_1 includes both this same resonance and a strong nonresonant contribution.

TABLE I. Transition strengths for ³He(³He, γ)¹²C resonance at $E(\text{lab}) = 2.55$ MeV, width $\Gamma(\text{c.m.}) = 1.6$ MeV.

	E_{res} (MeV)	J_{res}	E_{γ} (MeV)	σ_{TOT} (μb) ^a	Γ_{γ} (eV) ^b	$\Gamma_{\gamma}/\Gamma_{\text{res}}$
γ_0	0.0	0 ⁺	28.19	2.7	≥ 11.3	1.5×10^{-3}
γ_1	4.44	2 ⁺	23.75	1.1 ^c	≥ 4.6	0.9×10^{-4}
γ_2	7.56	0 ⁺	20.63	2.6	≥ 11.3	3.6×10^{-3}

^a The total resonant cross sections at 2.55 MeV are calculated assuming angular distributions identical to those measured at 3.5 MeV.

^b The lower limits on partial radiative widths are calculated assuming $J_{\text{res}} = 1$ and $\Gamma_{\text{3He}} = \Gamma$.

^c The nonresonant part of the cross section contributes an additional 0.9 μb at this energy, according to the fit to the data described in the text.





FIG. 5. Angular distributions measured at $E(^3\text{He}) = 3.5$ MeV. Curves of the form $W(\theta) = 1 + a_2 P_2(\cos\theta)$ are fitted to the data; for γ_1 an isotropic fit is also shown.

REF.

C. Brassard, H.D. Shay, J.P. Coffin, W. Scholz and D.A. Bromley
Phys. Rev. C6, 53 (1972)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
72 Br 4		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	29-35	D	14-24	NAI-D		DST

FIG. 3. Angular distributions of the γ_0 transition, in arbitrary units.




FIG. 11. Excitation functions for the total cross sections and angular-distribution coefficients for the γ_0 transition. The errors quoted here have a precise significance and are discussed in the text. The angular-distribution coefficients are as defined in the text (not as in Figs. 8-10).

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
72 G1 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
\$ P,G	ABX	22- 30	D	6-14	NAI-D		DST

$$W(\theta) = \frac{\sigma}{4\pi} \left[1 + \sum_{k=1}^{\infty} a_k P_k(\cos\theta) + \vec{P} \cdot \vec{n} \sum_{k=1}^{\infty} b_k \text{sink}\theta \right]$$

POLARIZED PROTONS

$$A_1(\theta) = \left[\sum_{k=1}^{\infty} b_k \text{sink}\theta \right] \left[1 + \sum_{k=1}^{\infty} a_k P_k(\cos\theta) \right]^{-1}$$

TABLE I. New limits on the configuration mixing in the GDR of ¹²C obtained with the (p,γ) reaction using polarized protons, as compared with the old limits derived from the unpolarized reaction.

Data	δ	min. s _{1/2} ²	max. d _{7/2} ²	min. d _{1/2} ²
Unpolarized	0	0.01	0.32	0.10
Polarized included	-15°	0.05	0.14	0.23
	0	0.06	0.26	0.21
	15°	0.02	0.49	0.07




FIG. 1 Measured angular distributions of the polarized-proton reactions ¹¹B(p,γ)¹²C and ¹¹B(p,γ)¹²C expressed as the analyzing power A₁(θ). The curves are fits of Eq. (2) with k=1, 2, 3.

(over)




FIG. 2. Summary of existing information on the reactions $^{11}\text{B}(p,\gamma_0)^{12}\text{C}$ and $^{11}\text{B}(p,\gamma_1)^{12}\text{C}$ in the giant dipole region of ^{12}C . The curves for σ_0 , a_1 , and a_2 are from Ref. 2. The values of b_2 are those obtained from the fitted curves in Fig. 1.

²R. G. Allas, S. S. Hanna, L. Meyer-Schutzmeister, R. E. Segel, Nucl. Phys. **58**, 122 (1964).




FIG. 3. The configurations in the γ_1 giant resonance of ^{12}C allowed by the data in Fig. 2. The paths in configuration space were obtained for $a_2 = -0.60$, $b_2 = -0.19$, and $\delta = 0, = 15^\circ$. The solid and dashed lines represent alternative solutions I and II, respectively. For each allowed value of $s_{1/2}|s_{1/2}|$, the allowed values of $d_{3/2}|d_{3/2}|$, $d_{3/2}|d_{7/2}|$, and $\phi_4 - \phi_3$ can be read from their respective graphs. Note that for solution I the paths for $d_{3/2}|d_{3/2}|$ and $d_{3/2}|d_{7/2}|$ are almost identical for $\delta = 0, = 15^\circ$.

METHOD				REF. NO.		hvm	
				72			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
S P,G	NOX	22- 29	D	6- 14	NAI-D		DST

POLARIZED PROTONS




Fig. 11. Summary of existing information on both $^{11}\text{B}(p,\gamma_0)^{12}\text{C}$ and $^{11}\text{B}(p,\gamma_1)^{12}\text{C}$




Fig. 10. Measured analyzing powers for $^{11}\text{B}(p,\gamma_0)^{12}\text{C}$ and $^{11}\text{B}(p,\gamma_1)^{12}\text{C}$.

REF. D. Hiramatsu, T. Kamae, H. Muramatsu, K. Nakamura, N. Izutsu,
and Y. Watase
PICNS-72, p.429 Sendai (see 73H15)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

72 Hi 8

hvm

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	ABX	0*70	D	700	MAG-D		DST

* SEP ENERGY RANGE




Fig. 2. Separation energy spectra for ^{12}C




Fig. 6. The separation energy of 1s and 1p states as a function of the mass number.




Fig. 3. Momentum distributions for ^{12}C

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				72 Ku 6		egf	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,N	ABX	19-30	D	20-30	ACT-I		4PI




Fig. 3. Absolute cross sections of the three reactions investigated. The error of the absolute scale is estimated to be $\pm 8\%$.




Fig. 4. The measured ratios of σ^-/σ^+ at an energy of 27 MeV compared with the straight line given by Herring *et al.*⁶⁾.

REF.

I. Linck and L. Kraus
C.R. Acad. Sc. Paris 275, 669 (1972)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

72 Li 4

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
HE,G	ABX	27- 35	D	1- 11	NAI-D		90

HE = HE3




Fig. 2 a. — Courbes d'excitation des transitions γ_0 et γ_1 .
L'échelle du haut donne l'énergie d'excitation dans ${}^{12}\text{C}$, au centre de la cible.




Fig. 2 b. — Courbes d'excitation des transitions γ_2 et γ_3 .

REF.

D.M. Skopik, Y.M. Shin, J.W. Horniachek, S.K. Kundu
Nucl. Phys. A197, 185 (1972)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

72 Sk 7

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E _e ,D	ABX	40-60	D	40-60	MAG-D		DST




Fig. 3. Angular distribution of deuterons near the kinematic threshold at an incident electron energy of 40 MeV.




Fig. 4. Angular distribution of deuterons ($E_d = 11.5 \text{ MeV}$) not restricted to ground or first excited state transitions at an incident electron energy of 40 MeV.

(over)

U.S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS




Fig. 2. Differential cross sections for the $^{12}\text{C}(\gamma, d)^{10}\text{B}$ reaction at three lab angles.




Fig. 6. Isochromats for deuterons of 11.5 MeV.




Fig. 5. Angular distribution of deuterons ($E_d = 11.5$ MeV) at an incident electron energy of 60 MeV.

ELEM. SYM.	A	Z
C	12	6

METHOD					REF. NO.		
					72 Sp 9	hvm	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	NOX	15	D	35- 71	MAG-D		DST

15=15.109 MEV

The analysis of the data has been performed in an iterative procedure with b and $\sqrt{B(M1, q)}$ as free parameters and $\rho = 0.16$ (corresponding to $\gamma = -1.4$). After each iteration the f_c 's are recalculated using the improved value of b . The best fit to the data (including the properly normalized results from Orsay (8)) is shown in fig. 7. The final results are $\sqrt{B(M1, 0)} = (1.737 \pm 0.012)10^{-1}$ fm, $b^2 = 3.542 \pm 0.078$ fm²; this corresponds to a radiative width of $\Gamma_Y^0 = (35.74 \pm 0.86)$ eV, where an error of 1% has been added linearly to the statistical error to account for the uncertainty in the f_c .




Fig. 7. Reduced transition probabilities for the 15.1 MeV level in ¹²C. The fit is obtained from an analysis performed for the combined set of Darmstadt data and Amsterdam data. The correction factors $\sigma_{BA}/\sigma_{DWBA}$ in the data reduction are shown in the insert.

REF.

S.G. Tonapetyan, N.V. Goncharov, A.I. Derebchinskii,
O.G. Kononov and V.M. Khvorostyan
Zh. Eksp. Teor. Fiz. 63, 1955 (1972)
Soviet Phys. JETP 36, 1033 (1973)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

72 To 9

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	86-240	C	250-999	BBL-D		90

999 = 1.2 GEV




FIG. 2




FIG. 3

FIG. 2. Comparison of experimental proton energy spectra for $E_p^{\max} = 250, 293, 400, 600,$ and 800 MeV with spectra calculated from Eq. (1), where L is taken respectively as 3, 4.4, 6.4, 7.7, and 9.1; solid curves—experiment, dashed curves—quasideuteron model.

FIG. 3. Differential cross section for photodisintegration of C^{12} at $\theta_{lab} = 90^\circ$ as a function of E_γ : Δ —results of the present work when integration over E_p is carried out in the region $E_p = 70-240$ MeV, \circ —integration over E_p in the region $E_p = 30-240$ MeV (curves 1 and 2 have been drawn through the experimental points). The errors are statistical. Curve 3—differential cross section for photodisintegration of the deuteron, increased by 30 times; curve 4—the sum of the differential cross sections for photoproduction of 1, 2, and 3 π mesons of all signs in nucleons of C^{12} .




FIG. 1. Energy spectra of photoprotons from C^{12} at an angle $\theta_{lab} = 90^\circ$ for bremsstrahlung photons with maximum energy from 10 to 1200 MeV. The errors are statistical. Hollow points—results of the present work, solid points—data from other sources: \blacksquare —ref. 1, \bullet —ref. 2, \blacktriangle —ref. 5.

⁴C. Levinthal et al. Phys. Rev. 82, 822 (1951)

⁵P. Dougan et al., Preprint LUSY-1003, 1970

¹²N.G. Afanas'ev et al., Prib. Tekh. Eksp. 5, 146 (1967)

REF.

R. E. Van de Vyver, H. Ferdinande, G. Knuyt, R. Garchon, J. Devos
Nucl. Phys. A198, 144 (1972)

ELEM. SYM. A Z

C 12 6

METHOD

REF. NO.

72 Va 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	ABX	18-32	C	18-32	BF3-I		4PI




Fig. 2. Comparison of the present total photoneutron cross section for ^{12}C with previous measurements. Our data are the results of a Cook analysis.

REF. J. Ahrens, H.B. Eppler, H. Gimm, H. Gundrum, M. Kroning,
 P. Riehn, G. SitaRam, A. Zieger, and B. Ziegler
 PICNS-73, Vol. I, p. 23 Asilomar

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	73 Ah 4 hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	10-140	C	140	MGC-D		4PI

Statistics may have been improved over those of 72Ah7.

See figure on other side.

(OVER)




Fig. 4. Same as Fig. 2 for C.

REF. S. Anami, K. Baba, I. Endo, H. Fujii, S. Kato, A. Murakami,
S. Ninomiya and Y. Sumi
PICNS-73, Vol. I, p.613 Asilomar

ELEM. SYM.	A	Z
C	12	6

METHOD					REF. NO.		
					73 An 8		hmg
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	ABX	140-824	D	323-824	MAG-D		28




Fig.

Table

<k> (MeV)	dσ/dΩ (μb/sr)	N _{eff}	2Γ (MeV/c)	p ₀ (MeV/c)	p _{free} (MeV/c)	χ ² /d.f.
323	30.8-30.9	1.4	47	242	279	0.76-0.77
373	39.2-42.3	2.0-2.2	47-50	286-289	329	1.26-1.76
423	44.5-47.8	2.7-2.9	51-54	333-336	378	1.46-1.69
473	57.7-61.6	3.2-3.4	62-64	380-384	425	0.65-1.28
524	57.8-58.9	3.1-3.2	69-70	424	473	0.91-1.04
574	62.0-66.6	3.1-3.3	72-76	468-471	518	1.03-1.34
623	61.5-66.8	2.7-3.0	72-77	519-523	561	1.09-1.43
674	66.4-68.1	2.7	82-83	567	606	1.35-1.42
724	72.9-78.5	3.1-3.3	94-98	600-605	649	0.45-0.76
774	65.3-67.1	3.4-3.5	105-106	646	692	0.46-0.53
824	61.9-70.9	3.9-4.5	114-133	672-682	734	0.36-0.52

REF. P.S. Ananien, I.V. Glavanakov, V.I. Kryshkin,
 V.N. Stibunov, A.G. Sterligov, Yu. P. Usov
 Lettere al Nuovo Cimento 8, 651 (1973)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	73 An 10
	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
\$ G,PI0p	NOX	137*235	C	800-900	TEL-D		90
\$ G,PI-P	NOX	155*235	C	800-900	TEL-D		90

\$P,*RECOIL P ENERGY

TABLE I. - Experimental results.

E_p (MeV)	$(\gamma p, \pi^0 p)$			$(\gamma n, \pi^- p)$		
	P	ΔP	Events	P	ΔP	Events
137	0.1	± 0.65	22			
155	-0.8	± 0.35	60	0.9	± 0.6	45
175	-0.19	± 0.3	71	-0.09	± 0.29	65
197	-0.63	± 0.28	63	-0.16	± 0.27	60
215	-0.06	± 0.31	57	-0.57	± 0.3	62
235	-0.25	± 0.25	59	-0.30	± 0.29	46

Table I presents the experimental results of the polarization measurements in reactions $(\gamma p, \pi^0 p)$ and $(\gamma n, \pi^- p)$ vs. recoil proton energy. The errors are one standard deviation evaluated from the maximum-likelihood procedure.

REF. B. T. Chertok, G. Sheffield, J.W. Lightbody, Jr., S. Penner,
and D. Blum
Phys. Rev. C8, 23 (1973)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

73 Ch 1

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	LFT	15	D	35- 56	MAG-D		DST

$$\Gamma_{\gamma} = 37.0 \pm 1.1 \text{ eV}$$

15.109 MEV LEVEL

TABLE III. Input data for fitting.

q^2 (fm^{-2})	$B(M1, q, \uparrow)$ (10^{-2} fm^2)	$[\sigma(B(M1))]^{-2} \text{ a}$
0.0338	0.9076	213
0.0454	0.9737	275
0.0591	0.9024	419
0.0920	0.8290	674
0.0560	0.8191	454
0.0773	0.8449	785
0.1298	0.6777	1570
0.1613	0.6505	1558
0.1295 ^b	0.6854	213
0.2730	0.4571	479
0.1185 ^c	0.728	385
0.1519	0.667	459
0.1880	0.519	758
0.1894	0.593	1138
0.2310	0.527	1441
0.2768	0.452	3058

^a 1% has been added linearly to the σ in Table II and to the data of Ref. 15.

^b 0.1295+0.2730 are from Ref. 20.

^c 0.1185 to 0.2768 inclusive are from Ref. 15.




FIG. 3. Fits using Eq. (11) are presented to the experimental reduced nuclear transition probability versus momentum transfer squared. The data from Yale (Ref. 15) and Orsay (Ref. 20) as reanalyzed here are included with our measurements.

¹⁵ G.A. Peterson, Phys. Lett. 25B, 549 (1967);
and private communication.

²⁰ B. Dudelzak and R.E. Taylor, J. Phys. Radium
22, 544 (1961).

REF. V. di Napoli, F. Salvetti, M.L. Terranova, H.G. de Carvalho,
and J.B. Martins
Phys. Rev. C8, 206 (1973)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		73 Di 4
		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, SPL	ABY	THR-999	C	999	ACT-I		4PI

999 = 1 GE




FIG. 3. Yields of ^{11}C and ^7Be versus the mass number of the target nucleus. Filled circle: ^{11}C , Ref. 16. Filled triangle: ^{11}C , Ref. 17. Filled square: ^{11}C , Ref. 8. Open circles: ^{11}C , present work. Reversed open triangles: ^7Be , present work.

⁸G. Andersson et al., Nucl. Phys. A197, 44 (1972).

¹⁶V. di Napoli et al., Nuovo Cimento 55B, 95 (1968).

¹⁷A. Masaïke, J. Phys. Soc. Japan 19, 427 (1964).

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XP	ABY	96-400	C	400	TET-D		DST




Fig. 9. In this figure, the straight lines show the experimental cross-sections at 40° and 74° for $E_p=150$ MeV. The other curves are the same cross-sections divided by atomic weight




Fig. 6. The ratios of the experimental cross-sections at 40 and 90 degrees for selected proton energies as a function of atomic number




Fig. 8. Experimental cross-sections at various angles for $E_p=150$ MeV divided by NZ/A plotted as a function of atomic weight

(over)

Table 4. Carbon. Bremsstrahlung endpoint energy: 400 MeV. Differential cross-sections in microbars/sterrad · MeV · eq. quantum. Quoted errors: statistical in percent

Energy	Angle								
	22	30	40	50	60	74	90	110	130
81.5		2.67 2.6	2.30 1.0	1.86 1.8	1.56 2.9	1.18 2.3	0.820 2.3	0.545 3.0	0.445 3.2
85.0	3.18 2.2	2.71 3.0	2.19 1.4	1.76 1.9	1.45 2.0	1.10 1.6	0.843 2.0	0.612 2.3	
99.4		2.19 3.1	1.72 2.0	1.31 2.3	1.13 1.7	0.891 2.9	0.575 3.0	0.393 3.8	0.266 4.5
102.5	2.59 2.6	2.04 2.1	1.68 1.7	1.27 2.4	1.09 2.5	0.849 2.1	0.587 2.6	0.410 3.1	
106.3	2.36 2.3		1.55 1.2	1.23 2.8	1.05 2.3	0.744 1.4	0.552 1.6	0.400 3.2	
109.2		1.82 2.4	1.39 0.9	1.28 2.4					
116.4		1.55 3.6	1.26 3.3		0.839 2.0	0.687 3.3	0.418 3.5	0.244 4.8	0.154 5.9
119.2	1.77 4.4	1.38 2.4	1.21 2.8	0.880 4.2	0.736 4.2	0.602 3.4	0.390 4.4	0.242 5.0	

Table 4 (continued)

Energy	Angle								
	22	30	40	50	60	74	90	110	130
128.9	1.65 2.9		1.04 1.1	0.903 3.6	0.718 3.0	0.529 2.1	0.355 2.6	0.207 4.6	
131.4		1.29 3.0	0.977 2.0	0.932 2.9					
150.0	0.952 3.8		0.719 1.0	0.626 4.4	0.472 3.7	0.347 3.2	0.205 5.2	0.112 6.5	
152.3		0.850 3.8	0.687 2.0	0.666 3.5					
171.6	0.695 3.1	0.600 2.3	0.513 1.3	0.416 2.1	0.341 4.6	0.216 2.7			
188.4	0.487 3.9	0.455 2.8	0.381 1.3	0.306 3.9	0.223 1.2	0.133 3.7			
205.1	0.363 4.4	0.302 3.4	0.264 2.8	0.203 2.9	0.142 3.4	0.0681 5.1			
221.0			0.211 7.2	0.146 9.7	0.0798 12.9				
223.8	0.298 3.8	0.267 3.7	0.208 2.6	0.146 5.7	0.0905 7.1	0.0398 10.2	0.0131 15.8		
238.4			0.169 8.4	0.118 11.3	0.0437 18.3				
240.1	0.192 5.0	0.185 4.7	0.135 3.4	0.0822 7.9	0.0546 9.6	0.0248 13.5	0.00471 27.7		
251.6	0.156 4.6	0.145 3.5	0.0805 3.4	0.0529 7.8					
254.8			0.0797 8.4	0.0492 17.1	0.0254 23.6				
256.4	0.152 5.5	0.110 5.9	0.0656 4.8	0.0492 10.1	0.0213 15.1	0.00481 30.2	0.00176 44.7		
266.8	0.0879 6.4	0.0744 5.5	0.0370 5.5	0.0275 11.3					
282.6	0.0703 7.0	0.0475 12.9	0.0218 6.4	0.0127 16.2					




Fig. 3. Energy distribution of protons at 40° from carbon irradiated with bremsstrahlung of 400 MeV maximum energy. The data not included in Table 4 are to be found in [32].

REF.

I. Endo, K. Baba, H. Fujii, Y. Sumi, S. Anami, S. Ninomiya,
S. Kato, and A. Murakami
Phys. Lett. 47B, 469 (1973)

ELEM. SYM. A Z

C 12 6

METHOD

REF. NO.

73 En 4 egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,PI+	ABX	319-819	C	300-850	MAG-D		28
E,PI-	ABX	319-819	C	300-850	MAG-D		28




Fig. 1. Typical examples of momentum spectrum at 28.4°. Bars on several points show typical statistical errors. The solid curve is the one fitted by the Gaussian form. The arrow p_{free} denotes the value of pion momentum corresponding to the free nucleon kinematics. The π^+ data (\times) at 63° and $\langle k \rangle = 575$ MeV are also shown [3].

³S. Arai et al., preprint, Nagoya University DPNU-9 (1972)




Fig. 2. (a) Values of $(d\sigma/d\Omega)_{carbon}$ (\bullet : π^+ , \square : π^-) and $6 \times (d\sigma/d\Omega)_{hydrogen}$ as a function of $\langle k \rangle$. (b) N_{eff} (\bullet : π^+ , \square : π^-) and N_{eff}^* (\triangle : π^+ , Δ : π^-). Errors are due to the fitting uncertainty; statistical errors of about 4% at all points are not shown.

over

Table 1
 The fitted values of parameters in the Gaussian distribution given by (4) and the values of effective nucleon numbers N_{eff} and N_{eff}^* . Errors quoted are only due to the fitting uncertainty; statistical errors are always about 4% and not shown here.

k (MeV)		$d\sigma/d\Omega$ ($\mu\text{b}/\text{sr}$)	2Γ (MeV/c)	$P_0 - P_{\text{free}}$ (MeV/c)	N_{eff}	N_{eff}^*
319	π^+	33.8 ± 1.4	51 ± 2	-34 ± 2	1.6 ± 0.1	3.6 ± 0.2
	π^-	30.9 ± 0.1	47 ± 1	-33 ± 1	1.4 ± 0.1	3.2 ± 0.1
369	π^+	39.4 ± 1.5	50 ± 1	-35 ± 2	1.8 ± 0.1	3.7 ± 0.2
	π^-	40.8 ± 1.6	49 ± 2	-38 ± 1	1.8 ± 0.1	3.6 ± 0.1
419	π^+	48.1 ± 6.1	54 ± 1	-39 ± 2	2.9 ± 0.4	5.3 ± 0.7
	π^-	46.2 ± 1.7	53 ± 2	-39 ± 2	2.7 ± 0.1	4.9 ± 0.2
469	π^+	56.5 ± 1.4	67 ± 2	-34 ± 1	3.3 ± 0.1	5.3 ± 0.2
	π^-	59.7 ± 2.0	63 ± 1	-40 ± 2	3.4 ± 0.1	5.5 ± 0.2
519	π^+	53.3 ± 4.2	62 ± 3	-28 ± 5	2.9 ± 0.3	4.3 ± 0.4
	π^-	58.4 ± 1.4	70 ± 1	-44 ± 1	3.4 ± 0.1	5.0 ± 0.1
569	π^+	65.4 ± 0.2	82 ± 2	-40 ± 1	3.3 ± 0.1	4.5 ± 0.1
	π^-	64.3 ± 2.3	74 ± 2	-44 ± 2	3.4 ± 0.1	4.8 ± 0.2
619	π^+	63.9 ± 3.2	65 ± 1	-27 ± 1	2.8 ± 0.1	3.7 ± 0.2
	π^-	64.2 ± 2.6	75 ± 3	-37 ± 2	3.1 ± 0.1	4.1 ± 0.2
669	π^+	88.8 ± 2.7	104 ± 5	-47 ± 4	3.5 ± 0.1	4.3 ± 0.2
	π^-	67.3 ± 0.9	83 ± 1	-35 ± 1	3.1 ± 0.1	3.7 ± 0.1
719	π^+	100.8 ± 1.1	106 ± 2	-42 ± 1	4.0 ± 0.1	4.5 ± 0.1
	π^-	75.5 ± 2.8	96 ± 2	-42 ± 3	3.6 ± 0.2	4.1 ± 0.2
769	π^+	72.6 ± 1.4	103 ± 2	-38 ± 1	3.7 ± 0.1	4.1 ± 0.1
	π^-	66.2 ± 0.9	106 ± 1	-41 ± 1	4.4 ± 0.1	4.8 ± 0.1
819	π^+	86.7 ± 3.3	122 ± 6	-43 ± 5	6.2 ± 0.3	6.5 ± 0.3
	π^-	66.4 ± 4.5	124 ± 10	-52 ± 5	6.4 ± 0.5	6.8 ± 0.5

REF.

H. J. v. Eyss, H. Schier and B. Schoch
PICNS-73, Vol. II, p. 989 (1973) Asilomar

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	
	73 Ey 2	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	64-123	D	64-123	TOF-D		DST




Fig. I: Cross-sections for photoneutrons from ^{12}C . The width of the peak observed, corresponds to the incident photon-spectrum. The errors shown are statistically.

(REV. 7-14-66)
USCOMM-NBS-DC

PHOTONUCLEAR DATA SHEET 330

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				73 Ey 3		egf	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	SPC	THR-234	C	234	TOP-D		90

NEUTS E ABV 12 MEV




Fig. 8. Comparison of the shape of the high-energy part of the photoneutron spectra from C, Al, Cu, Cd and Pb. These measurements were performed with the same γ -shower spectrum, produced in a 0.3 cm thick lead sheet (see Fig. 2b). All spectra were fitted to the value for carbon at $E_n = 100$ MeV. The values predicted by a quasi-deuteron model (solid lines), which are also fitted at $E_n = 100$ MeV, were calculated with the parameters (defined in the text): $E_b = -10$ MeV, $E_{well} = 30$ MeV and $C'L = 19.0$ for different impulse parameters $\alpha = 100, 120, 140, 160$ and 180 MeV/c




Fig. 9. Dependence of the production cross section on the mass number A with the neutron energy as parameter, measured at $E_e = 234$ MeV. The γ -quanta were produced in a 0.3 cm thick lead sheet (see Fig. 2b) in front of the target of mass number A . The solid lines are fit curves through the measured values. The dashed lines are values calculated using a quasi-deuteron model with the parameters (defined in the text): $E_b = -10$ MeV, $E_{well} = 30$ MeV, $\alpha = 140$ MeV/c and $C'L = 19.0$. The dotted curve represents the dependence NZ/A , fitted at $A = 12$. The error bars correspond to the statistical error

(over)




Fig. 10. Neutron yield from targets of mass number A relative to carbon, measured at $E_e = 234 \text{ MeV}$. The target arrangement is that of Fig. 2b. The solid lines are fit curves through the experimental values. The dashed curve shows the energy dependence of the ratio of the nuclear absorption factors $f_a(\text{Pb})/f_a(\text{C})$, taken from Fig. 6. The error bars correspond to the statistical error

N.V. Goncharov, A.I. Derebchinskii, O.G. Kononov,
 REF. S.G. Tonapetyan, and V.M. Khvorostyan
 Yad. Fiz. 17, 242 (1973)
 Sov. J. Nucl. Phys. 17, 124 (1973)

ELEM. SYM.	A	Z
G	12	6

METHOD			REF. NO.				
			73 Go 5		hmg		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABY	170-400	C	400	BBL-D		90
G,PI-	ABY	170-400	C	400	BBL-D		90
G,P	ABY	86-400	C	400	BBL-D		90




FIG. 2. Energy spectra of π^+ and π^- mesons, $E_7^{\max} = 400$ MeV, $\theta_{lab} = (90 \pm 7)^\circ$.




FIG. 4. Charged pion yield vs. the mass number of the nucleus: \bullet — $E_\pi = 105 \pm 10$ MeV, \circ — $E_\pi = 85 \pm 10$ MeV, \circ — $E_\pi = 65 \pm 10$ MeV.




FIG. 3. Energy spectra of protons, $E_7^{\max} = 400$ MeV, $\theta_{lab} = (90 \pm 7)^\circ$. Circles—present data, triangles—from [18].

¹⁸P. Dougan, W. Stiefler, LUSY Preprint, 1001-1003, 1970.

(over)




FIG. 5. Proton yields vs. mass number of the nucleus: \circ — $E_p = 100 \pm 10$ MeV, \bullet — $E_p = 125 \pm 15$ MeV, Δ — $E_p = 155 \pm 15$ MeV, \square — $E_p = 185 \pm 15$ MeV, \diamond — $E_p = 215 \pm 15$ MeV.




FIG. 6. Pion yield (1), proton yield (2), and summary pion and proton yield (3) vs. the mass number of the nucleus.

REF.

Stanley S. Hanna
PICNS-73, Vol. I, p.417 (1973) Asilomar

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		73 Ha 15
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	22 (22.5)	D	7 (7.20)	NAI-D		90

Table IV. Summary of cross section measurements for (γ, p_0) and (γ, n_0) reactions on ^4He , ^{12}C , and ^{16}O . Underlined values are essentially the consensus of several values.

	E_x (MeV)	$\sigma(\gamma, p_0)$ in mb		$\sigma(\gamma, n_0)$ in mb	
		New ^{a)}	Old	New	Old
^4He	26.0	1.95 ± 0.13	<u>1.84</u> ^{b)}	<u>2.0</u> ^{c)}	1.0 ^{d)}
^{12}C	22.5	12.2 ± 1.2	13.4 ^{e)} , 3.0 ^{f)}	<u>6.0</u> ^{g)}	6.0 ^{h)}
^{16}O	22.2	9.6 ± 1.5	3.8 ⁱ⁾ , 12.2 ^{j)}	~ 7.2 ^{k)}	9.1 ^{h)} , 7.2 ^{l)}

a) Ref. 25
b) Refs. 30-32
c) Refs. 22-24
d) Ref. 33

e) Ref. 34
f) Ref. 35
g) Ref. 36,37
h) Ref. 38

i) Ref. 39
j) Ref. 40
k) Ref. 41
l) Ref. 42

(over)

- 22 W. R. Dodge and J. J. Murphy, II, Phys. Rev. Letters 28, 839 (1972).
- 23 J. D. Irish, R. G. Johnson, B. J. Thomas, B. L. Berman, K. G. McNeill, and J. W. Jury, contribution to this conference.
- 24 D. V. Webb, C. K. Malcolm, Y. M. Smith, and D. M. Skopik, contribution to this conference.
- 25 J. R. Calarco, G. A. Fisher, E. M. Diener, C. C. Chang, and S. S. Hanna, to be published.
- 26 J. E. Brolley, Jr., T. M. Putman, and Louis Posen, Phys. Rev. 107, 820 (1957).
- 27 J. L. Detch, Jr., R. L. Hutson, N. Jarmie, and J. H. Jett, Phys. Rev. C4, 52 (1971).
- 28 M. Suffert, W. Feldman, J. Mahieux, and S. S. Hanna, Nucl. Instr. and Meth. 63, 1 (1968).
- 29 B. L. Berman, private communication.
- 30 M. Suffert, private communication.
- 31 J. E. Perry, Jr. and S. J. Bame, Jr., Phys. Rev. 99, 1368 (1955).
- 32 W. E. Meyerhof, M. Suffert, and W. Feldman, Nucl. Phys. A148, 211 (1970); the cross sections in this paper are based on Ref. 31.
- 33 B. L. Berman, S. C. Fultz, and M. A. Kelly, Phys. Rev. C4, 723 (1971); B. L. Berman, F. W. K. Firk, and C. P. Wu, Nucl. Phys. A179, 791 (1972).
- 34 R. G. Allas, S. S. Hanna, L. Meyer-Schützmeister, and R. E. Segel, Nucl. Phys. 58, 122 (1964).
- 35 C. Brassard, H. D. Shay, J. P. Coffin, W. Scholz, and D. A. Bromley, Phys. Rev. C6, 53 (1972).
- 36 J. W. Jury, J. S. Hewitt, and K. G. McNeill, contribution to this conference.
- 37 B. L. Berman et al., contribution to this conference.
- 38 C. P. Wu, F. W. K. Firk, and T. W. Phillips, Phys. Rev. Letters 20, 1182 (1968).
- 39 E. D. Earle and N. W. Tanner, Nucl. Phys. A95, 241 (1967).
- 40 J. E. E. Baglin and M. N. Thompson, Nucl. Phys. A138, 73 (1969).
- 41 D. B. C. B. Syme and G. I. Crawford, contribution to this conference.
- 42 J. T. Caldwell, R. L. Bramblett, B. L. Berman, R. R. Harvey, and S. C. Fultz, Phys. Rev. Letters 15, 976 (1965).

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
73 He 8		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	ABX	1*	D	2* 3	MAG-D		15




Fig.2 Momentum distribution of the residual nucleus extracted from quasi-elastic electron scattering from ^{12}C . The four momentum transfer to the proton was $q^2 = 10 \text{ fm}^{-2}$. The recoil protons were detected in the range of $52^\circ \leq \theta_p$ and $-15.5^\circ \leq \phi_p \leq +15.5^\circ$.

* E, GEV




Fig.1 Spectrum of the scattered electrons.

The solid curve is a fit to the experimental points using calculations by Devanathan for quasi-elastic scattering, the model of Gutbrod and Simon for the $\Delta(1236)$ resonance, and a Woods-Saxon wave function for the ^{12}C nucleus including short range correlations.

REF.

H. Hiramatsu, T. Kamae, H. Muramatsu, K. Nakamura, N. Izutsu,
and Y. Watase
Phys. Letters **44B**, 50 (1973)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

73 HI 5

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	SPG	0*70	D	700	MAG-D		UKN

*SEP ENERGY RANGE




Fig. 2. Proton momentum distributions. See fig. 1 for the meaning of the curves in a) and b).

5 G. Gioffi degli Atti, Nucl. Phys. **A106** (1968) 215.




Fig. 1. Separation energy spectra. The solid curves show the DWIA results with the Woods-Saxon potentials. In a), b) and c) the dot-dashed curves represent the DWIA calculation with the harmonic oscillator potential given in ref. [5], and the dashed curves show the multiple collision background calculated by the Monte Carlo method.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hmg
73 Ju 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	SPC	18- 28	C	23- 31	TOF-I		DST

Data fitted to Legendre polynomial $\sum_4 a_n P_n(\cos\theta)$,
 a_4 determined to be zero.

GND STATE NEUTRONS




Figure 2: Neutron energy spectra for an end-point of 30.5 MeV.




Figure 3: Ground state photo-neutron cross section ($4\pi\sigma_0$) & anisotropy coefficients.

2 D.E. Frederick and A.D. Sherick, Phys. Rev. 176, 1177 (1968).




Figure 4: Comparison of the a_2/a_0 coefficient with results of the coupled-channels reaction theory.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
73 Li 1		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
HE,G	ABX	27-35	D	1-11	NAI-D		DST

HE=HE-3




Fig. 2. Excitation curves for γ_0, γ_1 and γ_2 , at $\theta=90^\circ$ and for bombarding energies ranging from 1.5 to 11 MeV.





Fig. 3. Angular distributions for γ_0 .

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				73 Mo 9		hmg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	RLY	18- 39	C	18- 39	ACT-I		4 PT
E,N	RLY	18- 39	D	18- 39	ACT-I		4 PT

YIELD/PHOTO/ELECTRO



1. W.C. Barber, Phys. Rev. 111(1958)1642 (other ref. to previous work to be found in this article)

REF. R. Nath, F.W.K. Firk, R.J. Holt, H.L. Schultz
 PICNS-73, Vol.I, p.155 Asilomar

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	73 Na 5 hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
\$ G,N	NOX	22- 31	C	30,64	TOF-I		DST

POLARIZED G. S. NEUTRONS

See figure on reverse side.

(OVER)




FIG. 1

B.D. Anderson, M.R. Dwarakanath, J.S. Schweitzer, and
 A.V. Nero
 REF. Nucl. Phys. A233, 286 (1974)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		74 An 7
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P, G	LFT	16	D	1*	NAI-D		90

Abstract: The partial widths of the second $T = 1$ state of ^{12}C , at 16.11 MeV excitation energy, have been determined by measuring the $^{11}\text{B}(p, \gamma)$ and $^{11}\text{B}(p, \alpha)$ cross sections at the $E_p = 163$ keV resonance corresponding to this state. These measurements result in the new values of $\Gamma_p = 21.7 \pm 1.8$ eV and $\Gamma_\gamma = 21.6 \pm 3.3$ eV, for the partial widths of this state; approximately 3 times smaller and larger, respectively, than the present values in the literature. The new result for the proton width eliminates a serious discrepancy found in an earlier comparison of the partial widths of the $T = 1$ analogue states of the $A = 12$ system. Measurements were also made of the $^{11}\text{B}(d, n)^{12}\text{C}^*$ reaction to compare the proton widths of the 15.11 and 16.11 MeV $T = 1$ states; these measurements confirm the new, smaller proton width for the 16.11 MeV state. An attempt was also made to determine the γ -width of the 16.11 MeV state by measuring the γ -branching ratio in the $^{10}\text{B}(^3\text{He}, p)^{12}\text{C}^*(\gamma)^{12}\text{C}$ reaction.

* $l=220$ KEV

E NUCLEAR REACTIONS $^{11}\text{B}(d, n)$, $E = 6$ MeV; measured $\sigma(E_n, \theta)$; $^{11}\text{B}(p, \gamma)$, (p, α) , $E = 163$ keV; measured σ_R ; $^{10}\text{B}(^3\text{He}, p)$, $E = 2.2$ MeV; measured $p\gamma$ -coin. ^{12}C levels deduced I_p , I_γ , S .

TABLE 3
 $^{11}\text{B}(p, \gamma)$ results for 163 keV resonance

	$\sigma_R \Gamma_{c.m.}$	$\Gamma_{c.m.}$ taken (keV)	σ_R (μb)
Present work	836 ± 109	6.7	125 ± 16
Huus and Day ⁷⁾	788 ± 113	5.0	158 ± 24

⁷⁾ Ref. 7).

⁷⁾ T. Huus and R.B. Day, Phys. Rev. 91 (1953) 599.

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		74 Ba 15
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,FRG	RLY	THR - 999	C	800-999	TRK-D		DST

999=1GEV FRG A,6-11

TABLE 2
Measured Be/Li and B/Li yield ratios at 800 and 1000 MeV

Angle (deg.)	300 MeV		1000 MeV	
	Be/Li	B/Li	Be/Li	B/Li
10	0.30 ± 0.08	0.07 ± 0.04	0.33 ± 0.05	0.04 ± 0.01
20	0.49 ± 0.13	0.07 ± 0.04	0.32 ± 0.05	0.03 ± 0.01
30			0.44 ± 0.07	0.03 ± 0.01
40	0.45 ± 0.12	0.02 ± 0.02	0.38 ± 0.07	0.03 ± 0.01
50			0.32 ± 0.06	0.04 ± 0.02
60	0.23 ± 0.08	0.02 ± 0.02	0.28 ± 0.07	0.02 ± 0.01
70	0.29 ± 0.11	< 0.03	0.24 ± 0.05	0.01 ± 0.01
80			0.25 ± 0.05	0.01 ± 0.01
90	0.24 ± 0.10	< 0.03	0.19 ± 0.04	0.01 ± 0.01
mean	0.34 ± 0.04	0.036 ± 0.012	0.31 ± 0.02	0.025 ± 0.003

TABLE 3
Lab energy and momentum intervals of the identified nuclei and related yields for the 1000 MeV irradiation; fitted values of the mean velocity (v_{11}/c) of the hypothetical emitting nucleus and slopes (T_0^*) of the c.m. energy spectra

	$^6\text{Li}+^7\text{Li}$	$^8\text{Li}^*$	^7Be	^9Be	^8B	$^{10}\text{B}+^{11}\text{B}^b)$
$T(\text{MeV}/N)$	1.1 - 1.8	0.7 - 1.7	2.0 - 3.0	1.9 - 2.8	2.5 - 4.5	2.4 - 4.0
$P(\text{MeV}/c)$	300 - 380	290 - 450	430 - 530	540 - 650	550 - 740	670 - 870
Angle (deg)	yields (particles/msr)					
10	740 ± 30	76 ± 10	82 ± 10	18 ± 4	2 ± 1	22 ± 4
20	690 ± 30	87 ± 11	84 ± 11	16 ± 4	6 ± 2	12 ± 3
30	650 ± 30	65 ± 8	100 ± 13	10 ± 3	1 ± 1	9 ± 2
40	690 ± 40	65 ± 9	87 ± 14	20 ± 4	2 ± 1	12 ± 3
50	630 ± 30	54 ± 9	82 ± 12	16 ± 4	2 ± 1	12 ± 4
60	610 ± 40	49 ± 8	80 ± 14	9 ± 3	2 ± 1	6 ± 2
70	560 ± 30	49 ± 7	62 ± 10	12 ± 3	0.7 ± 0.7	9 ± 2
80	470 ± 30	47 ± 5	46 ± 8	8 ± 2	0.4 ± 0.4	4 ± 1
90	400 ± 20	35 ± 6	35 ± 6	3 ± 1	< 0.6	4 ± 2
$v_{11}/c(\times 10^3)$	4.5 ± 0.6	6.4 ± 1.1	11.5 ± 2.3	6.7 ± 2.0		8.5 ± 2.0
$T_0^*(\text{MeV}/N)^c)$	0.52 ± 0.02	0.48 ± 0.03	1.12 ± 0.12	0.48 ± 0.08		0.48 ± 0.07

^{a)} Corrected for efficiency.

^{b)} Assumed as ^{10}B in the calculations.

^{c)} Assuming the c.m. spectrum $d^2N/d\Omega^*dT^* \approx \exp(-T^*/T_0^*)$.

(over)




Fig. 4. Angular distributions of Li, Be and B at 1000 MeV; the data collected for each isotope correspond to the quoted segments of energy spectra.




Fig. 5. Energy spectra of the quoted isotopes in the c.m. of the emitting nucleus from the 1000 MeV irradiation. Each group of points marked with the same symbol refers to data collected at different angles (10° to 90°) in the same lab energy interval. The results of a best fit on the emitting nucleus velocities and on the slopes (T_0^*) of the energy spectra are given in table 3.




Fig. 6. Normalized laboratory energy spectra of ${}^6\text{Li}$ from carbon: (a) this work (10° - 90°); (b) ref. ²³); π^- at rest on C, N, O; (c) ref. ²⁶); 2.2 GeV protons (30° - 105°); (d) ref. ²²); 660 MeV protons (20° - 90°). Lines are the result of a fit on a spectrum $dN/dT \approx \exp(-T/T_0)$. Open circles have not been used.

REF.

M. Bernheim, A. Bussiere, A. Gillebert, J. Mougey, Phan Xuan Ho,
M. Priou, D. Royer, I. Sick, and G.J. Wagner
Phys. Rev. Letters 32, 898 (1974)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

74 Be 1

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,EP	ABX	10* 60	D	497	MAG-D		DST

The reaction $^{12}\text{C}(e, e'p)$ at 497 MeV in conjunction with a distorted-wave impulse-approximation analysis was used to determine kinetic and separation energies of bound protons. The spectral function for separation energies less than 74 MeV provides only half of the total binding energy; i.e., the data do not satisfy Koltun's sum rule. The momentum distributions are compatible with elastic electron scattering.

*MISSING ENERGY MEV




FIG. 1. A "spectrum" (see text) of the reaction $^{12}\text{C}(e, e'p)$ with (solid line) and without (dashed line) radiative corrections. The cross section is zero within the error bars for $64 < E_m < 74$ MeV.




FIG. 2. Recoil momentum distributions in different regions of separation energies. The DWIA (solid line) and PWIA results (dashed line) were normalized by least-squares fits to the data.

TABLE I. Occupation numbers n , average kinetic energies $\langle T \rangle$, and separation energies $\langle E_m \rangle$. All energies are in MeV.

Energy region (MeV)	Experimental results ^a			DWIA corrections			Corrected values		
	n	$\langle T \rangle$	$\langle E_m \rangle$	νl_j	η	ΔT	n	$\langle T \rangle$	$\langle E_m \rangle$
15-25	1.7	16.3	16.9	$1p_{3/2}$	0.66	2.14	2.6	18.4	16.9
25-74	0.56	11.5	39.7	$1s_{1/2}$	0.52	1.35	1.1	13.4	38.7
15-74							3.7	16.9	23.4

^aThe estimated relative error is 20% for n ; the statistical ones are 3% for $\langle T \rangle$ and 1.5% for $\langle E_m \rangle$.

METHOD					REF. NO.		
					74 Bo 14		
					hmg		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	SPC	150-345	C	345	EMU-D		DST
G,PI-	SPC	150-345	C	345	EMU-D		DST
G,P	SPC	46-345	C	345	EMU-D		DST

Energy spectra of charged pions and protons have been measured at angles of 30, 60, 90, 120, and 150° in the laboratory system for a bremsstrahlung maximum energy of 345 MeV. Charged pions with kinetic energies from 10 to 100 MeV and protons with energies from 30 to 160 MeV emitted from C¹² nuclei were detected by the emulsion technique. The differential cross sections for photoproduction of protons agree with calculations according to the cascade model. For pions the agreement is poorer.




FIG. 3. Energy spectra of photoprotons from C¹² for a bremsstrahlung maximum energy 345 MeV. Hollow circles - data of the present work, solid circles - data of Kim et al.^[13], triangles - ref. 14, squares - ref. 15. The stepped line is a calculation according to the cascade model, and the smooth curve according to the quasideuteron model; the dashed lines show calculations by Gabriel and Alsmiller^[1] according to the cascade model.




FIG. 1. Energy spectra of charged pions from C¹² for bremsstrahlung maximum energy 345 MeV: Circles - pi⁺ mesons, triangles - pi⁻ mesons, from the present work; squares - pi⁺ mesons from ref. 11. The stepped line shows the pi⁺ energy spectrum calculated with the cascade model.




FIG. 2. Angular distributions of photoprotons with kinetic energies $T_p = 155$ MeV emitted from C¹² for bremsstrahlung maximum energy $E_0 = 345$ MeV (hollow points); the solid points are data of Kim et al.^[15] at $T_p = 156$ MeV, $E_0 = 335$ MeV. The designations of the curves are the same as in Fig. 3.

(over)

- ⁷ T. A. Gabriel and R. G. Alsmiller, Jr., Phys. Rev. 182, 1035 (1969).
- ⁸ J. M. Peterson, W. S. Gilbert, and R. S. White, Phys. Rev. 81, 1003 (1951).
- ⁹ D. Luckey, Phys. Rev. 97, 469 (1955).
- ¹⁰ T. R. Palfrey, B. M. K. Nefkens, L. Mortara, and F. J. Loeffler, Phys. Rev. 122, 1323 (1961).
- ¹¹ S. Kabe, S. Kato, T. Kifune, Y. Kimura, et al., J. Phys. Soc. Jap. 19, 1800 (1964).
- ¹² N. V. Goncharov, A. L. Derebchinskiĭ, O. G. Kononov, S. G. Tonapetyan, and V. M. Khvorostyan, Zh. Eksp. Teor. Fiz. 64, 67 (1973) [Sov. Phys.-JETP 37, 38 (1973)].
- ¹³ B. T. Fedl, R. D. Godbole, A. Odian, F. Scherb, P. C. Stein, and A. Wattenberg, Phys. Rev. 94, 1000 (1954).
- ¹⁴ R. J. Cence and B. J. Moyer, Phys. Rev. 122, 1634 (1961).
- ¹⁵ Y. S. Kim, F. F. Liu, F. J. Loeffler, and T. R. Palfrey, Phys. Rev. 129, 1362 (1963).

METHOD				REF. NO.			
				74 Ce 4 hmg			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	12- 15	C	50	MAG-D		180

G-WIDTH OF 12.71 LEV

The ground-state magnetic dipole transition width of the 1^* level at 12.71 MeV in ^{12}C has been measured to be 0.35 ± 0.05 eV. This value is used to calculate the total width of the level based on a previous measurement of the relative ground-state γ width. A total width of 14.6 ± 2.6 eV is thus calculated. A model-dependent measurement of the isospin mixing between this level and the 15.11-MeV level yields the values $\beta = 0.19 \pm 0.01$ or 0.05 ± 0.01 .




FIG. 1. Spectrum of electrons inelastically scattered at 180° from ^{12}C .

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
74 Di 7		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,2P3N	ABY	THR* 1	C	300* 1	ACT-I		4PI

1=1 GEV




Fig. 1. Cross section per equivalent quantum σ_Q vs the bremsstrahlung maximum energy. Curve a is an eye-fit of the present work values (open circles) for the reaction $^{23}\text{Na}(\gamma, 2p 3n)^{18}\text{F}$. Curve b is an eye-fit of the present work values (open triangles) and the values taken from Ref. [2] (filled triangles) for the reaction $^{27}\text{Al}(\gamma, 2p 3n)^{22}\text{Na}$. Curve c is the best-fit of the present work values and values taken from Refs. [18, 20-22] for the reaction $^{12}\text{C}(\gamma, 2p 3n)^7\text{Be}$; for the sake of simplicity, experimental points have not been reported. Curve d is an eye-fit of the present work values (reversed open triangles) for the reaction $^{16}\text{O}(\gamma, 2p 3n)^{11}\text{C}$.

²V. di Napoli, A.M. Lacerenza, F. Salvetti, H.G. de Carvalho, and J.B. Martins, Nuovo Cimento Lettere 1, 835 (1971).

¹⁸V. di Napoli, D. Margadonna, F. Salvetti, H.G. de Carvalho, and J.B. Martins, Nucl. Inst. Method. 93, 77 (1971).

²⁰V. di Napoli, F. Dobici, O. Forina, F. Salvetti and H.G. de Carvalho, Lett. Nuovo Cimento 55B, 95 (1968).

²¹V. di Napoli, Nucl. Inst. Meth. 69, 155 (1969).

²²V.I. Noga, Yu.N. Ranyuk, P.V. Sorokin, and V.A. Tkachenko, Ukr. J. Phys. 16, 1850 (1971) in Russian.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
74 Do 5		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	ABX	100-999	C	125-999	TEL-D		DST

Tables of yield curves are given.

999=1 GEV




Fig. 3. Ratios of the experimental yields of protons of 82 MeV from carbon and 86 MeV from beryllium at 40° laboratory. The curves are the ratios of 4th-order fits σ_0 (full line) and σ_x (dashed line) as defined in the text

5 Antuf'ev, Yu.P., Agranovich, V.L., Ganenko, V.B., Kuz'menko, V.S., Miroshnichenko, I.I., Sorokin, P.V., Sanin, V.M.: Sov. Jour. Nucl. Phys. 9, 538 (1969).




Fig. 2. Comparison of the yields of protons of energy 99 MeV at 40° laboratory (this experiment: open circles) with the yields of proton of energy 97 MeV at 30° laboratory (Ref. 5: full circles). The vertical arrows indicate the mean photon energies for detecting protons from π^0 -photoproduction from hydrogen for identical kinematical conditions




Fig. 1. The results of unfolding 4th-order fits to the yields of proton at 40° and 70° lab from carbon. The vertical and horizontal bar indicate the mean photon energies and the photon energy resolutions which would be given by kinematically similar measurements of π^0 -photoproduction from hydrogen

(over)




Fig. 4. Ratios of the experimental yields at 40° and 70° laboratory from carbon for two proton energies. The curves are the ratios of 4th-order fits σ_Q (full line) and σ_L (dashed line) as defined in the text. The curves for the other proton energies are similar. The vertical and horizontal bars are as in Fig. 1

V. D. Epaneshnikov, V.M. Kuznetsov, and O.I. Stukov
 REF. Yad. Fiz. 19, 483 (1974)
 Sov. J. Nucl. Phys. 19, 242 (1974)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		74 Ep 2
		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	ABX	150-375	C	120-375	ACT-I		4PI




FIG. 2.

FIG. 2. Yield curve of the reaction $^{12}\text{C}(\gamma, \pi^-)^{12}\text{N}$ per equivalent quantum.




FIG. 3.

FIG. 3. Total cross sections for the reaction $^{12}\text{C}(\gamma, \pi^-)^{12}\text{N}$.

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	egf
	74 Fi 6	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XP	ABX	0- 20	D	60-100	MAG-D		DST




Figure 1. The proton spectrum from the $^{12}\text{C}(\gamma, p)$ reaction for $E_\gamma = 60 \pm 1$ MeV, $\theta_p(\text{lab}) = 45^\circ$. The excitation energy in the residual nucleus, ^{11}B , is shown, and arrows indicate the low lying states.




Figure 2. Angular distributions, $(d\sigma/d\Omega)_{\text{cm}}$ against $(\theta_p)_{\text{cm}}$, at: (a) $E_\gamma = 60$ MeV, (b) 80 MeV and (c) 100 MeV for the $^{12}\text{C}(\gamma, p)$ reaction leading to ^{11}B states below 7 MeV excitation. Full circles: ground state and 2.1 MeV state only; open circles: all states below 7 MeV. The errors shown are purely statistical; an additional systematic uncertainty of $\pm 10\%$ is estimated. The theoretical results of Weise (1972, private communication) are as follows. Curve 1: plane-wave approximation for outgoing protons. Curve 2: a continuum wavefunction in a complex energy-dependent Woods-Saxon potential for the outgoing protons. Curve 3(3'): initial- and final-state correlations simulated by a Jastrow correlation factor corresponding to a Gaussian momentum-exchange distribution of width $q_e = 100$ MeV/c centred at $q_e = 300$ (350) MeV/c (otherwise like curve 2). The initial-state wavefunctions are calculated in a Woods-Saxon well chosen to reproduce the experimental separation energies, except in the plane wave calculation for which harmonic oscillator wavefunctions were used.

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	74 Sc 8

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,N	ABX	50-150	D	63-150	TOF-D		DST

PHOTON DIFF.




Fig. 4. Measured neutron yield. The data selected here qualitatively show the energy and angle dependence of the processes under investigation.




Fig. 5. Measured cross sections for the photoproduction of neutrons on ^{12}C by quasimonochromatic photons. The energy values given correspond to the photon end energies. The arrows mark the neutron energies E_n calculated with formula (7) for the medium photon energy.

Fig. 6. Measured cross sections for the photoproduction of neutrons on ^{12}C by quasimonochromatic photons. The energy values given correspond to the photon end energies. The arrows mark the neutron energies E_n calculated with formula (7) for the medium photon energy.

METHOD

Page 3 of 3

REF. NO.	egf
74 Sc 8	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 10. Cross section for the $^{12}\text{C}(\gamma, n)$ reaction. The solid lines are calculated using the modified quasi-deuteron model of subsect. 3.5.

ELEM. SYM.	A	Z
C	12	6

METHOD			REF. NO.				
			74 Sh 3		hmg		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
HE,G	ABX	28-45	D	3-24	NAI-D		DST

HE=HE3, 4 LEVELS




FIG. 9. 90° differential cross sections for the reaction ${}^3\text{Be}({}^3\text{He}, \gamma_0){}^{12}\text{C}$. Points represented by x's were obtained with a thinner target. Error bars include only statistical errors.




FIG. 15. Angular distributions of the reaction ${}^3\text{Be}({}^3\text{He}, \gamma_0){}^{12}\text{C}$. Data are shown as bars, the height of which represents the statistical errors. The solid lines are Legendre polynomial fits, $d\sigma/d\Omega = \sum_{n=0}^4 A_n P_n(\cos\theta_{c.m.})$. Units are arbitrary.




FIG. 21. Total cross section and anisotropy coefficients for the reaction ${}^3\text{Be}({}^3\text{He}, \gamma){}^{12}\text{C}$. Shown are the results of fitting the ${}^3\text{Be}({}^3\text{He}, \gamma){}^{12}\text{C}$ angular distributions with $d\sigma/d\Omega = (\sigma/4\pi)[1 + \sum_{i=1}^4 A_i P_i(\cos\theta)]$. Errors are wholly statistical.

TABLE I. Levels identified in ${}^{12}\text{C}$. Experimental energies and widths extracted from ${}^3\text{Be}({}^3\text{He}, \gamma){}^{12}\text{C}$ excitation functions. The resonance parameters for each of the three decays, $\gamma_0, \gamma_1, \gamma_2$, are listed separately and, for those resonances appearing in several decays, the combined values are given. All values are in MeV.

Combined		γ_0 decay		γ_1 decay		γ_2 decay	
E	Γ	E	Γ	E	Γ	E	Γ
28.83 ± 0.04	1.54 ± 0.09	28.84 ± 0.04	1.55 ± 0.09	28.78 ± 0.07	1.19 ± 0.44
30.29 ± 0.03	1.96 ± 0.15	30.29 ± 0.03	1.96 ± 0.15
31.16 ± 0.03	2.10 ± 0.15	31.16 ± 0.03	2.10 ± 0.15
32.29 ± 0.04	1.32 ± 0.23	32.30 ± 0.04	1.12 ± 0.39	32.22 ± 0.11	1.43 ± 0.29
33.47 ± 0.21	1.93 ± 0.05	33.47	2.03 ± 0.06	33.47 ± 0.21	1.82 ± 0.11

6

C. Brassard, Ph.D. thesis, Yale Univ. 1970 (unpublished); C. Brassard, H.D. Shay, J.P. Coffin, W. Scholtz, D.A. Bromley, Phys.Rev. C3, 53 (1972).

7

R.G. Allas, S.S. Hanna, L. Meyer-Schutzmeister, R.E. Segel, Nucl. Phys. 58, 122 (1964).

TABLE II. Average cross section of radiative capture reactions in ${}^{12}\text{C}$ and their ratios.

Reaction	Energy range (MeV)	Average 90° differential cross section			
		γ_0 decay	γ_1 decay	γ_2 decay	γ_3 decay
${}^{11}\text{B}(p, \gamma){}^{12}\text{C}$	20-28 ^a	3.8	3.0	≈ 0.02 ^b	...
${}^9\text{Be}({}^3\text{He}, \gamma){}^{12}\text{C}$	28-35 ^c	0.93	1.63	$\ll 0.18$	0.32
${}^{10}\text{B}(d, \gamma){}^{12}\text{C}$	28-33 ^d	< 0.003	< 0.005

Ratios of cross sections over the energy interval 28-35 MeV

$$r_1 \equiv \frac{\sigma({}^9\text{Be}({}^3\text{He}, \gamma){}^{12}\text{C})}{\sigma({}^{11}\text{B}(p, \gamma_0){}^{12}\text{C})} = 1/11.8$$

$$r_{13} \equiv \frac{\sigma({}^{11}\text{B}(p, \gamma_1){}^{12}\text{C})}{\sigma({}^{11}\text{B}(p, \gamma_0){}^{12}\text{C})} = 0.34$$

$$r_2 \equiv \frac{\sigma({}^{10}\text{B}(d, \gamma_0){}^{12}\text{C})}{\sigma({}^{11}\text{B}(p, \gamma_0){}^{12}\text{C})} \approx 1/300$$

$$r_{21} \equiv \frac{\sigma({}^9\text{Be}({}^3\text{He}, \gamma_1){}^{12}\text{C})}{\sigma({}^9\text{Be}({}^3\text{He}, \gamma_0){}^{12}\text{C})} = 1.64$$

$$r_{11} \equiv \frac{\sigma({}^{11}\text{B}(p, \gamma_1){}^{12}\text{C})}{\sigma({}^{11}\text{B}(p, \gamma_0){}^{12}\text{C})} = 1.75$$

$$r_{22} \equiv \frac{\sigma({}^9\text{Be}({}^3\text{He}, \gamma_2){}^{12}\text{C})}{\sigma({}^9\text{Be}({}^3\text{He}, \gamma_0){}^{12}\text{C})} = 0.79$$

$$r_{12} \equiv \frac{\sigma({}^{11}\text{B}(p, \gamma_2){}^{12}\text{C})}{\sigma({}^{11}\text{B}(p, \gamma_0){}^{12}\text{C})} \leq 0.05$$

$$r_{23} \equiv \frac{\sigma({}^9\text{Be}({}^3\text{He}, \gamma_3){}^{12}\text{C})}{\sigma({}^9\text{Be}({}^3\text{He}, \gamma_0){}^{12}\text{C})} = 4.73$$

^a Data of Allas *et al.* (Ref. 7) normalized to those of Brassard (Ref. 6) by factor of 1/1.6.

^b Data of P. Paul, private communication.

^c Data of Brassard (Ref. 6).

^d This work.

R.R. Whitney, I. Sick, J.R. Ficenec, R.D. Kephart, and
 W.P. Trower
 REF. Phys. Rev. C9, 2230 (1974)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
74 Wh 3		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	0-300	D	500	MAG-D		60

See further analysis of this data in reference 79Zil

QUASIELASTIC SCAT




FIG. 1. The measured quasielastic peaks; the errors on the data points do not include an over-all 3% normalization uncertainty. The solid curve is a fit by the Fermi-gas model which yielded k_F (in MeV/c) and $\bar{\epsilon}$ (in MeV) as follows: (a) ${}^6\text{Li}$ (169, 17); (b) ${}^{12}\text{C}$ (221, 25); (c) ${}^{24}\text{Mg}$ (235, 32); (d) ${}^{40}\text{Ca}$ (249, 33); (e) ${}^{58}\text{Ni}$ (260, 36); (f) ${}^{89}\text{Y}$ (254, 39); (g) ${}^{118}\text{Sm}$ (260, 42); (h) ${}^{181}\text{Ta}$ (265, 42); (i) ${}^{208}\text{Pb}$ (265, 44). The fitting uncertainty in k_F is ± 5 MeV/c and in $\bar{\epsilon}$ it is ± 3 MeV. The small-amplitude dashed curve is the s-wave π -production contribution, the dot-dashed curve is the isobaric excitation, and the large-amplitude dashed curve is the total result.

(over)

TABLE I. Proton-normalized and radiative-corrected cross sections $d^2\sigma/d\Omega dE' = (N \pm \Delta N) \times 10^{-10}$ in mb/sr MeV, for $E = 500$ MeV and $\theta = 60^\circ$.

E' (MeV)	^6Li			^{12}C			^{24}Mg			^{40}Ca			^{82}Se			^{118}Sn			^{180}Pt			^{208}Pb					
	N	ΔN	n	N	ΔN	n	N	ΔN	n	N	ΔN	n	N	ΔN	n	N	ΔN	n	N	ΔN	n	N	ΔN	n			
480.0	1.79	0.19	7	3.03	0.42	7	1.22	0.17	6	1.71	0.19	6		
474.0	1.02	0.13	7		
470.0	1.72	0.18	7	5.75	0.52	7	1.55	0.15	6	3.90	0.29	6	5.85	0.41	6	7.09	0.67	6	7.00	0.68	6		
464.0	2.49	0.29	7	1.38	0.11	6	1.91	0.17	6	2.72	0.15	6	4.48	0.33	6	5.68	0.37	6	8.32	0.71	6	1.16	0.08	5	9.82	0.79	6
460.0	2.96	0.30	7	1.20	0.09	6	2.58	0.19	6	1.54	0.10	5	1.81	0.10	5	
454.1	5.02	0.17	7	9.21	0.71	7	2.96	0.20	6	1.20	0.17	6	7.00	0.41	6	1.07	0.05	5	1.85	0.09	5	1.70	0.11	5	2.31	0.11	5
450.0	8.92	0.47	6	1.03	0.05	5	1.83	0.09	5	
444.3	8.68	0.58	7	3.26	0.07	6	4.11	0.25	6	6.67	0.27	6	2.77	0.14	5	2.74	0.12	5		
440.0	1.11	0.06	6	2.59	0.13	6	5.23	0.26	6	3.43	0.16	5	3.24	0.13	5		
434.2	1.32	0.06	6	2.99	0.14	6	5.50	0.26	6	8.74	0.35	6	1.40	0.07	5	1.90	0.07	5	2.27	0.09	6	3.14	0.15	5	3.34	0.14	5
430.0	1.40	0.07	5	2.11	0.08	5	2.77	0.11	5	
424.3	2.12	0.08	6	3.75	0.15	6	7.31	0.29	6	1.12	0.04	5	1.54	0.08	5	2.31	0.09	5	2.88	0.12	5	4.43	0.18	5	4.74	0.19	5
414.4	2.88	0.12	6	1.75	0.19	6	8.78	0.35	6	1.32	0.05	5	1.78	0.10	5	2.88	0.11	5	3.40	0.14	5	4.98	0.20	5	5.64	0.23	5
404.5	3.51	0.14	6	5.46	0.22	6	1.02	0.01	5	1.56	0.06	5	2.09	0.08	5	3.39	0.12	5	3.90	0.10	5	5.89	0.24	5	6.57	0.26	5
400.0	6.35	0.25	6	1.09	0.01	5	2.35	0.09	5	3.34	0.13	5	4.29	0.17	5	6.56	0.27	5	7.00	0.28	5	
391.7	4.16	0.17	6	6.32	0.26	6	1.15	0.05	5	1.75	0.07	5	2.22	0.09	5	3.41	0.14	5	4.56	0.18	5	6.29	0.25	5	7.25	0.29	5
385.7	4.55	0.18	6	7.09	0.28	6	1.23	0.05	5	1.86	0.07	5	2.51	0.10	5	3.91	0.16	5	4.88	0.19	5	6.36	0.26	5	7.88	0.32	5
374.9	4.76	0.19	6	6.97	0.28	6	1.33	0.05	5	1.94	0.08	5	2.72	0.11	5	4.02	0.16	5	4.88	0.19	5	6.87	0.28	5	8.19	0.33	5
365.0	4.56	0.18	6	7.26	0.29	6	1.32	0.05	5	2.06	0.08	5	2.69	0.10	5	4.01	0.16	5	5.34	0.21	5	7.77	0.31	5	8.61	0.34	5
360.0	4.50	0.18	6	6.61	0.28	6	1.32	0.05	5	2.88	0.11	5	4.11	0.16	5	5.69	0.23	5	7.92	0.33	5	8.42	0.34	5	
355.2	4.35	0.17	6	6.97	0.28	6	1.16	0.05	5	2.08	0.08	5	2.69	0.11	5	4.23	0.17	5	5.22	0.21	5	7.51	0.30	5
345.3	3.68	0.15	6	6.51	0.26	6	1.35	0.05	5	1.85	0.07	5	2.72	0.11	5	4.02	0.16	5	5.37	0.21	5	7.59	0.30	5	8.83	0.35	5
345.4	2.90	0.12	6	5.91	0.21	6	1.29	0.05	5	1.87	0.08	5	2.48	0.10	5	4.08	0.16	5	4.92	0.19	5	7.44	0.29	5	8.68	0.35	5
335.5	2.59	0.10	6	5.23	0.21	6	1.05	0.01	5	2.48	0.11	5	3.78	0.15	5	4.83	0.19	5	6.93	0.28	5	
320.0	2.35	0.09	5	3.34	0.14	5	4.53	0.18	5	
315.7	2.16	0.10	6	4.13	0.18	6	9.11	0.38	6	1.47	0.06	5	2.26	0.09	5	3.43	0.14	5	4.34	0.17	5	6.51	0.26	5	7.76	0.31	5
305.8	1.84	0.09	6	3.79	0.15	6	8.61	0.32	6	1.23	0.05	5	2.01	0.08	5	3.27	0.13	5	4.32	0.17	5	6.11	0.24	5	6.92	0.28	5
300.0	1.97	0.08	5	3.11	0.12	5	4.03	0.16	5	
295.9	1.55	0.09	6	3.38	0.14	6	6.77	0.29	6	9.97	0.40	6	1.80	0.07	5	3.02	0.12	5	3.74	0.15	5	5.38	0.22	5	6.73	0.27	5
285.9	1.50	0.09	6	2.96	0.14	6	6.64	0.31	6	9.73	0.39	6	1.72	0.07	5	2.60	0.13	5	3.55	0.15	5	4.92	0.23	5	6.30	0.29	5
276.2	1.14	0.08	6	2.64	0.13	6	6.03	0.32	6	8.35	0.41	6	1.50	0.07	5	2.64	0.13	5	3.10	0.15	5	5.22	0.24	5	5.73	0.30	5
266.3	1.08	0.08	6	2.61	0.11	6	5.32	0.33	6	8.57	0.43	6	1.31	0.08	5	2.37	0.14	5	2.72	0.16	5	4.62	0.26	5	5.51	0.31	5
260.0	1.39	0.08	5	1.95	0.13	5	2.91	0.18	5	
256.1	1.28	0.09	6	2.13	0.15	6	5.71	0.35	6	8.33	0.45	6	1.27	0.08	5	2.27	0.14	5	2.87	0.18	5	4.57	0.28	5	6.03	0.33	5
246.5	1.20	0.09	6	2.55	0.16	6	5.17	0.36	6	8.55	0.46	6	1.39	0.09	5	2.14	0.13	5	2.95	0.19	5	4.33	0.28	5	5.40	0.35	5
236.7	1.15	0.10	6	2.51	0.16	6	5.18	0.36	6	8.71	0.51	6	1.31	0.09	5	2.21	0.15	5	3.02	0.20	5	4.35	0.30	5	4.90	0.34	5
226.8	1.27	0.11	6	2.88	0.19	6	5.62	0.42	6	8.72	0.51	6	1.29	0.10	5	2.29	0.16	5	2.73	0.20	5	4.57	0.30	5	5.88	0.37	5
216.9	1.41	0.14	6	2.91	0.21	6	6.15	0.49	6	9.81	0.56	6	1.11	0.10	5	2.38	0.17	5	3.26	0.22	5	5.42	0.36	5	5.76	0.38	5
207.0	1.60	0.16	6	2.91	0.21	6	6.59	0.52	6	1.02	0.06	5	1.43	0.11	5	2.51	0.18	5	3.24	0.22	5	5.58	0.37	5	6.11	0.40	5
197.2	1.78	0.17	6	3.12	0.24	6	7.01	0.59	6	1.59	0.12	5	2.77	0.20	5	3.43	0.24	5	5.67	0.38	5	5.99	0.41	5	

REF. K. Wienhard, K. Bangert, R. Stock, and H. Wolf
 Z. Phys. 270, 93 (1974)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
74 Wi 4		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XP	NOX	20- 29	C	31	SCD-D		90

P SPECTRUM




Fig. 2. Photoproton spectrum from ^{12}C at 90° , divided by the bremsstrahl number spectrum, which was normalized to unity at 22.3 MeV. It was assumed that ground state protons only are emitted. The experimental energy resolution is 100 keV at 20 MeV and 40 keV at 29 MeV.

REF.

J. Ahrens, H. Borchert, K.H. Czock, H.B. Eppler, H. Gimm,
 H. Gundrum, M. Kroning, P. Riehn, G. Sita Ram, A. Zieger,
 and B. Ziegler
 Nucl. Phys. A251, 479 (1975)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

75 Ah 3

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	10-160	C	140-275	MGC-D		4PI

920+

TABLE 2

The moments of the experimental nuclear cross section distributions integrated from 10 MeV to the energy E , and their statistical errors

	E (MeV)	Σ_{-2}		Σ_{-1}		Σ_0		Σ_{+1}		Σ_{+2}	
		(mb/MeV)	$\pm(\%)$	(mb)	$\pm(\%)$	(mb·MeV)	$\pm(\%)$	(b·MeV ²)	$\pm(\%)$	(b·MeV ³)	$\pm(\%)$
Li	100	0.196	1.1	4.64	1.0	143	1.7	5.82	3.1	305	5
	140	0.197	1.1	4.79	1.0	161	1.9	8.03	3.4	577	5
	210	0.198	1.1	5.03	1.0	206	2.0	16.60	3.7	2220	5
Be	100	0.192	2.5	5.19	1.5	173	2.0	7.11	3.4	362	5
	140	0.194	2.5	5.33	1.5	189	2.1	9.09	3.6	600	6
	210	0.195	2.5	5.58	1.5	236	2.1	17.80	3.5	2240	5
C	100	0.313	1.7	8.81	1.1	291	1.6	12.00	2.9	630	4
	140	0.316	1.7	9.18	1.2	334	2.2	17.10	5	1250	7
O	100	0.580	1.6	14.50	1.3	432	2.0	16.00	4	748	8
	140	0.585	1.6	15.10	1.3	508	2.5	25.20	5	1880	8
Al	100	1.10	1.8	25.70	1.5	739	2.6	27.9	5	1400	8
	140	1.11	1.8	26.3	1.7	807	3.9	36.4	9	2450	16
Ca	100	2.22	1.2	45.5	1.5	1120	3.6	34.9	9	1430	18
	140	2.23	1.2	46.8	1.7	1290	4.6	56.6	11	3710	19




Fig. 4. The same as fig. 2 for C.

Fig. 2. Total photonuclear cross section for natural Li. The error bars indicate one standard deviation of counting statistics from the main spectrometer. The dashed lines along the abscissa indicate the uncertainty due to counting statistics in the normalizing spectrometer. Oscillations of the base line within this area are possible, the period of these oscillations, however, must not be smaller than 10% in photon energy. The dashed and dotted lines through the cross section values have been drawn to guide the eye.

REF. U. Kneissl, E. A. Koop, G. Kuhl, K. H. Leister, A. Weller
 Nucl. Instrum. Methods 127, 1 (1975)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				75 Kn 8		egf	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	ABX	19- 32	D	19- 32	MOD-I		4PI

103E




Fig. 5. $\sigma(\gamma, \text{Tn})$ for ^{12}C .

REF.

H. Schier and B. Schoch
Nuovo Cimento Lett. 12, 334 (1975)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		75 Sc 7
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	62- 66	D	62- 66	TOF-D		DST

62.7-65.7 MEV

TABLE I. - Measured cross-sections obtained from Fig. 1-5. The photon energy corresponds to the maximum value of the difference spectrum.

Nucleus	Photon energy (MeV)	Excitation energy (MeV)	$d\sigma/d\Omega$ (40.2°) ($\mu\text{b}/\text{sr}$)	$d\sigma/d\Omega$ (90°) ($\mu\text{b}/\text{sr}$)
¹² O	63.0	ground state	5.0 ± 0.75	
		$0 \div 6.18$	20.7 ± 3.1	
¹² C	62.75	ground state	11.1 ± 0.8	7.8 ± 0.36
		$0 \div 4.79$	22.5 ± 1.1	17.1 ± 0.56
⁹ Be	61.25	ground state	0.8 ± 0.13	0.85 ± 0.1
		$0 \div (\gamma, np)$ threshold	3.4 ± 0.3	4.1 ± 0.22




Fig. 2.

Fig. 2. - Cross-section for the photoproduction of neutrons in ¹²C by quasi-monochromatic photons. The dashed lines show the photon spectrum fitted to the ground state, the 1.99 MeV level and two excited levels centred at 4.55 MeV in ¹²C. The solid line gives the sum of these levels. A calculation for the (γ, np) reaction leads to the dashed-dotted line. (65.6 - 62.6) MeV, $\theta_1 = 40.2^\circ$.




Fig. 3.

Fig. 3. - The same as Fig. 2 for $\theta_1 = 90^\circ$.

REF.

S. G. Tonapetyan, N. V. Goncharov, and V. M. Khvorostyan
 Yad. Fiz. 22, 440 (1975)
 Sov. J. Nucl. Phys. 22, 226 (1976)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	75 To 4	egf
----------	---------	-----

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABY	150-400	C	300,400	BBL-D		90
G,PI-	ABY	150-400	C	300,400	BBL-D		90
G,P	ABY	96-400	C	300,400	BBL-D		90




FIG. 1. Energy spectra of π^+ and π^- mesons. $E_{\pi}^{\max} = 300$ MeV, $\theta_{\text{lab}} = 90 \pm 7^\circ$. Points: \circ - ^{12}C , \triangle - ^{28}Si , \times - ^{40}Ca , \diamond - ^{93}Nb , \square - ^{181}Ta .




FIG. 3. Energy spectra of protons. $E_{\pi}^{\max} = 500$ MeV, $\theta_{\text{lab}} = 90 \pm 7^\circ$. Points: \circ - ^{12}C , \triangle - ^{28}Si , \times - ^{40}Ca , \diamond - ^{93}Nb , \square - ^{181}Ta , \circ -data from ref. 5, \square -data from ref. 6.




FIG. 2. Energy spectra of π^+ and π^- mesons. $E_{\pi}^{\max} = 400$ MeV, $\theta_{\text{lab}} = 90 \pm 7^\circ$. The points are the same as in Fig. 1.





FIG. 4. Energy spectra of protons. $E_{\pi}^{\max} = 400$ MeV, $\theta_{\text{lab}} = 90 \pm 7^\circ$. The points are the same as in Fig. 1; \square -data from ref. 7.

FIG. 5. Dependence of yields of π mesons 1, protons 2, and the sum of π -meson and proton yields 4 as a function of mass number of the nucleus. The dashed line 3 is the theory. Points: \circ -experimental differential cross sections for pions of all signs, \triangle -differential cross sections for protons emitted at the same angle $\theta_{\text{lab}} = 90^\circ$, \square -combined values of these differential cross sections. The statistical errors are shown.



⁵P.C.Murray et al., Phys.Rev. 94, 764 (54).

⁶C.Levinthal et al., Phys.Rev. 82, 822 (51).

⁷P.Dougan et al., LUSY Preprint 1002 (1970).

REF. E. Wolyneć, G. Moscati, J. R. Moreira, O. D. Goncalves,
M. N. Martins
Phys. Rev. C11, 1083 (1975)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

75 Wo 2

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	RLY	19- 40	C	19- 40	ACT-I		4PI

RATIO (G,N)/(E,N)




FIG. 1. Measured F for $^{12}\text{C}(\gamma, n)$. The full circles and triangles refer to our measurements with Teflon and PVC targets, respectively. Full squares are data from Ref. 7. Dashed curve is a polynomial fit to the points. Full curves are F_{PW} and F_{DW} predictions.

$$F_{PW}^{(\gamma, n)} = (N_r / Z_r^2 \gamma_e^2 N_r)$$

$$\times \frac{\int_0^{E_1 - m_e} \sigma_\gamma(\omega) \phi(E_1, \omega, Z_r) (d\omega/\omega)}{\int_0^{E_1 - m_e} \sigma_\gamma(\omega) N_{PW}^{E_1}(E_1, \omega) (d\omega/\omega)}, \quad (10)$$

$$F_{DW}^{(\gamma, n)} = (N_r / Z_r^2 \gamma_e^2 N_r)$$

$$\times \frac{\int_0^{E_1 - m_e} \sigma_\gamma(\omega) \phi(E_1, \omega, Z_r) (d\omega/\omega)}{\int_0^{E_1 - m_e} \sigma_\gamma(\omega) N_{DW}^{E_1}(E_1, \omega, Z_r) (d\omega/\omega)}, \quad (11)$$

REF. A.M. Bernstein, N. Paras, W. Turchinets, B. Chasan,
and E. C. Booth
Phys. Rev. Lett. 37, 819 (1976)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
76 Be 6		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	ABX	150-191	C	150-191	ACT-I		4PI

$$Y(191) = 0.9 \pm 0.2 \text{ub}/Q$$

B+ ACTIV-NITROGEN 12

The total cross section for the reaction $^{12}\text{C}(\gamma, \pi^-)^{12}\text{N}$ has been determined by observation of the residual ^{12}N radioactivity. The cross section was extracted from the bremsstrahlung excitation function which was measured in the region between 3.6 and 12.6 MeV above the threshold, with one point 33.5 MeV above the threshold. The variation of the measured cross section with energy is far more rapid than is predicted by calculations using the $\bar{e}\text{-}\bar{\nu}$ interaction. Even when the full interaction Hamiltonian is used, the experimental cross section rises somewhat more rapidly than predicted.




FIG. 2. Relative yield of the $^{12}\text{C}-^{12}\text{N}$ to the reaction $^{14}\text{N}(\gamma, 2\pi)^{12}\text{N}$ versus the end-point bremsstrahlung energy.




FIG. 3. (a) Yield and (b) cross section versus energy above threshold. The theoretical curves are from Koch (K) and from Ref. 19; curves 1 and 2 are from Nagl and Überall (Ref. 21). The best fit is a two-parameter fit with a step at the threshold and a linear increase with energy above the threshold. The shaded zone shows the errors in the best fit.

REF.

R. Carchon, R. Van de Vyver, H. Ferdinande, J. Devos,
E. Van Camp
Phys. Rev. C14, 456 (1976)

ELEM. SYM.

A

Z

C

12

6

METHOD

REF. NO.

76 Ca 5

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	18- 30	C	19- 30	SCD-D		DST

Expansion used: $d\sigma/d\Omega(E_\gamma, \theta) = A_0(E_\gamma) \left[1 + \sum_{i=1}^4 a_i(E_\gamma) P_i(\cos\theta) \right]$.




FIG. 4. $^{12}\text{C}(\gamma, p)^{11}\text{B}$ differential cross sections at seven angles in the laboratory system, derived from the proton spectra by assuming ground-state transitions only; values are plotted every 150 keV.

The energy spectra and angular distributions of photoprotons from the $^{12}\text{C}(\gamma, p)^{11}\text{B}$ reaction have been measured at seven angles simultaneously, using 30 MeV bremsstrahlung. The giant dipole resonance peaks at 22.3 MeV, and reaches an absolute cross section value of 13.1 ± 0.8 mb. The anisotropy parameter a_1 from the Legendre polynomial expansion of the angular distributions has an average value of about -0.55 but shows some structure especially around 25 MeV. On the other hand, the asymmetry coefficient a_1 is always positive and rises slowly with energy. Discussion of all coefficients leads to the conclusion that, although the photoneuclear absorption mechanisms in ^{12}C leading to photoproton emission is dominated by the E1 component, a nonnegligible E2 contribution (about 2%) might be present. A recent coupled-channel calculation by Hinkholz satisfactorily describes our results.

(over)




FIG. 7. Photoproton angular distribution coefficients deduced from the Legendre polynomial fit, as a function of center-of-mass excitation energy. The results from the theoretical calculations by Mshelia *et al.* (Ref. 26) (dashed line) and by Birkholz (Ref. 52) (solid line) are shown for comparison.




FIG. 5. The total photoproton cross section ($4\pi A_0$) assuming ground-state transitions only. Also shown are the theoretical predictions by Mshelia *et al.* (Ref. 26) and by Birkholz (Ref. 25).

- ²⁵J. Birkholz, Phys. Lett. 34B, 1 (1971); Nucl. Phys. A189, 385 (1972); F. Beck and J. Birkholz, in Proc. International Conf. on Photonuclear Reactions and Applications, Asilomar, 1973, Vol. 1, p.159.
- ²⁶E.D. Mshelia et al., Phys. Rev. Lett. 28, 847 (1972); E.D. Mshelia and R.F. Barrett, Nucl. Phys. A205, 581 (1973); Z. Phys. 261, 313 (1973).
- ⁵²J. Birkholz (private communication).

REF.

J. L. Matthews, D. J. S. Findlay, S. N. Gardiner,
R. O. Owens
Nucl. Phys. A267, 51 (1976)

ELEM. SYM.	A	Z
C	12	6
METHOD	REF. NO.	
	76 Ma 8	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	60-100	D	60-100	MAG-D		DST

TABLE I

Cross Section Table given.

Cross sections for the ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$ and ${}^{12}\text{C}(\gamma, p)$ reactions at $E_{\gamma}(\text{lab}) = 60 \text{ MeV}$, $\theta_{\text{lab}} = 45^\circ$ populating the excitation energy regions $E_x = 0-9 \text{ MeV}$ and $9-26 \text{ MeV}$ in the residual nuclei

Target nucleus	$(d\sigma/d\Omega)^{\text{stat.}}$ ($\mu\text{b}/\text{sr}$)	
	residual nucleus excitation energy E_x 0-9 MeV	9-26 MeV
${}^6\text{Li}$	5.5 ± 0.6	12.2 ± 2.0
${}^7\text{Li}$	9.5 ± 0.8	14.7 ± 3.0
${}^9\text{Be}$	31.8 ± 3.0	
${}^{12}\text{C}$	36.4 ± 3.9	

Only statistical errors are tabulated; there is an additional common systematic error of $\pm 22\%$.




Fig. 10. Angular distributions of photoprotons from ${}^{12}\text{C}$ at $E_{\gamma} = 60 \text{ MeV}$, populating states in ${}^{11}\text{B}$ as follows: solid circles, ground state; solid squares, 2.1 MeV state; open triangles, (4.4+5.0+6.7) MeV states.




Fig. 11. Angular distributions of photoprotons from ${}^{12}\text{C}$ at $E_{\gamma} = 80 \text{ MeV}$ and 100 MeV populating states in ${}^{11}\text{B}$ as follows: solid circles, ground state; open triangles, (4.4+5.0+6.7) MeV states.




Fig 13 The $^{12}\text{C}(\gamma, p)^{11}\text{B}$ differential and total cross sections as a function of photon energy: solid circles, present measurements for ground + 2.1 MeV states; open circles, corrected results from ref. 12) (see comments in caption to fig. 12); solid triangles, data from ref. 11); open triangles, estimated results obtained from the values of $d\sigma/d\Omega$ at 90° given in ref. 11); at these additional energies and the trend of the ratio of the 90° and 45° cross sections; open squares, results obtained from fig. 1 of ref. 13). The theoretical curves are discussed in sect. 6.

- 11 S. Penner et al., Phys. Rev. **114** (1959) 1101
- 12 G. Manuzio et al., Nucl. Phys. **A133** (1969) 225
- 25 G.G. Tarant et al., Sov. J. Nucl. Phys. **6** (1968) 816

TABLE 5

Total cross sections at $E_L(\text{lab}) = 60, 80$ and 100 MeV for the reactions $^6\text{Li}(\gamma, p)^5\text{He}$ (ground state + 2.6 MeV state), $^7\text{Li}(\gamma, p)^6\text{He}$ (ground state + 1.8 MeV state) and $^{12}\text{C}(\gamma, p)^{11}\text{B}$ (ground state, 2.1, 4.4, 5.0 and 6.7 MeV states in several combinations)

$E_L(\text{lab})$ (MeV)	Total cross section (μb)		
	$^6\text{Li}(\gamma, p)$ ($P_0 + P_1$)	$^7\text{Li}(\gamma, p)$ ($P_0 + P_1$)	$^{12}\text{C}(\gamma, p)$ ($P_0 + P_1$)
60	34.1 ± 1.6	43.8 ± 1.2	148.9 ± 4.6
80		11.39 ± 0.39	44.9 ± 2.1
100		17.2 ± 1.4	19.6 ± 1.1
			$P_0 + P_1 + \dots + P_n$
			238.1 ± 11.7
			65.4 ± 3.9

The error shown are statistical. There is an additional common systematic error of + 7.0.




Fig 8. Photoproton spectra for $E_L = 60 \pm 1$ MeV, $\theta = 45^\circ$ derived as described in the text from data such as those shown in fig. 7: (a) $^6\text{Li}(\gamma, p)^5\text{He}$; (b) $^7\text{Li}(\gamma, p)^6\text{He}$; (c) $^9\text{Be}(\gamma, p)^8\text{Be}$; (d) $^{12}\text{C}(\gamma, p)^{11}\text{B}$. The scale above each spectrum gives the excitation energy in the residual nucleus, with lines indicating the positions of known excited states. The location of the 1s shell peak observed in the (p, 2p) reaction is also marked.

REF.

D. Mehlig, K.H. Czock
Acta Physica Academiae Scientiarum Hungaricae 41, 55 (76)ELEM. SYM. A Z
C 12 6

METHOD

REF. NO.

76 Me 12

hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	15 (15.11)	C	UKN	NAI-D		135

The 15.11 MeV energy level in ^{12}C was excited by bremsstrahlung. The scattered photons were detected with a total γ absorption NaI(Tl) spectrometer set at 135° to the bremsstrahlung beam. The following level parameters were obtained: $\sigma_0^{\text{th}} = (17.9 \pm 0.6)$ barn, $\Gamma_{\text{tot}} = (69 \pm 4)$ eV and $\sigma_{\text{int}} = (1.36 \pm 0.12)$ mb MeV. The branching ratio to the first excited state and the ground state is $\Gamma_{\gamma 1}/\Gamma_{\gamma 0} = (3.6 \pm 1.0)\%$.

Table I

Parameters of the 15.11 MeV level

Reference	I_3 [mb. MeV]	σ_0^{th} barn	Γ_{tot} [eV]	$\frac{\delta}{\Gamma_{\text{tot}}}$	$\frac{\Gamma_{\text{exc MeV}}}{\Gamma_{\text{tot}}}$ %
FULLER [1]	1.90 ± 0.27	22.2 ± 2.2	79 ± 16	1	7
GARWIN [2]	2.33 ± 0.19	29.7 ± 1.1	64 ± 10	0.62 ± 0.1	5 ± 4
BUSSIERE [3]	2.45 ± 0.5	32	60 ± 8	0	11 ± 5
SCHMID [4]*	1.82 ± 0.12	32	45 ± 10	1	—
GUDDEN [5]*	1.79	32	35.5	—	—
KÜHNE [6]*	1.8 ± 0.2	32	39 ± 5	0.9 ± 0.3	—
present work	1.86 ± 0.12	17.9 ± 0.6	69 ± 4	0.48 ± 0.1	3.6 ± 1.0

REFERENCES

1. E. G. FULLER and E. HAYWARD, NBS-Report, 6255, 1959.
2. E. L. GARWIN, Phys. Rev., 114, 143, 1959.
3. A. BUSSIERE DE NERCY, Annales de Physique, 6, 1961.
4. H. SCHMID and W. SCHOLZ, Zeitschr. f. Physik, 175, 430, 1963.
5. F. GUDDEN, Thesis, Darmstadt 1964 — unpublished.
6. H. W. KÜHNE, P. AXEL and D. C. SUTTON, Phys. Rev., 163, 1278, 1967.

REF.

J. Mougey, M. Bernheim, A. Bussiere, A. Gillebert, Phan
 Xuan Ho, M. Priou, D. Royer, I. Sick, G. J. Wagner
 Nucl. Phys. A262, 461 (1976)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

76 Mo 5

egf

REACTION	RESULT*	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E, E/P$	ABX	$00^+ 57$	D	497	MAG-D		53

*MISSING ENERGY




Fig. 9. Missing energy spectra from $^{12}\text{C}(e, p)^{11}\text{B}$; (a) $0 \leq P \leq 36$ MeV/c, (b) $60 \leq P \leq 180$ MeV/c and (c) $0 \leq P \leq 60$ MeV/c for $20 \leq E \leq 60$ MeV.




Fig. 10. Momentum distribution from $^{12}\text{C}(e, p)^{11}\text{B}$; (a) $15 \leq E \leq 21.5$ MeV and (b) $30 \leq E \leq 50$ MeV. The solid and dashed lines represent DWIA and PWIA calculations respectively, with normalization obtained by a fit to the data.

REF. N. Nakamura, S. Hiramatsu, T. Kamae, H. Muramatsu,
N. Izutsu, Y. Watase
Nucl. Phys. A268, 381 (1976)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		76 Na 2
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	ABX	0* 60	D	700	MAG-D		DST

Abstract: The proton spectral function of ^{12}C was deduced by using the DWIA from the measurement of the reaction $^{12}\text{C}(e, e'p)$ at an incident electron energy of 700 MeV. The proton separation energy range of $E_s \leq 65$ MeV and the recoil momentum range of $K \leq 240$ MeV/c were covered in this experiment. The results were analyzed in terms of the independent-particle shell model using the proton single-particle wave functions consistent with the elastic electron scattering results, and the WKB approximation for the distorted proton wave. We also made a Monte Carlo study to estimate the background contribution from the multiple-collision processes to the spectral function. The results of this analysis indicate that the occupation probabilities given by the best fit to the data are smaller than the independent-particle shell-model values for both the 1p and 1s proton states, while the shapes of the distorted momentum distributions are well reproduced. Koltun's sum rule was tested taking the background contribution into account.

*SEPARATION ENERGIES




Fig. 3. Proton separation energy spectra for ^{12}C . The solid curves are the result of the DWIA fit with the momentum distributions calculated from the bound-state potentials given by Elton and Swift¹²⁾. The dashed curves show the contributions from the 1p and 1s states.




Fig. 9. Recoil momentum distributions for ^{12}C . For the meaning of the curves, see caption to fig. 3

¹²⁾ L.R.B. Elton et al., Nucl. Phys. A94 (1967) 52.

REF. M. Turk
Fizika 8, 173 (1976)

ELEM. SYM.	A	Z
C	12	6

METHOD
Remarks: Bremsstrahlung filtered by 40cm of aluminum. Cross section obtained by normalizing to Toms ($\gamma, 3\alpha$)

REF. NO. 76 Tu 3 egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PA	ABX	25- 42	C	42	EMU-D		4PI
G,DA	ABX	30- 42	C	42	EMU-D		4PI
G,3A	ABX	7- 42	C	42	EMU-D		4PI




Fig. 3. Cross section for the reaction $^{12}\text{C}(\gamma, 3\alpha)$ as function of gamma ray energy after normalizing the data (full points) to Toms results. Squares represent the spread of earlier measurements by various authors referenced in Ref.³.




Fig. 4. Measured cross sections for reactions (1-5) as functions of incident gamma ray energy.

³M.E.Toms, Nucl. Phys. 50, 561 (1964)

REF. V.G. Vlasenko, V.A. Gol'dshtein, A.V. Mitrofanova, V.I. Noga,
 Yu.N. Ranuuk, V.I. Startsev, P.V. Sorokin, Yu.N. Telegin
 Yad. Fiz. 23, 504 (1976)
 Sov. J. Nucl. Phys. 23, 265 (1976)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				76 V1 1		hmg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	100-600	D	812-999	MAG-D		14

999=1.396 GEV

Inelastic electron scattering has been used to measure the total hadronic cross sections for absorption of photons with energy 150-500 MeV by nuclei of C, Al, Ni, Mo, and W. The results obtained are compared with calculations carried out in the impulse approximation.




FIG. 1. Spectrum of electrons scattered by ^{12}C . The solid curve is elastic scattering distorted by radiation processes. The dashed curve is radiative quasielastic scattering.




FIG. 4. Total hadronic cross section for absorption of photons by the carbon nucleus. The curve is a calculation with Eq. (5). Points: — Ref. 1, ●—present work, ○—Ref. 4.




FIG. 3. Cross section for scattering of electrons by the carbon nucleus, assigned to one virtual photon. Points: ○—present work, △—Ref. 25, ●—data for hydrogen.




FIG. 6. The ratio $\sigma_T/\epsilon_\gamma^2$ as a function of A for $h=0.32$ GeV.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
76 Wa 3		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABY	140-250	C	250	MAG-D		90
G,PI-	ABY	140-250	C	250	MAG-D		90




Fig. 2. The energy spectra of photoproduced π^- from C, Al, Ca, and Cu at 90° in the laboratory system by 250-MeV bremsstrahlung. The data of Ca are normalized to $0.26 \mu\text{b sterad}^{-1} \text{MeV}^{-1} Q^{-1}$ for π^- at 35 MeV. The solid curves are the calculated spectra of π^- by a theoretical model.




Fig. 3. The π^-/π^+ ratio as a function of the kinetic energy of pions produced from C, Al, Ca, and Cu by 250-MeV bremsstrahlung. The solid curve in (a) is the calculated energy spectrum of π^-/π^+ ratio including the Coulomb potential for C. The dashed curve is the ratio calculated neglecting the Coulomb potential.




Fig. 4. The A -dependence of the $(\pi^+\pi^-)$ cross sections at the pion kinetic energies of ~ 20 MeV, ~ 35 MeV, ~ 50 MeV and ~ 70 MeV. The solid curves show the relative A -dependence obtained from the theoretical calculation. The dashed lines show $A^{2/3}$ dependence only for guiding eyes.

METHOD					REF. NO.		
					77 Ad 1		hmg
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P, G	LFT	16	D	0	SCD-D		45
		(16.11)		(.163)			

γ -ray decays of the 12.7 and 16.1 MeV states of ^{12}C are investigated in a coincidence study of the $^{10}\text{B}(^3\text{He}, p\gamma)$ reaction. We obtain $\Gamma_{\gamma_0}/\Gamma = (1.93 \pm 0.12)\%$, $\Gamma_{\gamma_1}/\Gamma_{\gamma_0} = (15.0 \pm 1.8)\%$ and $\Gamma_a/\Gamma = (97.8 \pm 0.1)\%$ for $^{12}\text{C}(12.7)$, and $\Gamma_{\gamma_1}/\Gamma = (2.42 \pm 0.29) \times 10^{-3}$ for $^{12}\text{C}(16.1)$. Relative γ -ray branching ratios of $^{12}\text{C}(16.1)$ were measured using the $E_p = 163$ keV $^{11}\text{B}(p, \gamma)$ resonance. We find $\Gamma_{\gamma_0}/\Gamma_{\gamma_1} = (4.6 \pm 0.7)\%$, $\Gamma_{\gamma}(16.1 \rightarrow 9.6)/\Gamma_{\gamma_1} = (2.4 \pm 0.4)\%$, and $\Gamma_{\gamma}(16.1 \rightarrow 12.7)/\Gamma_{\gamma_1} = (1.46 \pm 0.25)\%$. This information, together with existing data on M1 transitions and single nucleon transfer reactions, is used to determine the isospin mixing between the 12.7 and 15.1 MeV levels of ^{12}C . A charge dependent matrix element of 110 ± 30 keV is deduced.

STATE AT 16.11 MEV

TABLE I. Decay properties of the 12.7 and 16.1 MeV states of ^{12}C .

$^{12}\text{C}(12.7)$	$^{12}\text{C}(16.1)$
$\Gamma_{\gamma_0}/\Gamma = (1.93 \pm 0.12) \times 10^{-2}^a$	$\Gamma = 6.7 \pm 0.5$ keV ^a
$\Gamma_{\gamma_1}/\Gamma_{\gamma_0} = 0.150 \pm 0.018^a$	$\Gamma_{\gamma_1}/\Gamma = (2.42 \pm 0.29) \times 10^{-3}^a$
$\Gamma_a = (17.7 \pm 2.8)$ eV ^b	$\frac{\Gamma_{\gamma}(16.1 \rightarrow 0.0)}{\Gamma_{\gamma}(16.1 \rightarrow 4.4)} = (4.6 \pm 0.7)\%^a$
$\Gamma_{\gamma_0} = 0.35 \pm 0.05$ eV ^c = 0.008 W.u.	$\frac{\Gamma_{\gamma}(16.1 \rightarrow 9.6)}{\Gamma_{\gamma}(16.1 \rightarrow 4.4)} = (2.4 \pm 0.4)\%^a$
$\Gamma_{\gamma_1} = 0.053 \pm 0.010$ eV ^d = 0.005 W.u.	$\frac{\Gamma_{\gamma}(16.1 \rightarrow 12.7)}{\Gamma_{\gamma}(16.1 \rightarrow 4.4)} = (1.46 \pm 0.25)\%^a$
	$\Gamma_{\gamma}(16.1 \rightarrow 0.0) = 0.75 \pm 0.16$ eV = 0.53 W.u.
	$\Gamma_{\gamma}(16.1 \rightarrow 4.4) = (16.2 \pm 2.3)$ eV = 0.49 W.u.
	$\Gamma_{\gamma}(16.1 \rightarrow 9.6) = (0.39 \pm 0.09)$ eV = 4.0×10^{-2} W.u.
	$\Gamma_{\gamma}(16.1 \rightarrow 12.7) = (0.24 \pm 0.05)$ eV = 0.29 W.u.
	$\Gamma_p = (27.2 \pm 4.2)$ eV ^e

^aThis work.

^bAssuming $\Gamma_a + \Gamma_{\gamma_0} + \Gamma_{\gamma_1} = \Gamma$.

^cSee Ref. 6.

^dCombining Ref. 6 and present work.

^eSee Ref. 21.

^fSee Refs. 21, 22 and present work.

TABLE II. Comparison of A=12 electromagnetic observables with theory.

Transition	Expt. Γ_{γ} (eV)	Theory A ^a Γ_{γ} (eV)	Theory B ^a Γ_{γ} (eV)
$^{12}\text{B}(0.95 \rightarrow 0.00)$	$(2.19 \pm 0.24) \times 10^{-1}^b$	1.71 ± 10^{-1}	2.32 ± 10^{-1}
$^{12}\text{C}(16.11 \rightarrow 0.00)$	0.75 ± 0.16^c	0.607	0.607
$^{12}\text{C}(16.11 \rightarrow 4.44)$	16.2 ± 2.3^c	11.3	11.8
$^{12}\text{C}(16.11 \rightarrow 12.71)$	0.24 ± 0.05^c	0.215	0.215
$^{12}\text{C}(15.1 \rightarrow 0.0)$	37.0 ± 1.1^d	30.8	30.3
$^{12}\text{C}(15.1 \rightarrow 4.4)$	$0.92 \pm 0.36^{d,e}$	1.32	1.01
$^{12}\text{C}(15.1 \rightarrow 12.7)$	$0.56 \pm 0.16^{d,e}$	0.47	0.48
$^{12}\text{C}(12.71 \rightarrow 0.0)$	0.35 ± 0.05^f	0.11	0.11
$^{12}\text{C}(12.71 \rightarrow 4.4)$	$0.053 \pm 0.010^{f,g}$	0.015	0.015
State	Expt. ^h μ (μ_N)	Theory A ^a μ (μ_N)	Theory B ^a μ (μ_N)
$^{12}\text{B}(0.0)$	$+1.003 \pm 0.001$	-0.762	-0.948
$^{12}\text{N}(0.0)$	$+0.4571 \pm 0.0005$	-0.511	-0.413

(over)

- ⁶F.E. Cecil et al, Phys. Rev. C9, 798 (1974).
¹⁴D.E. Alburger et al., Phys. Rev. C5, 384 (1972).
²¹R.E. Segel et al., Phys. Rev. 124, 814 (1961).
²²T. Huus et al., Phys. Rev. 91, 599 (1953).
²⁷S. Cohen et al., Nucl. Phys. 73, 1 (1965);
ibid. A101, 1 (1967); D. Kurath, Phys. Rev.
C7, 1390 (1973); private communication
³¹B.T. Chertock et al., Phys. Rev. C3, 23 (1973).

REF. K.V. Alanakyan, M. Dzh. Amaryan, R.A. Demirchyan, K.Sh. Egiyan, M.S. Ogandzhanyan, & Yu.G. Sharabyan
 Yad. Fiz. 25, 545 (March 1977)
 Sov. J. Nucl. Phys. 25, 292 (March 1977)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	77 A1 9
	hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	90-999	C	2 *5 (4-5)	TEL-D	---	DST

COMMENTS: $f \sim \exp(-Bp^2)$
 $B = E$
 $p^4 (d^2\sigma/d\Omega dpQ)$

*E, GEV, 999=4.5 GEV

The A^n -dependence and momentum spectra of photoprotons in the nuclei ^{12}C , ^{27}Al , ^{63}Cu , ^{118}Sn , and ^{208}Pb have been studied experimentally for maximum bremsstrahlung energies of 2.0, 3.0, and 4.5 GeV. The A -dependence shows that the proton photoproduction mechanism for $E_\gamma > 400$ MeV is identical for the entire kinetic-energy region 65-280 MeV and the angle region 45° - 150° for the secondary protons studied. The dependence of the exponent n on the transverse momentum p_t is in good agreement with the same dependence for protons produced in nuclei by primary protons. In the momentum spectra of the invariant cross section $f = (E/p^2)(d^2\sigma/d\Omega dpQ) \sim \exp(-Bp^2)$ it is observed that the parameter B does not depend on the incident-photon energy and on the target nucleus, but depends on the proton-detection angle.




FIG. 1. Differential cross section for proton photoproduction as a function of atomic number A of the nucleus at $E_0 = 2$ GeV. The lines a correspond to $\beta_p = 60^\circ$, b to 90° ; and c to 150° . Points: \square — $E_p = 64$, Δ —80, \circ —101, \blacksquare —137, \blacktriangle —209, and \bullet —280 MeV.




FIG. 3. The same as Fig. 2. Experimental points: \blacktriangle — $E_0 = 0.13$, \circ —0.25, \triangle —0.4, \square —1.2, \blacksquare —2.0, \times —3.0, and \bullet —4.5 GeV.




FIG. 2. Dependence of the exponent n in the A^n dependence of the cross section for the reaction $\gamma A \rightarrow p A'$ as a function of proton transverse momentum: a— $E_0 = 2.0$ GeV, b— $E_0 = 3.0$ GeV, c— $E_0 = 4.5$ GeV. The points for a and b: Δ — $\beta_p = 60^\circ$, \circ — 90° , \square — 150° ; for c: Δ — $\beta_p = 46^\circ$, \circ — 86° , \square — 136° . The curves show the dependence of n on p_t for the reaction $A(p, p')A'$ taken from Ref. 9.




FIG. 4. Invariant cross section f as a function of p^2 . a—for $E_0 = 2.0$ GeV, b—for $E_0 = 3.0$ GeV. Experimental points: \bullet , \blacktriangle , \blacksquare —for $\beta_p = 60^\circ$ for the respective nuclei ^{12}C , ^{63}Cu , and ^{208}Pb ; \circ , Δ , \square —the same for $\beta_p = 90^\circ$; \times , \cdot , \diamond —the same for $\beta_p = 150^\circ$.

TABLE I. Differential cross section $d^2\sigma/d\Omega dTQ$ of the reaction $\gamma A \rightarrow pA'$ in $\mu\text{b}/\text{MeV}\cdot\text{sr}$.

A	E_γ , GeV	θ_p , deg	E_p , MeV						
			54	80	101	137	209	279	
^{12}C	2	80	3.720±0.056	2.630±0.052	1.907±0.057	1.425±0.038	0.725±0.220	0.429±0.016	
		90	2.274±0.045	1.587±0.047	1.130±0.039	0.783±0.022	0.256±0.008	0.117±0.007	
		150	1.152±0.032	0.690±0.014	0.505±0.028	0.218±0.007	0.071±0.005	0.021±0.002	
	3	80	4.240±0.100	3.424±0.063	1.897±0.043	1.829±0.046	0.653±0.024	0.462±0.019	
		90	2.440±0.056	2.031±0.040	1.145±0.029	0.807±0.028	0.243±0.009	0.088±0.005	
		150	1.360±0.042	0.877±0.029	0.438±0.016	0.300±0.010	0.057±0.003	-	
	2	60	3.460±0.127	6.014±0.120	4.083±0.109	3.233±0.097	1.513±0.046	-	
		90	5.920±0.107	3.750±0.088	2.502±0.084	1.718±0.052	0.603±0.018	-	
		150	3.127±0.078	1.797±0.035	1.189±0.060	0.844±0.019	0.164±0.011	-	
	^{27}Al	3	60	9.960±0.239	7.492±0.131	4.160±0.092	3.527±0.100	1.568±0.058	0.925±0.037
			90	6.090±0.130	4.843±0.107	2.688±0.078	1.995±0.065	0.596±0.021	0.239±0.013
			150	3.750±0.103	2.943±0.081	1.234±0.042	0.717±0.025	0.136±0.006	-
4.5		46	-	9.510±0.250	-	-	-	1.320±0.170	
		86	-	6.200±0.085	-	-	-	0.248±0.020	
		138	-	3.380±0.050	-	-	-	-	
^{63}Cu		2	60	23.500±0.379	15.170±0.299	10.931±0.269	8.163±0.240	3.939±0.110	2.115±0.064
			90	16.721±0.258	9.757±0.231	6.856±0.082	4.411±0.134	1.424±0.042	0.743±0.037
			150	10.592±0.212	5.217±0.103	3.662±0.165	1.697±0.050	0.342±0.021	0.115±0.011
	3	60	26.180±0.790	20.580±0.340	10.270±0.191	8.594±0.248	3.680±0.140	1.661±0.064	
		90	17.800±0.320	13.601±0.260	7.516±0.110	5.245±0.172	1.403±0.048	0.676±0.032	
		150	11.640±0.271	7.834±0.205	3.388±0.107	2.237±0.075	0.366±0.017	0.097±0.008	
	4.5	46	-	27.000±0.750	-	-	-	3.550±0.180	
		86	-	17.401±0.250	-	-	-	0.735±0.060	
		138	-	9.750±0.150	-	-	-	-	
^{119}Sn	2	60	45.601±0.538	30.050±0.593	19.970±0.587	13.102±0.380	7.137±0.210	-	
		90	32.550±0.553	18.890±0.466	13.840±0.428	8.287±0.320	2.588±0.078	-	
		150	19.571±0.391	10.289±0.203	6.548±0.321	3.022±0.090	0.585±0.041	-	

A	E_γ , GeV	θ_p , deg	E_p , MeV						
			54	80	101	137	209	279	
^{119}Sn	3	80	55.070±1.270	29.920±0.680	17.800±0.430	16.550±0.490	7.028±0.036	5.873±0.140	
		90	36.800±0.720	26.250±0.550	14.370±0.440	9.544±0.328	2.684±0.099	1.197±0.055	
		150	22.500±0.560	14.590±0.350	8.251±0.210	4.103±0.150	0.664±0.033	-	
	4.5	46	-	53.900±1.400	-	-	-	5.640±0.320	
		86	-	32.200±0.51	-	-	-	1.420±0.114	
		138	-	19.250±0.290	-	-	-	0.229±0.038	
	^{208}Pb	2	60	90.000±1.280	56.850±1.120	35.200±1.030	23.930±0.720	13.140±0.400	7.745±0.310
			90	60.990±0.970	34.080±0.800	23.690±0.720	14.220±0.400	4.522±0.135	2.453±0.120
			150	36.690±0.730	19.560±0.370	10.638±0.520	3.724±0.168	1.112±0.077	0.531±0.035
3		60	100.740±1.130	76.050±1.200	23.000±0.670	28.000±0.820	12.810±0.450	7.092±0.250	
		90	71.350±1.170	48.320±0.900	24.700±0.650	16.420±0.520	4.590±0.170	2.244±0.120	
		150	42.090±0.970	27.240±0.660	12.130±0.42	7.200±0.150	1.220±0.034	0.530±0.039	
4.5		46	-	45.000±0.350	-	-	-	11.600±0.560	
		86	-	28.780±0.320	-	-	-	4.050±0.214	
		138	-	23.600±0.430	-	-	-	0.465±0.084	

A	E_γ , GeV	θ_p , deg	80	119	166	231	281
^{12}C	4.5	46	5.210±0.280	3.670±0.086	2.520±0.064	1.190±0.046	0.755±0.042
		86	2.440±0.080	1.350±0.052	0.845±0.037	0.363±0.019	0.105±0.007
		138	1.320±0.029	0.427±0.029	0.196±0.015	0.045±0.005	0.018±0.002

TABLE II. Values of the parameter B in $(\text{GeV}/c)^{-2}$ in the relation $E_p/p_p^2 (d^2\sigma/d\Omega_p dT_p Q) = \sim \exp(-Bp^2)$.

Target	$E_\gamma = 2.0 \text{ GeV}$			3.0 GeV			4.5 GeV		
	$\theta_p = 46^\circ$	86°	138°	46°	86°	138°	46°	86°	138°
^{12}C	4.874±0.512	7.276±0.482	9.461±0.303	6.288±0.805	3.623±0.437	10.373±0.977	8.347±0.173	3.066±0.49	11.262±0.481
^{63}Cu	5.340±0.827	7.337±0.827	10.472±0.609	6.972±0.939	3.559±0.622	11.807±0.944	-	-	-
^{208}Pb	5.204±0.753	7.605±0.721	10.088±0.36	6.870±1.514	8.858±0.733	10.983±1.158	-	-	-




FIG. 6. Dependence of the parameter B from the relation $f \sim \exp(-Bp^2)$ on the atomic number of the target nucleus. The solid points refer to $E_\gamma = 2.0 \text{ GeV}$, and the hollow points to $E_\gamma = 3.0 \text{ GeV}$; the points \bullet and \circ are for $\theta_p = 80^\circ$, \blacktriangle and \triangle are for 90° , and \blacksquare and \square are for 150° .




FIG. 5. The same as Fig. 4 but for $E_\gamma = 4.5 \text{ GeV}$, in ^{12}C . Experimental points: \bullet — $\theta_p = 46^\circ$, \blacktriangle — 86° , \blacksquare — 138° .

REF.

K. Baba, I. Endo, H. Fujii, M. Fujisaki, S. Kodata, A. Murakami,
Y. Murata, S. Noguchi and Y. Sumi
J. Phys. Soc. Japan 42, 1049 (1977)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
77 Ba 8	egf	

REMARKS: Photon Differences

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	ABX	300-900	C	300-900	MAG-D		44

It is found that, even at a considerably large lab angle 44.2° , momentum spectra of pions in the reaction $^{12}\text{C}(\gamma, \pi^-)$ show a clear singly-peaked structure throughout a photon energy range between 300 and 850 MeV. The peak momentum is always about 40 MeV/c less than that determined by free nucleon kinematics. The width increases almost linearly with photon energy. Differential cross section does not exhibit a marked variation with energy. These data are quite well explained by an impulse-approximation calculation.




Fig. 1. Typical examples of pion momentum spectrum measured in the present experiment. Smooth curves are the ones calculated by an impulse-approximation method. The arrow means the value of pion momentum p_{free} corresponding to the elementary processes $\gamma-n \rightarrow \pi^- + p$.




Fig. 2(a). Values of the width 2Γ fitted by using the Gaussian form (1) and (b) those of the momentum shift $p_{free} - p_0$, as a function of $\langle k \rangle$. Smooth and dotted curves are the results from the impulse approximation calculation at $\theta = 28.4^\circ$ and 44.2° , respectively.




Fig. 3. Values of the fitted cross section $d\sigma/d\Omega$ as a function of $\langle k \rangle$. The symbols of data are the same as in Fig. 2. The curves shown for comparison are: Six times the elementary cross section $d\sigma/d\Omega$ ($\gamma n \rightarrow \pi^- p$) at 44.2° (solid line) and at 28.4° (broken line) taken from reference 3).

³T. Fujii, T. Kondo, F. Takasaki, S. Yamada, S. Homma, K. Huke, S. Kato, H. Okuno, I. Endo and H. Fujii: Phys. Rev. Lett. 28, (1972) 1672 and 29, (1972) 244(E), see also INS-Report-256 (1976)

REF. Z. Berant, R. Moreh and S. Kahane
Phys. Lett. 69B, 281 (1977)

ELEM. SYM	A	Z
C	12	6
REF. NO.		egf
77 Be 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	5 - 8	D	5-8	SCD-D		140

5.5-7.2 MeV

Table I
Differential cross sections ($\mu\text{b}/\text{sr}$) for elastic scattering from C and Mg targets at an angle of 140° . The theoretical values include the combined effect of NT, NR and D scattering; The R contribution was ignored. The values for pure NT scattering are also given.

E (keV)	C(Z = 6)			Mg(Z = 12)		
	Experiment	Theory	Thomson	Experiment	Theory	Thomson
5516	0.17 ± 0.04	0.16	0.17	0.64 ± 0.15	0.61	0.67
5752	0.20 ± 0.06	0.15	0.17	0.66 ± 0.15	0.61	0.67
6465	0.17 ± 0.04	0.15	0.17	0.64 ± 0.15	0.59	0.67
6517	0.16 ± 0.04	0.15	0.17	0.63 ± 0.15	0.59	0.67
6374	0.18 ± 0.04	0.15	0.17	0.59 ± 0.15	0.58	0.67
7163	0.16 ± 0.04	0.14	0.17	0.50 ± 0.15	0.57	0.67




Fig. 1. High-energy part of the elastic scattering spectrum: from a Mg target at 140° as measured by a 40 cm^3 Ge(Li) detector, and of the incident $V(n, \gamma)$ spectrum. Lines indicated by P and F denote photopeaks and first-escape peaks. Other lines denote double-escape peaks.

REF. F. Borowski, Ch. Schmitt, G. G. Simon, V. Walther, D. Drechsel,
W. Haxton, R. Rosenfelder
Phys. Rev. Lett. 38, 742 (1977)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
77 Bo 12		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,P+	RLX	150-280	D	280	MAG-D	-	DST
E,P-	ABX	150-280	D	280	MAG-D	-	DST

Double-differential cross sections for the electroproduction of pions of both charges have been measured. We compare the data obtained for production near threshold from ^{12}C and ^{16}O with theoretical treatments employing both shell-model and sum-rule nuclear descriptions with full inclusion of the final-state interaction.

SIG PI-/SIG PI+




FIG. 1. Double-differential cross sections for production of negative pions by 280-MeV electrons from (a) ^{16}O and (b) ^{12}C as a function of the pion angle. The shell-model results (solid line) are compared with the predictions of the sum-rule results when evaluated with (dashed line) and without (dotted line) the effective two-body Skyrme II (Ref. 14) interaction.




FIG. 2. The ratio of the double-differential cross sections for production of negative and positive 21.6-MeV pions from ^{12}C as a function of the pion angle. The dashed and solid curves are the shell-model results that follow on using a pion optical potential parametrized (Ref. 2) with and without the Lorentz-Lorenz effect, respectively. The dotted line is the Skyrme II sum-rule result with no Lorentz-Lorenz effect.

²J. Hüfner, Phys. Rep. 21C, 1 (1975)

¹⁴M. Beiner, H. Flocard, N. van Giai, and P. Quentin, Nucl. Phys. A238, 29 (1975)

METHOD				REF. NO.		hmg	
				77 Hi 5			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,n	ABI	18-30 (18.7)	C	18-30	ACT-I		49I

THICK BREMS TARGET

The integral experiment for a sensitivity check of photonuclear cross section data of C, Mn, Fe, In and Au was performed by using the bremsstrahlung produced in a thick iron target by 18, 22, 26 and 30 MeV electrons from a linear accelerator. The cross section data measured by the activation method showed better results for all incident electron energies than those by the photoneutron method, because the latter include the competing (γ, np) reaction above its threshold energy. It is necessary to obtain the cross section data of (γ, n), (γ, np), ($\gamma, 2n$), ($\gamma, p2n$) reactions etc., separately by the activation method.

The effective energy range and effective cross section in the giant resonance region were determined for C, Mn, Fe and Au. By using these quantities, the gross structure of the bremsstrahlung spectrum was obtained in good agreement with the theoretical calculation.

TABLE 3
Ratio of measured and calculated saturated activities at 90°, $R_s = A_s^{exp}/A_s^{cal}$

Reaction	Reference	Detector	Electron energy				Threshold energy of competing (γ, np) reaction (MeV)
			18 MeV	22 MeV	26 MeV	30 MeV	
$^{12}\text{C}(\gamma, n)^{11}\text{C}$	6	ACT ^a	-	1.205	1.09	0.824	27.4
	7	BF3 ^b	-	0.757	0.931	-	
	8	BF3	-	1.492	1.189	0.914	
	9	BF3	-	1.418	1.159	-	
$^{55}\text{Mn}(\gamma, n)^{54}\text{Mn}$	10	BF3	0.575	0.810	0.598	-	17.8
	11		0.595	0.875	0.654	0.609	
$^{54}\text{Fe}(\gamma, n)^{53}\text{Fe}$	12	ACT	0.542	0.783	0.722	1.12	20.9
$^{115}\text{In}(\gamma, n)^{114\text{m}}\text{In}$	13	BF3	0.448	0.511	-	-	15.9
	14	BF3	0.603	0.672	0.586	1.00	
$^{115}\text{In}(\gamma, \gamma)^{115\text{m}}\text{In}$	15	ACT	0.829	1.00	0.852	0.995	
$^{197}\text{Au}(\gamma, n)^{196}\text{Au}$	16	BF3	0.743	0.553	0.529	-	13.7
	17	BF3	0.842	0.628	0.608	0.476	

^a ACT: measurement of radioactivity of the target.
^b BF3: BF₃ neutron counter with moderator.

(OVER)

TABLE 4
Effective cross section and effective energy range.

Reaction	Effective energy range (MeV)	Effective cross section (mb)			Average
		22 MeV	26 MeV	30 MeV	
$^{12}\text{C}(\gamma, n)^{11}\text{C}$	20.5-24.5	-	5.98	5.28	5.63 ± 0.35
$^{55}\text{Mn}(\gamma, n)^{54}\text{Mn}$	15.0-20.5	-	56.2	52.7	54.5 ± 1.7
$^{54}\text{Fe}(\gamma, n)^{53}\text{Fe}$	17.0-24.0	-	33.6	47.9	40.8 ± 7.2
$^{197}\text{Au}(\gamma, n)^{196}\text{Au}$	12.5-15.0	553.5	532.5	415.0	500 ± 85




Fig. 3. Comparison of experimental and calculated bremsstrahlung spectra emitted from an iron target in the 9° direction by (a) 18 MeV electrons, (b) 22 MeV electrons, (c) 26 MeV electrons, (d) 30 MeV electrons.

METHOD

REF. NO.

77 Jo 3

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,2N	ABX	100 - 600	C	100-800	ACT-I		4PI




Fig. 1. The measured yield of $^{12}\text{C}(\gamma, 2n)^{10}\text{C}$ as a function of the maximum bremsstrahlung energy. The meaning of the solid and dashed curves is described in the text.




Fig. 3. The cross section for the reaction $^{12}\text{C}(\gamma, 2n)^{10}\text{C}$ as a function of photon energy. The present experimental result is given by the solid curve and the dashed curve shows the result from ref. 7. The hatched areas indicate the error. Open circles are calculated values.

7 H. Breuer and W. Pohlitz, Nucl. Phys. 30, 417 (1962)

REF. V.V. Kirichenko, A.F. Khodjachikh, Ju.M. Arkatov, P.I. Vatset,
I.V. Dogjust
Ukr. Fiz. Zh. 22, 959 (1977)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
77 Ki 11		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	40-120	C	150	CCH-D		DST

The reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$ is studied in the range of energy exceeding that corresponding to the giant resonance. The analysis of the obtained data shows that the electric dipole photon absorption is the main reaction mechanism as in the reactions of two particles photodisintegration of the lightest ^2He , ^3He and ^2H nuclei with the proton yield. The obtained values for the asymmetry coefficient is found to be in agreement within the experimental accuracy with the similar coefficient in the reactions of two particles photodisintegration of ^2He , ^3He but they differ from the asymmetry coefficient values for the case of ^2H photodisintegration.

Кириченко В.В., Ходячих А.Ф., Аркатов Ю.М., Ватсет П.И., Догжуст И.В. РЕАКЦИЯ $^{12}\text{C}(\gamma, p)^{11}\text{B}$ В ЭНЕРГЕТИЧЕСКОМ ИНТЕРВАЛЕ ОТ 40 ДО 120 МЭВ. Укр. физ. журнал, 22, 959-964.

С помощью диффузионной камеры, установленной на пути пучка тормозных фотонов от линейного ускорителя на 360 МэВ, измерены энергетическая зависимость полного сечения и угловые распределения протонов в интервале энергий от 40 до 120 МэВ. Из анализа данных получены полное сечение асимметричного дипольного поглощения и асимметрия угловых распределений. Результаты сравниваются с данными по исследованию реакций двухчастичного фоторасщепления легких ядер ^2He , ^3He , ^2H с выходом протонов.

Using a diffusion chamber in the way of a beam of bremsstrahlung photons from a 360-MeV linear accelerator, the energy dependence of the total cross section and the angular distributions of protons have been measured in the energy region from 40 to 120 MeV. From an analysis of the data, the total cross section for electric dipole absorption and the asymmetry of the angular distributions have been obtained. The results are compared with the data on the reactions of two-particle photodisintegration of the lightest nuclei ^2He , ^3He and ^2H with proton yield.




Рис. 1. Зависимость полного сечения реакции $^{12}\text{C}(\gamma, p)^{11}\text{B}$ от E_γ (МэВ): гистограмма — настоящий эксперимент, O — данные работы [1].




Рис. 2. Угловые распределения протонов реакции $^{12}\text{C}(\gamma, p)^{11}\text{B}$ в области энергий 40—50 МэВ (I), 50—60 (II), 60—80 (III), 80—120 (IV).

$$\frac{d\sigma}{d\Omega} = a \sin^2 \theta + b \sin^2 \theta \cos \theta + c \sin^2 \theta \cos^2 \theta + d, \quad (1)$$

E_γ , МэВ	a , $\frac{\text{мб}}{\text{ср}}$	b , $\frac{\text{мб}}{\text{ср}}$	c , $\frac{\text{мб}}{\text{ср}}$
40—50	$0,169 \pm 0,023$ $0,187 \pm 0,017$	$0,138 \pm 0,043$ $0,139 \pm 0,043$	$0,098 \pm 0,092$
50—60	$0,082 \pm 0,012$ $0,099 \pm 0,011$	$0,112 \pm 0,021$ $0,112 \pm 0,027$	$0,102 \pm 0,045$
60—80	$0,027 \pm 0,008$ $0,041 \pm 0,008$	$0,068 \pm 0,016$ $0,069 \pm 0,021$	$0,082 \pm 0,033$
80—120	$0,010 \pm 0,002$ $0,011 \pm 0,003$	$0,023 \pm 0,004$ $0,023 \pm 0,007$	$0,025 \pm 0,010$

ELEM. SYM.	A	Z
C	12	6

METHOD			REF. NO.				
			77 Kn 2		egf		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E-,N	ABX	19-32	D	26-32	ACT-I		4PI
E+,N	ABX	19-32	D	26-32	ACT-I		4PI

See figure on other side

(OVER)






Fig. 1. a Experimental absolute $(e, e'n)$ cross section for ^{12}C together with the results of DWBA and PW calculations. b Experimental absolute $(e, e'n)$ cross section for ^{63}Cu together with the results of DWBA and PW calculations. c Experimental absolute $(e, e'n)$ cross section for ^{107}Ag together with the results of DWBA and PW calculations

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,PI+	ABX	0-1 (0-.95)	D	195	MAG-D		DST

Three resonances in $^{12}\text{C} - ^{12}\text{C}$ with $E_{c.m.}(J^\pi)$ of 3.35(6⁺), 11.2(8⁺), and 13.75(10⁻) are reported. Together with earlier data, these resonances fit in a rotational band in ^{24}Mg . A model based on the rotation-vibration coupling is proposed to account for these data.




FIG. 2. Pion energy distribution in $^{12}\text{C}(e, e' \pi^+)^{12}\text{B}$. Solid curves are the theoretical estimates derived from Eqs. (3) and (4) and the Helm model using the electron scattering data (Refs. 2 and 3). Dashed curves are the theoretical result with a nuclear shell-model and pion wave with Coulomb correction (Ref. 5). Arrows indicate the maximum energy of pion leading to the ground- and first excited residual states.




FIG. 3. Pion angular distributions in $^{12}\text{C}(\gamma, \pi^+)^{12}\text{B}$ at $E_\gamma \approx 194$ MeV leaving the ground state (open circles) and the first excited state (closed circles) in ^{12}B . Solid curves and dashed curves are the theoretical estimates by the Helm model using electron scattering data (Refs. 2 and 3) and by the shell model (Ref. 4), respectively.

TABLE I. Comparison of cross sections of the $^{12}\text{C}(\gamma, \pi^+)^{12}\text{B}$ reaction relating to the ground and first excited residual states. R is the most probable value of the ratio $(d\sigma/d\Omega)_{\text{expt}} / (d\sigma/d\Omega)_{\text{theor}}$. σ_{expt} is calculated from σ_{theor} multiplied by R .

Residual state	Ground state ($J^\pi = 1^+$)	First excited state ($E_R = 0.95$ MeV, $J^\pi = 2^+$)
	Helm model theory ^a	
R	0.56 ± 0.03	1.42 ± 0.13
$\sigma_{\text{theor}} (\mu\text{b})$	27.04	16.55
$\sigma_{\text{expt}} (\mu\text{b})$	15.1 ± 0.3	23.5 ± 2.3
	Shell-model theory ^b	
R	0.14 ± 0.01	0.43 ± 0.04
$\sigma_{\text{theor}} (\mu\text{b})$	99.9	59.5
$\sigma_{\text{expt}} (\mu\text{b})$	14.0 ± 1.0	25.6 ± 2.4

^aRefs. 2 and 3.

^bRef. 4.

¹H. Überall, B. A. Lamers, C. W. Lucas, and A. Nagl, Phys. Lett. 44B, 334 (1972).

²F. Cannata, B. A. Lamers, C. W. Lucas, A. Nagl, H. Überall, C. Warnatz, and F. J. Kelly, Can. J. Phys. 52, 1405 (1974).

³J. B. Seaborn, V. Devanathan, and H. Überall, Nucl. Phys. A219, 461 (1974).

⁴S. Furui, private communication, and in Proceedings of the International Conference on High-Energy Physics and Nuclear Structure, Zürich, Switzerland, 29 August-2 September 1977 (to be published).

REF. K.A. Snover, P. Paul, and H.M. Kuan
Nucl. Phys. A285, 189 (1977)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
77 Sn 3		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	21-37	D	6- 23	NAI-D		DST

TABLE 1
Angular distribution coefficients for $^{11}\text{B}(p, \gamma_1)^{12}\text{C}$

E_p (MeV)	Transition	a_1	a_2	a_3	a_4
12.4	γ_0	0.30 ± 0.03	-0.65 ± 0.07	-0.08 ± 0.08	-0.04 ± 0.09
	γ_1	0.24 ± 0.02	-0.23 ± 0.04	-0.08 ± 0.05	-0.01 ± 0.06
	γ_3	0.22 ± 0.04	-0.06 ± 0.08	-0.01 ± 0.10	-0.10 ± 0.10
14.4	γ_0	0.37 ± 0.04	-0.74 ± 0.07	-0.22 ± 0.10	-0.04 ± 0.13
	γ_1	0.21 ± 0.05	-0.27 ± 0.08	-0.16 ± 0.10	-0.08 ± 0.13
	γ_3	0.33 ± 0.03	-0.28 ± 0.05	-0.10 ± 0.07	-0.07 ± 0.08

TABLE 2
Integrated photoproton cross section $\int \sigma(\gamma, p_0) dE_\gamma$

Transition	Integration limits (MeV)		Integrated cross section (MeV · mb)
	E_p	E_γ	
γ_0	1.0-23.0	16.9-37.0	64 ^{b)}
γ_1	1.0-23.0	12.4-32.6	19 ^{a)}
γ_2	6.0-23.0	13.8-29.4	5.5 ^{c)}
γ_3	6.0-23.0	11.8-27.4	5.6 ^{d)}

^{a)} $E_\gamma = \frac{11}{12} E_p + 15.96 - E_1$.
^{b)} $E_p = 1.0-13.7$ MeV from σ_{total} of ref. ³⁾; $E_p = 13.7-23.0$ from $\sigma(90^\circ)$ (fig. 2) assuming $\bar{a}_2 = -0.70$.
^{c)} Same as *b* except $\bar{a}_2 = 0.30$.
^{d)} From $\sigma(90^\circ)$ (fig. 2) assuming $\bar{a}_1 = 0$.

³64A12

(over)




Fig. 2. The 90° cross sections for the $^{11}\text{B}(p,\gamma)^{12}\text{C}$ reaction for transitions to the lowest-four energy levels in ^{12}C as a function of excitation energy (lower scale) and proton energy (upper scale). The error bars shown are statistical; systematic uncertainties due to the efficiency calibration are estimated to be $\pm \leq 20\%$. The γ_2 cross section has an additional uncertainty of $\pm 0.05 \mu\text{b}/\text{sr}$ (see text).

REF. I. Arai, H. Fujii, S. Homma, Y. Hoshi, H. Ikeda, T. Ishii, A. Itano, K. Maruyama, E. Ohshima, H. Okuno, A. Sasaki, & N. Yamashita J. Phys. Soc. of Japan, 45, 1 (July 1978)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.

78 Ar 7

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	SPC	360-600	D	510-750	MAG-D		41

The cross section of the photoproduction of pions from carbon at 41° in the laboratory system and in the incident energy region from 510 MeV to 750 MeV is measured by a magnetic spectrometer for pions and counter hodoscopes for recoil protons in coincidence with pions. A tagged photon beam is used. The results which show the production of pions from the quasi free nucleons inside the nucleus are analyzed in terms of the distorted wave impulse approximation.




Fig. 2. The momentum spectrum of the photoproduced pions from carbon as a function of the difference of momentum between the expected momentum from the kinematics of $\gamma + \pi^- \rightarrow \pi^- + p$ and the observed momentum. Solid line is the best fitted curve of eq. (3) in the text. Dotted curve is the quasi free peak calculated from DWIA.




Fig. 3. The percentage ratio of the number of protons detected by the hodoscope to the number of pions in the spectrometer as a function of the proton angle. The pions in this case are in the quasi free peak region ($p_{\pi} \leq 100$ MeV/c). Solid curve is the angular distribution calculated from DWIA.

Table II. Parameters of fittings.

E_{γ}	515~595 MeV	595~675 MeV	675~755 MeV
S	$51 \pm 5 \mu\text{b/str.}$	$44 \pm 5 \mu\text{b/str.}$	$41 \pm 3 \mu\text{b/str.}$
Γ	$51 \pm 4 \text{ MeV/c}$	$58 \pm 5 \text{ MeV/c}$	$67 \pm 5 \text{ MeV/c}$
p_0	$39 \pm 7 \text{ MeV/c}$	$40 \pm 9 \text{ MeV/c}$	$35 \pm 6 \text{ MeV/c}$
p_1	$74 \pm 24 \text{ MeV/c}$	$79 \pm 23 \text{ MeV/c}$	$99 \pm 13 \text{ MeV/c}$
1/R	295 ± 58	374 ± 58	411 ± 35
χ^2	1.60	1.13	1.19

$$\frac{d^2\sigma}{d\Omega_{\pi} dp_{\pi}} = \frac{S}{\sqrt{2\pi}\Gamma} \exp\left\{-\frac{(p-p_0)^2}{2\Gamma^2}\right\} + R(p-p_1)\theta(p-p_1) \quad (3)$$

$$\theta(x) = 0 \quad \text{for } x < 0$$

$$\theta(x) = 1 \quad \text{for } x \geq 0$$

METHOD

REF. NO.

78 Ar 9

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	THR-30		12*30	NAI-D	---	4PI

The total cross section of hadron photoproduction on C, Cu and Pb nuclei is measured for six energy values in the range 12-30 GeV. The obtained cross-section values for C and Cu nuclei have a weak energy dependence at high energies (above 20 GeV). The cross section for the Pb nucleus is somewhat higher in comparison with that expected, and energy dependence is not observed. The A-dependence of the effective number of hadrons agrees with VDM predictions.

*Energy in GeV




Fig. 3. Energy dependence of A_{eff}/A for C, Cu, Pb nuclei. For comparison the data of DESY and SLAC-UCSB, and also the theoretical curves, corresponding to VDM and to the case when the photon is ~20% of the time in a "pure" state, without shadowing. Δ , DESY; \circ , SLAC-UCSB; \times , Serpukhov; \square , Serpukhov [4]; — VDM; --- 0.8 VDM + 0.2 pointlike interaction; -.- general VDM.

Table 2
Value of A_{eff} for nuclei C, Cu, Pb for different energies of γ quanta. Only statistical errors are given.

γ -quanta energy (GeV)	^{12}C	^{64}Cu	^{207}Pb
12.6-15.0	0.79 ± 0.04	0.77 ± 0.04	0.73 ± 0.05
15.0-17.7	0.81 ± 0.04	0.72 ± 0.05	0.75 ± 0.06
17.7-21.0	0.87 ± 0.04	0.80 ± 0.05	0.85 ± 0.07
21.0-24.6	0.80 ± 0.05	0.74 ± 0.07	0.72 ± 0.08
24.6-27.9	0.79 ± 0.05	0.69 ± 0.05	0.79 ± 0.09
27.9-30.0	0.71 ± 0.05	0.68 ± 0.07	0.75 ± 0.13

$$\frac{A_{eff}}{A} = \frac{\sigma_t(\gamma, A)}{Z\sigma_t(\gamma, p) + (A-Z)\sigma_t(\gamma, n)}$$

where

$$\sigma_t(\gamma, p) = (98.7 \pm 3.6) + (65 \pm 10)E^{-1/2} \mu\text{b},$$

$$\sigma_t(\gamma, n) = \sigma_t(\gamma, p) - (18.3 \pm 6.1)E^{-1/2} \mu\text{b}.$$

Table 1
Hadron photoproduction cross sections (in μb) for C, Cu, Pb nuclei for different energies of γ -quanta. Only statistical errors are given.

γ -quanta energy (GeV)	^{12}C	^{64}Cu	^{207}Pb
12.6-15.0	1084 ± 48	5600 ± 240	17140 ± 1170
15.0-17.7	1100 ± 43	5200 ± 310	17480 ± 1140
17.7-21.0	1175 ± 34	5740 ± 340	19680 ± 1720
21.0-24.6	1058 ± 53	5220 ± 460	16400 ± 1720
24.6-27.9	1047 ± 55	4870 ± 350	17920 ± 1920
27.9-30.0	930 ± 66	4730 ± 510	16840 ± 2810

REF. K. Baba, I. Endo, M. Fujisaki, S. Kadota, Y. Sumi, H. Fujii
 Y. Murata, S. Noguchi and A. Murakami
 Nucl. Phys. A306, 292 (1978)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		78 Ba 4
		rs

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABX	300-850	D	300-850	MAG-D		DST
G,PI-	ABX	300-850	D	300-850	MAG-D		DST

Abstract: The momentum spectra of charged pions produced in the reaction $^{12}\text{C}(\gamma, \pi^\pm)$ have been measured for incident photon energies in the interval between 300 and 850 MeV in steps of 50 MeV. Pions with relatively high momenta have been detected by a magnetic spectrometer set at the lab angles 28.4° and 44.2° . All these spectra exhibit a clear singly peaked structure. Detailed features of the structure are quantitatively investigated and compared with a plane-wave impulse-approximation calculation. Calculated results are found to reproduce the spectral shape very well and the absolute magnitude is also rather well fitted by introducing the pion absorption effect properly.




Fig. 5. The fitted values of parameters in the Gaussian distribution for pion momentum spectra: (a) The momentum shift $p_{r,rel} - p_0$, (b) the width 2Γ , and (c) the cross section $d\sigma/d\Omega$ as a function of $\langle k \rangle$. Curves in (a) and (b) are the result of the PWIA calculation for $\theta_L = 28.4^\circ$ (solid and dotted curves for π^- and π^+ , respectively) and 44.2° (dashed one for π^-). Curves in (c) show the differential cross sections (multiplied by a factor of 6) for elementary processes (solid and dotted ones for π^- and π^+ at 28.4° , respectively, while dashed one is for π^- at 44.2°). Symbols in (b) and (c) are the same as in (a).




Fig. 7 Effects of scattered-in pions and double pion photoproduction. Dotted curves represent the contribution of scattered-in pions, which takes into account also the charge-exchange scattering of photoproduced π^0 . The dot-dash curve shows the contribution from double pion production, which is appreciable only above $\langle k \rangle = 650$ MeV. Dashed curves are calculated $1/4$ times PWIA results. Solid curves are the sums of these contributions.

REF. K. Baba, I. Endo, M. Fujisaki, S. Kadota, Y. Sumi, H. Fujii,
Y. Murata, S. Noguchi & A. Murakami
Nucl. Phys. A306, 292 (1978)

ELEM. SYM.	A	Z
C	12	6

METHOD		REF. NO.		hmg			
		78 Ba 12					
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABX	150-700	D	300-850	MAG-D		DST
G,PI-	ABX	150-700	D	300-850	MAG-D		DST

COMMENT: Subtracted bremsstrahlung spectra used

Abstract: The momentum spectra of charged pions produced in the reaction $^{12}\text{C}(\gamma, \pi^\pm)$ have been measured for incident photon energies in the interval between 300 and 850 MeV in steps of 50 MeV. Pions with relatively high momenta have been detected by a magnetic spectrometer set at the lab angles 28.4° and 44.2° . All these spectra exhibit a clear singly peaked structure. Detailed features of the structure are quantitatively investigated and compared with a plane-wave impulse-approximation calculation. Calculated results are found to reproduce the spectral shape very well and the absolute magnitude is also rather well fitted by introducing the pion absorption effect properly.

E NUCLEAR REACTIONS $^{12}\text{C}(\gamma, \pi^+), (\gamma, \pi^-)$, $E = 300\text{--}850$ MeV; measured $\sigma(E, p_\pi, \theta)$. PWIA calculations.




Fig. 4. Double differential cross sections for $^{12}\text{C}(\gamma, \pi^\pm)$ at $\theta_\pi = 28.4^\circ$ and 44.2° as a function of $p_\pi - p_{free}$, where p_{free} denotes the pion momentum satisfying free nucleon kinematics. Solid (28.4°) and dashed (44.2°) curves show the results obtained from the PWIA calculation, while the dotted curve represents the experimental resolution when the target nucleon is at rest.

(OVER)




Fig. 5. The fitted values of parameters in the Gaussian distribution for pion momentum spectra: (a) The momentum shift $p_{free} - p_0$, (b) the width 2Γ , and (c) the cross section $d\sigma/d\Omega$ as a function of $\langle k \rangle$. Curves in (a) and (b) are the result of the PWIA calculation for $\theta_a = 28.4^\circ$ (solid and dotted curves for π^- and π^+ , respectively) and 44.2° (dashed one for π^-). Curves in (c) show the differential cross sections (multiplied by a factor of 6) for elementary processes (solid and dotted ones for π^- and π^+ at 28.4° , respectively, while dashed one is for π^- at 44.2°). Symbols in (b) and (c) are the same as in (a).

REF. S. V. Dementii, V. I. Ogurtsov, N. G. Afanas'ev
 Yad. Fiz. 28, 865 (1978)
 Sov. J. Nucl. Phys. 28, 443 (1978)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.			
				78De11		hg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	50-200	D	1*2	MAG-D		DST
				(799-1178)			

Measurements were made of quasielastic scattering of electrons in the (e, e') reaction by the nuclei ${}^9\text{Be}$, ${}^{12}\text{C}$, ${}^{14}\text{N}$, ${}^{16}\text{O}$, and ${}^{27}\text{Al}$. An experimental estimate is obtained of the effective mass of an intranuclear nucleon. At excitation energies up to 80 MeV, a ratio $M^*/M = 0.6$ is obtained, corresponding to a linear potential $V(E) = V_0 + 0.4E$ and in good agreement with the data on proton scattering by nuclei [C. M. and F. G. Perey, Atomic Data and Nuclear Tables 13, 294 (1974)]. At excitation energies above 120 MeV the nucleon effective mass turned out to be close to that of the free nucleon, $M^*/M = 0.9$.

*GEV, QUASIELASTIC

PACS numbers: 25.30.Cg, 27.20.+n, 27.30.+t




FIG. 4. Reduction coefficient M^*/M as a function of energy transfer: \bullet — ${}^9\text{Be}$, \blacktriangle — ${}^{12}\text{C}$, \circ — ${}^{14}\text{N}$, \triangle — ${}^{16}\text{O}$, \square — ${}^{27}\text{Al}$. The curve has been drawn through the experimental points by hand.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	4, 16	D	57-215	MAG-D		180

The transverse electromagnetic form factors squared of the $^{12}\text{C } 2^+$ levels at 4.439 MeV ($T=0$) and at 16.107 MeV ($T=1$) have been measured by means of 180° electron scattering over a momentum-transfer range from $q=0.51$ to 2.05 fm^{-1} . Evidence is presented for appreciable contributions of nuclear convection currents to the transverse 4.439-MeV form factor at low q , and spin magnetization contributions to the transverse 16.107-MeV form factor at higher q .

4=4.439; 16=16.107




FIG. 1. Squared form factors for the $T=1$, 16.107 MeV and the $T=0$, 4.439-MeV 2^+ states in ^{12}C . Curves A and B are the transverse form factors squared of the $T=1$ and $T=0$ states calculated with $b=1.64$ and 1.76 fm , respectively. Curve C is the longitudinal form factor squared for the $T=0$ state calculated with $b=1.76 \text{ fm}$. (Triangles, data of Ref. 15; unfilled squares, data of Ref. 10; and filled squares, data of Ref. 11.)




FIG. 2. Deduced convection-current part of the $T=0$, 4.439-MeV 2^+ transverse form factor squared. The dashed curve was obtained by using Cohen and Kurath configurations with Woods-Saxon wave functions normalized to the $B(E2)$ value at the photon point, and the solid curve by using a pure $(p_{1/2}, p_{3/2})$ configuration. (Δ , datum point of Ref. 15.)

¹⁰Tohoku University, Sendai, Japan, Research Report of the Laboratory of Nuclear Science, 1969 (unpublished), Vol. 2, p.1.

¹¹J.S. McCarthy et al., Ref. 10.

¹⁵P. Strehl, Z. Phys. 234, 416 (1970).

REF. A. Friebel, P. Manakos, A. Richter, E. Spamer, W. Stock and
O. Titze
Nucl. Phys. A294, 129 (1978)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	
78 Fr 2	rs

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E_e E'$	LFT	16	D	33- 62	MAG-D		DST

16=16.11 MeV

Abstract: The electroexcitation of the 2^+ , $T = 1$ state at 16.11 MeV in ^{12}C has been investigated for momentum transfers $q < 0.5 \text{ fm}^{-1}$. Longitudinal and transverse contributions were separated by means of the angular dependence of the cross section. The ground-state radiative width has been determined to $\Gamma_0^0 = 0.346 \pm 0.041 \text{ eV}$ and the contribution of the magnetization density to the current has been derived from the q -dependence of the reduced transition probability. These quantities are compared to $1p$ - $1h$ shell-model and intermediate coupling model calculations which are both in disagreement with the experimental numbers. This discrepancy is, however, partly removed when collective states with $2\hbar\omega$ excitations are included.




Fig. 3. Angular distributions of the measured inelastic cross sections at various constant momentum transfers. The symbols on the ordinate and abscissa are explained in the main text. The angle associated with each data point is also indicated. This figure is needed to determine $B(C2, q)$ from the slopes of the curves fitted to the data and $B(E2, q)$ from their intersects with the ordinate.




Fig. 5. Longitudinal and transverse form factors for the electroexcitation of the 16.11 MeV state in ^{12}C . Besides the Darmstadt data of the present experiment a data point from Mainz ²¹ is also shown. The curves compared to the data are calculated within the shell model including collective states with $2h\omega$ excitations and using free nucleon charges and magnetic moments.

²¹H.D. Wohlfahrt, Thesis, University of Mainz, 1969, unpublished

TABLE 2
Experimental $B(C2, k)\uparrow$ value compared to various theoretical predictions

Configuration	$B(C2, k)\uparrow$ (fm^4)		
Experiment	1.80 ± 0.21		
$(p_{3/2})^{-1}p_{1/2}$	7.13		
$(p)^8$	3.87 ^{a)}	4.22 ^{b)}	3.14 ^{c)}
$(pf)(p)^7(s)^4 + (p)^8$	2.60 ^{c)}		
$(pf)(p)^7(s)^4 + (p)^8 + (sd)(p)^8(s)^3$	2.18 ^{c)}	2.80 ^{a)}	

^{a)} With p-shell interaction (8-16) POT from ref. ¹⁵).

^{b)} With p-shell interaction (8-16) 2BME from ref. ¹⁵).

^{c)} With central p-shell interaction and Rosenfeld exchange mixture. $L:K = 6:8$ and $a:K = 4:5$.

¹⁵S. Cohen and D. Durath, Nucl. Phys. 73 1 (1965)

REF. V.V. Kirichenko, Yu.M. Arkatov, P.I. Vatset, I.V. Dogyust,
A.F. Khodyachikh
Yad. Fiz. 27, 588 (1978)
Sov. J. Nucl. Phys. 27, 314 (1978)

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		78 Ki 8
		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	16-120	C	150	CCH-D		DST

A diffusion chamber in a magnetic field has been used to measure total and differential cross sections for the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$ in the energy interval from threshold to 120 MeV. The angular distributions are analyzed in order to identify the principal transitions. It is shown that for $E_\gamma > 43$ MeV an important contribution to the reaction is provided by the channel with the residual nucleus in the $1/2^-$ state and excitation energy 6.793 MeV. The contribution from this channel is estimated. The results are compared with theoretical calculations.

$$\int_{16}^{120} \sigma(\gamma, p) dE = 189 \pm 9 \text{ MEV}\cdot\text{mb} \text{ (statistical and systematical errors included)}$$

PACS numbers: 25.20.+y, 27.20.+n

δ represents contribution of excited-state transitions to assumed ground state cross section




FIG. 1. Total cross section for the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$ as a function of energy. The histogram shows the results of the present experiment; the hollow circles are from Ref. 4, the solid squares are from Ref. 2, and the solid circles are from Ref. 7.




FIG. 2. Differential cross sections for protons from the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$ for several intervals of γ -ray energy. Points: \bullet —present experiment, \circ —Ref. 10, \blacksquare —Ref. 2, \times and Δ —Ref. 3.

USCOMM-DC 26910-P64




FIG. 3. The coefficients $\alpha_{i,k}$ as a function of energy. The solid curve is from Ref. 1, and the hollow squares from Ref. 4; the remaining designations are the same as in Fig. 2.

$$\frac{d\sigma}{d\Omega} = \sum_{i=0}^3 A_i P_i(\cos\theta)$$

$$A_i/A_k = \alpha_{ik}$$

TABLE I. Relative contribution δ of the channel with an excited state of the residual nucleus ($E_\gamma' = E_\gamma + \epsilon$).

E_γ , MeV	E_γ' , MeV	δ	E_γ , MeV	E_γ' , MeV	δ
22-24	29-31	0.05±0.14	34-36	41-43	0.47±0.14
24-26	31-33	-0.11±0.15	36-40	43-47	0.65±0.17
26-28	33-35	0.03±0.15	40-50	47-57	0.61±0.15
28-30	35-37	0.03±0.15	50-60	57-67	0.60±0.15
30-32	37-39	0.04±0.17	60-90	67-87	0.44±0.11
32-34	39-41	0.04±0.22	90-120	87-127	0.14±0.10

ELEM. SYM.	A	Z
C	12	6
REF. NO.		egf
78 K1 5		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
(E,N)	ABX	18-29	D	30	ACT-I		4PI

The cross section for the production of the ¹¹C activity by bombarding ¹²C with 30 MeV electrons has been measured absolutely. The result, $11.9 \pm 0.2 \mu\text{b}$, has a smaller absolute error than any previous measurement. This value corresponds to a (γ, n) cross section integrated to 30 MeV of $41.0 \pm 0.6 \text{ MeV mb}$, based on a distorted-wave Born-approximation electric dipole virtual photon spectrum.

[NUCLEAR REACTIONS ¹²C(e,n), $F_e = 30 \text{ MeV}$, $\sigma(e,n)$ measured absolutely.]

The weighted average cross section for 30 MeV electrons is then

$$\sigma(e, n) = 11.92 \pm 0.15 \mu\text{b}.$$

The systematic errors common to all of the measurements, 1.5% for the absolute ²²Na source calibration and 0.5% for the target thickness, must be combined with the above to yield

$$\sigma(e, n) = 11.9 \pm 0.2 \mu\text{b}.$$

TABLE I. The cross section for the production of ¹¹C by 30 MeV electrons.

Run	Cross section
I	12.15 ± 0.31
II	11.73 ± 0.47
III	11.84 ± 0.32
IV	11.99 ± 0.34
V	11.76 ± 0.32
Weighted average	11.92 ± 0.15

⁷G. Kuhl and U. Kneissl, Nucl. Phys. A195, 559 (1972).
⁸J. Müller, C. Schule, G. Tamas, and C. Tzara, J. Phys. (Paris) 27, 3 (1966).
⁹W. A. Lochstet and W. E. Stephens, Phys. Rev. 141, 1002 (1966).
¹⁰B. C. Cook, J. E. E. Baglin, J. N. Bradford, and J. E. Griffin, Phys. Rev. 143, 724 (1966).
¹¹S. C. Fultz, J. T. Caldwell, B. L. Berman, R. L. Bramblett, and R. R. Harvey, Phys. Rev. 143, 790 (1966).

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	78 Ma 10

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABY	19-68	C	30-68	ACT - I		4PI
G,AN	ABY	26-68	C	30-68	ACT - I		4PI

Analysis is made of reactions interfering with photon activation analysis procedures.

The activation yield curves have been presented for a number of photonuclear reactions in the energy range from 30 to 68 MeV, in order to evaluate quantitatively the interferences due to competing reactions in multielement photon activation analysis. The general features of the yields as functions of both target mass number and excitation energy were elucidated from the data obtained, discussion being given on the results in terms of the reaction mechanism.

Simultaneous neutron activation due to appreciable neutron production from the converter and surrounding materials has also been studied, and, finally, the magnitudes of interferences in real multielement analysis were given in the form of their energy dependences.




Fig. 2. Yield per equivalent quanta versus target neutron number.




Fig. 3. Activation yield curves for the reactions on C, Na, Mg and Al
 □ ¹²C(p,n)¹¹C, ■ ⁹Be(p,n)⁹Be, ▲ ²³Na(p,n)²³Na,
 ▢ ²⁴Mg(p,n)²⁴Na, ● ²⁴Mg(p,n)²⁴Na, ▣ ²⁷Al(p,n)²⁷Al,
 ○ ²⁷Al(p,n)²⁷Na.

(over)




Fig. 9. Yields of the (γ, n) reactions as a function of bremsstrahlung maximum energy and target mass number. The numerical values in the figure are yields per equivalent quanta in mb.

REF. J. Mougey, M. Bernheim, D. Royer, D. Tarnowski, S. Turck,
 P.D. Zimmerman, J.M. Finn, S. Frullani, D.B. Isabelle,
 G.P. Capitani, E. De Sanctis and I. Sick
 Phys. Rev. Lett. 41, 1645 (1978)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		rs
78 Mo 4		

METHOD			SOURCE		DETECTOR		ANGLE
DEFINE: $R(E_0, \omega) = \sigma \text{ measured} / \sigma \text{ proton mott}(E_0, \theta)$			TYPE	RANGE	TYPE	RANGE	
REACTION	RESULT	EXCITATION ENERGY	D	160-520	MAG-D		DST
E, E/	ABX	30-350					

A systematic study of the deep-inelastic electron-scattering response function of ^{12}C has been carried out at scattering angles of 60° and 130° and electron energies between 160 and 520 MeV. A pronounced transverse strength, the origin of which is not understood, is found in the region between the quasielastic and the N peak.




FIG. 1. Inelastic response function for ^{12}C at 60° and electron incident energies between 240 MeV (topmost curve) and 520 MeV in steps of 40 MeV. Where not shown, the error bar is smaller than the dot.




FIG. 2. Inelastic response function for ^{12}C , as in Fig. 1 but for 130° .

over




FIG. 3. Response function for ^{12}C compared to a calculation for one-nucleon-knockout process (solid line), coherent pion production (dash-dot line) (Ref. 16), Δ excitation (dashed line) (Ref. 17), and meson-exchange-current contribution (dotted line) (Ref. 19).

- ¹⁶E. Borie, in "Proceedings of the Seventh International Conference on High-Energy Physics and Nuclear Structure, Zurich, 1977" (Schweizerisches Institut für Nuklearforschung, Villigen, 1977), p. 247.
- ¹⁷G. Do Dang, Phys. Lett. 69B, 425 (1977), and private communication.
- ¹⁹T.W. Donnelly, J.W. van Orden, T. de Forest, Jr., and J.C. Hermans, Phys. Lett. 76B, 393 (1978); T. de Forest, Jr., private communication.

REF. N. G. Shevchenko, A. Yu. Buki, B. V. Mazan'ko, V. N. Polishchuk,
 A. A. Khomich
 Yad. Fiz. 28, 12 (1978)
 Sov. J. Nucl. Phys. 28, 5 (1978)

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.		hg	
				78Sh8			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,E/	SPC	18 (18.1)	D	140	MAG-D		DST

Energy spectra of 140-MeV electrons are measured after scattering by ^{10}B and ^{12}C nuclei. Peaks above the nucleon emission threshold are observed in the ^{10}B spectra at energies 8.26, 9.0, and 9.7 MeV, and at 13.1 MeV in the ^{12}C spectra. The existence of these peaks and their energy location is predicted by calculations based on inclusion of the contribution of triangle Feynman diagrams to the reaction amplitude.

EX 18.1 MEV

Examines levels above the nuclear threshold

PACS numbers: 25.30.Cg, 27.20.+n




FIG. 3. Spectra of 140-MeV electrons after scattering by ^{12}C at angles 90 and 100°. We have indicated in the figure the energies of the discrete levels of ^{11}B and ^{11}C , and also the proton and neutron binding energies in the ^{12}C nucleus. For each spectrum we have indicated the exposure in monitor units of the second range R2 and of the fourth range R4, equal to 20.3×10^{-4} Coulomb.

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		79 Bo 3
		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	ABX	150-360	C	150-360	ACT-I		90
E,PI-	RLX	200-360	D	150-360	ACT-I		90

Total cross sections for $^{12}\text{C}(\gamma, \pi^-)^{12}\text{N}$ and $^7\text{Li}(\gamma, \pi^-)^7\text{Be}$ have been measured from threshold to 360 MeV photon energy by detecting the radioactivity of the residual nuclei, thereby singling out the ground state of ^{12}N and the ground and first excited states of ^7Be . The cross sections are found to peak at about 40 MeV pion energy and then to fall gradually. In contrast to pion charge exchange and other photopion experiments, these results are well reproduced both in shape and in magnitude by distorted-wave impulse-approximation calculations.




FIG. 2. Yields for (a) ^{12}C and (b) ^7Li as a function of incident electron energy. The theoretical curves are those calculated from the corresponding cross sections shown in Fig. 3 with the two-step background added back in. (c) Real-to-virtual ratios per equivalent quantum (i.e., photoproduction-to-electroproduction ratios) for ^{12}C vs endpoint energy compared with Dalitz-Yennie predictions.




FIG. 3. Extracted cross sections for (a) ^{12}C and (b) ^7Li vs photon energy above threshold (shaded regions). Calculation of EST (see Ref. 13) shown as solid lines (full interaction) and dotted line (Coulomb only). Calculation of NU (see Ref. 15) shown as dash-dotted line.

REF. K. Sh. Egiyan
 Yad. Fiz. 30, 890 (1979)
 Sov. J. Nucl. Phys. 30, 461 (1979)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
79 Eg 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	RLY	16-250	C	130,250	MAG-D		DST

Experimental data are presented on the inclusive photoproduction of protons in the nuclei ^{12}C , ^{24}Mg , ^{63}Cu , ^{118}Sn , and ^{208}Pb irradiated by bremsstrahlung with maximum energies 0.13 and 0.25 GeV. The regions of angles $30-90^\circ$ and of photoproton momenta $0.24-0.48 \text{ GeV}/c$ were studied.

PACS numbers: 25.20. + y




FIG. 2. Relative yields of photoprotons as a function of emission angle; experimental points: \circ , \bullet —for $p_p = 0.29 \text{ GeV}/c$; Δ , \blacktriangle — $0.34 \text{ GeV}/c$; \square , \blacksquare — $0.40 \text{ GeV}/c$. The hollow points are for $E_{\gamma, \text{max}} = 0.25 \text{ GeV}$ and the solid points for $E_{\gamma, \text{max}} = 0.13 \text{ GeV}$: a—for ^{12}C , b—for ^{24}Mg , c—for ^{208}Pb .




FIG. 3. Momentum spectra of protons. The experimental points are as follows: \circ and \bullet —for ^{12}C , Δ —for ^{63}Cu , \square —for ^{208}Pb . The hollow points are for $E_{\gamma, \text{max}} = 0.25 \text{ GeV}$ and the solid points are for $E_{\gamma, \text{max}} = 0.13 \text{ GeV}$; a—for $\vartheta_p = 30^\circ$, b—for $\vartheta_p = 60^\circ$, c—for $\vartheta_p = 90^\circ$. The lines have been drawn through the experimental points by the method of least squares.

TABLE II. Values of the exponent π in the A^π dependence of the proton yield in reactions (2) and (3).

ϑ_p , deg	$E_{\gamma, \text{max}} = 0.25 \text{ GeV}$			$E_{\gamma, \text{max}} = 0.13 \text{ GeV}$	
	$p_p, \text{ GeV}/c$			$p_p, \text{ GeV}/c$	
	0.29	0.34	0.40	0.29	0.34
30	1.15 ± 0.04	1.17 ± 0.04	1.20 ± 0.05	0.59 ± 0.18	0.62 ± 0.06
60	—	1.17 ± 0.02	1.22 ± 0.03	—	—
90	1.02 ± 0.03	1.11 ± 0.03	1.24 ± 0.03	—	—

(over)




FIG. 5. A-dependence of the photoproton yield in reactions (2) and (3). Experimental points for $E_{\gamma \text{ max}} = 0.25$ GeV: circles— $p_p = 0.29$ GeV/c; triangles— $p_p = 0.34$ GeV/c, squares— $p_p = 0.40$ GeV/c; half-open symbols—for $\vartheta_p = 30^\circ$, open symbols— $\vartheta_p = 60^\circ$, solid symbols— $\vartheta_p = 90^\circ$, for $E_{\gamma \text{ max}} = 0.13$ GeV: \diamond — $p_p = 0.29$ GeV/c, $\vartheta_p = 30^\circ$; \blacklozenge — $p_p = 0.34$ GeV/c, $\vartheta_p = 30^\circ$. The lines have been drawn through the experimental points by the method of least squares.

REF. V.N. Eponeshnikov, Yu.F. Krechetov
 JETP Lett. 29, 401 (1979)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
79 Ep 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, PIOP	ABY	0*600	C	450	CKV-D		DST

*MOM, MEV/C, COIN

The cross sections of reaction $(\gamma, \pi^+ p)$ on Li^7, C^{12} nuclei were measured in the 0-600 MeV/c range of momentum transfer to the residual nucleus. For large values of momenta, cross section values disagree with calculations carried out within the framework of a shell model and a model of quasi-free meson photoproduction on nuclei.

PACS numbers: 25.20. + y




FIG. 1. Dependence of reaction cross section on the proton angle of emission: O—experiment, with total measurement errors; --- calculations using plane waves; — final state interactions taken into account; effect of S-shell shown separately; — calculations with momentum distribution from Ref. 3 with allowance for correlation by the Jastrow model. Oscillator parameters $\alpha_1 = \alpha_2 = 115$ MeV/sec for Li^7 nucleus, 120 MeV/sec for C^{12} and 113 MeV/sec for O^{16} . Arrow indicates angle of proton emission in the case of reaction $\gamma + p \rightarrow \pi^0 + p$.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.E/	FMF	12.15	D	0*3	MAG-D		DST

The structure and the degree of isospin mixing for the 1^+ levels in ^{12}C at 12.71 MeV ($T=0$) and 15.11 MeV ($T=1$) are determined from measured 180° -electron-scattering form factors. The resulting charge-dependent isospin-mixing matrix element ranges from 130 to 165 keV depending on the theoretical model of the isoscalar form factor.

*MOM FM-1, 2 LEVELS




FIG. 2. The 15.11-MeV form factor. Solid circles are the data of this experiment, and open circles those of other experiments (Ref. 5). The solid curve was obtained as described in the text, and includes meson-exchange currents. The inset shows the 180° -Coulomb distortion correction used to obtain the points shown.




FIG. 3. Shell-model predictions for the 12.71-MeV form factor compared with the present data. The solid lines show the (S-16) 2BME prediction and the dashed lines the CK-KB-G prediction. The upper curves include isospin mixing with amplitude β as given in Table I but the lower curves do not. The inset shows the 180° -Coulomb-distortion corrections used to obtain the points shown.

REF. I.V. Glavanakov, V.N. Stibunov
 Yad. Fiz. 29, 1455 (1979)
 Sov. J. Nucl. Phys. 29, 746 (1979)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.	hg
	79 G1 5	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-P	ABY	150-390	C	390	MAG-D		120

The differential yield of the reaction $^{12}\text{C}(\gamma, \pi^- p)$ has been measured for a bremsstrahlung maximum energy 390 MeV. The results of the measurement are analyzed in terms of the impulse approximation and the shell model. Inclusion of the final-state interaction leads to satisfactory quantitative agreement of the theoretical reaction yields and the experimental data.

PACS numbers: 25.20.+y, 27.20.+n, 13.60.Kd




FIG. 1. Differential yield of the reaction as a function of proton energy for $e_\gamma = 350$ MeV.




FIG. 2. Differential yield of the reaction as a function of the invariant mass of the (πp) pair at $p_\pi = 70$ MeV/c.




FIG. 3. Differential yield of the reaction as a function of proton energy for $W = 1200$ MeV.

REF. I.V. Glavanakov, V.N. Stibunov
 Yad. Fiz. 30, 897 (1979)
 Sov. J. Nucl. Phys. 30, 465 (1979)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
79 G1 6		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-P	ABY	150-390	C	340-390	MAG-D		DST

COINC PI- WITH P

We have measured the differential cross section for the reaction $^{12}\text{C}(\gamma, \pi^- p)$ at photon energies 340, 360, and 380 MeV in two ranges of residual-nucleus excitation energy (0-10 MeV and 10-40 MeV). The results of the measurements are analyzed in terms of the impulse approximation and the shell model. It is shown that the main contribution to the reaction yield is due to the quasifree photoproduction mechanism. The interaction in the final state exerts a substantial effect on the value of the cross section. An estimate is obtained for the parameter of the oscillator wave function of the neutrons bound in the ^{12}C nucleus.

PACS numbers: 25.20. + y




FIG. 2. Energy dependence of the cross section for $e_x < 10$ MeV: a— $E_\gamma = 340$ MeV, b— $E_\gamma = 360$ MeV, c— $E_\gamma = 380$ MeV.




FIG. 3. Energy dependence of the cross section for $10 \text{ MeV} < e_x < 40 \text{ MeV}$: a— $E_\gamma = 360$ MeV, b— $E_\gamma = 380$ MeV.

e_x = excitation energy of residual nucleus

REF. S. Hartwig, F. H. Heimlich, G. Huber, E. Rössle, J. Bleckwenn, M. Köbberling, J. Moritz, K. H. Schmidt, D. Wegener, D. Zeller, P. David, H. Mommsen
Phys. Lett. 82B, 297 (1979)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
79Ha2		

METHOD			SOURCE		DETECTOR		ANGLE
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE	
E, E/	RLX	1*2	D	2*3	MAG-D		13
		(.95-2.16)		(2.18, 3.06)			

*GEV, RLX E+/E-

The ratio $R = \sigma^+/\sigma^-$ of the cross sections for inelastic positron and electron scattering on ^{12}C and ^{27}Al has been measured for four momentum transfers $0.08 (\text{GeV}/c)^2 \leq q^2 \leq 0.45 (\text{GeV}/c)^2$ of the virtual photon and invariant masses $0.95 \text{ GeV} < W < 3.3 \text{ GeV}$ of the hadronic system. The mean value of the ratio is $R = (1.005 \pm 0.027)$. No q^2 , respectively, W dependence of the ratio is observed.

Table 1
Ratio $R = \sigma_+/\sigma_-$ for inelastic lepton scattering on ^{12}C . E_1 , primary electron energy, θ_e , electron scattering angle, q^2 , four-momentum transfer of the virtual photon, W , invariant mass of the excited hadronic system, A_2 , amplitude of two-photon exchange, A_1 , amplitude of one-photon exchange. All errors include the systematic error contributions given in the text.

$E_1 = 2.68 \text{ GeV}$, $\theta_e = 13^\circ$				$E_1 = 3.08 \text{ GeV}$, $\theta_e = 13^\circ$			
W (GeV)	q^2 ((GeV/c) ²)	R	$\text{Re}(A_2/A_1)$	W (GeV)	q^2 ((GeV/c) ²)	R	$\text{Re}(A_2/A_1)$
0.95	0.342	1.027 ± 0.06	(0.7 ± 1.5)%	0.928	0.450	1.041 ± 0.031	(1.0 ± 0.8)%
1.247	0.297	1.004 ± 0.037	(0.1 ± 0.9)%	1.275	0.391	1.032 ± 0.029	(0.8 ± 0.7)%
1.478	0.254	1.019 ± 0.037	(0.5 ± 0.9)%	1.520	0.337	1.025 ± 0.029	(0.6 ± 0.7)%
1.670	0.213	1.013 ± 0.037	(0.3 ± 0.9)%	1.702	0.292	1.045 ± 0.032	(1.1 ± 0.8)%
1.855	0.168	0.948 ± 0.04	(-1.3 ± 1.0)%	1.866	0.247	0.992 ± 0.032	(-0.2 ± 0.8)%
		0.985 ± 0.025	(-0.4 ± 0.6)%	2.022	0.199	0.983 ± 0.031	(-0.4 ± 0.8)%
				2.163	0.154	0.957 ± 0.031	(-1.1 ± 0.8)%
						1.019 ± 0.022	(0.5 ± 0.5)%




Fig. 1. Ratio $R = \sigma_+/\sigma_-$ of inelastic positron and electron scattering on ^{12}C and ^{27}Al as a function of the invariant mass W of the excited hadronic system, calculated for a free target nucleus.




Fig. 2. Ratio $R = \sigma_+/\sigma_-$ of inelastic positron and electron scattering on ^{12}C and ^{27}Al as a function of the four-momentum transfer q^2 of the virtual photon.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
79 Ki 2		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, PA	ABX	27-120	C	120	CCH-D		DST
G, NA	ABX	27-120	C	120	CCH-D		DST

A diffusion chamber in a magnetic field has been used to measure the total and differential cross sections as a function of energy in the interval from threshold to 120 MeV. Investigations were made of the energy correlations of the reaction products, the distributions in the average energy of the particles, and the excitation energy of the various intermediate states. It is shown that at energies above the giant resonance the two reactions occur through intermediate excited states of the nuclei ^{11}B and ^{11}C . The excitation energies and widths of the resonances of these states have been measured. An estimate is made of the relative contribution of electric quadrupole transitions. Possible mechanisms for interaction of the γ rays with the nucleus are discussed.

Photon energy was calculated on the assumption that the residual nuclei are in the ground state.

PACS numbers: 25.20. + y, 24.30. - v, 27.20. + n




FIG. 1. Total cross sections for reaction (1) and (2) and their ratio η as a function of γ -ray energy. a)—total cross section for reaction (1), b)—for reaction (2), c)— $\eta = \sigma(\gamma, p\alpha) / \sigma(\gamma, n\alpha)$. The solid curve shows the data of Ref. 9.




FIG. 2. Energy correlations of the products of reactions (1) and (2) in various γ -ray energy intervals. The solid curves represent phase space.³




FIG. 6. Asymmetry coefficient β of nucleon angular distributions as a function of γ -ray energy. Points: ●—reaction (1), ▲—reaction (2), ○— $^4\text{He}(\gamma, p)^3\text{H}$ (Ref. 16), Δ and ◀— $^4\text{He}(\gamma, n)^3\text{He}$ (Refs. 5 and 6).

(OVER)

TABLE I. Comparison of integrated cross sections

$$\sigma_{int} = \int_{E_{\gamma 1}}^{E_{\gamma 2}} \sigma(E_{\gamma}) dE_{\gamma}$$

and yield of ${}^7\text{Be}$ per equivalent quantum

$$\sigma_Q = \int_{E_{\gamma 1}}^{E_{\gamma 2}} \sigma(E_{\gamma}) dE_{\gamma} / E_{\gamma}$$

with the results of other studies.

E_{γ} , MeV	${}^6\text{C}(\gamma, \text{pa}){}^7\text{Li}$		${}^6\text{C}(\gamma, \text{na}){}^7\text{Be}$			
	σ_{int} , MeV-mb		σ_{int} , MeV-mb		σ_Q , mb	
	a	b	a	b	a	b
25-40	5.9±0.4	4.9 (*)	3.8±0.3	4.1 (*)	0.105±0.017	0.083±0.007 (*)
25-55	7.9±0.5		5.6±0.4	8.5±1.0 (*)	0.142±0.097	0.143±0.011 (*)
25-80	9.3±0.8	4.17 (*)	6.4±0.4		0.170±0.010	0.158±0.027 (*)
25-100	9.8±0.6		6.8±0.5		0.176±0.010	0.180±0.020 (*)
25-120	9.7±0.6	10.2±1.0 (*)	6.9±0.5	8.6±1.7 (*)	0.173±0.011	0.190±0.030 (*)
40-120	3.8±0.3	5.3 (*)	3.1±0.3	4.5 (*)		
60-120	1.4±0.2	1.8 (*)	1.2±0.2	1.5 (*)		

Note. In the columns, a is the result of the present experiment and b are the data cited.

TABLE II. Results of fitting the nucleon angular distributions by a sum of Legendre polynomials: $d\sigma/d\Omega = \sum_{l=0}^2 A_l P_l(\cos\theta)$.

E_{γ} , MeV	${}^6\text{C}(\gamma, \text{na}){}^7\text{Li}$			${}^6\text{C}(\gamma, \text{na}){}^7\text{Be}$		
	A_0 , $\frac{\text{mb}}{\text{sr}}$	A_1 , $\frac{\text{mb}}{\text{sr}}$	A_2 , $\frac{\text{mb}}{\text{sr}}$	A_0 , $\frac{\text{mb}}{\text{sr}}$	A_1 , $\frac{\text{mb}}{\text{sr}}$	A_2 , $\frac{\text{mb}}{\text{sr}}$
25-32	2.57±0.13	0.56±0.24	-0.08±0.35	1.26±0.13	-0.23±0.21	-0.26±0.25
32-40	3.07±0.17	0.53±0.28	0.85±0.36	2.47±0.16	-0.16±0.23	-1.05±0.36
40-50	1.22±0.11	0.54±0.19	0.43±0.23	0.99±0.10	0.00±0.15	-0.40±0.19
50-70	0.54±0.15	0.43±0.11	0.55±0.12	0.43±0.05	0.29±0.09	-0.02±0.10
50-120	0.33±0.03	0.33±0.06	0.28±0.07	0.31±0.03	0.23±0.05	0.14±0.06

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
79 Ko 1		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	SPC	53-89	D	40-80	NAI-D		60

Observations are reported from the first systematic studies of proton radiative capture at intermediate energies. In addition to captures to the ground and first few excited states, the reactions $^{11}\text{B}(p,\gamma)^{12}\text{C}$ and $^{27}\text{Al}(p,\gamma)^{28}\text{Si}$ reveal unexpectedly strong transitions to isolated high-lying states. These latter transitions could take place via "second-harmonic" giant resonances. Protons on ^{12}C produce a rich spectrum of γ rays which may arise from captures to high-lying single-particle states.




FIG. 1. γ -ray spectra from the bombardment of ^{11}B with 40-, 60-, and 80-MeV protons, observed at $\theta_{\text{lab}} = 60^\circ$. Labeled arrows indicate the expected positions of captures to the ground state (γ_0) and low-lying excited states of ^{12}C . The peak labeled "19.2" is identified as radiative capture to a state at ~ 19.2 -MeV excitation energy. *Inset*: time-of-flight spectrum, at $E_p = 60$ MeV, for events leaving more than 30 MeV in the NaI(Tl) crystal. The time interval between the peaks is 12 ns. Following beam bursts provide the "stop" signal.




FIG. 3. The spectrum from $^{12}\text{C}(p,\gamma)$ at 40-MeV bombarding energy, $\theta_{\text{lab}} = 60^\circ$.

REF. F.L. Milder, E.C. Booth, B. Chasan, A.M. Bernstein, J. Comuzzi,
G. Franklin, A. Nagl, H. Uberall
Phys. Rev. C19, 1416 (1979)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
79 Mi 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABX	150-175	C	175	TEL-I		4PI

The π^+ photoproduction on ^{12}C has been measured relative to the proton from 0-20 MeV above threshold. Total cross sections from 0-12 MeV, summed over states in ^{12}B have been extracted with accuracies between 5% and 8% relative to the proton photoproduction cross section. Distorted wave impulse approximation calculations have been performed and show good agreement with the data. The yield from 0-3 MeV above threshold is well described by a one parameter fit for the ground state cross section using only the $\sigma \cdot \vec{\epsilon}$ term of the Hamiltonian.

DETECT PI-MU-E DECAY

TABLE II. Comparison of the total summed ^{12}C cross section with DWIA theoretical calculation at several photon energies above threshold.

ΔE_{LSD}	$\sigma_{exp} (\mu b)$	$\sigma_{DWIA} (\mu b)$
2	1.02 ± 0.08	1.23
4	2.33 ± 0.15	2.47
6	5.07 ± 0.24	5.04
8	9.42 ± 0.45	8.75
10	14.09 ± 0.69	13.13

[NUCLEAR REACTIONS $^{12}\text{C}(\gamma, \pi^+)^{12}\text{B}$, bremsstrahlung endpoint energies to 175 MeV, deduced $\sigma(E)$; calculated $\sigma(E)$, DWIA.]




FIG. 3. (a) Experimental yields per equivalent quanta for $p(\gamma, \pi^+)n$ and fit generated from $\sigma_H = 201(p^2/k)$ ($1-0.0063E$). The fit provides the absolute normalization for the experiment. (b) Experimental yields per equivalent quanta for $^{12}\text{C}(\gamma, \pi^+)^{12}\text{B}$. The solid curve is obtained by folding the bremsstrahlung spectrum with the total summed cross section shown in Fig. 4.

(over)




FIG. 4. Unfolded total summed $^{12}\text{C}(\gamma, \pi^+)^{12}\text{B}$ cross section from a fit to the measured yields. The width of the line represents the total estimated error. The upper dashed line is the theoretical summed cross section from Table II. Also shown as a dashed line is the ground state calculation $\sigma_{\text{BWA}}^{\text{I}}$ from Table III.


REF. J. Arends, J. Eyink, H. Hartmann, A. Hegerath, B. Mecking, G. Noldeke, H. Rost
Z. Phys. A298, 103 (1980)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
80 Ar 8		hg


REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,XP	ABX	200-385	D	200-385	MAG-D		DST
G,PPI	ABX	200-385	D	200-385	MAG-D		54
G,PN	ABX	200-385	D	200-385	MAG-D		DST
G,PP	ABX	200-385	D	200-385	MAG-D		DST

Data are presented for proton emission off ^{12}C using tagged photons in the energy range 200-385 MeV. The protons are detected in a magnetic spectrometer. In addition, charged or neutral particles can be measured with a scintillation counter setup. Contributions to the proton energy spectra from primary pion production and quasideuteron reactions are separated using the coincidence data. The total $^{12}\text{C}(\gamma, p)X$ cross section is derived as a function of the photon energy. The data are compared to the predictions of an intranuclear cascade calculation. Final state interactions have to be taken into account to describe the data.

(G,PPI) COINC P/CHARGED PION
(G,PN) COINC PROTON/NEUTRON
(G,PP) COINC PROTON/PROTON



* Fig. 5. Double differential cross section for the photoemission of a proton off ^{12}C by photons of energies between 210 and 381 MeV. The laboratory angular range is 44-54 degrees



* Fig. 10. Double differential cross section for the photoemission of protons off ^{12}C by photons with (315-353) MeV as a function of the proton energy. The proton angular interval is 44°-54°. a) Single arm data (no particle in the scintillation counter arm is required). b-d) The particle detected in the scintillation counter arm is required to be a charged pion b, a neutron c, a proton d

*Additional plots of type given in Figs. 5 and 10 are in Figs. 6-10, 11 and 12 of text.




Fig. 13. Ratio of pp to pn pairs as a function of the photon energy. The spectrometer angle was restricted to 44° - 54° while the full acceptance of the scintillation counter arm was used for the second particle




Fig. 14. Distribution of the missing energy E_m for NN coincidence events. The photon energy range is 353 - 391 MeV. Full line: pn coincidences, dashed line: pp coincidences




Fig. 15. Total inclusive cross sections for proton emission off ^{12}C as a function of the photon energy. \bullet total (γ, p) cross section. Δ total (γ, p) cross section without a pion in the final state. The extrapolation of the data was done using the cascade calculation PICA




Fig. 16. Total cross section for (γ, p) with a pion in the final state. The dashed curve shows the sum of the elementary cross sections for $\gamma p \rightarrow \pi^0 p$ and $\gamma n \rightarrow \pi^- p$. The full line represents the result of a Fermi gas model including pion final state interactions and Pauli blocking

REF. P. Argan, G. Audit, A. Bloch, N. de Botton, J.-L. Faure, C. Schuhl, G. Tamas, C. Tzara, E. Vincent, J. Deutsch, D. Favart, R. Prieels, B. Van Oystaeyen
Phys. Rev. C21, 662 (1980)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
80 Ar 9		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	RLY	151-159	C	153-159	ACT-I		4PI

We performed a measurement of the $\gamma + {}^{12}\text{C} \rightarrow \pi^+ + {}^{12}\text{B}_{gs}$ reaction up to 4 MeV above threshold relative to the $\gamma + p \rightarrow \pi^+ + n$ reaction. The resulting cross section is in agreement, at the accuracy level of $\pm 5\%$ with the available theoretical predictions and with another experimental result.

[NUCLEAR REACTIONS ${}^{12}\text{C}(\gamma, \pi^+){}^{12}\text{B}_{gs}$, E_{γ} : 0-4 MeV, measured σ relative to proton.]




FIG. 1. Experimental yields per nucleus versus the excess energy above threshold in the laboratory system. The curves are best fits to the data using Eqs. (1) and (2).

TABLE I. Photoproduction yield per nucleus for the proton and carbon targets at different values of the nominal bremsstrahlung endpoint energy E_e .

E_e (MeV)	A_p (a.u.)	E_e (MeV)	A_c (a.u.)
153.0	55.7 ± 3.7	153.5	0.68 ± 0.14
154.0	146.9 ± 6.0	156.0	1.41 ± 0.21
155.0	267.0 ± 9.7	156.5	2.45 ± 0.27
155.5	333.9 ± 11.3	157.0	3.76 ± 0.45
156.0	429.5 ± 10.3	158.0	9.19 ± 0.55
		158.5	11.47 ± 0.73

ELEM. SYM.	A	Z
C	12	6
METHOD		REF. NO.
		80 Co 6
		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	34-53	D	20-40	NAI-D		90

Cross section absolute scale obtained by normalization to the cross section of the 15.11 MeV (p,p', γ) transition in ^{12}C .




Fig. 2. - a) Differential cross-section for the $^{11}\text{B}(p, \gamma)^{12}\text{C}$ reaction at $\theta_{lab} = 90^\circ$. b) Differential cross-section for the $^{11}\text{B}(p, \gamma)^{12}\text{C}$ reaction at $\theta_{lab} = 90^\circ$. \bullet ref. (1), Δ ref. (2), \cdot present work.

REF. W.R. Dodge, E. Hayward, R.G. Leicht, B.H. Patrick, R. Starr
 Phys. Rev. Lett. 44, 1040 (1980)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
80 Do 1		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	23-39	D	23-59	NAI-I		DST

The elastic-photon-scattering cross section for ^{12}C has been measured at 90° and 135° in the energy range from 23.5 to 39 MeV. These data disagree with the predicted scattering, derived from the measured photonuclear absorption cross section, if only $E1$ transitions are assumed. To explain the difference in these cross sections, a large component of electric quadrupole absorption between 24 and 40 MeV is inferred.

PACS numbers: 23.20.Js, 25.20.+y, 27.20.+n, 24.30.Cz




FIG. 1. The photon scattering cross section at 135° . The errors on the experimental points are standard deviations based only on the number of counts. The dashed curve was calculated by using the total photonuclear absorption cross section (Ref. 1) in the optical theorem and the dispersion relation and assuming that only electric dipole absorption occurs according to Eq. (6). The solid curve labeled $E1$ shows the effect of applying a form factor to the Thomson scattering amplitude according to Eq. (9). The solid curve labeled $E1 + E2$ represents the best fit to the data assuming that part of the absorption cross section results from $E2$ excitations. The two solid curves cross near 25 MeV with the onset of the $E2$ strength. The introduction of $E2$ excitations results in destructive interference above 25 MeV and constructive interference below that energy, depending on the relative signs of the real parts of their scattering amplitudes.




FIG. 2. The total photonuclear absorption cross section and its $E2$ part. The upper solid curve is the published total photonuclear absorption cross section decreased by 0.12% of the total attenuation cross section and the dashed curves represent the upper and lower limits resulting from the systematic errors. The $E2$ cross section obtained from the best fit to the photon scattering data is also shown. The cross-hatched area represents 2 standard deviations about the best fit and corresponds to $2.7^{+2.9}_{-1.3}$ total energy-weighted $E2$ sums.

METHOD				REF. NO.			
				80 Ga 8		hg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, N	ABY	18-999	D	320-999	ACT-I		4PI
G, N	ABY	18-999	D	320-999	ACT-I		4PI

999=1.2 GEV

The reactions $^{12}\text{C} \rightarrow ^{11}\text{C}$ and $^{63}\text{Cu} \rightarrow ^{62}\text{Cu}$ have been studied by the induced-activity method at electron and photon energies from 0.32 to 1.2 GeV. Activation of the targets was carried out directly by an electron beam. The use of targets in the form of stacks permitted cross sections to be obtained for photodisintegration and electrodisintegration of the nuclei. Comparison of the experiment with theoretical calculations in the plane-wave approximation indicates a dominant role of E1 transitions of photons in these reactions.

PACS numbers: 25.20. + y, 25.30.Cg, 27.20. + n, 27.50. + e




FIG. 1. Yield and cross section of the reactions $^{12}\text{C}(j, n)^{11}\text{C}$, and also the ratio of these quantities, as functions of electron energy. The data presented were obtained with allowance for the E_0 -dependence of the radiator thickness in radiation lengths. Points: \circ —results of the present work and of Refs. 3 and 5, \bullet —Ref. 4, ∇ —Ref. 6, \square —Ref. 7, \swarrow —Ref. 3.

REF. H. Göringer, B. Schoch
Phys. Lett. 97B, 41 (1980)


ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
80 Go 7		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,NO	ABX	60	C	60	TOF - D		DST

$$\Delta E_n / E_n = 4\%$$

BREMS. TIP

Differential (γ, n_0) cross sections on ^{12}C and ^{16}O have been measured for photon energies $60 \text{ MeV} < E_\gamma < 160 \text{ MeV}$. These results combined with the corresponding (γ, p_0) cross sections support an absorption mechanism of the photon by neutron-proton pairs.



[13] J.L. Matthews, D.J.S. Findlay, S.N. Gardiner and R.O. Owens, Nucl. Phys. A267 (1976) 51.

Fig. 5. As fig. 4 for ^{12}C . Proton data: ref. [13].

ELEM. SYM.	A	Z
C	12	6

METHOD				REF. NO.		hg	
				80 Is 4			
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	ABX	15-32	C	32	NAI-D		90

Cross sections have been measured for elastic scattering of photons by the nuclei ^{12}C and ^{16}O in the region of the giant resonance. The energy resolution achieved in the experiment is commensurate with the energy resolution in the total photoabsorption cross section. It is shown that study of the cross sections for total absorption and elastic scattering of photons permits spectroscopic information to be obtained on the high-lying states of nuclei.

PACS numbers: 25.20. + y




FIG. 1. Cross section for elastic scattering of photons by ^{12}C obtained in the present work (a) and in Ref. 9 (b); cross section for total absorption of photons by ^{12}C from Ref. 10 (c).

TABLE II.

Energy of levels, MeV		J ^π , T	σ _{tot} , mb (Ref. 10)	σ _{el} , mb, present work	σ _{tot} /σ _{el} , b	σ _{el} , b
Ref. 10, total absorption	Present work					
^{12}C						
18.4	18.7±0.2	3 ⁻ , 1*	3	4±2	1±0.6	23.2
20.5	20.5±0.2	3 ⁻ , 1*	3	8±3	0.7±0.4	17.3
22-24	23-24	1 ⁻ , 1*	25	50±5	12.6±1	13.7
26.2	26.5±0.4	-	3.5	5±2	2.5±1	10.3
27.5	-	-	1.5	-	-	-
29.5	29.5±0.3	-	-	3±1	1.3±0.5	3.3
^{16}O						
17.1	-	1 ⁻ , 1**	3	-	-	-
19.5	19.5±0.2	1 ⁻ , 1**	3	10±3	0.9±0.4	19.1
21	-	1 ⁻ , 1**	6	-	-	-
22.3	22.0±0.2	1 ⁻ , 1**	30	65±10	10.7±4	14.4
24-25	24-25	1 ⁻ , 1**	30	85±10	10.7±4	12.1
27.2	27.2±0.2	2 ⁻ , 1**	3	10±3	0.9±0.5	10.3
-	28.1±0.2	-	-	-	-	-
-	31.0±0.3	-	-	-	-	-

**Data of the present work and also of Refs. 16 and 17 on the position of the maximum of the giant dipole resonance.

¹A. S. Penfold and E. L. Garwin, Phys. Rev. 116, 120 (1959).
²J. Ahrens *et al.*, Nucl. Phys. A251, 479 (1975).

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	80 Is 5 hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, p	ABX	16-31	C	21-31	TEL-D		90

A semiconductor counter telescope in a bremsstrahlung beam has been used to measure photoproton spectra from ^{12}C . The bremsstrahlung maximum energy took on the following values: 21.7, 23.4, 25.0, 26.6, 28.2, 29.6, and 31.0 MeV. The photoprotons were detected at an angle 90° . From the measured spectra we obtained the energy dependence of the cross sections for the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$ with formation of the final nucleus in the ground state and a number of excited states. In the region up to 30.5 MeV the integrated cross section for transitions to the ground state amounts to 73.7% of the total photoproton cross section.

SIG FOR GND, 1, 2 STATE

PACS numbers: 25.20. + y




FIG. 2. Proton spectra with correction for energy loss in the target (the spectra have been normalized to unit dose).




FIG. 4. Partial cross sections for the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$ at an angle 90° .




FIG. 3. Residual spectrum of protons for $E_\gamma^{\text{max}} = 31.0$ MeV with subtraction of the contributions of transitions to the ground state and the first three excited states of the ^{11}B nucleus (with energies 0.0, 2.12, and 4.44 + 5.02 MeV).

TABLE I. Relative yields (in percent) of partial channels of the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$.

E_γ^{max} , MeV	Data of Ref. 3			Relative errors, %	Data of the present work		
	E_γ^{max} , MeV				E_γ^{max} , MeV		
	21.7	25.0	31.0		21.7	25.0	31.0
7.20: $^{11}\text{B}^*$	0.5	0.5	0.5	± 30			
6.79: $^{11}\text{B}^*$	0.3	0.7	0.7	± 30			
6.71: $^{11}\text{B}^*$	0.4	0.3	0.3	± 30			
5.02: $^{11}\text{B}^*$	1.5	1.6	2.0	± 15			
4.44: $^{11}\text{B}^*$	0.7	0.8	1.4	± 15			
2.12: ^{11}B	8.0	8.0	5.4	± 10	17.2 \pm 0.5	19.0 \pm 0.6	20.8 \pm 0.6
0.0: ^{11}B	0.18	0.24	0.3		82.8 \pm 1.6	51.0 \pm 1.3	76.9 \pm 1.1

(OVER)

TABLE II. Integrated cross sections ($\int_{E_1}^{E_2} \sigma(E) dE$, MeV-mb/sr) of partial channels of the reaction $^{12}\text{C}(\gamma, p)^{11}\text{B}$.

E_p^0, MeV J^π	Data of Ref. 3			Data of the present work		
	$E_1 - E_2, \text{MeV}$			$E_1 - E_2, \text{MeV}$		
	Threshold -4.5	24.5-27.0	27.0-33.0	Threshold -4.5	24.5-27.0	27.0-33.0
3.02; $1/2^-$	0.1	0	0.1			} 0.4
4.44; $1/2^-$	0.06	0	0.15			
2.12; $1/2^-$	0.2(5)	0.05	0.2	1.1	0.3	
Ground state $3/2^-$	5.0	1.2	1.8	4.3	1.3	1.2
All excited states	0.4	0.2(5)	0.6	1.1	0.3	1.4

REF. A.F. Khodyachik, P.I. Vatset, I.V. Dogyust, V.A. Zolenko,
V.V. Kirichenko
Sov. J. Nucl. Phys. 32, 453 (1980)

ELEM. SYM.	A	Z
C	12	6

METHOD					REF. NO.		
					80 Kh 2	hg	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, NP	ABX	34-150	C	UKN	CCH-D		4PT

A diffusion chamber in a magnetic field has been used to measure the cross section for the reaction $^{12}\text{C} + \gamma \rightarrow p + n + ^{10}\text{B}$ as a function of γ -ray energy, and also the distribution in relative energy of np pairs in the intermediate energy region. It is concluded that the main mechanism of the reaction is the interaction of the γ ray with a correlated np pair from the p shell of the carbon nucleus.

NP PAIRS

PACS numbers: 25.20. + γ , 27.20. + n




FIG. 2. Total cross section as a function of energy. The hollow circles are from the present experiment, and the curve from Ref. 1.

TABLE II. Dependence of total cross section on energy.

E_γ , MeV	$\sigma \cdot 10^{10}$, cm ²	$\sigma \cdot 10^{10}$, cm ² *	E_γ , MeV	$\sigma \cdot 10^{10}$, cm ²	$\sigma \cdot 10^{10}$, cm ² *
34-36	0.33±0.05	1.5±0.5	63-68	0.30±0.04	-
36-38	0.30±0.06	0.9±0.3	68-73	0.29±0.04	0.7±0.4
38-40	0.62±0.07	0.7±0.2	73-78	0.35±0.04	0.3±0.15
40-42	0.65±0.08	0.60±0.2	78-83	0.15±0.03	0.3±0.15
42-44	0.60±0.06	0.45±0.2	83-88	0.21±0.03	-
44-46	0.53±0.06	0.40±0.2	88-98	0.18±0.03	-
46-48	0.52±0.07	0.7±0.2	98-108	0.18±0.03	0.23±0.15
48-51	0.4±0.05	0.6±0.2	108-118	0.116±0.02	0.10±0.05
51-54	0.28±0.05	0.85±0.3	118-128	0.089±0.02	0.25±0.12
54-57	0.32±0.05	0.4±0.2	128-138	0.083±0.019	0.15±0.07
57-60	0.27±0.04	0.3±0.1	138-148	0.073±0.02	0.20±0.10
60-63	0.29±0.04	0.25±0.1			

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
80 Mi 1		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,PI+	ABX	4 (4.5)	D	200	MAG-D		DST

Strong spin-isospin mode, $T=1, T_z=1$ analog states at 4.5 MeV in ^{12}B were studied by the reaction $^{12}\text{C}(e, e'\pi^+)^{12}\text{B}$ at the electron energy $E_e = 200$ MeV. The photoproduction cross sections at seven angles ranging from 30° to 150° were obtained with use of virtual-photon theory and an experimentally determined real-to-virtual photon ratio. The results are compared with theoretical calculations and also with the available data on inelastic scattering from the 19.5-MeV $T=1, T_z=0$ analog complex in ^{12}C .

VIRT PHOT ANAL

PACS numbers: 24.30.Cz, 25.30.Cg, 27.20.+n




FIG. 1. The pion energy spectra from ^{12}C at $\theta = 50^\circ$ and 120° . The solid curves are least-squares fits to the data using virtual-photon spectrum shapes (Ref. 9), with the lower lines representing the background due to transitions to the ground and 0.95-MeV states in ^{12}B .




FIG. 2. Photopion cross sections compared with the shell-model calculations using the $\vec{\sigma} \cdot \vec{\epsilon}$ production amplitude (Ref. 10).

(OVER)




FIG. 3. Photopion cross sections compared with the shell-model calculations (Ref. 11) using the full Blomqvist-Laget amplitude (Ref. 12), and also with the Helm-model calculations (Ref. 13).

METHOD

REF. NO.

80 Sh 9

hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E,PI+	ABX	150-195	D	195	MAG-D		DST

Abstract: Energy distributions of π^- produced from ^{12}C by electrons of total energy 195 MeV were measured at various angles. The results show large contributions from transitions leaving the residual nucleus in the ground (1^+), first (2^-) excited state and states at around 4.5 MeV. The angular distributions of $^{12}\text{C}(\gamma, \pi^-)^{12}\text{B}$ leading to these residual states are deduced from the energy distributions by the unfolding method with the virtual photon theory. Theoretical results with the Helm model and the shell model are compared with the experimental results. Their relative shapes are in good agreement. A better agreement in the absolute value is found for the theoretical results which include the final-state interaction estimated with a pion optical potential. The surface production model shows better agreement with the experimental (γ, π^-) cross sections than the volume production model.

VIRT PHOTON ANALYSIS

E NUCLEAR REACTIONS $^{12}\text{C}(\gamma, \pi^-)$, $E = 195$ MeV; measured $\sigma(E_\gamma, \theta)$; ^{12}B deduced levels, transition strength. Natural target.




Fig. 10. Angular distributions of $^{12}\text{C}(\gamma, \pi^-)^{12}\text{B}$. Open circles: present result for the residual ground state; closed circles: that for the first excited residual state; crosses: for the residual states at around 4.5 MeV.

34) M.I. Adamovich, V.G. Lurionova, A.I. Lebedev, S.P. Khartamov and F.R. Yaegudina, in photomeson and photonuclear processes, ed. D.V. Skobel'tsyn (Consultants Bureau, NY 1967) p. 49

(OVER)




Fig. 11. Comparison of cross sections of the $H(\gamma, \pi^-)$ reaction. Open circle: present result; closed circles: previous result²⁴.




Fig. 16. Comparison of angular distributions for $^{12}\text{C}(\gamma, \pi^-)^{12}\text{B}$ leaving the residual states at around 4.5 MeV. Open circles: present result; broken curves: theoretical estimates with a generalized Heilm model for the residual states 4.5 MeV (2^-) and 5.0 MeV (4^-) as indicated^{23,24}; solid curve: sum of the above estimates.




Fig. 17. Comparison between the photon spectrum in radiative π^- capture of ^{12}C [ref. 15] and the transition strengths of the $^{12}\text{C}(\gamma, \pi^-)^{12}\text{B}$ reaction. The strengths are normalized at $E_1 = 0$.

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
81 A1 8		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABY	16-999	C	999	TEL-D		DST

Abstract: The angular dependences of proton photoproduction from the nuclei ^{12}C , ^{63}Cu and ^{208}Pb irradiated by bremsstrahlung γ -quanta with maximum energy 4.5 GeV, both in the cumulative region (i.e. in the kinematical region in which the production of protons in the collision of γ -quanta of the given energy with the quasi-free nuclear nucleon is forbidden) and in the non-cumulative region, are investigated. The experimental data obtained are compared with the results of theoretical calculations of cumulative proton photoproduction according to the following models: the "quasi-two-body" scaling model, the low-nucleon correlation model, the fluctuon model and the cluster model.

999=4.5 GEV

E NUCLEAR REACTIONS ^{12}C , ^{63}Cu , $^{208}\text{Pb}(\gamma, p)$, $E = 4.5$ GeV bremsstrahlung; measured $\sigma(E_p, \theta_p)$; deduced reaction mechanism. Natural target.




Fig. 3. The invariant cross section of the reaction $\gamma - A - p - X$ as a function of the detection angle of protons on the nucleus ^{12}C irradiated with bremsstrahlung γ -quanta with the maximum energy 4.5 GeV. \bullet - the proton energy 80 MeV, \blacktriangle - 100 MeV, \blacksquare - 136 MeV, \circ - 180 MeV, \triangle - 210 MeV, \square - 290 MeV. Statistical errors do not exceed the symbol sizes. The lines are drawn through experimental points by eye.

REF. K.V. Alanakyan, M.Dzh. Amaryan, R.A. Demirchyan, K.Sh. Egiyan,
Dzh.V. Karumyan, Zh.L. Kocharova, M.S. Ogandzhanyan, Yu.G. Sharabyan
Yad. Fiz. 34, 1494 (1981)
Sov. J. Nucl. Phys. 34, 828 (1981)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	81 A1 14

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PX	ABY	130*4	C	*4	MAG-D		DST

Experimental energy and angular distributions are presented for inclusive protons emitted from the ^{12}C nucleus bombarded by bremsstrahlung with maximum energy 4.5 GeV. The protons were detected in the range of angles $20\text{--}120^\circ$ and momenta $0.4\text{--}1.3\text{ GeV}/c$. It is shown that the energy distributions of photoprotons in the case of cumulative production are satisfactorily described by an exponential, while for noncumulative protons the spectrum is flattened out in the high-energy region. The experimental data obtained are analyzed from the point of view of a number of theoretical models proposed for description of cumulative particle production.

*4=4.5 GeV

PACS numbers: 25.20. - y

TABLE I. Invariant yield of protons in the reaction $\gamma^{12}\text{C} \rightarrow pX$ at $E_{\gamma, \text{max}} = 4.5\text{ GeV}$ in units of $\mu\text{b} \cdot \text{c}^3/\text{GeV}^2 \cdot \text{sr} \cdot \Omega$.

T, GeV	θ_p, deg				
	20	40	60	90	120
0.223	4040 ± 130	2920 ± 60	1320 ± 30	408 ± 13	155 ± 4
0.317	2650 ± 70	1110 ± 30	280 ± 10	87.0 ± 3.0	48.6 ± 0.7
0.425	1340 ± 40	444 ± 14	118 ± 3	11.4 ± 0.6	1.66 ± 0.17
0.530	704 ± 23	222 ± 7	33.8 ± 1.4	1.36 ± 0.23	0.173 ± 0.028
0.62	483 ± 14	93.7 ± 3.0	11.2 ± 0.6	0.247 ± 0.051	-




FIG. 2. Proton energy spectra for various values of the angle θ_p (\bullet, \circ —for $\theta_p = 20^\circ$; ∇, ∇ — 40° ; \blacksquare, \square — 60° ; $\blacktriangle, \triangle$ — 90° ; \bullet, \circ — 120° ; \circ — 160°). The solid points are the experimental data of the present work for the region $0.7 \leq p_p \leq 1.3\text{ GeV}/c$, and the hollow points are the data of Ref. 15 for $p_p \leq 0.9\text{ GeV}/c$; $E_{\gamma, \text{max}} = 4.5\text{ GeV}$.




FIG. 7. Angular distributions of photoprotons for various values of their momenta: \blacklozenge —proton momentum $p_p = 0.4\text{ GeV}/c$; \bullet — $0.44\text{ GeV}/c$; \blacksquare — $0.52\text{ GeV}/c$; \blacktriangle — $0.608\text{ GeV}/c$; \square — $0.66\text{ GeV}/c$; \circ — $0.69\text{ GeV}/c$; ∇ — $0.79\text{ GeV}/c$; ∇ — $0.84\text{ GeV}/c$; \square — $0.98\text{ GeV}/c$; \blacktriangle — $1.13\text{ GeV}/c$; \circ — $1.25\text{ GeV}/c$. The hollow points show the experimental data of the present work for the region $0.7 \leq p_p \leq 1.3\text{ GeV}/c$. The remaining points are the data of Ref. 15 for $p_p \leq 0.9\text{ GeV}/c$. The arrows indicate the boundaries between the cumulative and noncumulative regions. The solid curves are the results of calculations according to the cluster model (a), the FNC model (b), and the two-particle scaling model (c). The solid and dashed curves (see the text) marked by identical numbers correspond to identical momentum values p_p .

REF. J. Arends, J. Eyink, A. Hegerath, K.G. Hilger, B. Mecking,
G. Nöldeke, H. Rost
Phys. Lett. 98B, 423 (1981)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
81 Ar 1		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, MU-T	ABX	215-386	D	215-386	TOF-D		4PI

DATA ALSO IN 81AR3

Double differential cross sections for the photo-emission of protons and charged pion production were investigated for a number of target nuclei (He, Be, C, O, Al, Ti, Cu, Sn, Pb) in the photon energy range $k = (215-386)$ MeV. On the basis of these experimental results the total hadronic cross section was determined.




Fig. 7. Parameter σ_0 as a function of photon energy (data points) compared to the mean cross section for a free nucleon (solid line).

The total hadronic cross sections for all measured elements can be parametrized in the form

$$\sigma(k, A) = \sigma_0(k) \cdot A^x,$$

A being the atomic number, with a constant exponent $x = 1.1$. The photon energy dependence of σ_0 is shown in fig. 7. Compared to the mean cross section for a free nucleon (the solid line in fig. 7) the excitation of the Δ -resonance is suppressed. Such a suppression is expected in the Δ -hole model [11].

REF. J. Arends, J. Eyink, A. Hegerath, K.G. Hilger, B. Mecking,
G. Nöldeke, H. Rost
Nucl. Phys. A358, 367c (1981)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
81 Ar 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,MU-T	ABX	215-386	D	215-386	TOF-D		4PI

Abstract: Double differential cross sections for the photoemission of protons and charged pion photoproduction were investigated for a number of target nuclei (He, Be, C, O, Al, Ti, Cu, Sn, Pb) using the tagged bremsstrahlung beam at the Bonn 500 MeV-Synchrotron in the photon range $k = (215-386)$ MeV. On the basis of these experimental results the total hadronic cross section was determined.




Fig. 2. Cross section for the process: $\gamma + \text{Pb} \rightarrow p + X$. The proton threshold is 53 MeV.




Fig. 3. Total hadronic cross section for Be. The data are compared to the cross section taken from ref. 2).

The photon energy dependence of the total cross sections for heavier nuclei are similar to the Be results. The complete data set can be parametrized in the form

$$\sigma(k, A) = \sigma_0(k) \cdot A^x.$$

The exponent is constant $x = 1.1$. The photon energy dependence of σ_0 is shown in fig. 4. Compared to the mean cross section for a free nucleon, the excitation of the Δ -resonance is suppressed.




Fig. 4. Parameter σ_0 compared to the cross section for a free nucleon (full line).

METHOD	REF. NO.
	81 Av 10
	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
S G, XP	RLX	0*2	D	0*2	TEL-D		100

We report the results of a study of the reaction $\gamma A \rightarrow pX$ at an angle $\theta_{\gamma} = 100^\circ$ lab in a beam of quasimonochromatic polarized photons. The measurements were made for three values of photon energy ($E_{\gamma} = 0.69, 1.40, \text{ and } 1.95 \text{ GeV}$) in the nuclei ^{12}C , ^{64}Cu , and ^{207}Pb . The range of kinetic energies of the protons was $\approx 100\text{--}230 \text{ MeV}$. It is shown that the slope parameter B for the invariant cross section $f = C \exp(-Bp^2)$ is a weak function of A and does not depend on E_{γ} , but the parameter $C_{\gamma} = C/A\bar{\sigma}$, increases with increase of E_{γ} , the slope of the lines $C_{\gamma}(E_{\gamma})$ being greater for larger A .

COH-BRMS .69*1.95 GEV

PACS numbers: 25.20.+y, 13.60.Rj




FIG. 2. Invariant cross sections f for photoproduction of cumulative protons as a function of their momentum squared for three target nuclei (^{12}C —lower family of points, ^{64}Cu —middle family of points, ^{207}Pb —upper family of points). The solid, hollow, and combined points correspond respectively to the values $E_{\gamma} = 0.69, 1.40 \text{ and } 1.95 \text{ GeV}$. The curves are described in the text.




FIG. 3. The dependence of the parameter C_{γ} on E_{γ} . Points: \circ — ^{12}C , \square — ^{64}Cu , \triangle — ^{207}Pb .

TABLE II.

$E_{\gamma}, \text{ GeV}$	Parameter	Nucleus		
		^{12}C	^{64}Cu	^{207}Pb
0.69	B	9.50 ± 0.51	11.01 ± 0.32	10.19 ± 0.38
	C	$71.9^{+11.9}_{-11.7}$	663^{+61}_{-55}	3252^{+312}_{-302}
1.40	B	10.45 ± 0.31	11.21 ± 0.22	9.63 ± 0.45
	C	$105.4^{+7.3}_{-9.0}$	973^{+61}_{-58}	3358^{+305}_{-317}
1.95	B	10.36 ± 0.49	11.14 ± 0.37	10.07 ± 0.31
	C	$138.0^{+29.6}_{-18.0}$	1317^{+150}_{-116}	8201^{+342}_{-330}

Note. The parameter B is given in units of $(\text{GeV}/c)^{-2}$, while C is given in relative units.

TABLE I. Invariant cross section f (in relative units)

Nucleus	$E_{\gamma}, \text{ GeV}$	$T_p, \text{ MeV}$						
		116	135	153	173	190	210	229
^{12}C	0.69	7.30 ± 0.13	4.80 ± 0.10	3.40 ± 0.08	1.88 ± 0.05	1.55 ± 0.05	1.05 ± 0.01	0.74 ± 0.03
	1.40	9.18 ± 0.34	8.04 ± 0.30	4.04 ± 0.24	2.89 ± 0.19	1.68 ± 0.16	1.07 ± 0.12	0.45 ± 0.04
	1.95	13.12 ± 1.00	7.24 ± 0.71	4.91 ± 0.58	3.02 ± 0.46	2.34 ± 0.38	1.43 ± 0.20	1.00 ± 0.22
^{64}Cu	0.69	54.73 ± 0.39	39.66 ± 0.68	28.10 ± 0.55	15.97 ± 0.43	11.54 ± 0.33	8.49 ± 0.25	4.59 ± 0.20
	1.40	69.54 ± 2.32	59.45 ± 2.14	37.70 ± 1.73	24.12 ± 1.41	14.23 ± 1.05	8.37 ± 0.60	5.21 ± 0.61
	1.95	125.0 ± 6.65	77.9 ± 5.53	48.82 ± 4.11	31.61 ± 3.30	21.32 ± 2.44	13.42 ± 1.57	8.37 ± 1.42
^{207}Pb	0.69	392.7 ± 3.3	242.1 ± 3.4	172.5 ± 4.8	119.2 ± 3.7	85.3 ± 2.6	51.4 ± 2.2	31.2 ± 1.2
	1.40	529.3 ± 13.2	365.3 ± 17.3	253.2 ± 13.2	147.2 ± 9.9	106.2 ± 8.7	76.8 ± 7.0	42.8 ± 5.0
	1.95	734.0 ± 68.0	525.0 ± 50.8	333.6 ± 38.7	213.4 ± 29.3	137.4 ± 21.1	108.5 ± 16.9	62.4 ± 12.5

REF. R.O. Avakyan, A.É. Avetisyan, N.Z. Akopov, S.S. Danagulyan, I.Kh. Kosakov, A.A. Oganessian, Zh.V. Petrosyan, S.P. Taroyan, G.M. Élbakyan
Sov. J. Nucl. Phys. 33, 448 (1981)
Yad. Fiz. 33, 858 (1981)

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
81 Av 13		

METHOD

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
\$ G,XP	ASM	0*2	C	0*2	UKN		100

COH-BRMS .69*1.95 GEV

At the present time it is rather well established that the experimental values of the invariant cross section $f = (E/p^2)(d^2\sigma/d\Omega dp)$ of the reaction

$$aA \rightarrow bX \quad (1)$$

in the cumulative region^{1,2} are described by an exponential dependence of the form $f = C \exp(-Bp^2)$. Most of the experiments in which reaction (1) induced by various particles (π, p, γ, \dots), has been studied were designed to study the energy, angular, and A dependence of the parameters B and C .³⁻⁹ As a result of the investigations it has been established that the parameter B does not depend on the mass number A of the target nucleus, on the type of incident particle, or on its energy, beginning with $E_a \approx 1$ GeV, while the parameter $C_p = C/\sigma_{tot}$ (σ_{tot} is the total cross section for the aA interaction) does not depend on the type of particle a . In addition to the established properties of the quantities B and C it would be interesting to check the dependence of the parameters B and C on the direction of polarization of the initial particle. For this purpose it is necessary to measure the asymmetry Σ of the cross section for reaction (1) as a function of the direction of the initial-particle polarization vector.

In the present work we report the results of a study of the photoproduction of cumulative protons at an angle $\vartheta_{\gamma p} = 100^\circ$ in the laboratory system in the nuclei ^{12}C , ^{64}Cu , and ^{207}Pb for three photon energy values ($E_\gamma = 0.69, 1.40, \text{ and } 1.95$ GeV). The possibility of measurement at a definite photon energy was based on the use of the method of subtraction of the coherent peak^{10,11} in the spectrum of quasimonochromatic polarized photons emitted by electrons in passing through a diamond crystal.¹² The existence of a significant degree of polarization of the photons in the coherent

peak has enabled us to measure the value of the cross-section asymmetry Σ of the reaction $\gamma A \rightarrow pX$. The asymmetry was calculated from the relation

$$\Sigma = \frac{1}{\bar{P}_\gamma} \frac{y^+ - y^-}{y^+ + y^- - 2y^*}$$

where y^+, y^- are the reaction yields in the case of perpendicular and parallel orientation of the photon polarization vector with respect to the reaction plane in the coherent bremsstrahlung spectrum; y^* is the reaction yield for an ordinary bremsstrahlung spectrum; \bar{P}_γ is the average value of photon polarization in the subtracted coherent peak.

Measurements of Σ were made in the nuclei ^{12}C , ^{64}Cu , and ^{207}Pb for protons with kinetic energy respectively $T_p = 173, 164, \text{ and } 163$ MeV. The energy bin was $\Delta E = 60$ MeV.

The experimental apparatus and measurement technique have been described in detail elsewhere.¹³

Numerical values of Σ with their standard deviations $\sigma(\Sigma)$ are given in the table.

The values of $\sigma(\Sigma)$ contain both the statistical error and the error in determination of the quantity \bar{P}_γ .¹³

From the figure, where we have shown Σ as a function of E_γ for the three nuclei it can be seen that the absolute values of the asymmetry in the region investigated are insignificant and depend weakly on E_γ . We note that Σ for carbon is close to zero for all E_γ , and the maximum value 0.29 ± 0.16 is achieved in the case of lead for $E_\gamma = 1.95$ GeV. The data show that within experimental error the asymmetry is almost indepen-

TABLE I.

$E_\gamma, \text{ GeV}$	Nucleus					
	^{12}C		^{64}Cu		^{207}Pb	
	Σ	$\sigma(\Sigma)$	Σ	$\sigma(\Sigma)$	Σ	$\sigma(\Sigma)$
0.69	-0.0784	± 0.0233	-0.1533	± 0.0351	-0.1192	± 0.0323
1.40	0.0402	± 0.0258	0.0831	± 0.0309	-0.0798	± 0.0310
1.95	0.0580	± 0.0235	0.1526	± 0.0331	0.2931	± 0.1633




FIG. 1. Asymmetry Σ of the cross section for photoproduction of cumulative protons as a function of the photon energy E_γ for target nuclei ^{12}C (○), ^{64}Cu (□), and ^{207}Pb (△).

ELEM. SYM.	A	Z
C	12	6
REF. NO.		hg
81 Be 11		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
\$ G, XP	SPC	0*2	D	0*2	TEL-D		100

We report the results of a study of the reaction $\gamma A \rightarrow pX$ at an angle $\theta_{\gamma} = 100^\circ$ lab in a beam of quasimonochromatic polarized photons. The measurements were made for three values of photon energy ($E_{\gamma} = 0.69, 1.40, \text{ and } 1.95 \text{ GeV}$) in the nuclei ^{12}C , ^{64}Cu , and ^{207}Pb . The range of kinetic energies of the protons was $\approx 100\text{--}230 \text{ MeV}$. It is shown that the slope parameter B for the invariant cross section $f = C \exp(-Bp^2)$ is a weak function of A and does not depend on E_{γ} , but the parameter $C_{\gamma} = C/A\bar{\sigma}$, increases with increase of E_{γ} , the slope of the lines $C_{\gamma}(E_{\gamma})$ being greater for larger A .

COH-BRMS .69*1.95GE

PACS numbers: 25.20. + y, 13.60.Rj




FIG. 2. Invariant cross section f for photoproduction of cumulative protons as a function of their momentum squared for three target nuclei (^{12}C —lower family of points, ^{64}Cu —middle family of points, ^{207}Pb —upper family of points). The solid, hollow, and combined points correspond respectively to the values $E_{\gamma} = 0.69, 1.40$ and 1.95 GeV . The curves are described in the text.




FIG. 3. The dependence of the parameter C_{γ} on E_{γ} . Points: \circ — ^{12}C , \square — ^{64}Cu , \triangle — ^{207}Pb .

TABLE II.

$E_{\gamma}, \text{ GeV}$	Parameter	Nucleus		
		C	^{64}Cu	^{207}Pb
0.69	B	3.50 ± 0.41	11.01 ± 1.02	10.17 ± 1.76
	C	71.9 ± 11.7	563 ± 63	3252 ± 312
1.40	B	10.10 ± 0.31	11.21 ± 0.22	9.05 ± 0.47
	C	105.4 ± 4.3	973 ± 26	3853 ± 217
1.95	B	10.50 ± 0.40	11.14 ± 0.37	13.07 ± 0.51
	C	138.0 ± 15.6	1317 ± 116	6201 ± 213

Note. The parameter B is given in units of $(\text{GeV}/c)^{-2}$, while C is given in relative units.

TABLE I. Invariant cross section f (in relative units)

Nucleus	$E_{\gamma}, \text{ GeV}$	$T_p, \text{ MeV}$						
		116	125	133	173	190	210	221
^{12}C	0.69	7.35 ± 0.13	4.80 ± 0.10	3.40 ± 0.08	1.89 ± 0.05	1.53 ± 0.05	1.05 ± 0.04	0.53 ± 0.03
	1.40	9.18 ± 0.34	6.04 ± 0.30	4.04 ± 0.25	2.60 ± 0.19	1.83 ± 0.16	1.07 ± 0.12	0.53 ± 0.04
	1.95	13.12 ± 1.00	7.24 ± 0.71	4.91 ± 0.53	3.02 ± 0.46	2.01 ± 0.38	1.18 ± 0.21	1.00 ± 0.22
^{64}Cu	0.69	61.73 ± 0.89	39.86 ± 0.66	28.10 ± 0.55	15.97 ± 0.43	11.74 ± 0.33	6.49 ± 0.25	4.59 ± 0.20
	1.40	62.71 ± 2.02	39.40 ± 2.14	37.70 ± 1.79	21.10 ± 1.11	14.33 ± 1.05	9.37 ± 0.60	5.21 ± 0.61
	1.95	125.0 ± 6.65	77.14 ± 5.53	46.82 ± 4.11	31.81 ± 3.50	21.82 ± 2.41	13.42 ± 1.37	8.37 ± 1.42
^{207}Pb	0.69	392.7 ± 6.3	272.1 ± 5.4	172.5 ± 4.6	119.2 ± 3.7	66.8 ± 2.6	31.4 ± 2.2	31.2 ± 1.2
	1.40	513.0 ± 11.2	263.8 ± 17.3	228.2 ± 15.1	147.2 ± 9.9	106.2 ± 8.7	75 ± 7.0	42.8 ± 5.6
	1.95	734.0 ± 63.0	525.0 ± 59.6	358.0 ± 36.7	213.4 ± 29.3	137.4 ± 21.1	108.5 ± 16.9	62.4 ± 12.5

REF. R. Leicht, K.P. Schelhaas, M. Hammen, J. Ahrens, B. Ziegler
Nucl. Inst. Meth. 179, 131 (1981)

ELEM. SYM.	A	Z
C	12	6

METHOD					REF. NO.		
					81 Le 5		egf
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G _s G	RLX	15	D	15	NAI-D		DST

A facility for the measurement of absolute cross sections and angular distributions of photons scattered elastically from nuclei has been set up in the Mainz Linac Laboratory. Quasimonoenergetic photons of 8–100 MeV are produced by the method of the positron annihilation in flight. The scattered photons can simultaneously be detected in four NaI(Tl) spectrometers, which are installed at different scattering angles. The performance of the system and test measurements are discussed.

15=15.11 MEV




Fig. 11. Measured angular distribution of photons scattered from the 15.11 MeV level in ¹²C (points with error bars) in comparison with the dipole angular distribution (full line). The scattering cross section has been adjusted at 90° to be 0.5.

METHOD						REF. NO.		hg
						81 Lo 6		
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE	
			TYPE	RANGE	TYPE	RANGE		
E,p	ABX	16-200	D	200	MAG-D		DST	

Double differential cross sections at six angles ranging from 45° to 143° have been measured for the ¹²C(e,p)e' reaction. The proton energy ranged from 15.6 to 17.2 MeV at an incident electron energy of 200 MeV. At the backward angles our results are in good agreement with data reported by Vysotskaya and Afanas'ev but for forward angles the results are lower.

On a mesuré les sections efficaces différentielles doubles, à six angles allant de 45° à 143°, pour la réaction ¹²C(e,p)e'. L'énergie des protons variait de 15,6 à 17,2 MeV, pour une énergie des électrons incidents de 200 MeV. Pour les angles vers l'arrière nos résultats sont en bon accord avec les données rapportées par Vysotskaya et Afanas'ev, mais ils sont plus bas dans le cas des angles vers l'avant.

Can. J. Phys., 59, 271 (1981) [Traduit par le journal]




FIG. 2. Angular distributions of 15.6 (●), 16.2 (+), 16.7 (○), and 17.2 (■) MeV protons. The data of Vysotskaya and Afanas'ev (×, ref. 3) are also plotted for comparison purposes. The error bars reflect the statistical uncertainties only.

TABLE 2. Double differential cross sections for ¹²C(e,p)e' in μb/(MeV sr)*

E _p (MeV)	θ _p = 45°	θ _p = 65°	θ _p = 90°	θ _p = 105°	θ _p = 120°	θ _p = 143°
15.6	0.219 ± 0.002	0.290 ± 0.003	0.214 ± 0.002	0.199 ± 0.002	0.150 ± 0.001	0.108 ± 0.001
16.2	0.226 ± 0.002	0.273 ± 0.003	0.227 ± 0.002	0.183 ± 0.002	0.135 ± 0.001	0.098 ± 0.001
16.7	0.215 ± 0.002	0.258 ± 0.003	0.211 ± 0.002	0.168 ± 0.002	0.121 ± 0.001	0.089 ± 0.001
17.2	0.199 ± 0.002	0.233 ± 0.002	0.194 ± 0.002	0.154 ± 0.002	0.108 ± 0.001	0.083 ± 0.001

* The uncertainties reflect the statistics only.

REF. R.M. Sealock, H.S. Caplan, G.J. Lolos, W.C. Haxton
 Phys. Rev. C23, 1293 (1981)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	hg
81 Se 2	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E, \pi^+ +$	ABX	6*13	D	200	MAG-D		DST

Angular distributions for near threshold electroproduction of positive pions from ^{12}C are reported and compared to theory. These data provide tests of recently developed second-order optical potentials and of the nuclear response to electroproduction in the giant resonance region.

*PION ENERGY IN MEV

NUCLEAR REACTIONS $^{12}\text{C}(e, \pi^+ e')$; measured $\sigma(E_\pi, \theta)$, $E_e = 200$ MeV.
 $E_\pi = 6-13$ MeV; calculated σ , distorted wave Born approximation; giant resonances.




FIG. 1. Double differential cross sections for $^{12}\text{C}(e, \pi^+ e')$ as a function of θ_π for 6, 9, and 13 MeV pions. Theoretical results are given for transitions to $0h\omega$ final states in ^{12}B (dashed line), to $0h\omega$ and $1h\omega$ states (dash-dotted line), and to all states kinematically allowed (solid line).




FIG. 2. Double differential cross sections for $^{12}\text{C}(e, \pi^+ e')$ at $\theta_\pi = 45^\circ, 90^\circ,$ and 143° as a function of pion energy. The 50° and 90° data of Shoda (Ref. 13) are shown for comparison. The theoretical curves are as in Fig. 1.

(OVER)

TABLE I. Double differential cross sections for $^{12}\text{C}(e, \pi^+ e')$ (nb/MeV sr).

E_{ν} (MeV) \ θ_{ν}	45°	65°	90°	105°	120°	143°
6.0	0.304 ± 0.023	0.324 ± 0.035	0.300 ± 0.024		0.278 ± 0.035	0.278 ± 0.023
7.0	0.324 ± 0.021	0.345 ± 0.032	0.329 ± 0.033	0.268 ± 0.032	0.310 ± 0.021	0.297 ± 0.021
8.0	0.336 ± 0.020	0.361 ± 0.040	0.340 ± 0.021	0.334 ± 0.020	0.295 ± 0.020	0.304 ± 0.020
9.0	0.347 ± 0.020	0.376 ± 0.030	0.353 ± 0.030	0.339 ± 0.020	0.284 ± 0.020	0.305 ± 0.020
10.0	0.353 ± 0.020	0.382 ± 0.029	0.357 ± 0.030	0.328 ± 0.020	0.278 ± 0.020	0.295 ± 0.020
11.0	0.356 ± 0.029	0.384 ± 0.029	0.351 ± 0.029	0.313 ± 0.019	0.273 ± 0.024	0.283 ± 0.019
12.0	0.358 ± 0.029	0.384 ± 0.029	0.343 ± 0.029	0.297 ± 0.019	0.269 ± 0.029	0.275 ± 0.019
13.0	0.359 ± 0.028	0.381 ± 0.028	0.331 ± 0.029	0.281 ± 0.028	0.265 ± 0.038	0.266 ± 0.019

REF. M. Bernheim, A. Bussièrè, J. Mougey, D. Royer, D. Tarnowski,
S. Turck-Chieze, S. Frullani, S. Boffi, C. Giusti, F.D. Pacati,
G.P. Capitani, E. De Sanctis, G.J. Wagner
Nucl. Phys. A375, 381 (1982)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	82 Be 3
	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/P	NOX	312*572	D.	500	MAG-D		DST

Abstract: The measurement of recoiling nucleus momentum distributions in (e,e'p) reactions has been performed for p-hole states in ¹²C and ¹⁶O under extended kinematical conditions. The analysis of the experimental data has been performed without requiring the usual factorization of the cross section. The sensitivity of the data and of the analysis for deducing bound and scattering proton states is discussed.

DLTQ MEV/C

E NUCLEAR REACTIONS ¹²C, ¹⁶O(e,e'p), E = 500 MeV; measured σ (missing energy, recoil momentum); deduced proton-hole spectral functions. Unfactorized DWIA analysis.

TABLE 7

Experimental values of the momentum distributions (without corrections for radiative effects) for ¹²C(e, e'p)¹¹B in perpendicular (a) and parallel (b) kinematics

p_B (MeV/c)	Perpendicular kinematics $S(p)(\text{GeV}/c)^{-3} = \Delta S(p)$		Parallel kinematics $S(p)(\text{GeV}/c)^{-3} = \Delta S(p)$	
	$p_B < 0$	$p_B > 0$	$p_B < 0$	$p_B > 0$
	15	7.3 ± 2.5	11.0 ± 3.4	13.2 ± 1.5
25	11.7 ± 2.6	14.1 ± 3.8	16.2 ± 1.6	8.6 ± 1.0
35	22.2 ± 3.9	20.6 ± 5.4	25.0 ± 2.1	11.3 ± 1.1
45	29.0 ± 3.0	29.0 ± 3.1	31.3 ± 2.4	16.8 ± 1.5
55	36.3 ± 3.2	30.7 ± 3.0	35.7 ± 2.7	22.3 ± 2.0
65	44.5 ± 3.9	36.8 ± 4.0	38.6 ± 3.0	27.5 ± 2.7
75	50.8 ± 4.8	43.6 ± 6.1	41.0 ± 3.2	34.0 ± 4.0
85	65.2 ± 6.5		50.2 ± 4.3	44.1 ± 4.8
95	54.8 ± 4.5	50.1 ± 4.3	51.4 ± 4.2	47.8 ± 4.8
105	56.0 ± 4.1	48.1 ± 4.0	47.4 ± 3.8	46.0 ± 4.2
115	48.3 ± 3.8	48.4 ± 4.4	41.8 ± 3.3	48.5 ± 4.1
125		44.8 ± 5.1	36.6 ± 2.9	52.8 ± 5.5
135			31.4 ± 2.4	53.6 ± 4.6
145	38.9 ± 3.0	29.2 ± 2.3	24.8 ± 2.0	45.6 ± 4.2
155	33.5 ± 2.5	26.1 ± 2.1	21.3 ± 1.9	
165	31.8 ± 2.4			

ELEM. SYM.	A	Z
C	12	6
REF. NO.		
82 Co 2		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	21-30	D	5-14	NAI-D		90

We have performed two independent measurements of the $^{11}\text{B}(p,\gamma_0)^{12}\text{C}$ reaction over the energy range of $E_p = 5-14$ MeV. The two measurements are in good agreement with each other and indicate that the previously accepted results are in error. The new values for the γ_0 cross section reported here resolve several outstanding conflicts. Their implications are discussed.

[NUCLEAR REACTIONS $^{11}\text{B}(p,\gamma)^{12}\text{C}$, $5 \leq E_p \leq 14$ MeV; measured E_γ ,]
 absolute cross sections for $d\sigma/d\Omega$ (90° , E_γ), $d\sigma/d\Omega$ (90° , E_γ).]




FIG. 6. The $^{11}\text{B}(p,\gamma_1)^{12}\text{C}$ differential cross section at $\theta_{lab}^{ab} = 90^\circ$. The solid curve is taken directly from Ref. 1.




FIG. 7. The ratio of the 90° differential cross section for $^{12}\text{C}(\gamma,\rho_0)^{11}\text{B}$, deduced from the proposed result shown in Fig. 5(a), and the cross section for $^{12}\text{C}(\gamma,\rho_0)^{11}\text{C}$ taken from Ref. 14. The calculations of Ref. 13 are shown as open circles.




FIG. 5. (a) The $^{11}\text{B}(p,\gamma_0)^{12}\text{C}$ differential cross section at $\theta_{lab}^{ab} = 90^\circ$. The proposed result (solid curve) is obtained by applying the correction factor shown in (b) to the data of Ref. 1 (see Ref. 11). The total cross section, shown in (c) as the solid line, was constructed from the proposed result in (a) and the a_2 and a_4 angular distribution coefficients of Ref. 1. The dashed curve in (c) is the total cross section from Ref. 1.

(OVER)




FIG. 8. The $E2$ cross section obtained from $^{11}\text{B}(\bar{p}, \gamma_0)^{12}\text{C}$ measurements (Ref. 16) using the present results to correct the absolute scale.

REF. G.W. Dodson, E.C. Booth, F.L. Milder, B.E. Parad, B.L. Roberts,
D.R. Tieger, J. Comuzzi
Phys. Rev. C26, 2548 (1982)

ELEM. SYM.	A	Z
C	12	6

METHOD

REF. NO.	egf
82 Do 3	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI0	ABY	THR*20	C	140-155	CKV-I		1PI

Photoproduction of π^0 mesons off targets of ${}^6\text{Li}$, ${}^{12}\text{C}$, ${}^{28}\text{Si}$, ${}^{40}\text{Ca}$, natural Cd, and natural Pb was studied using a bremsstrahlung beam with endpoint energies of 140, 145, 150, and 155 MeV. Photoproduction from a liquid hydrogen target was employed as a normalization. The measured yields were found to be in disagreement with published theoretical cross sections for ${}^6\text{Li}(\gamma, \pi^0){}^6\text{Li}$ and also in disagreement with a simple schematic model which assumed only coherent contributions from the M_{1+} multipole. The schematic model, however, did approximately predict the relative magnitudes of the yield curves for the energy range 14–20 MeV over threshold.

*MEV ABOVE THR

NUCLEAR REACTIONS ${}^6\text{Li}$, ${}^{12}\text{C}$, ${}^{28}\text{Si}$, ${}^{40}\text{Ca}$, Cd, Pb, (γ, π^0) ;
 $E_\gamma = 140-155$ MeV; measured σ ; test of reaction model.




FIG. 6. The experimental and calculated yields for (γ, π^0) off a range of complex nuclei. The data were scaled so that the ${}^{12}\text{C}(\gamma, \pi^0){}^{12}\text{C}$ experimental yield fit the calculated yield at 9.7 MeV over threshold (see text).

REF. M.C. Wright, N.R. Roberson, H.R. Weller, D.R. Tilley, Dean Halderson
 Phys. Rev. C25, 2823 (1982)

ELEM. SYM.	A	Z
C	12	6

METHOD	REF. NO.
	82 Wr 2
	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,G	ABX	18-20	D	3-5	NAI-D		DST

PROTON E 2.9-4.6 MEV

The 60° and 90° yield curves for the $^{11}\text{B}(p,\gamma)^{12}\text{C}$ reaction have been measured for $E_p = 2.9$ to 4.6 MeV in 100 keV steps for γ rays leading to the ground and first and third excited states. Recent (π,π') data suggest the existence of an isospin mixed doublet near 19.5 MeV in ^{12}C . A continuum shell model calculation which successfully describes the (π,π') data predicts an observable effect in the γ_3 channel. However, no evidence for these states is found in our γ_3 data.

NUCLEAR REACTIONS $^{11}\text{B}(p,\gamma)^{12}\text{C}$; measured $\sigma(E_p)$, $E_p = 2.9$ to 4.6 MeV, $\theta = 60^\circ, 90^\circ$, for $\gamma_0, \gamma_1, \gamma_3$. Compared γ_3 data to continuum shell model calculations. No evidence for 19.25 and/or 19.65 MeV states is found.

(OVER)




FIG. 2. The cross section at $\theta_{\text{lab}} = 60^\circ$ for $^{11}\text{B}(p,\gamma)^{12}\text{C}$. The error bars represent the statistical errors associated with each data point. The arrows on the γ_3 plot indicate the previously proposed location of the 4^- states.




FIG. 3. Same as Fig. 2, for $\theta_{\text{lab}} = 90^\circ$.



C
A=13

C
A=13

C
A=13

Method ρ detector; N detector; activation
 Ref. No. 57 Co 1
 EGF

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
$C^{13}(\gamma, xn)$ (561)	5-37	13.5 \pm 1	5 \pm 1	$\int_0^{17} = 22$ MeV-mb		$E_{th} = 4.95$ MeV; $\sigma_{max} = 3.7$ mb.
		24 \pm 2	10 \pm 2	$\int_{17}^{38} = 70-95$		$\sigma_{max} = 7.9$ mb.
$C^{13}(\gamma, p)$ (297)	18-38	25.5	6	$\int_{18}^{24} = 55$ MeV-mb		$E_{th} = 17.54$ MeV; $\sigma_{max} = 8.8$ mb.
$C^{13}(\sigma_{\tau})$	5-38	13.5	5	$\int_5^{17} = 22$ MeV-mb		$\sigma_{max} = 3.7$ mb.
		25.5	8.5	$\int_{17}^{30} = 125-140$ MeV- mb		$\sigma_{max} = 16$ mb; $\int_5^{30} \frac{\sigma}{E} dE = 7.54$ mb.

Per Figure 12, values for cross section of Halpern and Mann [Phys. Rev. 83, 370 (1951)]:

$C^{12}(\gamma, p)$	19-23	21.5	1.8	$\int_{20}^{24} = 63$ MeV-mb		$E_{th} = 15.96$ MeV; $\sigma_{max} = 34$ mb.
$C^{12}(\sigma_{\tau})$	18.5-24.5	21.5	2.5	$\int_{18}^{24} = 100$ MeV-mb		$\sigma_{max} = 41$ mb; $\int_{18}^{21} \frac{\sigma}{E} dE = 4.75$ mb.

Thresholds:

	C^{12}	C^{13}
(γ, n)	18.72	4.95
(γ, p)	15.97	17.54
(γ, np)	27.42	20.90
($\gamma, 2n$)	32.04	23.67
($\gamma, 2p$)	27.19	---




Fig. 12. The C^{12} - C^{13} cross section. The dotted curve for the C^{13} - C^{12} cross section is taken from Halpern and Mann reference [1].

[For other figures, see page 2]



Ref.

Elem. Sym.	A	Z
C	13	6

Method

Ref. No.	EGF
57 Co 1	

Reaction	E or ΔE	E ₀	Γ	∫σdE	Jπ	Notes
----------	---------	----------------	---	------	----	-------




FIG. 13. The total photon absorption cross section for C¹³ and C¹².




Figure 11: The cross sections for C¹³(γ, xn) and C¹²(γ, xn). The solid curve is the experimental cross section. The dotted curves for the cross sections C¹³(γ, n)C¹² and C¹³(γ, pn)B¹¹ + C¹³(γ, 2n)C¹¹ are based on reasonable assumptions about the neutron multiplicity. The curve "C¹³(γ, n)C¹² + C¹³(γ, pn)B¹¹ + C¹³(γ, 2n)C¹¹ [hypothetical]" is the assumed cross section for all processes yielding neutrons. The assumptions made are discussed in text.

Elem. Sym.	A	Z
C	13	6
Ref. No.		JHH
60 Ed 1		

Method γ 's from $F^{19}(p, \alpha \gamma)$ reaction, using 874-kev protons from 3-MeV VandeGraff; BF_3 ctrs. in 4 geom.

Reaction	E or ΔE	E_0	Γ	$\int \sigma dE$	$J\pi$	Notes
(γ, n)	6.4					$\sigma_{\gamma, n}(6.4 \text{ MeV}) = 94.1 \pm 10 \mu\text{b}$, using $\sigma_{H^2(\gamma, n)p}(6.4 \text{ MeV}) = (21.2 \pm 3) \times 10^{-28} \text{ cm}^2$ as reference standard. Weighted average of: 6.13 MeV 65% 6.9 MeV 24% 7.1 MeV 11%

METHOD				REF. NO.			
Linac; neutron spectrum, time-of-flight				61 Sa 2		NVB	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	SPC	7-14	C	14 (13.9)	TØF-D	2-9	79




Figure 1
 Energy spectrum of neutrons from the $C^{13}(\gamma, n)$ reaction.

METHOD				REF. NO.			
Synchrotron				64 De 2		NVB	
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,P	ABX	18 - 50	C	18 - 50	ACT-I		4PI

31.3




FIG. 1. Differential cross section for the reaction $C^{13}(\gamma, p)B^{12}$. The hollow circles indicate average values of three independent series of measurements; the solid circles refer to a separate series of measurements obtained at intermediate energy values.

Energy of peaks, MeV:	18.5	20.0	23.5	26.0	29.0	32-50
$\int \sigma(E) dE$, MeV-mb:	1.1	4.4	22.8	7.0	8.4	28.0




FIG. 2. Comparison of our experimental differential cross section curve for the reaction $C^{13}(\gamma, p)B^{12}$ (solid curve) with the curve obtained by Cook^[1] (dashed curve), and with the curve obtained by Dodge and Barber^[2] (dotted curve) for the cross section of the $C^{13}(\gamma, p)B^{12}$ reaction.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	6-11	D	6-11	BF3-I		4PI

Enriched target, ~ 50% C¹³.

TABLE II. Summary of measured cross sections (millibarns).

Source	Energy* (MeV)	Ta ¹⁸¹	Li ⁷	Targets Li ⁶	C ¹³	B ¹⁰
Aluminum	7.72	4.1 ± 0.4	0.06 ± 0.01	1.13 ± 0.12	1.7 ± 0.2	...
Copper	7.91	10.8 ± 1.0	0.07 ± 0.01	1.1 ± 0.2	0.97 ± 0.13	...
Chlorine	8.56	29 ± 6	0.17 ± 0.12
Nickel	9.00	44 ± 6	0.16 ± 0.06	1.6 ± 0.3	0.6 ± 0.1	0.11 ± 0.01
Nitrogen	10.83	121 ± 12	1.07 ± 0.25	...	4 ± 2	0.9 ± 0.2
Chromium	9.72	84 ± 25	0.55 ± 0.25	0.23 ± 0.05
Iron	7.64	0.0 ± 0.9	0.079 ± 0.014	1.3 ± 0.2	0.23 ± 0.05	...
Iron	9.30	0.09 ± 0.03
Lead	7.38	...	0.068 ± 0.035	1.2 ± 0.2	0.3 ± 0.3	...
Sulphur	5.45	0.42 ± 0.07
Sodium	6.41	0.6 ± 0.1
Titanium	6.75	1.3 ± 0.2
Titanium	6.61 ^b	0.32 ± 0.04	...
Manganese	7.16 ^c	0.9 ± 0.1	0.4 ± 0.1	...
Zinc	7.38	1.0 ± 0.2	1.2 ± 0.2	...

* Energies taken from Reiss, 4 and 5.
^b Weighted average of 6.75-, 6.35-, and 6.41-MeV γ rays.
^c Weighted average of 7.16-, 7.15-, and 7.05-MeV γ rays.




FIG. 5. Energy versus cross section, C¹³(γ ,n).

ELEM. SYM.	A	Z
C	13	6
REF. NO.		NVB
64 Ko 2		

METHOD			SOURCE		DETECTOR		ANGLE
Betatron; proton spectrum, yield; CSI scintillator; ion chamber monitor					SCI-D		90
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.P	SPC	20 - 32	C	32	SCI-D	3-13	90

SEPARATED ISOTOPES

Tabelle 1. Überblick über die Messungen

Bei der Berechnung des integrierten Wirkungsquerschnittes wurde isotrope Winkelverteilung angenommen.

E_{min} : untere Abschneidegrenze bei der Bestimmung der Protonenausbeute.

Target	Isotop	Anreicherungsgrad des seltenen Isotops %	Druck des Targetgases mm Hg	$(\frac{dY}{d\Omega})_{90^\circ}$ $\frac{\mu b}{MeV \cdot ster}$	registrierte integrale Protonenzahl	E_{min} MeV	$E_0 \int \sigma dE_p$ MeV mb
Methan CH_4	C^{13}	55,2	700	$2,5 \pm 15\%$	8327	3,5	$37 \pm 20\%$
Knallgas $H_2 + O_2$	O^{18}	90	a) 1560	—	9560	3,5	—
			b) 750	$4,6 \pm 15\%$	7376	2,5	$33 \pm 20\%$




Fig. 3. Energieverteilung der Photoprotonen aus C^{13} . Das Histogramm wurde aus dem der Fig. 1 gewonnen, indem der Anteil der Protonenemission aus C^{12} abgezogen wurde. Nähere Erläuterungen im Text




Fig. 5. Differentieller Wirkungsquerschnitt für Photoprotonenemission aus C^{13} . a) Unter 90° zum γ -Strahl unter der Annahme, daß nur Grundzustandsübergänge vorliegen, diese Arbeit (Histogramm —). b) Ergebnisse von COOK* (Kurve).
 Differentieller Wirkungsquerschnitt für Photoprotonenemission aus C^{12} nach (1) (Histogramm - - -)

REF. W. Bertozzi, P.T. Demos, S. Kowalski, F.R. Paolini, C.P. Sargent,
and W. Turchinets
Nucl. Instr. & Methods 33, 199 (1965)

ELEM. SYM.	A	Z
C	13	6
REF. NO.		egf
65 Be 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	6-14	C	8-14	TOF-D		DST




Fig. 11. Differential cross section for $C^{13}(\gamma, n)C_0^{12}$ for neutrons populating the ground state of C^{12} . The open circles were derived from runs using bremsstrahlung of maximum energy 13.7 MeV, the solid circles 11.9 MeV and the crosses 7.85 MeV.

REF. K. Fukuda, S. Okabe, Y. Sato
 Annu. Rep. Radiat. Center Osaka Prefect. 7, 38 (1966)

ELEM. SYM.	A	Z
C	13	6

METHOD	REF. NO.	
	66 Fu 3	egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	RLY	THR-15	C	5-15	BF3-I		4PI

NSA 13943
BREAKS

13943 FINE STRUCTURE OF THE $^{13}\text{C}(\gamma, n)$ REACTION.
 Fukuda, Kyue; Okabe, Shigeru; Sato, Yuji. Annu. Rep. Radiat.
 Center Osaka Prefect., 7: 38-40(1966).

The neutron yield curve for the $^{13}\text{C}(\gamma, n)$ reaction was studied in detail for the energy region between 4.9 and 15 MeV using an 18 MeV electron linear accelerator. Several breaks (fine structures of the cross section) appeared as changes in the slope of the yield curve at energies of 7.1, 10.4, 11.6 and 14.6 MeV distinctly, and less distinctly at 6.2 and 9.2 MeV. The existence of these breaks means that many resonances of the $^{13}\text{C}(\gamma, n)$ reaction exist below the giant resonance region. (auth)

REF.

G. A. Peterson
Phys. Letters 25B, 549 (1967)

ELEM. SYM.

A

Z

C

13

6

METHOD

REF. NO.

67 Pe 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	LFT	15	D	40-65	D	20-65	DST

$$\Gamma_Y(C^{+3}) = 25 \pm 7 \text{ eV}$$




Fig.1. Ratio of differential cross sections for the excitation of the 15.11 MeV state in ¹³C to that of the 15.11 MeV state in ¹²C versus momentum transferred to the nucleus in units of fm⁻¹.

ELEM. SYM.	A	Z
C	13	6
REF. NO.		HMG
68 Ri 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
P,P'G	LFT	3	D	5	C		
		(3.09)		(4.5)			
		(3.68)					

D,PG also used

DOPPLER SHIFT METHOD

TABLE I. Measurements on the lifetime of the 3.09-MeV state in ¹²C. Slowing down times of $\alpha = 330$ fsec for ¹²C ions in gold and 295 fsec for ¹²C ions in nickel were used to compute the lifetime limits. The attenuation factor F_a is the ratio of the stopped shift to the vacuum shift.

Reaction	E_x (MeV)	Stopping medium	Target thickness ($\mu\text{g}/\text{cm}^2$)	F_a	τ (fsec)	No. of measurements
¹² C(p,p') ¹² C*	4.10	Au	25	1.064 ± 0.056	<33	2
	4.125	Ni	25	0.982 ± 0.024	<38	3
	4.55	Ni	25	0.973 ± 0.020	<35	5
¹² C(d,p) ¹² C*	1.79	Au	15	0.944 ± 0.015	<29	1
	2.715	Au	18	0.988 ± 0.011	<22	3
	3.60	Au	25	0.951 ± 0.025	<40	3
	2.715	Au	15	1.007 ± 0.015	<18	1

TABLE II. Measurements on the lifetime of the 3.68-MeV level in ¹²C. Slowing-down times of $\alpha = 330$ fsec for ¹²C ions in gold and 295 fsec for ¹²C ions in nickel were used to compute the lifetime limits.

Reaction	E_x (MeV)	Stopping medium	Target thickness ($\mu\text{g}/\text{cm}^2$)	F_a	τ (fsec)	No. of measurements
¹² C(p,p') ¹² C*	4.55	Ni	25	0.976 ± 0.018	<35	4
¹² C(d,p) ¹² C*	1.79	Au	15	0.987 ± 0.027	<25	2

H. S. Caplan, R. K. Gupta, E. L. Tomusiak, C. S. Yang

Contributions Montreal Conference 134 (1969)

C

13

6

METHOD

REF. NO.

69 Ca 1

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E, E/$	SPC	0-20	D	140	MAG-D	100-140	DST

ELECTRON SCATTERING FROM CARBON-13

H. S. Caplan, R. K. Gupta, E. L. Tomusiak and C. S. Yang, Saskatchewan Accelerator Laboratory, Saskatoon, Canada.

In the simplest j-j coupling shell model, the ground state of Carbon-13 has one nucleon outside a closed shell. We have used this nucleus to initiate a program of electron scattering experiments to study the effects of an extra nucleon outside a closed shell.

Crannell et al (Nucl. Phys. A103 (1967) 677) measured the elastic scattering of electrons from Carbon-13 and concluded that the rms radius of the charge distribution was smaller than that of Carbon-12. Using the harmonic oscillator shell model they obtained the ratio 0.96 ± 0.01 for the rms radii. Peterson (Phys. Lett. 25B (1967) 549) measured the ratio of the strengths of the transitions to the 15.1 MeV levels in Carbon-12 and Carbon-13. At the time of this writing the above two experiments were the only published (e,e') experiments performed on Carbon-13.

We have measured the elastic and inelastic scattering (up to 20 MeV excitation) in the range of momentum transfer $0.6 < q < 1.2 \text{ fm}^{-1}$ using a 84% enriched Carbon-13 target. The isotopic abundance was checked by measuring the peak at 4.43 MeV due to the Carbon-12 in the Carbon-13 target. Our elastic scattering results confirm Crannell's conclusion that the charge radius of Carbon-13 is smaller than that of Carbon-12. The inelastic spectra show strong excitation of the negative parity states and virtually no excitation of the positive parity states. We observed peaks at 3.06 ($3/2^-, 1/2^-$), 7.55 ($5/2^-, 1/2^-$), 8.35 ($1/2^-, 1/2^-$), 11.7 ($1/2^-, 1/2^-$), 13.76 ($1/2^-, 3/2^-$), 15.1 ($1/2^-, 3/2^-$) and 19.12 ($7/2^-, 1/2^-$) MeV (Ajzenberg-Selove $\lambda = 13$, Lemon Aid Preprint May 1969). We have obtained the form factors for these levels and using the angular dependence of the form factors we have deduced some of the longitudinal and transverse matrix elements.

We have adopted the point of view that if a model of Carbon-13 (or any other nucleus for that matter) is capable of predicting the correct level spacings and strengths for γ -transitions then the electron scattering form factors are logically the next test on the model. At the time of this writing we are calculating the form factors predicted by the various p-shell intermediate coupling wave functions that have been published in the last few years. Presumably this will show us which model is the best and also indicate the need for higher configurations as (speculation) the experimental form factor deviates from the theoretical one at higher momentum transfers.

V. K. Rasmussen and C. P. Swann
Phys. Rev. 183, 918 (1969)

C

13

6

METHOD

REF. NO.

69 Ra 3

hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,G	LFT	3	C	3	NAI-D		DST
		(3.68)		(3.79)			(96,128)

$\Gamma = 0.44 \pm .04$ eV. The width of the level was determined by comparing the scattering with that by the ^{31}P 3.51 and ^6Li 3.56 MeV levels.


REMARKS3 = 3.68 MEV

FIG. 5. Plot of the angular distribution coefficient A_2/A_0 for the spin sequence $\frac{1}{2}-\frac{1}{2}$. The A_4 term is identically zero for this sequence. The experimental limits for the C^{13} and P^{31} distributions are indicated.

REF.

M. Tomaselli, L. Grünbaum, G. A. Beer, H. G. Clerc and G. Wittwer
Phys. Letters 29B, 579 (1969)

ELEM. SYM.

A

Z

C

13

6

METHOD

REF. NO.

69 To 1

EGF

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	3	D	34-65	MAG-D	30-65	DST
		(3.03)					(105-165)


Angular distribution at const. $q = .35, .45, .56 \text{ fm}^{-1}$.3.08 MeV

Fig. 1. Longitudinal and transverse reduced transition probabilities $B(C1, q)$ and $B(E1, q)$ for excitation of the 3.08 MeV level in ^{13}C as a function of momentum transfer.

Solid lines: present theory; the parameter μ is the magnetic moment of the odd neutron as used in the calculations. In the upper part of the figure, the photon point $q = k$ is not shown for reasons of scale; the calculations give $B(E1, k) = B(C1, k) = 2.1 \times 10^{-2} \text{ fm}^2$.
Open circles: present (e, e')-experiments;
full circle \bullet : from measured lifetime [9].

METHOD

REF. NO.

69 Wi 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E.E./	LFT	3-15	D	36-65	MAG-D	15-65	DST

7 LEVELS




Fig. 1. Spectrum of 57 MeV electrons scattered inelastically at 165° from a carbon target (91.7% ^{13}C). The peak at about 5.6 MeV excitation energy is due to electrons scattered elastically from the hydrogen contamination in the target.

Table 1.

Experimental results

E_x (MeV)	I^π	$X\lambda$	R_{E2} (fm)	Γ^2 (eV)	Γ^2/Γ_w	δ (E2/M1)
3.69 = 0.02	$3/2^-$	C2	3.50 = 0.37	$(3.61 = 0.39) \times 10^{-3}$ a)	3.52	0.100 = 0.008 ^{b)}
		M1	2.76 = 0.16			
6.85 = 0.06	$5/2^+$	M2	(3.27)	$(6.0 = 3.6) \times 10^{-5}$	0.055	
7.34 = 0.02	$5/2^-$	C2	3.01 = 0.25	0.1150 = 0.0062 ^{d)}	3.15	
5.56 = 0.02	$1/2^-$	M1	2.50 = 0.19	3.06 = 0.46	0.230	
		C0	(4.60)	2.09 = 0.36 ^{e)}		
9.90 = 0.02	$3/2^-$	C2	(3.46)	$(6.3 = 2.1) \times 10^{-3}$	0.045	0.44 = 0.08
		M1	2.93 = 0.25			
11.07 = 0.02	$1/2^-$	M1	3.03 = 0.22	1.02 = 0.20	0.0359	
		C0	5.27 = 0.43	2.62 = 0.26 ^{e)}		
		or **): C2	4.01 = 0.27	0.256 = 0.026		
15.11 = 0.02	$3/2^-$	C2	(3.46)	0.59 = 0.11	0.50	0.161 = 0.017
		M1	2.55 = 0.20	22.7 = 2.6 ^{e)}		

*) Monopole matrix element in fm^2 .

**) The electric quadrupole classification is an alternative to the M1/C0 classification: in this case, the electric quadrupole transition accounts for at least 2/3 of the observed transverse cross section; therefore an M1 width is not given.

a) 3.27×10^{-3} eV according to ref.5.

c) 0.44 = 0.04 eV according to ref.15.

b) 0.096 = 0.020 according to ref.13.

d) 0.119 eV according to ref.5.

e) 25 = 7 eV according to ref.3.

For the energy levels indicated in the first column, the third column gives the character and multipolarity of the ground state transitions: $C\lambda$ denotes a longitudinal electric, $M\lambda$ a magnetic transition of multipolarity λ , and $C0$ a monopole transition. Column 4 lists the transition radii as defined, e.g. in ref.2. Transition radii R_{E2} in parenthesis were used in the analysis, but not determined by this experiment. The last two columns give the transition strengths in Weisskopf units as defined in ref.14 and the E2/M1 mixing ratios. Experimental data from the literature are given for comparison.

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	6-14	C	5-14	BF3-I		90




Fig. 4. Neutron yield curves from ¹³C target measured with and without the X-ray target as a function of the maximum energy of bremsstrahlung spectrum. Plots on the yield curve from 7.5 to 12.5 MeV are measured without the X-ray target. A dotted line from 10 to 13 MeV shows the neutron yield resulting from photons below 10 MeV.




Fig. 5. Photoneutron cross sections for ¹³C. The solid line is the cross-section curve determined in the present experiment. In addition to the present data, the results of Cook ¹⁾, Edge ²⁾ and Green and Donahue ³⁾ are shown for comparison.

TABLE I

Observed peaks, integrated cross sections and angular distribution functions

Energy (MeV)	$\int \sigma(k) dk$ (MeV · mb)	$W(\theta)$
6.2	0.02	$1 - (1.7 \pm 1.7) \sin^2 \theta$ $1 - (1.5 \pm 0.4) \sin^2 \theta$ $1 - (0.00 \pm 0.30) \sin^2 \theta$ $1 - (0.46 \pm 0.20) \sin^2 \theta$
7.5	0.3	
8.2	0.6	
9.1	1.2	
13	14 ^{*)}	

- 1) B.C. Cook, Phys. Rev. 106, (1957) 300.
- 2) R.D. Edge, Phys. Rev. 119, (1960) 1643.
- 3) L. Green and D.J. Donahue, Phys. Rev. 135, B701 (1964).

*) Integrated cross section from 10 to 14 MeV.

ELEM. SYM.	A	Z
G	13	6
METHOD		REF. NO.
		70 Fu 3
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N	ABX	6-14	G	5-14	BF3-I		DST




Fig. 1 Schematic diagram of the counter assembly and the target chamber. ¹³C and natural carbon targets are set in the target chamber on a target changing device. They move up and down in a vertical line on the plane of the diagram. The electron beam is brought to the center of the target by viewing the beam spot on a zinc sulfide screen with an ITV.




Fig. 7 Photoneutron cross sections for ¹³C. The solid line is the cross-section curve determined in the present experiment. In addition to the present data, the results of Cook¹⁾, Edge¹⁰⁾ and Green and Donahue¹⁷⁾ are shown for comparison.

¹B. C. Cook et al., Phys. Rev. 106, 300 (1957).

¹⁶R. D. Edge, Phys. Rev. 119, 1643 (1960).

¹⁷L. Green et al., Phys. Rev. 135, B701 (1964).




Fig. 6 Angular distributions of photoneutrons for the incident bremsstrahlung with different values of the maximum energy. On the left are photoneutron yields versus angles in the lab system at energies of 6.8, 8.7, 10.1 and 12.1 MeV. Broken lines are least-squares fits of the experimental data to a curve of the form $a + b \sin^2 \theta$. Also shown in the right of the figure are the photoneutron yield curve of ¹³C and its extrapolation corresponding to each peak. Arrows show the energies at which angular distributions are measured.

Table I
Observed peaks, integrated cross sections and angular distribution functions

k_i	$\int \sigma(k) dk$ (MeV.mb)	$W(\theta)$
6.2	0.02	$1 + (1.7 \pm 1.7) \sin^2 \theta$
7.5	0.3	$1 + (1.5 \pm 0.4) \sin^2 \theta$
8.2	0.6	
9.1	1.2	$1 + (0.0 \pm 0.3) \sin^2 \theta$
13	14 ^{a)}	$1 + (0.46 \pm 0.20) \sin^2 \theta$

a) Integrated cross section from 10 to 14 MeV 477

METHOD			REF. NO.		egf		
			70 Wi 1				
REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E _e /	LFT	3-15	D	36-65	MAG-D		DST

Tabelle 2. Longitudinale und transversale Anteile (T_L und T_T) des Wirkungsquerschnitts aus Winkelverteilungen bei konstanter Impulsübertragung q . Die Fehler sind in Prozent angegeben und stehen in Klammern

E_x [MeV]	q^2 [fm ⁻²]	T_L [10 ⁻⁹]	T_T [10 ⁻⁹]
3,08	0,119	14,4 (10)	0,55 (22)
	0,198	23,8 (12)	0,85 (18)
	0,308	36,9 (11)	1,24 (14)
3,69	0,117	45,5 (16)	17,7 (6)
	0,305	22,5 (9)	23,6 (7)
7,54	0,106	54,5 (5)	1,67 (20)
	0,181	145,7 (4)	2,7 (42)
	0,273	293,7 (3)	7,05 (17)
7,83	0,105	2,99 (18)	0,464 (12)
	0,180	2,73 (28)	0,584 (16)
	0,271		1,98 (20)
8,86	0,102	4,59 (35)	5,92 (6)
	0,176	19,6 (14)	8,17 (5)
	0,267	33,8 (5)	9,51 (5)
9,90	0,099	0,68 (85)	0,778 (15)
	0,173	0,40 (83)	0,980 (10)
	0,262	2,62 (21)	1,09 (10)
10,86	0,097	1,97 (8)	0,081 (25)
	0,170	4,0 (42)	< 0,5
	0,257	2,5 (170)	0,77 (100)
11,07	0,096	8,84 (7)	0,776 (12)
	0,169	23,6 (5)	1,05 (25)
	0,256	43,8 (3)	1,04 (18)
15,11	0,085	3,08 (17)	1,429 (3)
	0,154	8,80 (13)	2,180 (2)
	0,238	25,4 (12)	2,440 (4)

9 LEVELS




Fig. 3. Winkelabhängigkeit der Wirkungsquerschnitte in Bornscher Näherung ($f_c = 1$, σ_E in B. N.). An die Meßpunkte konstanter Impulsübertragung q wurden Gerade gemäß Gl. (2) angepaßt; q^2 ist an den jeweiligen Geraden angegeben




Fig. 4. Quadratwurzeln der reduzierten Übergangswahrscheinlichkeiten $B(\lambda, q)$ in Bornscher Näherung ($f_c=1$, σ_E in B. N.) als Funktion von q^2 . Die eingezeichneten Kurven wurden gemäß Gl. (4) an die Meßpunkte angepaßt. Die in Tabelle 3 angegebenen Ergebnisse wurden nicht aus diesen Anpassungen gewonnen, sondern erst in einem zweiten Schritt unter Benutzung von Korrekturfaktoren (s. Abschnitt 4.1). Für $E_x=7,54$ MeV wurden longitudinaler und transversaler Anteil zusammen aufgetragen, um die Abweichungen von der Abschätzung Gl. (6) darstellen zu können

Tabelle 3. Ergebnisse. Der Charakter und die Multipolarität für die Grundzustandsübergänge der in Spalte 1 aufgeführten Niveaus sind in Spalte 3 angegeben; $C\lambda$ bedeutet einen longitudinalen elektrischen, $M\lambda$ einen magnetischen Übergang der Multipolarität λ , $C0$ einen elektrischen Monopol-Übergang. Spalte 4 enthält die Übergangsradien; Übergangsradien in Klammern wurden in der Auswertung benutzt, aber nicht durch die Messungen bestimmt. Die folgenden drei Spalten geben die reduzierten Übergangswahrscheinlichkeiten am Photonenpunkt ($q=k$), die Strahlungsbreiten sowie die Übergangsstärken in Weißkopfeinheiten (Definition nach Wilkinson²⁸). In der letzten Spalte ist der Fehler ϵ in Prozent angegeben; er gilt sowohl für Γ_0^2/Γ_W und $B(\lambda, k)$ als auch für Γ_0^2 , wenn man den Fehler der Anregungsenergien unberücksichtigt läßt

E_x [MeV]	I^π	$X\lambda$	R_{1r} [fm]	$B(k, k)$ [fm ²]	Γ_0^2 [eV]	Γ_0^2/Γ_W	ϵ [%]
$3,08 \pm 0,03$	$1/2^-$	$C1$	(2,84)	0,0221	0,68	0,62	34
$3,69 \pm 0,02$	$3/2^-$	$C2$	$3,50 \pm 0,37$	13,1	$3,61 \times 10^{-3}$	3,52	11
		$M1$	$2,76 \pm 0,16$	$13,6 \times 10^{-3}$	0,358	0,339	13
$6,85 \pm 0,06$	$5/2^+$	$M2$	(3,27)	0,0171	$6,9 \times 10^{-5}$	0,055	52
$7,54 \pm 0,02$	$5/2^-$	$M3$	(3,27)	58	$1,01 \times 10^{-5}$	35	60
		$C2$	$3,01 \pm 0,25$	17,53	0,115	3,15	6
$8,86 \pm 0,02$	$1/2^-$	$M1$	$2,50 \pm 0,19$	$4,61 \times 10^{-3}$	3,36	0,230	14
		$C0$	(4,60)	2,09*			9
$9,90 \pm 0,03$	$3/2^-$	$C2$	(3,46)	0,165	$6,3 \times 10^{-3}$	0,045	33
		$M1$	$2,83 \pm 0,25$	$0,643 \times 10^{-3}$	0,324	0,0159	15
$11,07 \pm 0,02$	$1/2^-$	$M1$	$3,03 \pm 0,22$	$0,721 \times 10^{-3}$	1,02	0,0359	19
		$C0$	$5,27 \pm 0,43$	2,62*			5
	oder $3/2^-$	$C2$	$4,01 \pm 0,27$	3,82	0,256	1,03	11
		$M1$	(2,64)	$0,242 \times 10^{-3}$	0,172	0,0060	33
11,50	$3/2^-$	$M1$	(2,64)	$4,0 \times 10^{-3}$	3,45	0,100	25
$15,11 \pm 0,02$	$3/2^-$	$C2$	(3,46)	1,86	0,59	0,50	18
		$M1$	$2,55 \pm 0,20$	$12,6 \times 10^{-3}$	22,7	0,313	12

* Monopolmatrixelement (ME) in fm².

METHOD

REF. NO.

71 Be 2 hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	8-44	D	77-106	MAG-D		DST

Figures also given for: 77 MeV, 75°
55.4 MeV, 145.7°
81 MeV, 145.7°

963

TABLE I. Resonances observed in this experiment above 15-MeV excitation in ^{13}C .

E_x (MeV)	Width (keV)	
16.2 (16.0)	300	$E_x \leq 17.0$
17.7	300	
18.7	460	
18.7	1200	$17.0 \leq E_x \leq 21.7$
19.3	700	
20.1	400	
20.5	400	
21.2	400	
22.2	1100	
24.7	600	$21.7 \leq E_x \leq 26.7$
25.5	500	
27.3	600	
28.1 (29.4)	500 1200	$E_x > 26.7$ MeV




FIG. 11. The ratio of the ^{13}C to ^{12}C form factors (squared) in the 10-29-MeV region (circles) and the 17-29-MeV region (boxes). The form factors were obtained by integrating the differential form factors over the indicated energy regions. The errors reflect counting statistics and the elastic peak tail uncertainty (see text). The tail uncertainty tends to cancel in the formation of the ratios.

TABLE III. Comparison of the excitation energies of resonances seen in this experiment with the energies calculated by Mashev (Ref. 21).

Present experiment (MeV)	Mashev (MeV)
21.7	24.0, 24.5, 21.0 (unresolved?)
25.5	25.7
20.5	20.2
21.0	20.0
19.3	18.0

(over)




FIG. 3. Similar to Fig. 2, but incident energy is 106 MeV.

ELEM. SYM.	A	Z
C	13	6
REF. NO.		
71 Mu 1		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G.NG	ABY	9-28	C	21,28	SCD-D	2-16	90




Fig. 1. - Energy level diagram showing levels and transitions pertinent to this experiment.

TABLE I. - Gamma-ray strengths from $^{12}\text{C}(\gamma, n\gamma)$.

Bremsstrahlung endpoint (MeV)	E_γ (MeV)	Strength (*)
21	4.444	$(10.1 \pm 3.2) \cdot 10^{-20}$
	15.11	Not seen
28	4.444	$(30.6 \pm 0.7) \cdot 10^{-20}$
	15.11	Clearly seen

(*) Strength is given in units of events per target nucleus roentgen of bremsstrahlung hardened by 20.6 cm of graphite.

(*) H. USERRALL: private communication.

[over]




Fig. 2. - Parts of the pulse-height spectrum taken with 23 MeV bremsstrahlung. The sharp line at 2.615 MeV is due to neutrons scattering inelastically in the lead shield.

reactions. Since the 15.11 MeV level could be fed by the lower T resonance, its observation does not necessarily constitute a signature of the T upper resonance. However, the isospin coupling coefficients give a factor of 4 for the population of this level by the T upper to that due to T lower. It is reasonable therefore to conjecture that the 15.11 MeV level is fed to a large extent by neutrons emitted from the $T = \frac{1}{2}$ component of the giant resonance in ^{12}C .

REF.

Y. M. Shin, C. F. Wong, H. S. Caplan
Nucl. Phys. A166, 162 (1971)

ELEM. SYM.

A

Z

C

13

6

METHOD

REF. NO.

71 Sh 1

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, p	SPC	18-31	D	43	MAG-D	3-13	DST




Fig. 1. Proton energy spectra from the reactions $^{12}\text{C}(e, e'p)^{11}\text{B}$ and $^{13}\text{C}(e, e'p)^{12}\text{B}$ with 43 MeV electrons.

[over]




Fig. 2. Differential cross sections of the $^{12}\text{C}(\gamma, p)^{11}\text{B}$ and $^{13}\text{C}(\gamma, p)^{11}\text{B}$ reactions at 90° with $E = 43$ MeV. The solid line is from the results of Allas *et al.*




Fig. 4. Energy dependence of coefficients in the angular distribution of the reaction $^{13}\text{C}(e, e'p)^{12}\text{B}$.

METHOD

REF. NO.

[Page 1 of 3]

71 Ya 1

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
$E_1 E_2$	FMF	3-20	D	40-125	MAG-D		DST

TABLE 2

10 LEVELS

Total inelastic form factors of the runs performed

E_1 (MeV)	θ (°)	$F^2(q, \theta) \times 10^3$ Excitation energy (MeV)									
		3.69	7.55	3.86	9.50	11.72	11.97	12.32	13.76	15.11	19.15
125.1	155	5.7 ± 0.4	12.7 ± 1.2	^{b)}	^{b)}	3.8 ¹⁾ ± 0.3	^{c)}	^{d)}	4.9 ± 0.5	^{e)}	
95.2	155	5.5 ± 0.3	9.2 ± 0.4	1.5 ± 0.2	^{d)}	1.5 ¹⁾ ± 0.2	^{c)}	5.3 ± 0.4	3.8 ± 0.3	1.6 ± 0.3	
104.7	125	4.3 ± 0.3	7.4 ± 0.4	0.73 ± 0.06	0.22 ± 0.04	0.89 ± 0.09	0.44 ± 0.07	0.39 ± 0.06	0.55 ¹⁾ ± 0.08	0.91 ± 0.09	0.4 \leq
70.0	125	2.77 ± 0.17	3.0 ± 0.4	0.5 ± 0.6	^{d)}	^{d)}	^{d)}	^{d)}	0.78 ¹⁾ ± 0.12	0.15 ± 0.09	^{c)}
63.5	155	4.2 ± 0.3	4.2 ± 0.3	3.0 ± 0.2	1.12 ^{b)} ± 0.11	1.05 ¹⁾ ± 0.11	^{c)}	^{d)}	3.4 ± 0.2	3.4 ± 0.2	^{c)}
97.1	110	4.7 ± 0.4	5.3 ± 0.3	^{b)}	^{b)}	^{b)}	^{b)}	^{b)}	1.92 ± 0.17	1.92 ± 0.17	^{b)}
85.0	140	6.6 ± 0.7	6.4 ± 0.6	^{b)}	^{b)}	^{b)}	^{b)}	^{b)}	2.4 ± 0.4	2.4 ± 0.4	^{b)}

^{a)} The Tsai¹⁰⁾ correction and the Meister and Griffy¹²⁾ correction have been applied to the elastic and inelastic peaks respectively.

^{b)} Not investigated in this run. ^{c)} Very weakly excited in this run. ^{d)} Peak not well defined in this run.

^{e)} Not seen in this run. ^{f)} Peak appeared at 13.9 MeV. ^{g)} Peak appeared at 13.5 MeV.

^{h)} The sum of 9.50 MeV and 9.89 MeV peaks. ⁱ⁾ The sum of 11.72 MeV and 11.97 MeV peaks.

* We use the notation and definitions of DeForest and Walecka¹³⁾.

TABLE 3

Comparisons of experimental observations and theoretical predictions for the 3.69, 7.55 and 15.11 MeV levels

Energy level (MeV)		Theory ^{a)}			
		CK	GHN	B	
3.69	Γ_7^0 (eV)	0.44 ± 0.04 ^{b)}	0.62	0.67	0.52
	$\chi(E2/M1)$	-0.096 ± 0.021 ^{c)}	-0.047	-0.053	-0.052
7.55	Γ_7^0 (eV)	0.119 ^{d)}	0.065	0.061	0.023
	$ \chi(E2/M3) \times 10^{-4}$	1.1 ± 0.7 ^{e)}	0.20	0.38	0.45
15.11	Γ_7^0 (eV)	23.3 ± 2.7 ^{f)}	26-29 ^{f)}		22.4
	$ \chi(E2/M1) $	0.161 ± 0.017 ^{g)}			0.147

^{a)} B, CK and GHN stand for our calculated predictions using wave functions of refs. 1-3). No charge renormalization has been used in our calculations.

^{b)} Ref. 14), $\Gamma_7^0 = 0.36 \pm 0.05$ eV according to ref. 7).

^{c)} Ref. 15), $|\chi(E2/M1)| = 0.10 \pm 0.09$ according to ref. 7).

^{d)} Ref. 16), $\Gamma_7^0 = 0.115 \pm 0.006$ eV according to ref. 7).

^{e)} Ref. 7).

^{f)} Ref. 5).




Fig. 5. Comparisons of the form factor squared between theory and experiment for the 7.55 MeV level in ^{13}C . See caption of fig. 4 for curves. (a) Longitudinal component (M_2^{Coul}), all three curves are indistinguishable in this case; (b) Transverse component (the sum of T_1^{mag} and T_2^{el}).




Fig. 4. Comparison of the form factor squared between theory and experiment for the 3.69 MeV level in ^{13}C . The solid, dash and dot-dash curves correspond to our computed prediction curves using wave functions of Boyarkina ¹⁾, Cohen and Kurath ²⁾ and Goldhammer, Hill and Nachamkin ³⁾. (a) Longitudinal component (M_2^{Coul}); (b) Transverse component (the sum of T_1^{mag} and T_2^{el}).

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	




Fig. 6. Comparisons of the form factor squared between theory and experiment for the 15.11 MeV level in ^{13}C . The wave functions of Boyarkina ¹⁾ were used. (a) Longitudinal component ($M_2^{C^01}$); (b) Transverse component (the sum of T_2^{01} and T_1^{00}).

References

- 1) A. N. Boyarkina, Izv. Akad. Nauk SSSR (ser. fiz), 28 (1964) 337
- 2) S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1
- 3) P. Goldhammer, J. R. Hill and J. Nachamkin, Nucl. Phys. A106 (1967) 62
- 4) H. Crannell, L. R. Suelzle, F. J. Uhrhane and M. R. Yearian, Nucl. Phys. A103 (1967) 677
- 5) G. A. Peterson, Phys. Lett. 25B (1967) 549
- 6) M. Tomascelli, L. Grünbaum, G. A. Beer, H.-G. Clerc and G. Wittwer, Phys. Lett. 29B (1969) 579
- 7) G. Wittwer, H.-G. Clerc and G. A. Beer, Phys. Lett. 30B (1969) 634; preprint from Institut für Technische Kernphysik der Technischen Hochschule Darmstadt, Germany (1970)
- 8) C. S. Yang, Ph. D. thesis, University of Saskatchewan (1970)
- 9) L. Katz, G. A. Beer, D. E. McArthur and H. S. Caplan, Can. J. Phys. 45 (1967) 3721
- 10) Y. S. Tsai, Phys. Rev. 122 (1961) 1398
- 11) L. C. Maximon and D. B. Isabelle, Phys. Rev. 133 (1964) B1344
- 12) N. T. Meister and T. A. Griffy, Phys. Rev. 133 (1964) B1032
- 13) T. DeForest and J. D. Walecka, Adv. in Phys. 15 (1966) 1
- 14) J. L. Friar, Stanford University rep. ITP311 (1968)
- 15) P. Stichel and E. Werner, Nucl. Phys. A145 (1970) 257
- 16) V. K. Rasmussen and C. P. Swann, Phys. Rev. 183 (1969) 918
- 17) A. R. Poletti, J. W. Olness and E. K. Warburton, Phys. Rev. 151 (1966) 312
- 18) B. G. Harvey, J. R. Meriwether, J. Mahoney, A. Bussiére deNercy and D. J. Horen, Phys. Rev. 146 (1966) 712

REF. H. R. Weller and R. A. Blue
Nucl. Phys. A211, 221 (1973)

ELEM. SYM.	A	Z
C	13	6
REF. NO.		egf
73 We 3		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
D,G	ABX	19- 22	D	1- 4	NAI-D		DST




Fig. 4. The measured excitation curve for the $^{11}\text{B}(d, \gamma)^{13}\text{C}$ reaction at 90° . The error bars represent the statistical errors associated with the data points.





Fig. 5. The measured $^{11}\text{B}(d, \gamma)^{13}\text{C}$ angular distributions at $E_d = 2.03$ and 3.61 MeV. The error bars represent the statistical errors associated with the data points.

- 15 P. S. Fisher, D. F. Measday, F. A. Nikolaev, A. Kalmykov, A. B. Glegg, Nucl. Phys. 45 (1963) 113.
- 16 V. P. Denisov, A. V. Kulikov, L. A. Kul'chitskii, JETP (Sov. Phys.) 19 (1964) 1007.
- 17 B. R. Easlea, Phys. Lett. 1 (1962) 163.



489 Fig. 6. Comparison of the (d, γ) resonance with the (p, p) and (p, γ) data for the $A = 13$ nuclei [refs. ^{15, 16}]. The theoretical calculations of Easlea ¹⁷ are also shown.

REF. E. D. Arthur, D. M. Drake, I. Halpern
Phys. Rev. Lett. 35, 914 (1975)

ELEM. SYM.	A	Z
C	13	6
METHOD		REF. NO.
		75 Ar 5
		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
N,G	NOX	18 (18.9)	D	14	NAI-D		DST

Relative yields of capture photons have been observed for four nuclei at angles of 55°, 90°, and 125° in bombardments with 14-MeV neutrons. The yields from ¹⁰B, ²⁹Si, and ⁴⁰Ca show smaller fore-aft anisotropies than those observed in corresponding proton captures. This suggests that the forward peaking in (p, γ) reactions is due mainly to direct rather than collective capture amplitudes. Photons from ¹²C(n, γ)¹³C peak backward, but this peaking cannot be straightforwardly accounted for in terms of the interference between the collective excitations dominant in this energy region.

TABLE I. Angular distribution coefficients.

Reaction	E* (MeV)	a ₂	R _n = 0.57a ₁ - 0.39a ₂	R _p
¹⁰ B(n, γ) ¹¹ B	25	-0.44 ± 0.23	0.05 ± 0.08	¹⁰ B(p, γ) ¹¹ C ~ 0.3 ^a
¹² C(n, γ) ¹³ C	18	-0.08 ± 0.13	-0.15 ± 0.06	¹² C(p, γ) ¹³ N ~ 0.3 ^b
²⁹ Si(n, γ) ³⁰ Si	24	0.2 ± 0.24	0.02 ± 0.1	
⁴⁰ Ca(n, γ) ⁴¹ Ca	22	0.03 ± 0.20	-0.06 ± 0.08	³⁹ K(p, γ) ⁴⁰ Ca ~ 0.2 ^c

^aRef. 13.

^bRef. 14.

^cRef. 15.




FIG. 3. Angular distributions obtained for 14-MeV neutron capture leading to the designated final states. The horizontal bars show the angular widths subtended by the capture targets.

¹³H.M. Kuan, Nucl. Phys. A151, 129 (1970).

¹⁴M. Hasinoff, private communication.

¹⁵E.M. Diener et al., Phys. Rev. C7, 695

(1973); E. M. Diener et al., Phys. Rev. C7, 705 (1973).

$$W(\theta_{n\gamma}) - 1 = \sum_{n=1} a_n P_n.$$

REF.

B. H. Patrick, E. M. Bowey, E. J. Winhold, J. M. Reid,
and E. G. Muirhead
J. Phys. (London) G1, 874 (1975)

ELEM. SYM.	A	Z
C	13	6
METHOD		REF. NO.
		75 Pa 2
		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,NG	ABX	10- 36	C	15- 44	SCD-D		112
G,PG	ABX	19- 37	C	15- 44	SCD-D		112
G,2N	RLY	24- 40	C	35, 40	ACT-I		4PI.

$^{13}\text{C}(G,2N)$ yield: $E_0 = 35$ MeV $1.9 \pm 0.5\%$ of $^{12}\text{C}(G,N)$
 $= 40$ MeV $1.7 \pm 0.5\%$ of $^{12}\text{C}(G,N)$

Table 1. Integrated cross sections.

Reaction	Energy of state to which particle emitted (MeV)	E_{max} (MeV)	$\int_0^{E_{max}} \sigma dE$ (MeV mb)	Total (MeV mb)	Source of data	Reference
γ, n	ground state	38	35	121	Deduced from $^{12}\text{C}(p, \gamma)$ data	Fisher <i>et al</i> (1963)
	4.44	38	8			
	15.1	38	30			
	16.1	38	48			
γ, xn		38		117	Measured	Cook (1957)
		29		62	Measured	McKenzie (1974)
γ, p	ground state	38	25	55	Deduced	Present experiment
	0.95	38	30			
γ, p		38		73	Measured	Cook (1957)
		38		50	Measured	Denisov <i>et al</i> (1964)




Figure 10. The solid curves are the cross sections for neutron emission to the ground, 4.44, 15.1 and 16.1 MeV states in ^{12}C and for proton emission to the ground and 0.95 MeV states in ^{12}B as measured or deduced in the present experiment. The broken curves are the corresponding results from the calculations of Kissener *et al* (1974), multiplied by 0.7.

Kissener *et al.*, Nucl. Phys. A219, 601 (1974)
 Marangoni *et al.*, Phys. Lett. 49B, 253 (1974)
 Fisher *et al.*, Nucl. Phys. 45, 113 (1963)
 Cook, Phys. Rev. 106, 300 (1957)
 McKenzie, Thesis, U. of Melbourne, 1974
 Denisov *et al.*, Sov. Phys. JETP 19, 1007 (1964)

(over)

U.S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS

Table 2. Percentage population of final states in ^{12}C and ^{12}B .

Reaction	Final state (energies in MeV)	Present experiment	Kissener <i>et al</i> (1974)	Marangoni <i>et al</i> (1974)
$^{13}\text{C}(\gamma, n)^{12}\text{C}$	gs	20	2.5 (4.5)	15
	4.44	5	8 (15)	14
	10.3	—	4 (7)	—
	12.7	—	— (—)	4
	15.1	17	8 (15)	15
	16.1	27	13 (24)	26
	Total	69	35.5 (65)	74
$^{13}\text{C}(\gamma, p)^{12}\text{B}$	gs	14	3 (15)	9
	0.95	17	11 (20)	17
	Total	31	19 (35)	26

REF. R. Koch and H. H. Thies
Nucl. Phys. A272, 296 (1976)

ELEM. SYM.	A	Z
C	13	6
REF. NO.		egf
76 Ko 5		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, XN	ABX	5- 25	C	5- 25	BF3-I		4PI

Abstract: The photoneutron cross section in ^{13}C was measured using bremsstrahlung from an electron synchrotron operated in energy sweeping mode. Neutrons were detected by a multi-BF₃-counter detection system. Decorrelated cross-section ordinates were calculated directly using a modified Leiss-Penfold analysis. The cross section from threshold to 25 MeV shows detailed structure, some nine maxima, in agreement with recent continuum shell-model calculations.




Fig. 2. Photoneutron cross section of ^{13}C , present experiment.

REF. J.G. Woodworth, K.G. McNeill, J.W. Jury, P.D. Georgopoulos,
R.G. Johnson
Can. J. Phys. 55, 1704 (1977)

ELEM. SYM.	A	Z
C	13	6
REF. NO.		
77 Wo 4		egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,N0		6-37	C	11-38	TOF-D		98
G,N1		10-36	C	11-38	TOF-D		98

The differential cross sections at 98° for the reactions $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ and $^{13}\text{C}(\gamma, n_1)^{12}\text{C}$ were measured over the range of 6.0 to 37.0 MeV and 10.5 to 35.5 MeV, respectively. The cross sections are compared with previous results on the same nucleus or results from the similar reaction $^{12}\text{C}(p, \gamma_0)^{13}\text{N}$. In the region of the so-called pygmy resonance (about 13.5 MeV) there are many sharp resonances which correspond very well with the level structure of ^{13}C as deduced from several other studies using various reactions. For the full energy region studied, further evidence for the isospin splitting of the ^{13}C photoneutron cross section is obtained. In comparison of the cross sections with recent theoretical results, general agreement is obtained. However, several details of this comparison are in disagreement; the implications of this disagreement are discussed.




FIG. 3. (a) The 98° differential cross section for the reaction $^{13}\text{C}(\gamma, n_1)^{12}\text{C}$ and (b) for the reaction $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ as a function of excitation energy (full energy range).

(over)




FIG. 6. (a) The 98° differential cross section for the reaction $^{13}\text{C}(\gamma, n_1)^{12}\text{C}$ and (b) for the reaction $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ as a function of excitation energy (excitation energies below 20 MeV).

TABLE 2. ^{13}C differential cross sections integrated over energy

Energy region (MeV)	Ground state $\int \frac{d\sigma_p}{d\Omega} dE$ (MeV mb/sr)	First excited state $\int \frac{d\sigma_p}{d\Omega} dE$ (MeV mb/sr)	Ratio $\frac{\int \frac{d\sigma_p}{d\Omega} dE}{\int \frac{d\sigma_0}{d\Omega} dE}$
6.00-37.00	4.13 ± 0.16	2.00 ± 0.38	0.48
6.00-19.00	2.08 ± 0.08	1.19 ± 0.15	0.57
7.59-8.38	0.066 ± 0.003		
11.51-12.03	0.114 ± 0.004	0.364 ± 0.013	0.32
13.13-14.10	0.295 ± 0.008	0.134 ± 0.013	0.45
14.85-15.51	0.166 ± 0.005	0.147 ± 0.015	0.88
17.98-18.55	0.086 ± 0.004	0.075 ± 0.014	0.86
20.13-22.64	0.554 ± 0.016	0.200 ± 0.017	0.36
25.83-28.70	0.371 ± 0.015	0.171 ± 0.036	0.46

TABLE 1. Summary of resonant structure below 20 MeV excitation energy

Center of mass neutron energy E_n (MeV)	Excitation energy E_x (MeV)	Natural width Γ (keV)	Ajzenberg-Selove (1970)		
			E_x (MeV)	Γ (keV)	J^π
2.55	7.71	60	7.68 ± 12 ^a	72 ± 10	$\frac{1}{2}^-$
2.71	7.88	375			
3.61	3.87	175	8.358 ± 14	161 ± 13	$\frac{1}{2}^-$
4.18	9.49	< 90 ^a	9.499 ± 4	≤ 5	($\frac{1}{2}^-$)
4.37	9.69	< 100 ^a			
5.57 } 1.49 ^c }	11.00	< 150 ^a	11.000 ± 20	37	($\frac{1}{2}^-$)
6.56 } 2.48 ^c }	12.08	150			
7.99 } 3.89 ^c }	(13.62)	500	13.55	≈ 500	
9.34	(15.09)	400	14.95 ± 50		
5.30 ^c	15.13	< 135 ^a	15.1087 ± 2.5	5.9 ± 0.9	$\frac{1}{2}^-, T = \frac{1}{2}$
11.04	16.94	< 400 ^a	16.96 ± 50	330	
11.97	17.95	< 450 ^a	(17.99)		

^aError in keV.
^bThe natural width is less than the system resolution. The numbers quoted are the system resolution.
^cFirst excited state transition.

REF. J.W. Jury, B.L. Berman, D.D. Faul, P. Meyer, K.G. McNeill,
J.G. Woodworth
Phys. Rev. C19, 1684 (1979)

ELEM. SYM.	A	Z
C	13	6

METHOD	REF. NO.
	79 Ju 3

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G, 1N	ABX	5-42	D	7-42	BF3-I		4PI
G, 2N	ABX	23-42	D	23-42	BF3-I		4PI

The photoneutron cross sections for ^{13}C have been measured from near threshold to over 40 MeV using monoenergetic photons from positron in-flight annihilation. Several sharp features below the giant resonance were distinguished. The results both for this "pygmy-resonance" region and for the giant resonance near 24 MeV differ markedly from previously reported measurements and provide a much better quantitative comparison with recent theoretical calculations of the photoneutron reaction in ^{13}C . Comparison of the measured total photoneutron cross section with recent data on the ground-state photoreaction and with average photoneutron energies provides evidence for the isospin splitting of the giant resonance for this nucleus.

[NUCLEAR REACTIONS: $^{13}\text{C}(\gamma, n)$, $E_\gamma = 7.6-41.9$ MeV; measured 4π neutron yield for monoenergetic photons; $\sigma(E_\gamma, 1n)$, $\sigma(E_\gamma, 2n)$, integrated cross sections; isospin splitting of the giant resonance.]

TABLE II. Integrated cross sections for ^{13}C .^a

Reaction	$\sigma_{\text{int}} = \int \sigma dE$ (MeV mb)	$\tau_{-1} = \int \sigma E^{-1} dE$ (mb)	$\sigma_{-2} = \int \sigma E^{-2} dE$ (mb MeV ⁻²)
(γ, n)	121.3	5.58	0.301
$(\gamma, 2n)$	4.7	0.14	0.004
(γ, n_{opt})	126.1 ^b	5.72	0.306

^a From threshold to $E_{\gamma, \text{max}} = 41.9$ MeV.

^b TRK sum rule is $60NZ/A = 193.3$ MeV mb.

Uncertainties in the values shall not exceed 10%.

Systematic uncertainties vary from 7% near or below the giant-resonance region to about 20% at the highest energy measured.

TABLE III. Comparisons of integrated photoneutron cross sections for ^{13}C .

Energy interval (MeV)	Present results (MeV mb)	Kissener <i>et al.</i> ^a (MeV mb)	Marangoni <i>et al.</i> ^b (MeV mb)	Other experiments (MeV mb)
5-10	2.4	1		2 ^c
10-14	10.6	10		14 ^c
5-17	21.3	22		20, 122 ^e
17-38	97.7	158		35 ^d
5-42	126.1		157	

^a Reference 9.

^b Reference 11.

^c Fukuda, Ref. 15.

^d Bergstrom *et al.*, Ref. 17.

^e Cook, Ref. 13.

TABLE IV. Isospin sum-rule comparisons for ^{13}C .

Reference	$[\sigma_{-1}(1/2^+) - 0.5\sigma_{-1}(1/2^-)]$ (mb)
Present (γ, n) work plus (γ, p) work of Cook, Ref. 13 Albert <i>et al.</i> , Ref. 12	1.2 ± 0.2 ^a
Soper interaction	1.12
Tabakin interaction	0.56
Marangoni <i>et al.</i> , Ref. 11	1.75

^a See text.

(over)




FIG. 1. Photoneutron cross sections for ^{13}C . Part (a) shows the total photoneutron cross section $\sigma(\gamma, n) + \sigma(\gamma, pn) + \sigma(\gamma, 2n)$, part (b) shows the single photoneutron cross section $\sigma(\gamma, n) + \sigma(\gamma, pn) + \sigma(\gamma, 2n)$, and part (c) shows $\sigma(\gamma, 2n)$. The plotted error bars reflect the statistical uncertainties only.

¹⁴M. Marangoni, P. L. Ottaviani, and A. M. Saruis,
Nucl. Phys. A277, 229 (1977).

Ref. 9: H.R. Kissener, et. al., Nucl. Phys. A219, 601 (1974) Theory

13: 57 Co 1

27: 71 Be 2

¹⁵K. Fukuda, Nucl. Phys. A156, 10 (1970).

ELEM. SYM.	A	Z
C	13	6
REF. NO.		hg
79 Wo 2		

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
(G,NO)	ABX	7-24	C	12-25	TOF-D		DST
		(7.6-24)		(12.0-24.6)			

Abstract: The angular distribution of photoneutrons for the reaction $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ was measured in the region of excitation energy between 7.6 and 24 MeV. The anisotropy coefficients over many of the sharp resonances seen have been interpreted using a simple single-particle description, and information on the spin and parity of these excited ^{13}C states has been found to be in agreement with the known level scheme. New levels at 7.95 MeV and at 12.05 MeV are seen and both are given the tentative J^π assignment of $\frac{1}{2}^-$. Comparison is made with results for the similar $^{12}\text{C}(p, \gamma_0)^{13}\text{N}$ reaction. The measured angular distribution coefficient, a_2/a_0 , displays general agreement in magnitude with the predictions of a recent calculation but the gross energy-dependent features are in marked disagreement. The observed a_2/a_0 coefficient between 14 and 18 MeV has the unexpectedly large average value of $+0.38 \pm 0.05$.

NUCLEAR REACTIONS $^{13}\text{C}(\gamma, n_0)$, $E = 7.6-24$ MeV; measured photoneutron angular distributions with time of flight; deduced $\sigma(E_\gamma, n_0)$.




Fig. 5. Angular distribution coefficients. N.B. The $4\pi a_2$ coefficient has a minimum energy resolution of 75 keV while 120 keV resolution was imposed on the relative Legendre coefficients. The a_3/a_0 data in the region indicated by the horizontal bar are possibly contaminated by non-ground-state neutrons.

Table I

Formulae for the Legendre coefficients describing photoneutron angular distributions from E1, E2 or M1 absorption[†] from ref. (13)

$$a_0 = 21E^2(0, \frac{1}{2}) + 4E^2(1, \frac{1}{2}) + 2M^2(1, \frac{1}{2}) + 4M^2(1, \frac{1}{2}) + 6E^2(1, \frac{1}{2})$$

$$a_1 = -4.000E(1, \frac{1}{2}) + 6.928E(1, \frac{1}{2})E^2(1, \frac{1}{2}) - 4.000E(1, \frac{1}{2})M(1, \frac{1}{2}) + 12.471E(1, \frac{1}{2})E^2(1, \frac{1}{2}) + 4.000E(1, \frac{1}{2})M(1, \frac{1}{2}) + 1.386E(1, \frac{1}{2})E^2(1, \frac{1}{2})$$

$$a_2 = -2.000E^2(1, \frac{1}{2}) + 6.928M(1, \frac{1}{2})E^2(1, \frac{1}{2}) - 2.000M(1, \frac{1}{2})E^2(1, \frac{1}{2}) + 2.000E^2(1, \frac{1}{2}) + 3.429E^2(1, \frac{1}{2}) - 4.000E(1, \frac{1}{2})E(1, \frac{1}{2}) - 4.000M(1, \frac{1}{2})M(1, \frac{1}{2}) + 6.928M(1, \frac{1}{2})E^2(1, \frac{1}{2}) + 6.928M(1, \frac{1}{2})E^2(1, \frac{1}{2})$$

$$a_3 = -5.541E(1, \frac{1}{2})E^2(1, \frac{1}{2}) - 6.928E(1, \frac{1}{2})E^2(1, \frac{1}{2}) - 8.341E(1, \frac{1}{2})E^2(1, \frac{1}{2})$$

[†] For example, E1(0, $\frac{1}{2}$) is the matrix element for the formation via electric dipole absorption of a state with $J = \frac{1}{2}$ and $I = 0$.

ELEM. SYM.	A	Z
C	13	6
REF. NO.		.
80 Ho 3		hg

METHOD			SOURCE		DETECTOR		ANGLE
REACTION	RESULT	EXCITATION ENERGY	TYPE	RANGE	TYPE	RANGE	
G,NO	LFT	6-10	C	10	TOF-D		DST

The angular distribution for the $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ reaction was observed in the energy region 6.5 to 9.3 MeV and at angles of 90° and 135° . The photoneutron measurements were analyzed in terms of a multilevel R-matrix formalism. The $^{12}\text{C}(n, n)^{12}\text{C}$ reaction channel was explicitly included in this analysis. The effects of potential capture were directly observed in the photoneutron spectra. The ground-state radiative widths for resonances in this energy region were deduced from the R-matrix interpretation of the results. The ground-state transition probabilities for E1 excitations at 7.69 and 8.19 MeV were found to be in good agreement with the predictions of the weak-coupling model.

TABLE IV. Deduced ground-state radiative widths for the $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ reaction.

E_γ (MeV)	πL	J^π	$\Gamma_{\gamma 0}$ (eV)	
			Present work	Darmstadt
7.56	E2	$\frac{5}{2}^-$	$0.11 = 0.015$	$0.1150 = 0.0062$
7.69	E1	$\frac{3}{2}^+$	$0.6 = 0.1$	
8.19	E1	$\frac{3}{2}^+$	$7.0 = 0.9$	
8.39	M1	$\frac{1}{2}^-$	$5.4 = 0.5$	$3.36 = 0.46$

[NUCLEAR REACTIONS $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$, $E_{exc} = 6.5 - 9.3$ MeV, measured $\sigma(\theta)$,
 $\theta = 90^\circ, 135^\circ$; deduced $\Gamma_{\gamma 0}$.]




FIG. 1. High-resolution spectra for the $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ reaction at angles of 90° and 135° . The curves represent the results of an R-matrix analysis of the data.

TABLE II. R-matrix parameters for $^{13}\text{C}(\gamma, n_0)^{12}\text{C}$ reaction: $R = 4.61$ fm, $\delta_{1/2} = -0.524$.

πL	J^π	$E_{\gamma\lambda}$ (MeV)	$(\Gamma_{n\lambda})^{1/2}$ (MeV) ^{1/2}	$(\Gamma_{\gamma\lambda})^{1/2}$ (eV) ^{1/2}
E2	$\frac{5}{2}^-$	7.534	0.045 ^a	0.346
E1	$\frac{3}{2}^+$	7.636	0.412	2.068
E1	$\frac{3}{2}^+$	8.151	1.053	1.664
M1	$\frac{1}{2}^-$	8.361	0.412	-2.258
E1	$\frac{1}{2}^+$	11.000 ^b	1.414	-2.966
E1	$\frac{3}{2}^+$	11.000 ^b	1.414	-2.230

^aThis width was chosen to be 0.1 of the energy resolution spread at this energy.

^bThese resonances were arbitrarily placed here in order to account for the effects of distant levels.

TABLE V. Comparison of the deduced internal capture widths with theoretical predictions.

E_γ (MeV)	πL	Present experiment	$\Gamma_{\gamma 0}$ (eV)			
			Barker	Jäger <i>et al.</i>	Klissener <i>et al.</i>	Kurath ^a
7.56	E2	0.11 ± 0.015				0.13
7.69	E1	0.6 ± 0.1	0.71	0.60	1.54	
8.19	E1	7.0 ± 0.9	5.49	0.53	0.39	3.73 ^b
8.39	M1	$5.4 = 0.5$			6.13	13.37-14.73

^aReferences 9 and 22.

^bOnly the integrated strength from both E1 excitations is quoted.

REF. G. Kajrys, W. Del Bianco, J. Kim, S. Landsberger, R. Lecomte,
S. Monaro, P. Paradis
Can. J. Phys. 59, 781(1981)

ELEM. SYM.	A	Z
C	13	6
REF. NO.		
81 Ka 1		hg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
D,G	DST	20-29	D	2-12	NAI-D		90

The 90° differential cross section of the $^{11}\text{B}(d,\gamma_0)^{12}\text{C}$ reaction has been measured at deuteron energies from 1.95 to 12.0 MeV in steps varying from 100 to 200 keV. The γ -rays have been detected by a 23 cm long \times 23 cm diameter NaI(Tl) crystal spectrometer enclosed in a plastic scintillator anticoincidence shield. The yield curve shows a resonant structure at the energies $E_{\text{exc}} = 20.4 \pm 0.1$ and 22.0 ± 0.2 MeV.

La section efficace différentielle de la réaction $^{11}\text{B}(d,\gamma_0)^{12}\text{C}$ a été mesurée pour des énergies des deutons allant de 1.95 à 12.0 MeV. Les rayons gamma ont été détectés par un cristal de NaI(Tl) entouré d'un scintillateur plastique en anticoincidence. La courbe d'excitation montre de la structure aux énergies $E_{\text{exc}} = 20.4 \pm 0.1$ et 22.0 ± 0.2 MeV.

Can. J. Phys., 59, 781 (1981)




FIG. 2. The measured 90° differential cross section of the $^{11}\text{B}(d,\gamma)^{12}\text{C}$ reaction between $E_{\text{exc}} = 20.3$ and 28.9 MeV compared with the results of Weiler and Blue (5) (solid line).

REF. J.J. LeRose, K. Min, D. Rowley, B.O. Sapp, P. Stoler, E.J. Winhold, P.F. Yergin, A.M. Bernstein, K.I. Blomqvist, S. Dytman, G. Franklin, M. Pauli
Phys. Rev. C25, 1702 (1982)

ELEM. SYM.	A	Z
C	13	6

METHOD

REF. NO.	egf
82 Le 1	

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI-	ABX	17*42	C	31*41	MAG-D		90

Differential cross sections for the reaction $^{13}\text{C}(\gamma, \pi^-)^{13}\text{N}_{\text{g.s.}}$ were measured at 90° (lab) for three pion energies: 17, 29, and 42 MeV. The experimental cross sections are anomalously low compared with distorted wave impulse approximation calculations and contrary to the predicted enhancement due to pion condensation precursor effect.

*PION ENERGY

[NUCLEAR REACTIONS $^{13}\text{C}(\gamma, \pi^-)^{13}\text{N}_{\text{g.s.}}$; $E_\pi = 17, 29, \text{ and } 42 \text{ MeV}$; measured $\sigma(E_\pi, \theta = 90^\circ)$; compared with DWIA calculations.]




FIG. 1. Pion yield as a function of pion energy in $^{13}\text{C}(\gamma, \pi^-)^{13}\text{N}$. The threshold energies for exciting the residual nuclear levels are indicated by arrows. The solid line is the least-squares fitted yield curve, using flat background plus calculated photon spectrum (see text), from which the cross section of Fig. 2 is obtained.




FIG. 2. Comparison of experimental cross sections with theoretical calculations. (●) experiment, (○) Heim model, (Δ) Maleki, and (+) Singham, Tabakin, and Dytman. See text.

REF. J. LeRose, K. Min, D. Rowley, B.O. Sapp, P. Stoler, P.-K. Teng, E.J. Winhold, P.F. Yergin, A.M. Bernstein, K.I. Blomqvist, H.S. Caplan, S.A. Dytman, G. Franklin, M. Pauli, K. Shoda, M. Yamazaki Phys. Rev. C26, 2554 (1982)

ELEM. SYM.	A	Z
C	13	6

METHOD

REF. NO.

82 Le 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
G,PI+	ABX	18*42	C	195-250	MAG-0		90

Differential cross section measurements were made for $^{13}\text{C}(\gamma, \pi^+)^{13}\text{B}(\text{g.s.})$ at 90° (lab) for pion energies of 18, 29, and 42 MeV, and for $^{13}\text{C}(\gamma, \pi^+)^{13}\text{B}$ ($E_x = 3.5$ MeV) at 42 MeV. The ground-state results are compared to several distorted-wave impulse approximation calculations and to Helm model calculations. There are significant discrepancies between experiment and theory and among the theoretical results.

*PION ENERGY, TO G.S.

TABLE II. Experimental results for $^{13}\text{C}(\gamma, \pi^+)$. Errors quoted are statistical only (see text).

T_π (MeV)	q_{lab} (fm^{-1})	$d\sigma/d\Omega$ (90° lab)	
		$E_x = 0$	$E_x = 3.5$ MeV
18	0.94	178 ± 9 nb/sr	
29	1.03	230 ± 4	
42	1.15	325 ± 10	166 ± 18

[NUCLEAR REACTIONS $^{13}\text{C}(\gamma, \pi^+)^{13}\text{B}$ ($E_x = 0, 3.5$ MeV), $\theta_\pi = 90^\circ$ (lab), $E_\gamma = 18, 29, 42$ MeV, measured $d\sigma/d\Omega$, compared with DWIA calculations.]




FIG. 3. Experimental isochromat (relative number of pions per MeV versus electron energy) obtained at 42 MeV pion energy. The solid curve is the fit to the data as described in the text, and includes contributions from background and the transitions to the ground state and 3.5 MeV region.




FIG. 5. The solid circles are the present experimental results and the solid square is the 90° result of Shoda *et al.* (Ref. 13). Error bars denote statistical errors only (see text). The curves labeled H1 and H2 are the results of Helm model calculations using the code of Nagl and Überall (Ref. 31) and the parameters of Table III. The points labeled O1(cp) and O2(cp) are as calculated by Sato, Koshigiri, and Ohtsubo (Ref. 22) using HM and CK wave functions, respectively as in Fig. 4, but now including core polarization effects.

(OVER)




FIG. 4. The solid circles are the present experimental results. The solid square experimental point is due to Shoda *et al.* (Ref. 13). The remaining curves and points are the results of several DWIA calculations. The curve labeled M is the result of a calculation by Maleki (Ref. 20), the point labeled C is Cheon's calculation (Ref. 21), and the curves labeled STD1 and STD2 are the results of the calculation of Singham, Tabakin, and Dytman (Ref. 23) using the 1979 and 1982 pion optical potentials of Stricker *et al.*, respectively. The open diamond points labeled O1 and O2 are due to Sato, Koshigiri, and Ohtsubo (Ref. 22), using Hauge-Maripuu (HM) and Cohen-Kurath (CK) wave functions, respectively.

C
A=14

C
A=14

C
A=14

REF. H. Grannell, P.L. Hallowell, J.T. O'Brien, J.M. Finn, F.J. Kline,
S. Penner, J.W. Lightbody, Jr., and S.P. Fivozinsky
PICNS-72, 375 (1972) Sendai

ELEM. SYM.	A	Z
C	14	6
METHOD		REF. NO.
		72 Cr 5
		hvm

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	FMF	7, 8	D	60-120	MAG-D		DST

7= 7.01, 8=8.32 MEV


Table 1. Form Factors Squared for ^{14}C 2^+ States

q (fm^{-1})	θ (Degrees)	$F^2(q)$ (7.01 MeV, 2^+)	$F^2(q)$ (8.32 MeV, 2^+)
0.445	92.9	$(7.10 \pm 1.78) \times 10^{-4}$	---
0.671	93.2	$(2.73 \pm 0.26) \times 10^{-3}$	$(6.86 \pm 1.37) \times 10^{-4}$
0.742	92.9	$(4.68 \pm 0.47) \times 10^{-3}$	$(1.17 \pm 0.24) \times 10^{-3}$
0.750	127.6	$(4.54 \pm 0.44) \times 10^{-3}$	$(1.14 \pm 0.23) \times 10^{-3}$
0.754	145.9	$(5.42 \pm 0.54) \times 10^{-3}$	$(1.64 \pm 0.33) \times 10^{-3}$
0.790	145.6	$(4.52 \pm 0.45) \times 10^{-3}$	$(1.45 \pm 0.30) \times 10^{-3}$
0.888	93.2	$(5.01 \pm 0.50) \times 10^{-3}$	$(1.54 \pm 0.30) \times 10^{-3}$
0.905	127.6	$(6.79 \pm 0.68) \times 10^{-3}$	---
0.906	145.9	$(6.50 \pm 0.65) \times 10^{-3}$	$(1.71 \pm 0.34) \times 10^{-3}$

Table 2. Transition Strengths and Radii for 2^+ Levels in ^{14}C

Excitation Energy (MeV)	$\sqrt{B(C2,0)}$ † ($e \text{ fm}^2$)	R_{tr} (fm)	M^2 (W. U.)	Γ ($eV \times 10^{-3}$)
7.01	4.32 ± 0.29	3.22 ± 0.22	1.79 ± 0.24	50.8 ± 7.1
8.32	2.02 ± 0.38	3.05 ± 0.69	0.39 ± 0.15	4.1 ± 1.7

(over)






Fig. 3. $F^2(q)$ as a function of $(\frac{1}{2} + \tan^2 \theta/2)$ for the 7.01 MeV and 8.32 MeV levels in ^{14}C for $q = 0.75 \text{ fm}^{-1}$. The dashed line shows the least square straight line fit to the data.




Fig. 4. $\sqrt{B(C_2, q)}$ as a function of q^2 for the 2^+ states in ^{14}C . The solid lines show the fit to the data obtained with the restriction that $R_{tr}^4 = (R_{tr}^2)^2$.

ELEM. SYM.	A	Z
C	14	6
REF. NO.		
73 Fa 5		hmg

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	SPC	0- 16	D	50	MAG-D		180
				(50.4)			

PEAKS 7.3, 7.9, 9.3




Fig. 12. Preliminary 180° electron scattering spectrum from ¹⁴C. Vertical scale unavailable at publication time

F.J. Kline, H. Crannell, J.M. Finn, P.L. Hallowell, J.T. O'Brien,
C.W. Werntz, S.P. Fivozinsky, J.W. Lightbody, Jr. and S. Penner
Nuovo Cimento 23, 137 (1974)

C

14

6

METHOD

REF. NO.

74 Kl 2

egf

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	SPC	20- 24	D	61, 81	MAG-D		146

NO T=2 STATE FOUND

Summary. — A search for the lowest-energy $T=2$ state in ^{14}C has been attempted employing inelastic electron scattering techniques. Spectra obtained at 61 and 81 MeV incident electron energy and 143.7° scattering angle fail to reveal significant structure at the predicted resonance energy. The implications of this result are discussed in terms of the particle-hole model and the ground-state structure of the ^{14}B analogue of this level.





Fig. 1. — Bin-sorted spectra of electrons scattered from ^{14}C with excitation energy between 20.0 and 25.0 MeV. The smooth peaks shown illustrate the experimental resolution, determined from elastic scattering, and correspond in height to the assumed upper limit of cross-section for the $T=2$ state: a) incident energy of 61.50 MeV and scattering angle of 143.7° , b) incident energy of 80.70 MeV and scattering angle of 145.7° .

TABLE 1. — Upper limits for excitation of a nonrot $T=2$ state in ^{14}C .

Incident energy (MeV)	g (fm $^{-1}$)	$d\sigma/d\Omega$ (cm 2 /sr)	$F_2^2(g)$
61	0.46	$1.4 \cdot 10^{-33}$	$7 \cdot 10^{-3}$
81	0.67	$3 \cdot 10^{-33}$	$2.7 \cdot 10^{-3}$

REF. H. Crannell, J.M. Finn, P. Hallowell, J.T. O'Brien, N. Ensslin,
L.W. Fagg, E.C. Jones, Jr., and W.L. Bendel
Nucl. Phys. **A278**, 253 (1977)

ELEM. SYM.	A	Z
C	14	6

METHOD	REF. NO.
	77 Cr 1

REACTION	RESULT	EXCITATION ENERGY	SOURCE		DETECTOR		ANGLE
			TYPE	RANGE	TYPE	RANGE	
E, E/	ABX	7 - 16	D	37-60	MAG-D		180

8 STATES

Abstract: Inelastic scattering of 37, 50, and 60 MeV electrons at 180° from ¹²C has been studied. Cross sections for the excitation of eight states in ¹²C with excitation energy less than 16 MeV have been observed. Most of the strength is observed to be concentrated in one transition at 11.31 MeV which is assigned a spin and parity of 1⁻. The total width of this state is observed to be 207 ± 13 keV, while the electromagnetic transition width is determined to be 6.8 ± 1.4 eV.

TABLE 2
Measured cross sections for levels in ¹²C

E _x	J ^π	Cross section (10 ⁻³¹ cm ² sr)		
		37 MeV	50.5 MeV	60.5 MeV
7.01 ^{a)}	2 ⁻	0.90 ± 0.2	1.61 ^{b)} ± 0.24 ± 0.27	1.62 ± 0.3
7.34	2 ⁻	< 0.3	0.72 ± 0.33	0.66 ± 0.4
8.32	2 ⁻	< 0.3	0.66 ± 0.18	1.02 ± 0.24
9.80	(1)	0.63 ± 0.3	0.84 ± 0.24	1.32 ± 0.24
10.5		< 0.4	0.72 ± 0.24	0.99 ± 0.3
11.31 ^{a)}	1 ⁻	11.95 ^{b)} ± 0.72 ± 1.20	10.98 ^{b)} ± 0.6 ± 1.08	9.01 ± 0.48 ± 0.87
12.97	(3, 4)	0.9 ± 0.3		1.41 ± 0.6
14.67		0.9 ± 0.5		0.57 ± 0.3
15.11 ^{b)}	1 ⁻	22.4	19.1	14.7
DWBA PWBA ^{c)}		1.13	1.09	1.06

^{a)} Smaller uncertainty due to peak-area ratio measurement only. Larger uncertainty includes uncertainties due to target composition effects (¹²C, ¹³C isotopic composition).

^{b)} The ¹²C 15.11 MeV cross sections used for normalization.

^{c)} Interpolated from the work of Chertok *et al.*³⁾

DEFINITIONS OF ABBREVIATIONS AND SYMBOLS

Note: In this list definitions are given for various photoneutron reactions in which the following symbols are used: N, NL, nN, SN and XN. Corresponding definitions apply for reactions involving other nuclear particles where the symbols N (neutron) is replaced by, e.g. P, D, T, HE, A etc. Where unknown reactions result in the production of a specific radionuclide, the chemical symbol and mass number is listed as the reaction product, e.g. a G,NA22 reaction in ⁵⁹Co.

A	alpha particle		response function. Contrast with D = discrete.
ANAL	analysis		
ABI	absolute integrated cross-section data	CCH	cloud chamber
ABX	absolute cross-section data	CF	compared with
ABY	absolute yield data. Often means cross-section per equivalent quantum is listed.	CHRGD	charged
ACT	measurement of induced radio-activity of the target	CMPT	Compton
ASM	asymmetric, asymmetry	COIN COINC	coincidence, coincide
AVG	average	COH	coherent
BBL	bubble chamber	CK	Cerenkov
BEL B(EL)	reduced electric radiative transition probability	D	deuteron or discrete. When discrete, it is used to describe a photon source or a detector response function. Contrast with C = continuous.
BF3	BF ₃ neutron counter with moderator e.g., Halpern detector, long counter	DLTE	energy loss
BML	reduced magnetic radiative transition probability, B(ML)	DLTQ	momentum transfer
BREAKS	levels located by "breaks" in the yield curve	DST	distribution
BRKUP	breakup	DT BAL	detailed balance
BRMS	bremstrahlung	E	electron
BTW	between	E/	inelastically scattered electron
C	continuous. Used to describe a photon source or a detector	E+	positron
		EDST	energy distribution or spectrum
		E/N	used only to indicate a coincidence experiment as in (E,E/N).

	N stands for any outgoing particle measured in coincidence with an inelastically scattered electron. Distinguish from eg., (E,N) which is used to represent an electron induced reaction when only the outgoing particle N is detected.	KE	kinetic energy
EMU	emulsions (photographic plates)	L	may be an integer or zero that always follows a reaction product symbol. This is used to indicate transitions to specific states in the residual nuclide. When the letter is used as in (G,NL) the cross section given is that for the sum of transitions to two or more specific final states.
EXCIT	excited	LFT	excited state lifetime
F	fission	LIM	limit
FMF	form factor	LV,LVS	level, levels
FM-1	inverse femtometers	LQD	liquid
FRAG	fragment	MAG	magnetic spectrometer
G	photon	MEAS	measurement(s)
G/	inelastically scattered photon	MGC	magnetic Compton spectrometer
G-WIDTH	gamma-ray transition width	MGP	magnetic pair spectrometer
HAD	hadrons, hadron production	MOD	moderated neutron detector <u>not</u> employing a BF ₃ counter, e.g. rhodium foil, Szilard-Chalmers reaction, ³ He, ⁶ Li reactions, GD loaded liquid scintillator, etc.
HE He3	³ He particle	MSP	mass spectrometer
INT	interaction, integral, intensity	MULT	multiple, multipole, multiplicity
INC	includes	MU-T	used only in combination with G to indicate a total photon absorption cross section measurement, i.e. (G,MU-T)
ION	ionization chamber	N	neutron (see also XN and SN). The notation (G,N) is used to indicate a reaction in which only a single neutron is emitted, i.e. the reaction that can, in many cases, be measured by observing the radioactive decay of the residual nuclide.
ISOB	isobaric		
ISM	isomer		
J	multiplicity of particle defined by following symbol e.g. (G,PJN) with remark J = 2,3,5,7		
JPI J-PI	spin and parity of a nuclear state		
K	second multiplicity index, e.g. (G,JPKN) with both J & K positive integers greater than 1		

nN	where n is any integer. (G,nN) indicates the sum over all reaction cross sections in which n neutrons are emitted.	SN	sum of neutron producing reactions, $\sigma(\gamma,SN) = \sigma(\gamma,N) + \sigma(\gamma,NP) + \sigma(\gamma,2N) + \sigma(\gamma,3N) + \text{etc.}$
NAI	NaI(Tl) spectrometer	SPC	photon or particle energy spectrum
NEUT	neutron(s)	SPK	spark chamber
NOX	no cross-section data	SPL	spallation
P	proton (see also XP)	STAT	statistical
PART	particle(s)	SYM	symetric, symmetry
PHOT	photon(s)	T	triton
PI	pion, usually written as PI+, PI-, PIO to indicate charge	TEL	counter telescope
POL	polarized or polarization	THR	threshold for reaction or threshold detector, e.g., $^{29}\text{Si}(n,p)^{29}\text{Al}$.
Q-SQUAR	momentum transfer squared (q^2)	TOF	time-of-flight detector
RCL	recoil	TRK	tracks of particles or fragments observed in solid materials (glass, mylar, etc.)
REL	relative	TRNS	transition
RLI	relative integrated cross-section data	UKN UNK	unknown
RLX	relative cross-section data	VIB	vibrational
RSP	reaction spectrometer	VIR PHOT	virtual photon(s)
RLY	relative yield data	XN	all neutrons, total neutron yield, $\sigma(\gamma,XN) = \sigma(\gamma,N) + 2\sigma(\gamma,2N) + 3\sigma(\gamma,3N) + \sigma(\gamma,NP) + \text{etc.}$
SCTD	scattered	XP	all protons, total proton yield $\sigma(\gamma,XP) = \sigma(\gamma,P) + \sigma(\gamma,NP) + 2\sigma(\gamma,2P) + \text{etc.}$
SCD	semiconductor (solid state) detector	XX XXX	reaction products defined in REMARKS
SCI	scintillator detector other than NaI, e.g., CsI, KI, organic (liquid or solid), stilbene, He	YLD	yield
SEP	separation		
SEP ISOTP	separated isotope used		
SIG	SIGMA (cross section)		

4PI	a 4π geometry was used or a method like radioactivity or a total absorption measurement		products was determined. The polarized particle is indicated in REMARKS.
999	energy defined in REMARKS	* or @	symbols used to indicate that the units associated with the numerals on one or both sides of the symbol in a specific column are not MeV. The units are defined in REMARKS.
\$	indicates the measurement involved beams or targets that were either polarized or aligned, or that the polarization of the reaction		

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET <i>(See instructions)</i>	1. PUBLICATION OR REPORT NO. NBSIR 83-2742	2. Performing Organ. Report No.	3. Publication Date October 1983
4. TITLE AND SUBTITLE Photonuclear Data-Abstract Sheets 1955-1982			
5. AUTHOR(S) E.G. Fuller and Henry Gerstenberg			
6. PERFORMING ORGANIZATION <i>(If joint or other than NBS, see instructions)</i> NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, D.C. 20234		7. Contract/Grant No.	
		8. Type of Report & Period Covered	
9. SPONSORING ORGANIZATION NAME AND COMPLETE ADDRESS <i>(Street, City, State, ZIP)</i>			
10. SUPPLEMENTARY NOTES <input type="checkbox"/> Document describes a computer program; SF-185, FIPS Software Summary, is attached.			
11. ABSTRACT <i>(A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here)</i> These abstract sheets cover most classes of experimental photonuclear data leading to information of the electromagnetic matrix element between the ground and excited states of a given nucleus. This fifteen volume work contains nearly 7200 abstract sheets and covers 89 chemical elements from hydrogen through americium. It represents a twenty-seven year history of the study of electromagnetic interactions. The sheets are ordered by target element, target isotope, and by an assigned bibliographic reference code. Information is given on the type of measurement, excitation energies studied, source type and energies, detector type, and angular ranges covered in the measurement. For a given reference, the relevant figures and tables are mounted on a separate sheet for each nuclide studied.			
12. KEY WORDS <i>(Six to twelve entries; alphabetical order; capitalize only proper names; and separate key words by semicolons)</i> data-abstract sheets, elements, experimental, isotopes, nuclear physics, photonuclear reactions			
13. AVAILABILITY <input type="checkbox"/> Unlimited <input checked="" type="checkbox"/> For Official Distribution. Do Not Release to NTIS <input type="checkbox"/> Order From Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402. <input type="checkbox"/> Order From National Technical Information Service (NTIS), Springfield, VA. 22161		14. NO. OF PRINTED PAGES @ 520 pp	
		15. Price	

