

A11102 145725

NAT'L INST OF STANDARDS & TECH R.I.C.



A11102145725

/NSRDS-NBS
QC100 .U573 V57;1976 C.1 NBS-PUB-C 1964

NBS
PUBLICATIONS



NSRDS-NBS 57

U.S. DEPARTMENT OF COMMERCE / National Bureau of Standards



Yields of Free Ions Formed in Liquids by Radiation

QC
100
.U573
No. 57
1976
c.2

NATIONAL BUREAU OF STANDARDS

The National Bureau of Standards¹ was established by an act of Congress March 3, 1901. The Bureau's overall goal is to strengthen and advance the Nation's science and technology and facilitate their effective application for public benefit. To this end, the Bureau conducts research and provides: (1) a basis for the Nation's physical measurement system, (2) scientific and technological services for industry and government, (3) a technical basis for equity in trade, and (4) technical services to promote public safety. The Bureau consists of the Institute for Basic Standards, the Institute for Materials Research, the Institute for Applied Technology, the Institute for Computer Sciences and Technology, and the Office for Information Programs.

THE INSTITUTE FOR BASIC STANDARDS provides the central basis within the United States of a complete and consistent system of physical measurement; coordinates that system with measurement systems of other nations; and furnishes essential services leading to accurate and uniform physical measurements throughout the Nation's scientific community, industry, and commerce. The Institute consists of the Office of Measurement Services, the Office of Radiation Measurement and the following Center and divisions:

Applied Mathematics — Electricity — Mechanics — Heat — Optical Physics — Center for Radiation Research: Nuclear Sciences; Applied Radiation — Laboratory Astrophysics² — Cryogenics² — Electromagnetics² — Time and Frequency².

THE INSTITUTE FOR MATERIALS RESEARCH conducts materials research leading to improved methods of measurement, standards, and data on the properties of well-characterized materials needed by industry, commerce, educational institutions, and Government; provides advisory and research services to other Government agencies; and develops, produces, and distributes standard reference materials. The Institute consists of the Office of Standard Reference Materials, the Office of Air and Water Measurement, and the following divisions:

Analytical Chemistry — Polymers — Metallurgy — Inorganic Materials — Reactor Radiation — Physical Chemistry.

THE INSTITUTE FOR APPLIED TECHNOLOGY provides technical services to promote the use of available technology and to facilitate technological innovation in industry and Government; cooperates with public and private organizations leading to the development of technological standards (including mandatory safety standards), codes and methods of test; and provides technical advice and services to Government agencies upon request. The Institute consists of the following divisions and Centers:

Standards Application and Analysis — Electronic Technology — Center for Consumer Product Technology: Product Systems Analysis; Product Engineering — Center for Building Technology: Structures, Materials, and Life Safety; Building Environment; Technical Evaluation and Application — Center for Fire Research: Fire Science; Fire Safety Engineering.

THE INSTITUTE FOR COMPUTER SCIENCES AND TECHNOLOGY conducts research and provides technical services designed to aid Government agencies in improving cost effectiveness in the conduct of their programs through the selection, acquisition, and effective utilization of automatic data processing equipment; and serves as the principal focus within the executive branch for the development of Federal standards for automatic data processing equipment, techniques, and computer languages. The Institute consists of the following divisions:

Computer Services — Systems and Software — Computer Systems Engineering — Information Technology.

THE OFFICE FOR INFORMATION PROGRAMS promotes optimum dissemination and accessibility of scientific information generated within NBS and other agencies of the Federal Government; promotes the development of the National Standard Reference Data System and a system of information analysis centers dealing with the broader aspects of the National Measurement System; provides appropriate services to ensure that the NBS staff has optimum accessibility to the scientific information of the world. The Office consists of the following organizational units:

Office of Standard Reference Data — Office of Information Activities — Office of Technical Publications — Library — Office of International Relations — Office of International Standards.

¹ Headquarters and Laboratories at Gaithersburg, Maryland, unless otherwise noted; mailing address Washington, D.C. 20234.

² Located at Boulder, Colorado 80302.

JUL 7 1976

not acc.

QC100

.U573

.D57

.D57

.D57

Yields of Free Ions Formed in Liquids by Radiation

t. National Standard Reference Data Series, NSRDS-NBS

A. O. Allen

Department of Chemistry
Brookhaven National Laboratory
Upton, New York 11973



U.S. DEPARTMENT OF COMMERCE, Elliot L. Richardson, Secretary

James A. Baker, III, Under Secretary

Dr. Betsy Ancker-Johnson, Assistant Secretary for Science and Technology

U.S. NATIONAL BUREAU OF STANDARDS, Ernest Ambler, Acting Director

Issued April 1976

Library of Congress Cataloging in Publication Data

Allen, Augustine O.

Yields of free ions formed in liquids by radiation.

(Nat. stand. ref. data ser.; NSRDS-NBS 57)

"CODEN: NSRDAP."

Includes bibliographical references.

Supt. of Docs. No.: C 13.48:57

1. Ions. 2. Liquids, Effect of radiation on. I. Title. II. Series:
United States. National Bureau of Standards. National standard
reference data series; NSRDS-NBS 57.
QC100.U573 No. 57 [QD561] 602'.1s [541'.372] 75-619361

NSRDS-NBS 57

Nat. Stand. Ref. Data Ser., Nat. Bur. Stand. (U.S.), 57, 25 pages (Apr. 1976)

CODEN: NSRDAP

© 1976 by the Secretary of Commerce on Behalf of the United States Government

U.S. GOVERNMENT PRINTING OFFICE
WASHINGTON: 1976

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402
(Order by SD Catalog No. C13.48:57). Price 55 cents. (Add 25 percent additional for other than U.S. mailing).

Foreword

The National Standard Reference Data System provides access to the quantitative data of physical science, critically evaluated and compiled for convenience and readily accessible through a variety of distribution channels. The System was established in 1963 by action of the President's Office of Science and Technology and the Federal Council for Science and Technology, and responsibility to administer it was assigned to the National Bureau of Standards.

NSRDS receives advice and planning assistance from a Review Committee of the National Research Council of the National Academy of Sciences-National Academy of Engineering. A number of Advisory Panels, each concerned with a single technical area, meet regularly to examine major portions of the program, assign relative priorities, and identify specific key problems in need of further attention. For selected specific topics, the Advisory Panels sponsor subpanels which make detailed studies of users' needs, the present state of knowledge, and existing data resources as a basis for recommending one or more data compilation activities. This assembly of advisory services contributes greatly to the guidance of NSRDS activities.

The System now includes a complex of data centers and other activities in academic institutions and other laboratories. Components of the NSRDS produce compilations of critically evaluated data, reviews of the state of quantitative knowledge in specialized areas, and computations of useful functions derived from standard reference data. The centers and projects also establish criteria for evaluation and compilation of data and recommend improvements in experimental techniques. They are normally associated with research in the relevant field.

The technical scope of NSRDS is indicated by the categories of projects active or being planned: nuclear properties, atomic and molecular properties, solid state properties, thermodynamic and transport properties, chemical kinetics, and colloid and surface properties.

Reliable data on the properties of matter and materials are a major foundation of scientific and technical progress. Such important activities as basic scientific research, industrial quality control, development of new materials for building and other technologies, measuring and correcting environmental pollution depend on quality reference data. In NSRDS, the Bureau's responsibility to support American science, industry, and commerce is vitally fulfilled.

ERNEST AMBLER, *Acting Director*

Preface

This report is one of a series of data publications on radiation chemistry; the aim of the series is to compile, evaluate, and present the numerical results on processes occurring in systems which have been subjected to ionizing radiation. Various kinds of data are important in radiation chemistry. The quantities which were measured first were the observed radiation yields or G values (molecules formed or destroyed per 100 eV). Various indirect methods based on G values have been used to determine yields of transient species and relative rates of reactions. The spectral properties (optical, electron spin resonance) of transients have provided a direct method for their identification, and rates of the very fast reactions of transients which occur in irradiated systems have been measured directly by spectroscopic methods. Conductivity and luminescence methods have also provided a means of measuring properties of transients and their kinetics. Some reactions which occur in irradiated systems have also been studied by other methods, such as photochemistry, electric discharge, ultrasonics, chemical initiation, electron impact, etc. The emphasis in these publications is on the data of radiation chemistry, but where other pertinent data exist, they are included.

The data of radiation chemistry are voluminous; thousands of systems have been investigated. As a result there are certain collections, *e.g.* rate constants of particular types of reactions or certain properties of transients, for which tabulations of the data are considered essential, but for which critical assessment of each value is impossible. On the other hand, certain systems and properties have been studied so extensively that critical examination of these data is desirable and timely. Authors of this series of data publications have been asked to evaluate the extent to which the data can be critically assessed, to describe their criteria for evaluation, and to designate preferred values whenever possible.

Yields of Free Ions Formed in Liquids by Radiation.*

A. O. Allen

Department of Chemistry
Brookhaven National Laboratory
Upton, N.Y. 11973

Free ions are those produced by ionizing radiation which escape initial recombination. Yields of free-ion pairs are tabulated for liquid alkanes, alkenes, alkynes, dienes and aromatic hydrocarbons, and other non-polar and polar compounds, including alcohols, ethers, esters, halides, amines, nitriles, etc. Total ion yields for a few liquids are compared with gas phase ion yields. Theoretical treatments of these data are discussed.

Key words: Electrons; free ions; ion yields; liquid; non-polar systems; polar systems; radiation chemistry; radiation physics.

1. Definitions

When X-rays, gamma rays or fast charged particles traverse a *gas*, positive and negative ions are formed in equal numbers, and (in the absence of an applied electric field) become uniformly distributed through the volume by diffusion, and eventually disappear by pairwise *general recombination*. If a D.C. electric field is applied, the ions drift to the electrodes and give up their charge to the external circuit. At ordinary radiation intensities, a relatively modest field strength suffices to collect all the ions. The number of ions of either sign produced by unit energy input is called the *ion yield* and is a characteristic property of each gaseous substance.

In a *liquid* under radiation, the charge collected by an applied field is usually only a small fraction of that produced by the same radiation absorbed in the same amount of the same material in gaseous form. The cause of this difference was correctly described nearly seventy years ago by G. Jaffé [080002]¹. In a gas, ionization knocks out an electron with sufficient momentum to escape from the electric field of the positive ion it leaves behind; but in a liquid, collisions with surrounding molecules rapidly reduce its initial outward momentum so that it has only a small escape probability. Usually the electron rapidly returns to its parent ion and recombines with it. This process is called *initial recombination* or *geminate recombination*. The number of electrons that escape initial recombination, per unit energy input by the radiation, is called the *free-ion yield*. Since an applied electric field tends to pull the electron out of the field of the positive ion, the free-ion yield increases with the applied field. The field strength that a liquid can stand is limited by dielectric breakdown; in most liquids, the free-ion yield at the maximum usable field is still only a small fraction of the ion yield found in the gaseous form of the same substance. Only in monatomic liquids (liquefied rare gases) has complete collection of the ions been reported.

The escaped electron can (1) remain as a *quasi-free electron* in a delocalized state in a conduction band; (2) aggregate solvent molecules about itself to form a *solvated electron*; (3) react with a solvent molecule, or an impurity or solute molecule to form a new negatively charged molecule, or *negative ion*.

* This is a data review prepared for, and in cooperation with, the Radiation Chemistry Data Center of the Radiation Laboratory, University of Notre Dame, Notre Dame, Indiana 46556. The Laboratory is operated under contract with the Energy Research and Development Administration. The work of the Center is supported in part by the National Bureau of Standards, Office of Standard Reference Data.

¹ Numbers in brackets indicate the literature references at the end of this paper.

The term "negative ion" is however sometimes used to indicate any negatively charged entity, including the electron.

Although attempts are made to extrapolate the free-ion yield to infinite field strength to obtain the *total ion yield*, the zero-field free-ion yield is precisely determinable and has been the focus of much recent work.

The free-ion yield depends strongly on the "radiation quality" or linear energy transfer (LET). At high LET, the positive charges are grouped in columns, the electrons are pulled back by a large group of positive ions instead of by just one or a few, and the escape probability becomes much lower. Most of the data reported here were taken with low LET or near-minimum ionization radiations such as γ rays, 1–2 MeV X-rays or megavolt electron beams, and this condition may be assumed unless otherwise indicated.

The free-ion yield is of practical significance in determining the chemical changes that occur in liquid and amorphous systems under radiation. The scientific significance of free-ion yield measurements lies in the fact that no adequate theory exists for the behavior of electrons in non-crystalline condensed systems, and free-ion yields offer one type of data which may point the way to the development of such a theory.

The purpose of this review is to assemble currently available data on zero-field free-ion yields in pure liquids for the convenience of those who wish to use these quantities. No attempt is made here to assess probable errors or to ascertain probable "best" values. Before this can be done, the many differences in results reported from different laboratories will have to be resolved in the light of independent determinations.

2. Notation and Units

The ion yield in gases is usually expressed as its reciprocal, denoted W , which is always expressed in eV per ion pair. In liquids, the yield is usually given as $G \equiv 100/W$, the number of ions of either sign produced per 100 eV energy input. In this paper, the zero-field free-ion yield in liquids is denoted by G_{fi}° , while the total ion yield is G_{fi}^∞ . The electron is represented by e^- .

3. Methods of Measurement

3.1. Scavenging

The scavenging method involves adding a substance which reacts directly with the electron (or in some cases with the positive ion or its deprotonated successor radical) to form a chemical product which is determined quantitatively. Examples of scavengers are anthracene, biphenyl, pyrene and the free radical galvinoxyl, which readily add an electron to form a colored radical anion of known optical absorptivity, and N_2O , which reacts with an electron to form O^- and a molecule of nitrogen, which is determined. There are two varieties of the method: (1) the *steady-state* method, in which a permanent chemical product is determined, following an exposure to a continuous source such as γ rays; (2) the *pulse* method, in which the optical absorption of a transient product such as pyrene ion gives a measure of the electron yield.

There are many sources of error. If the scavenger concentration is too high, some of the initially recombining electrons or ions may be counted as well as the free ions; results should always be extrapolated to zero scavenger concentration. A less obvious error occurs if some of the electrons react with the scavenger to form other products, which are not determined; or the electron may also be reacting with impurities, or with free radicals or other radiation products; these errors make the reported yield too low. If, however, the determined product also is formed by the reaction of the scavenger with other free radicals formed by the radiation, the reported yield might be too high. If the radiation dose is too large, the product may begin to compete with the scavenger. It is generally difficult or impossible to estimate the accuracy of a determination of a free-ion yield by the scavenger method except by comparison with results of other methods.

3.2. Conductivity

The specific conductance of a system is proportional to the product of the number of ions formed and their mobility. If the mobility is known, or guessed, then a determination of the conductivity produced by a known intensity of radiation can be used to calculate the free-ion yield. For non-polar liquids, the method is difficult and not too reliable and has been superseded by the charge collection method. Used with a pulse radiolysis system in aqueous and alcoholic media, this method has given valuable results.

3.3. Charge Collection

Charge collection methods comprise steady-state and pulse methods.

In the steady-state method, the current of ions collected by the field is measured with the radiation intensity so low and the field so high that essentially no general ion recombination occurs as the ions are swept out by the field.

In the pulse or "clearing field" method the charge produced by a short pulse of radiation is collected by the field and measured. In one version, the field is turned on immediately after the radiation pulse to give a direct measurement of the zero-field free-ion yield; a ballasting condenser takes care of the charge necessary to bring the electrodes to voltage when the field is switched on directly after the radiation pulse. In another version, the field is maintained during the pulse and the zero-field value obtained by extrapolation.

In all the electrical methods, the collected currents or charges must be kept very small to avoid recombination errors, and it is difficult to avoid systematic errors in the electrical measurement systems. Also, difficulties in measuring the energy put into the liquid may occur at these very low dose rates.

Although careful determinations of G_{fi}° by charge collection should be accurate to $\pm 5\%$, values reported by different laboratories for the same pure liquid at the same temperature unfortunately often differ by many times this amount.

4. Tabulations of G_{fi}°

Tables 1-4 give values of G_{fi}° for a variety of pure liquids. When values were determined by a particular worker over a series of temperatures, only one value is usually given in the tables; values obtained over a range of temperatures are shown graphically in figures 1-6. For temperature effects in aromatic hydrocarbons, see [741110]. The tables give the name of the liquid, temperature of measurement, dielectric constant of the liquid, method used, reported value of G_{fi}° , and literature reference. The dielectric constants are omitted from the hydrocarbon tables because the values nearly all lie about 1.8-2. When no temperature is reported, "room temperature" may be assumed. Under "Methods" is indicated whether conductivity (condy.), charge collection (charge coll.) or scavenging (scav., followed by the formula or name of the scavenger used) was employed.

Data on mixtures of non-polar compounds are given in [680376] and [700055], but are not reproduced here. In mixtures of hydrocarbons, G_{fi} changes gradually with composition; but addition of a little CCl_4 to neopentane results in an abrupt drop of G_{fi}° . Relative values of G_{fi}° in mixtures of hexane and alcohols are reported in [710464] and [731090], mixtures of benzene and alcohols in [741084], mixtures of ethanol and water in [730143], and mixtures of alcohols and alkanes in [741035].

TABLE 1. Zero-field free-ion yields in saturated hydrocarbon liquids

Compound	Temp. ^{a,b} K	Method ^c	G_{H}^{d}	Reference
Methane	140	charge coll.	1.13	741012
Ethane	200	charge coll.	0.158	741012
Propane	238	charge coll.	0.166	741012
	233	charge coll.	0.143	720166
Cyclopropane	234	charge coll.	0.049	741012
Butane	293	charge coll.	0.225	741012
	296	charge coll.	0.193	700055
Isobutane	294	charge coll.	0.31	720166
<i>n</i> -Pentane	296	charge coll.	0.17	731054
	296	charge coll.	0.145	700055
Isopentane	296	charge coll.	0.170	700055
Neopentane	296	condy.	1.1	700090
	296	charge coll.	0.86	700055
	294	charge coll.	1.09	720269
	294	scav., N ₂ O	0.95	700650
Cyclopentane	296	charge coll.	0.155	700055
<i>n</i> -Hexane	296	condy.	0.10	650216
	297	condy.	0.10	660149
	—	condy.	0.137	680821
				recalc. in 720405
	296	condy.	0.11	690240
	296	scav., (C ₆ H ₅) ₃ CCl	0.12	690348
	—	scav., CH ₃ Br	0.12	700047
	296	charge coll.	0.131	700055
	298	scav., <i>c</i> -C ₆ F ₁₂	0.08	700299
	—	condy.	0.13	720405
	292	charge coll.	0.18	731054
	—	scav., C ₂ H ₅ OT	0.12	730242
2-Methylpentane	296	charge coll.	0.148	700055
3-Methylpentane	296	charge coll.	0.146	700055
	298	scav., <i>c</i> -C ₆ F ₁₂	0.17	700299
2,3-Dimethylbutane	296	charge coll.	0.192	700055
2,2-Dimethylbutane	296	charge coll.	0.304	700055
Cyclohexane	296	charge coll.	0.148	700055
	—	scav., CH ₃ Br	0.13	700047
	298	scav., <i>c</i> -C ₆ F ₁₂	0.14	700299
	296	scav., anthracene	0.17	700360
	296	scav., (C ₆ H ₅) ₃ CCl	0.13	690348
	296	scav., galvinoxyl	0.15	700019
	294	charge coll.	0.19	731054
Methylcyclohexane	293	scav., pyrene	0.122	730124
<i>n</i> -Heptane	296	charge coll.	0.131	700055
2,4-Dimethylpentane	296	charge coll.	0.178	700055
2,2,3-Trimethylbutane	296	charge coll.	0.290	700055
<i>n</i> -Octane	296	charge coll.	0.124	700055
2,3,4-Trimethylpentane	296	charge coll.	0.174	700055
2,2,4-Trimethylpentane	296	charge coll.	0.332	700055
	—	scav., CH ₃ Br	0.36	700047
	—	scav., <i>c</i> -C ₆ F ₁₂	0.37	700299
	296	scav., (C ₆ H ₅) ₃ CCl	0.36	690348
	296	scav., galvinoxyl	0.39	700019
	—	scav., CH ₃ Br	0.13	720032
	—	scav., C ₂ H ₅ OT	0.22	730242
	—	scav., CH ₃ Br	0.3	740092
2,2,3,3-Tetramethylbutane	379°	charge coll.	0.80	720269
<i>n</i> -Nonane	296	charge coll.	0.117	700055
2,2,4,4-Tetramethylpentane	295	charge coll.	0.83	720269
2,2,3,3-Tetramethylpentane	295	charge coll.	0.42	720269
<i>n</i> -Decane	296	charge coll.	0.117	700055

TABLE 1. Zero-field free-ion yields in saturated hydrocarbon liquids—Continued

Compound	Temp. ^{a,b} K	Method ^c	G_{fi}^d	Reference
2,2,5,5-Tetramethylhexane	293	charge coll.	0.67	720269
2,2,6,6-Tetramethylheptane	293	charge coll.	0.47	720269
2,2,7,7-Tetramethyloctane	316	charge coll.	0.34	720269
<i>n</i> -Tetradecane	296	charge coll.	0.120	700055
Squalane (C ₃₀ H ₆₂)	296	charge coll.	0.119	700055

^aWhen no temperature is given in the reference, room temperature may be assumed.

^bWhen determinations were made at several temperatures on the same compound, only one result is given here; the others are displayed in the accompanying figures.

^cAbbreviations: condy. = conductivity under irradiation combined with measured or assumed ion mobility values
charge coll. = charge collection by clearing field,
scav. = scavenging of the negative or positive charge by the compound indicated.

^dFree-ion yield in molecule/100 eV.

^eThis octane isomer melts at 374 K.

TABLE 2. Zero-field free-ion yields in unsaturated and aromatic hydrocarbon liquids

Compound	Temp. K	Method	G_{fi}^d	Reference
Ethylene	170	charge coll.	0.010	741012
Propylene	234	charge coll.	0.042	741012
Butene-1	293	charge coll.	0.093	731054
<i>cis</i> -Butene-2	293	charge coll.	0.23	731054
<i>trans</i> -Butene-2	293	charge coll.	0.080	731054
Isobutene	293	charge coll.	0.25	731054
2-Methylbutene-2	292	charge coll.	0.26	731054
Hexene-1	296	charge coll.	0.062	700055
	293	charge coll.	0.10	731054
Hexene-2 ^a	296	charge coll.	0.076	700055
<i>trans</i> -Hexene-2	293	charge coll.	0.092	731054
<i>cis</i> -Hexene-3	293	charge coll.	0.13	731054
<i>trans</i> -Hexene-3	293	charge coll.	0.10	731054
3,3-Dimethylbutene-1	296	charge coll.	0.169	700055
2,3-Dimethylbutene-2	293	charge coll.	0.44	731054
Cyclohexene	296	charge coll.	0.15	700055
	293	charge coll.	0.20	731054
Propadiene(Allene)	282	charge coll.	0.050	731054
Butadiene-1,3	269	charge coll.	0.038	731054
Pentadiene-1,4	293	charge coll.	0.067	731054
Cyclopentadiene	195	scav., NH ₃	0.05	652067
Hexadiene-1,5	292	charge coll.	0.066	731054
Heptadiene-1,6	292	charge coll.	0.067	731054
Octadiene-1,7	292	charge coll.	0.065	731054
Acetylene	198	charge coll.	0.02	731054
Propyne	260	charge coll.	0.17	731054
Butyne-2	293	charge coll.	0.32	731054
Hexyne-1	253	charge coll.	0.10	731054
Hexyne-2	293	charge coll.	0.19	731054
Hexyne-3	293	charge coll.	0.21	731054
Benzene	—	scav., tritiated water or alc.	0.077	690650
	—	scav., tritiated water or alc.	0.098	700349
	298	scav., galvinoxyl	0.055	700019
	296	charge coll.	0.053	700055
	292	charge coll.	0.081	741110
Toluene	298	scav., galvinoxyl	0.051	700019
	292	charge coll.	0.093	741110
<i>tert</i> -Butylbenzene	296	charge coll.	0.089	700055
<i>o</i> -Xylene	292	charge coll.	0.094	741110

TABLE 2. Zero-field free-ion yields in unsaturated and aromatic hydrocarbon liquids—Continued

Compound	Temp. K	Method	G_{fi}^0	Reference
<i>m</i> -Xylene	292	charge coll.	0.082	741110
<i>p</i> -Xylene	292	charge coll.	0.073	741110
1,2,3-Trimethylbenzene	292	charge coll.	0.12	741110
1,2,4-Trimethylbenzene	292	charge coll.	0.083	741110
1,3,5-Trimethylbenzene	293	charge coll.	0.087	741110
1,2,3,4-Tetramethylbenzene	292	charge coll.	0.11	741110
1,2,4,5-Tetramethylbenzene	358	charge coll.	0.15	741110
Pentamethylbenzene	353	charge coll.	0.25	741110
Hexamethylbenzene	448	charge coll.	0.6	741110
Naphthalene	376	charge coll.	0.11	741110
Anthracene	500	charge coll.	~0.1	741110
Biphenyl	373	pulse radiolysis, ion spectra	<0.35	690431

* *Cis-trans* mixture.

TABLE 3. Zero-field free-ion yields in miscellaneous non-polar liquids

Compound	Temp. K	ϵ	Method	G_{fi}^0	Reference
Tetramethylsilane	296	1.84	charge coll.	0.74	700055
Tetramethyltin	296	2.07	charge coll.	0.62	700055
Carbon disulfide	296	2.63	charge coll.	0.314	700055
	296		scav., (C ₆ H ₅) ₃ CCl	0.26	690348
Triethylamine	296	2.44	charge coll.	0.148	700055
Germanium tetrachloride	296	2.44	charge coll.	0.127	700055
Carbon tetrachloride	296	2.232	charge coll.	0.096	700055
	296		scav., (C ₆ H ₅) ₃ CCl	0.093	690348
	296		condy.	0.068	660148
1,4-Dioxane	295	2.21	condy.	0.05	650216
	296		condy.	0.038	660148
	—		scav., benzophenone	0.1	690521
	296		charge coll.	0.046	700055
	296		scav., anthracene	0.10	700360
	293	2.20	charge coll.	0.08	751047
	335	2.15	charge coll.	0.12	751047
	374	2.08	charge coll.	0.16	751047
Carbon tetrafluoride	143	1.64	charge coll.	0.07	730095
Perfluoro- <i>n</i> -pentane	296	1.68	charge coll.	0.035	700055
Perfluoromethylcyclohexane	296	1.85	charge coll.	0.028	700055
Oxygen	87	1.50	charge coll.	0.013	720034

TABLE 4. Zero-field free-ion yields in polar liquids ($\epsilon > 3$)

Compound	Temp. K	ϵ	Method	G_{fi}^0	Reference
Ethers:					
Di- <i>n</i> -butyl ether	—	3.1	condy.	0.11	650216
	187	4.50	charge coll.	0.21	751047
	220	3.99	charge coll.	0.23	751047
	255	3.52	charge coll.	0.26	751047
	293	3.14	charge coll.	0.30	751047
	332	2.83	charge coll.	0.36	751047
	373	2.55	charge coll.	0.42	751047
	406	2.35	charge coll.	0.48	751047
	436	2.20	charge coll.	0.54	751047

TABLE 4. Zero-field free-ion yields in polar liquids ($\epsilon > 3$) — Continued

Compound	Temp. K	ϵ	Method	G_{fi}^∞	Reference
Di- <i>n</i> -propyl ether	169	5.64	charge coll.	0.23	751047
	202	4.76	charge coll.	0.25	751047
	254	3.90	charge coll.	0.30	751047
	293	3.43	charge coll.	0.34	751047
	342	2.91	charge coll.	0.38	751047
	376	2.60	charge coll.	0.44	751047
Diisopropyl ether	296	3.9	charge coll.	0.305	700055
Diethyl ether	—	4.3	condy.	0.19	650216
	296		charge coll.	0.35	700055
	—		scav., CH ₃ Br	0.15	700249
	296		scav., anthracene	0.28	700360
	195	8.0	charge coll.	0.53	751047
	223	6.45	charge coll.	0.50	751047
	252	5.40	charge coll.	0.49	751047
	293	4.39	charge coll.	0.54	751047
	328	3.70	charge coll.	0.57	751047
	165	10.7	charge coll.	0.64	751047
	195	9.0	charge coll.	0.73	751047
	226	7.4	charge coll.	0.74	751047
Tetrahydropyran	231	7.28	charge coll.	0.45	751047
	254	6.60	charge coll.	0.48	751047
	293	5.60	charge coll.	0.51	751047
Tetrahydrofuran	296	7.5	scav., pyrene, anthracene	0.68	700406
	—		scav., benzophenone	0.5	690521
	—		scav., maleic anhydride	0.6 ^a	701042
	—		pulse radiolysis	0.39 ^b	731069
	—		scav., pyrene	0.3	741030
	205	11.6	charge coll.	0.72	751047
	296	7.9	scav., anthracene	0.23	700360
Halides:					
<i>n</i> -Butyl bromide	295	6.9	condy.	0.27	650216
<i>n</i> -Butyl chloride	295	7.2	condy.	0.39	650216
Fluoroform	183	21.4	charge coll.	1.1	730095
Methyl fluoride	183	27.9	charge coll.	1.6	730095
Difluoromethane	183	38.9	charge coll.	1.9	730095
Ketones:					
Methyl ethyl ketone	293	18.5	condy.	0.84	720153
Acetone	296	21	scav., anthracene	1.18	700360
	—		scav., anthracene	1.5	710186
	—		scav., stilbene and anthracene	1.1	710189
	—		condy.	1.2	720002
	296	46.6	scav., anthracene	1.62	700360
Dimethyl sulfoxide	—		scav., N ₂ O	1.8	710215
	—		scav., biphenyl	1.8	720232
	—		scav., anthracene and Br ⁻	1.2–1.8	731016
Esters:					
Methyl formate	296	8.5	scav., anthracene	0.32	700360
1,2-Propanediol carbonate	296	64.5	scav., anthracene	2.25	700360
Nitriles:					
Benzonitrile	296	25.2	scav., anthracene	1.40	700360
	296		scav., <i>trans</i> -stilbene	1.4	741100
Acetonitrile	296	37.5	scav., anthracene	1.55	700360
Pyridine	296	12	scav., anthracene	0.40	700360
Amines:					
Aniline	—	6.9	scav., pyrene	0.95	680227
Tributylamine	293	5.	condy.	0.25	720153
Butylamine	293	5.	condy.	0.27	720153
Diethylamine	—	3.6	scav., pyrene	0.1	741030
Ethylamine	—	6.2	scav., pyrene and (C ₆ H ₅) ₂	1.8	741094

TABLE 4. Zero-field free-ion yields in polar liquids ($\epsilon > 3$) — Continued

Compound	Temp. K	ϵ	Method	G_{fi}^0	Reference
Methylamine	—	9.4	scav., pyrene and $(C_6H_5)_2$	2.2	741094
Ammonia	258	20.	scav., pyrene and $(C_6H_5)_2$	3.3	741094
	296	17.	pulse radiolysis	3.2	741062
	225	25	pulse radiolysis	3.0	740159
	225		scav., Cu^{2+}	2.9	740159
Ethanolamine	—	57.7	scav., anthracene	0.2	731106
Alcohols:					
Neopentyl alcohol	337	8.3	scav., galvinoxyl	0.21	700019
<i>tert</i> -Butyl alcohol	333	7.8	scav., anthracene and pyrene	0.7	720574
	296	12.3	scav., galvinoxyl	0.67	700019
Hexyl alcohol	293	13.3	scav., anthracene and pyrene	1.1	720574
<i>n</i> -Pentyl alcohol	298	14.2	scav., galvinoxyl	0.45	700019
	337		scav., galvinoxyl	0.46	700019
Isopentyl alcohol	293	15.	scav., anthracene and pyrene	0.35	720574
	293		condy.	0.40	720153
<i>n</i> -Butyl alcohol	298	17.1	scav., galvinoxyl	0.63	700019
	298		scav., acetone	0.55	670004
	293		scav., anthracene and pyrene	0.56	720574
	293		condy.	0.58	720153
Isobutyl alcohol	293	17.7	scav., anthracene and pyrene	0.5	720574
Isopropyl alcohol	296	18.3	scav., anthracene	0.92	700360
	—		scav., $(C_6H_5)_2$	1.0	650499
	298		scav., N_2O	1.2	680042
	—		scav., NO_3^-	1.7	690655
	298		condy.	2.2	710064
	291		scav., tetranitromethane	1.2	710564
	293		scav., anthracene and pyrene	1.0	720574
<i>n</i> -Propyl alcohol	—	20.1	scav., $(C_6H_5)_2$	1.0	650499
	298		scav., N_2O	1.2	690387
	298		scav., acetone	0.65	670004
	298		condy.	2.2	710064
	150–298		pulse radiolysis	1.45	751054
Ethyl alcohol	—	24.3	scav., $(C_6H_5)_2$	1.0	640113
	298		scav., N_2O	1.5	680047
	296		scav., galvinoxyl	1.0	700019
	294		scav., anthracene	0.98	700360
	298		condy.	1.95	710064
	290		condy.	3.9°	710531
	296		scav., acetone, nitrobenzene	1.7	720199
	—		scav., $(C_6H_5)_2$	1.7	700172
Methyl alcohol	—	32.6	scav., $(C_6H_5)_2$	1.1	650499
	298		scav., acetone	1.3	670004
	—		scav., N_2O	2.0	670018
	296		scav., anthracene	1.1	700360
	298		scav., N_2O	2.0	680610
	298		condy.	2.0	710064
	—		condy.	1.9	731086
	296		scav., nitrobenzene and acetone	1.7	730292
Ethylene glycol	—	38.	scav., $(C_6H_5)_2$	1.2	650499
	293		scav., anthracene and pyrene	1.2	720574
Nitro compounds:					
Nitrobenzene	298	34.8	scav., galvinoxyl	0.43	700019
Nitromethane	298	37.8	scav., galvinoxyl	0.31	700019
Amides:					
Hexamethylphosphoramide	—	30	scav., biphenyl and anthracene	2.4	720105
	293		condy.	1.2	720153
	—		scav., anthracene and pyrene	2.3	731091
Dimethylformamide	296	37.	scav., anthracene	2.0	700360
Dimethylacetamide	296	38.	scav., anthracene	1.7	700360

TABLE 4. Zero-field free-ion yields in polar liquids ($\epsilon > 3$) – Continued

Compound	Temp. K	ϵ	Method	G_{fi}°	Reference
Formamide	—	109.	scav., N_2O	3.3	700173
<i>N</i> -Methylpropionamide	296	164.	scav., anthracene	2.6	700360
Water ^d (H_2O)	—	80	scav., $KMnO_4$	$G(e_{aq}^-) = 2.7$	670311
	—	—	review	$G(e_{aq}^-) = 2.7$	730418
	—	—	condy.	$G(OH^-) = 0.7$; $G_{fi}^\circ = G(H^+) = 3.4$	710138
	—	—	condy.	$G(OH^-) = 0.8$; $G_{fi}^\circ = G(H^+) = 3.5$	700243
	—	—	condy.	$G(OH^-) < 0.3$; $G_{fi}^\circ = 2.85 \pm 0.15$	720404
(D_2O)	—	—	scav., $KMnO_4$	$G(e_{aq}^-)(D_2O) = 2.9$	680061

^aMuch lower values were obtained with high-LET radiations.

^bAbsorption due to $e^- \cdot Na^+$ ion pair, extrapolated to zero Na^+ .

^cYield includes $G(\text{anions}) = 2.1$, in addition to solvated electrons, as deduced from decay kinetics; not observed by other workers.

^dYields change very little with temperature up to 523 K, C.J. Hochanadel, [550048].




FIGURE 1. G_{fi}° for *n*-pentane, *n*-hexane and cyclohexane.
Open symbols from [680376]; filled symbols, [720269].




FIGURE 2. G_{fi}° for low-molecular-weight saturated hydrocarbons.
Data from [741012, 720034 and 720166].




FIGURE 3. G_{fi}° for some saturated hydrocarbons, from [720269].




FIGURE 5. G_{fi}° for some dienes, from [731054].




FIGURE 4. G_{fi}° for some singly-unsaturated hydrocarbons, from [741012].




FIGURE 6. G_{fi}° for some alkynes, from [731054].

5. Remarks on G_{fi}° Data

5.1. Hydrocarbons

The chief property determining G_{fi}° in hydrocarbons is molecular shape. By and large, the more nearly spherical the molecules, the bigger G_{fi}° . Despite variations in G_{fi}° by a factor of up to 100, the temperature coefficients are quite comparable. The effect of double and triple bonds is relatively minor, although the difference between *cis*- and *trans*-butene-2 is surprising. Comparing results from the two laboratories responsible for most of the data, we find that the more recent data from Freeman's laboratory are consistently about 30% higher than the data on the same liquids by Schmidt and Allen. Further work is required to settle this and other discrepancies.

5.2 Other Non-polar Compounds

The near-spherical tetramethides have large values of G_{fi}° , but equally spherical carbon tetrachloride has a low value. This is attributed to the reactivity of CCl_4 with e^- , which seems to convert e^- to Cl^- before it emerges from the positive ion field. However, CF_4 , which does not ordinarily react readily with electrons [690179], also has a very low G_{fi}° . This seems to be an outstanding anomaly.

5.3. Polar Liquids

The yield of free solvated electrons in water has been the subject of dozens of papers, and only in recent years has agreement been obtained. The two papers cited in table 4 for $G(e_{aq}^-)$ in water contain references to earlier work. For effects of radiation quality (LET) on the yields see [610164] and [690185]. The free-ion yield also includes some OH^- which is shown by pulse conductivity studies; therefore, G_{fi}° is actually larger than G_e^- .

There is some controversy about yields of solvated electrons and free ions in methyl and ethyl alcohols. Earlier scavenger studies gave $G(e^-)$ about 1.0, but pulse conductivity gave values near 2. In the light of later work, it now appears that electron reaction with biphenyl and anthracene may have been underestimated because of the short lifetime of the product ions in alcohol, while in other cases too low a scavenger concentration was used. The two conductivity studies differ widely, however, on the total ion pair yield; [710531] found the yield of other anions to be larger than that of solvated electrons while [710064] found the anion yield less than 8% of the electron yield. Another disagreement involves the yield in tetrahydrofuran.

G_{fi}° should increase with increasing dielectric constant, which shortens the range of the Coulomb forces. Table 4 shows a strong tendency in this direction, but with many exceptions. The yield in ammonia, which is much larger than that in liquids of similar dielectric constant, seems connected with the slowness of reaction of the solvated electron with the conjugate acid ion NH_4^+ .

6. Total Ion Yields

Only in two polyatomic liquids has G_{fi} been determined to high enough fields to allow a reasonable extrapolation to infinite field. Table 5 shows that the extrapolated total yield for γ rays in neopentane and CS_2 is close to that measured in the gas phase. In the monatomic liquids Ar, Kr and Xe, nearly complete collection of the ions is possible, and the total ion yield in the liquid is found to be greater than in the gas. In the solid forms of these elements, optical absorptions are found by Baldini at longer wave-lengths than in the corresponding gas [629023] and it is believed that the resulting excited states spontaneously ionize, in the liquid as well as the solid state [719437]. If this is true, Baldini's absorptions give the effective ionization potential in the liquid, I_{liq} . Takahashi *et al.* [740153] found reasons for assuming that the ratio of W to I should be the same in the liquid and gas, and obtained the following calculated values for W_{liq} : Ar, 24.0; Kr, 20.1; Xe, 16.7. These calculated values agree with their measured values within the experimental error (3 to 8%).

TABLE 5. Total ion yields reported for selected liquids

Compound	Temp. K	Radiation Source	% satn. obtained	G_{fi}^*	W_{liq} (eV)	G_{liq}/G_{gas}	Reference
Carbon disulfide	Room	γ rays from Ra	80	3.8	26	~ 1	340008
Neopentane	296	^{60}Co γ rays	84	4.03	24.8	0.94	700090 ^a
Argon	87	20-kV or 50-kV X rays	95-100	3.9	26 ± 3	~ 1	660847
	87	Internal conversion e^- , from ^{207}Bi	~ 100	4.2	23.6	1.11	740029
	87	^{63}Ni β -rays	98	4.22	23.7	1.11	740153
	87	^{60}Co γ rays	100	4.5	22.5	1.18	690429
	87	1.7 MeV X-rays	100	7.3	13.7	1.9	730059
Krypton	148	1.7 MeV X-rays	98	13.0	7.7	3.2	730059
	120	^{63}Ni β -rays	91	4.9	20.5	1.17	740153
Xenon	183	1.7 MeV X-rays	~ 100	13.7	7.3	3.0	730059
	183	Po α -rays		5.8	17.3	1.27	730735
	166	^{63}Ni β -rays	90	6.1	16.4	1.33	740153

^a G_{gas} from 710122.

7. Theoretical Treatments

No good quantitative theory exists of the free-ion yield. Jaffé's theory [080002] was based on a columnar model which is unrealistic for low LET radiations and has not proved useful quantitatively. Recent treatments start with Onsager's formula for the probability $P(r)$ of escape by thermal agitation of a pair of ions separated by a distance r :

$$P(r) = (1 + f(E))\exp(-r_c/r)$$

where $r_c = e^2/\epsilon kT$, e is electronic charge, ϵ dielectric constant, k Boltzmann's constant, T the absolute temperature, and $f(E)$ is a function of the applied field expressed as a double summation of terms again involving ϵ and T . The model for hydrocarbons assumes that an electron very quickly loses energy to electronic and vibronic modes and attains a level of energy which is less than that of molecular vibrations but still greater than thermal. It then loses energy much more slowly by elastic collisions, and how far it travels before it is thermalized depends on how smooth a potential surface it finds, which in turn depends on how nearly spherical the liquid molecules are. Eventually it attains thermal energy at a distance r from the positive ion, which of course is not the same for all electrons. Some distribution function $D(r)$ must exist which gives the probability of the electron becoming thermalized at r , and the escape probability is then $\int D(r)P(r)dr$. If this model holds, the same function should determine both the field dependence and the temperature dependence of G_{fi} . Every worker who has in recent years seriously considered G_{fi} has expressed his results in terms of an assumed $D(r)$, which usually includes two parameters, the total yield G_{fi}^* and a "mean penetration range" called b or r_0 . Both the temperature coefficient of the zero-field free-ion yield and the field effect on the yield may be used to obtain the best value of the range r_0 defined by an assumed $D(r)$. Mozumder [748021] fitted experimental data on G_{fi} for 19 hydrocarbons and other organic liquids with five different assumed forms of $D(r)$ to obtain values of the range r_0 . For any particular liquid, the five values of r_0 ranged over a factor of 1.6 to 1.8. But all the calculated "mean" ranges are many times longer than anyone would have supposed prior to the analysis of the free-ion yields. Thus, the smallest r_0 values in Mozumder's results were obtained from a "truncated power law" $D(r) = (2.7/4\pi r_0^3) (r/r_0)^{-5.7}$ and ranged from 28 Å for 1,4-dioxane to 133 Å for neopentane. The majority of electrons formed in the ionization of liquids by α , β , γ or X-rays have initial energies below 100 eV. Extrapolation of the usual "electron ranges" obtained by experiments on kilovolt electrons suggests maximum ranges < 1 Å for 100-eV electrons in hydrocarbons [660148]. The usual "electron range" determination consists in finding how thick a foil must be inserted in an electron beam to prevent ionization in air beyond the foil; the foil is thus only sufficient to reduce the electron energy below the ionization threshold (15-20 eV). The extra distance traveled in liquids is required to reduce the electron energy from a few eV to thermal levels. This distance depends strongly on the shape of the molecules.

The ranges in dielectric liquids deduced from free-ion yields agree in order of magnitude with values obtained by photoionization of solute molecules [710638] and metal surfaces [720439]. Recent work on electron scattering in metals has shown similar effects; the "mean free path" in the metal, which decreases with decreasing energy at higher energies, shows a sharp upturn at electron energies below 40 eV [749139].

The trouble with the Onsager model is that it does not conform well with physical reality, for only about 40% of the ions are formed in isolated pairs. The remainder occur in groups, usually known as "spurs", which contain from 2 to 50 or more ionizations, located close together. The electrons formed by ionization in such a group are attracted back by the collective charge of all the ions in the group, partially shielded by the charge on all the other electrons. Obviously the Onsager formula cannot apply to such a vague and complex situation. When the field is not too high and the escape probability from a spur remains low, the parameter r_0 may be rationalized as the mean distance between the positive and negative charges in the last pair to remain in a spur at the moment that all the other pairs have recombined. When the field is high enough so that an appreciable number of spurs contribute more than one free-ion pair, the use of the Onsager formula cannot be justified. Dodelet and Freeman [758020] have proposed a model of the spur which, together with an assumed $D(r)$, gives a reasonable fit to the data on field effects in such diverse liquids as xenon and water. The values obtained for the parameters G_{fi} and r_0 are not however very different from those obtained (much more simply) by not taking the spurs into account. The physical significance of such values remains doubtful as long as $D(r)$ is taken arbitrarily, which it must be until the process of electron slowing-down in the low energy range is better understood.

We do not include any quoted values of the "penetration range" in our tables of data, because the numbers have no precise quantitative significance. Nevertheless, the r_0 values listed in any paper on free-ion yields, have great qualitative meaning. It is clear that electrons of somewhat superthermal energy must lose their momentum more slowly, in passing through a liquid, the more nearly isotropic are the molecules of the liquid. The same behavior of thermal electrons is indicated by mobility measurements of thermal electrons in hydrocarbon liquids, although quantitative correlation between G_{fi} and mobility has not yet been made. A great deal of information remains to be extracted from G_{fi} data.

References

- [080002] Jaffé, G., Ueber die Ionisation fluessiger Dielektrika durch Radiumstrahlen, ANN. PHYSIK. 25: 257-84 (1908).
- [340008] Mohler, F.L.; Taylor, L.S., Ionization of liquid carbon disulphide by X-rays, J. RES. NAT. BUR. STD. 13: 659-64 (1934).
- [550048] Hochanadel, C.J., Proc. of the Int. Conf. Peaceful Uses Atomic Energy, Geneva, Aug. 8-20, 1955, Vol. 7, p.521-5, N.Y., United Nations, 1956.
- [610164] Allen, A.O., The radiation chemistry of water and aqueous solutions, D. Van Nostrand Co., Inc., New York, 1961, p.58.
- [629023] Baldini, G., Ultraviolet absorption of solid argon, krypton, and xenon, PHYS. REV. 128(4): 1562-7 (1962).
- [640113] Taub, I.A.; Harter, D.A.; Sauer, M.C.; Dorfman, L.M., Pulse radiolysis studies. IV. The solvated electron in the aliphatic alcohols, J. CHEM. PHYS. 41(4): 979-85 (1964).
- [650216] Freeman, G.R.; Fayadh, J.M., Influence of the dielectric constant on the yield of free ions produced during radiolysis of a liquid, J. CHEM. PHYS. 43(1): 86-92 (1965).
- [650499] Sauer, M.C., Jr.; Arai, S.; Dorfman, L.M., Pulse radiolysis studies. VII. The absorption spectra and radiation chemical yields of the solvated electron in the aliphatic alcohols, J. CHEM. PHYS. 42(2): 708-12 (1965).
- [652067] Bonin, M.A.; Busler, W.R.; Williams, F., The polymerization of cyclopentadiene by free ions. Determination of the propagation rate constant, J. AMER. CHEM. SOC. 87(2): 199-207 (1965).
- [660148] Hummel, A.; Allen, A.O.; Watson, F.H., Jr., Ionization of liquids by radiation. II. Dependence of the zero-field ion yield on temperature and dielectric constant, J. CHEM. PHYS. 44(9): 3431-6 (1966).
- [660149] Hummel, A.; Allen, A.O., Ionization of liquids by radiation. I. Methods for determination of ion mobilities and ion yields at low voltage, J. CHEM. PHYS. 44(9): 3426-31 (1966).
- [660847] Ullmaier, H.A., Measurement of the ionization in liquid argon due to X-rays, PHYS. MED. BIOL. 11(1): 95-101 (1966).
- [670004] Basson, R.A.; van der Linde, H.J., Polarity effects in the radiolysis of n-alcohols, J. CHEM. SOC. PT. A (1): 28-32 (1967).
- [670018] Seki, H.; Imamura, M., Radiolysis yields from γ -irradiated liquid methanol containing nitrous oxide and the effect of acid, J. PHYS. CHEM. 71(4): 870-75 (1967).
- [670311] Fielden, E.M.; Hart, E.J., Primary radical yields in

- pulse-irradiated alkaline aqueous solution, *RADIAT. RES.* 32(3): 564-80 (1967).
- [680042] Russell, J.C.; Freeman, G.R., Reactions of the primary reducing species in the radiolysis of liquid 2-propanol, *J. PHYS. CHEM.* 72(3): 808-15 (1968).
- [680047] Russell, J.C.; Freeman, G.R., The yields of the primary reducing species in the radiolysis of liquid ethanol, *J. PHYS. CHEM.* 72(3): 816-21 (1968).
- [680061] Fielden, E.M.; Hart, E.J., Primary radical yields and some rate constants in heavy water, *RADIAT. RES.* 33: 426-36 (1968).
- [680227] Cooper, R.; Thomas, J.K., Nanosecond-pulse radiolysis of aniline, *J. CHEM. PHYS.* 48(11): 5103-9 (1968).
- [680376] Schmidt, W.F.; Allen, A.O., Yield of free ions in irradiated liquids: determination by a clearing field, *J. PHYS. CHEM.* 72(11): 3730-6 (1968).
- [680610] Jha, K.N.; Freeman, G.R., I. Kinetics of reactions of electrons during radiolysis of liquid methanol. II. Reaction of electrons with liquid alcohols and with water, *J. CHEM. PHYS.* 48(12): 5480-90 (1968).
- [680821] Jacobi, W.; Jahns, A., Ionization and recombination in dielectric fluids after γ -irradiation, *Molecular Structure and Radiation Effects*, Glubrecht, H. (ed.), Georg Thieme Verlag, Stuttgart, 1968, p.25-35.
- [690179] Rajbenbach, L.A., Radiolysis of solutions of perfluorocarbons in n-hexane, *J. PHYS. CHEM.* 73(2): 356-60 (1969).
- [690185] Appleby, A.; Schwarz, H.A., Radical and molecular yields in water irradiated by γ rays and heavy ions, *J. PHYS. CHEM.* 73: 1937-41 (1969).
- [690240] Conrad, E.E.; Silverman, J., Short-lived transient species in irradiated n-hexane, *J. CHEM. PHYS.* 51: 450-1 (1969).
- [690348] Capellos, C.; Allen, A.O., Ionization of liquids by radiation studied by the method of pulse radiolysis. II. Solutions of triphenylmethyl chloride, *J. PHYS. CHEM.* 73(10): 2364-8 (1969).
- [690387] Jha, K.N.; Freeman, G.R., Effect of temperature on the reactions of electrons during the γ radiolysis of liquid n-propanol, *J. CHEM. PHYS.* 51(7): 2839-46 (1969).
- [690429] Klassen, N.V.; Schmidt, W.F., Ionization of liquid argon by X-rays: W-value, *CAN. J. CHEM.* 47(22): 4286-8 (1969).
- [690431] Theard, L.M.; Peterson, F.C.; Holroyd, R.A., Nanosecond pulse radiolysis of liquid naphthalene and biphenyl, *J. CHEM. PHYS.* 51(9): 4126-32 (1969).
- [690521] Beaumont, D.; Rodgers, M.A.J., Pulse radiolysis studies of ion association, *TRANS. FARADAY SOC.* 65: 2973-80 (1969).
- [690650] Kroh, J.; Hankiewicz, E., Ionic processes in the radiolysis of liquid benzene as studied by tritium substitution, *INT. J. RADIAT. PHYS. CHEM.* 1(4): 451-6 (1969).
- [690655] Hecquet, M.F.; Roux, J.C.; Simonoff, G.N., Radiolyse de l'isopropanol et des melanges eau-isopropanol, *INT. J. RADIAT. PHYS. CHEM.* 1(4): 529-40 (1969).
- [700019] Capellos, C.; Allen, A.O., Ionization of liquids by radiation studied by the method of pulse radiolysis. III. Solutions of galvinoxyl radical, *J. PHYS. CHEM.* 74(4): 840-5 (1970).
- [700047] Warman, J.M.; Rza, S.J., Relevance of the concentration dependence of methyl radical formation from methyl bromide solutions to the distribution of ion-pair separation distances in irradiated hydrocarbons, *J. CHEM. PHYS.* 52(2): 485-90 (1970).
- [700055] Schmidt, W.F.; Allen, A.O., Free-ion yields in sundry irradiated liquids, *J. CHEM. PHYS.* 52(5): 2345-51 (1970).
- [700090] Schmidt, W.F., Radiation-induced conductivity and ion yields in neopentane at high electric fields, *RADIAT. RES.* 42: 73-8 (1970).
- [700172] Fletcher, J.W.; Richards, P.J.; Seddon, W.A., Pulse radiolysis of ethanol, *CAN. J. CHEM.* 48(11): 1645-50 (1970).
- [700173] Head, D.A.; Walker, D.C., Yield of free-ions in the radiolysis of formamide, a liquid of very high dielectric constant, *CAN. J. CHEM.* 48(11): 1657-63 (1970).
- [700243] Barker, G.C.; Fowles, F.; Sammon, D.C.; Stringer, B., Pulse radiolytic induced transient electrical conductance in liquid solutions. Part 1. Technique and the radiolysis of water, *TRANS. FARADAY SOC.* 66(6): 1498-508 (1970).
- [700249] Bansal, K.M.; Rza, S.J., Electron scavenging in the γ radiolysis of liquid diethyl ether, *J. PHYS. CHEM.* 74(15): 2888-92 (1970).
- [700299] Sagert, N.H.; Reid, J.A., γ -Radiolysis of hydrocarbons using perfluorocyclohexane as an electron scavenger: determination of the free ion yields, *CAN. J. CHEM.* 48: 2429-32 (1970).
- [700349] Kroh, J.; Burzynska, E., The influence of LET on the radiation yield of free ions in benzene as measured by tritium substitution, *RADIOCHEM.-RADIOANAL. LETT.* 4(3): 119-22 (1970).
- [700360] Hayon, E., Yield of ions and excited states produced in the radiolysis of polar organic liquids, *J. CHEM. PHYS.* 53(6): 2353-8 (1970).
- [700406] Baxendale, J.H.; Beaumont, D.; Rodgers, M.A.J., Electrons, ions and excited states in the pulse radiolysis of tetrahydrofuran, *TRANS. FARADAY SOC.* 66: 1996-2003 (1970).
- [700650] Horacek, K.; Freeman, G.R., Kinetics of electron and ion reactions during the γ radiolysis of liquid neopentane: dependence of charge scavenging kinetics on hydrocarbon molecular structure, *J. CHEM. PHYS.* 53(12): 4486-9 (1970).
- [701042] Matsui, M.; Imamura, M.; Karasawa, T., LET effect on the hydrogen formation in the radiolysis of liquid tetrahydrofuran, *IPCR Cyclotron Progress Report*, 1970, Vol. 4, Dec., p.126-7.
- [710064] Fowles, P., Pulse radiolytic induced transient electrical conductance in liquid solutions. Part 4. The radiolysis of methanol, ethanol, 1-propanol and 2-propanol, *TRANS. FARADAY SOC.* 67(2): 428-39 (1971).
- [710122] Stoneham, T.A.; Ethridge, D.R.; Meisels, G.G., Precise measurements of W, the mean energy required for ion-pair formation in gases. I. Alkanes, *J. CHEM. PHYS.* 54(9): 4054-9 (1971).
- [710138] Rabani, J.; Graetzel, M.; Chaudhri, S.A.; Beck, G.; Henglein, A., The radiation chemical yield of OH⁻

- as determined by conductometric pulse radiolysis, J. PHYS. CHEM. 75(11): 1759-60 (1971).
- [710186] Arai, S.; Kira, A.; Imamura, M., Pulse radiolysis studies of acetone solutions of biphenyl and anthracene: Formation of ions and excited singlet state, J. CHEM. PHYS. 54(12): 5073-81 (1971).
- [710189] Rodgers, M.A.J., Pulse radiolysis studies of acetone solutions, TRANS. FARADAY SOC. 67(4): 1029-40 (1971).
- [710215] Cooper, T.K.; Walker, D.C., Free-ion yield in the radiolysis of dimethylsulfoxide, CAN. J. CHEM. 49(13): 2248-53 (1971).
- [710464] Brown, B.J.; Barker, N.T.; Sangster, D.F., The behavior of the solvated electron in ethanol-n-hexane mixtures, J. PHYS. CHEM. 75(23): 3639-40 (1971).
- [710531] Rabani, J.; Graetzel, M.; Chaudhri, S.A., Conductometric pulse radiolysis determination of ionic yields and neutralization kinetics in liquid ethanol, J. PHYS. CHEM. 75(25): 3893-4 (1971).
- [710564] Asmus, K.-D.; Chaudhri, S.A.; Nazhat, N.B.; Schmidt, W.F., Pulse radiolysis studies of tetranitromethane in iso-propanol, TRANS. FARADAY SOC. 67(9): 2607-17 (1971).
- [710638] Jarnagin, R.C., Photoionization processes in organic solids and fluids, ACC. CHEM. RES. 4(12): 420-7 (1971).
- [719437] Raz, B.; Jortner, J., Energy of the quasi-free electron state in dense neon, CHEM. PHYS. LETT. 9(3): 222-24 (1971).
- [720002] Chaudhri, S.A.; Asmus, K.-D., Ion yields and ion neutralization processes in pulse-irradiated acetone, J. PHYS. CHEM. 76(1): 26-31 (1972).
- [720032] Rzađ, S.J.; Bansal, K.M., Radiolysis of liquid 2,2,4-trimethylpentane. Kinetics of scavenging processes, J. PHYS. CHEM. 76(17): 2374-80 (1972).
- [720034] Fuochi, P.G.; Freeman, G.R., Molecular structure effects on the free-ion yields and reaction kinetics in the radiolysis of the methyl-substituted propanes and liquid argon: Electron and ion mobilities, J. CHEM. PHYS. 56(5): 2333-41 (1972).
- [720105] Nauta, H.; van Huis, C., Pulse radiolysis of hexamethylphosphoric triamide, J. CHEM. SOC. FARADAY TRANS. I 68(4): 647-52 (1972).
- [720153] Vannikov, A.V.; Mal'tzev, E.I.; Zolotarevsky, V.I.; Rudnev, A.V., Transient charged radiolysis products of organic liquid studied by electrical and optical methods, INT. J. RADIAT. PHYS. CHEM. 4: 135-48 (1972).
- [720166] Dodelet, J.-P.; Fuochi, P.G.; Freeman, G.R., Effect of electric field strength on the free ion yields in the X-radiolysis of liquids: Influence of molecular structure and temperature, CAN. J. CHEM. 50(11): 1617-26 (1972).
- [720199] Jha, K.N.; Freeman, G.R., Effect of pressure on the rates of reaction of solvated electrons in liquid ethanol, J. CHEM. PHYS. 57(4): 1408-14 (1972).
- [720232] Bensasson, R.; Land, E.J., Transient species in the pulse radiolysis of dimethyl sulphoxide, CHEM. PHYS. LETT. 15(2): 195-8 (1972).
- [720269] Dodelet, J.-P.; Freeman, G.R., Mobilities and ranges of electrons in liquids: Effect of molecular structure in C₅-C₁₂ alkanes, CAN. J. CHEM. 50(16): 2667-79 (1972).
- [720404] Schmidt, K.H., Electrical conductivity techniques for studying the kinetics of radiation-induced chemical reactions in aqueous solutions, INT. J. RADIAT. PHYS. CHEM. 4(4): 439-68 (1972).
- [720405] Terlecki, J.; Fiutak, J., Onsager's recombination theory applied to liquid alkanes, INT. J. RADIAT. PHYS. CHEM. 4(4): 469-78 (1972).
- [720439] Holroyd, R.A.; Dietrich, B.K.; Schwarz, H.A., Ranges of photoinjected electrons in dielectric liquids, J. PHYS. CHEM. 76(25): 3794-800 (1972).
- [720574] Rudnev, A.V.; Vannikov, A.V.; Bakh, N.A., Optical absorption spectra and mobility of solvated electrons in aliphatic alcohols, HIGH ENERGY CHEM. 6(5): 416-7 (1972), Transl. from Khim. Vys. Energ. 6(5): 473-5 (1972).
- [730059] Robinson, M.G.; Freeman, G.R., X-radiolysis ion yields and electron ranges in liquid xenon, krypton, and argon: Effect of electric field strength, CAN. J. CHEM. 51(5): 641-9 (1973).
- [730095] Robinson, M.G.; Freeman, G.R., Radiolysis free ion yields and electron ranges in polar and nonpolar liquids: Fluoromethanes, CAN. J. CHEM. 51(7): 1010-5 (1973).
- [730124] Baxendale, J.H.; Bell, C.; Wardman, P., Observations on solvated electrons in aliphatic hydrocarbons at room temperature by pulse radiolysis, J. CHEM. SOC. FARADAY TRANS. I 69(4): 776-86 (1973).
- [730143] Roux, J.C.; Baquey, Ch.; Sutton, J., The radiolysis of binary ethanol-water mixtures, INT. J. RADIAT. PHYS. CHEM. 5: 309-22 (1973).
- [730242] Kroh, J.; Piekarska, J.; Karolczak, S., Ionic reactions during the radiolysis of liquid n-hexane and 2,2,4-trimethylpentane, BULL. ACAD. POL. SCI. SER. SCI. CHIM. 21(4): 279-83 (1973).
- [730292] Jha, K.N.; Freeman, G.R., Effect of pressure on the rates of reaction of solvated electrons and hydrogen atoms in liquid methanol, J. AMER. CHEM. SOC. 95(18): 5891-6 (1973).
- [730418] Sehested, K.; Bjerbakke, E.; Fricke, H., The primary species yields in the ⁶⁰Co γ-ray radiolysis of aqueous solutions of H₂SO₄ between pH 7 and 0.46, RADIAT. RES. 56(2): 385-99 (1973).
- [730735] Konno, S.; Kobayashi, S., Ionization of liquid xenon and argon by α-particles, SCI. PAPERS INST. PHYS. CHEM. RES. (TOKYO) 67(2): 57-62 (1973).
- [731016] Cooper, T.K.; Walker, D.C.; Gillis, H.A.; Klassen, N.V., The pulse radiolysis of dimethylsulfoxide and binary mixtures with water, CAN. J. CHEM. 51(13): 2195-206 (1973).
- [731054] Dodelet, J.-P.; Shinsaka, K.; Kortsch, U.; Freeman, G.R., Electron ranges in liquid alkenes, dienes, and alkynes: Range distribution function in hydrocarbons, J. CHEM. PHYS. 59(5): 2376-86 (1973).
- [731069] Bockrath, B.; Dorfman, L.M., Kinetics of electron transfer from aromatic radical anions to alkyl halides in tetrahydrofuran. Effects of sodium cation pairing, J. PHYS. CHEM. 77(22): 2618-22 (1973).
- [731086] Busi, F.; Ward, M.D., On the molar extinction coefficient of the solvated electron and the G(free

- ion) yield in liquid methanol, INT. J. RADIAT. PHYS. CHEM. 5: 521-3 (1973).
- [731090] Brown, B.J.; Barker, N.T.; Sangster, D.F., Electron solvation processes in alcohol-hydrocarbon mixtures, AUST. J. CHEM. 26(10): 2089-99 (1973).
- [731091] Shaede, E.A.; Dorfman, L.M., Flynn, G.J.; Walker, D.C., Spectrum, kinetics, and radiation chemical yield of solvated electrons in hexamethylphosphoric triamide, CAN. J. CHEM. 51(23): 3905-13 (1973).
- [731106] Marevtsev, V.S.; Vannikov, A.V., Optical absorption spectra of the solvated electron in irradiated monoethanolamine, HIGH ENERGY CHEM. 7(2): 104-8 (1973), Transl. from Khim. Vys. Energ. 7(2): 119-24 (1973).
- [740029] Miyajima, M.; Takahashi, T.; Konno, S.; Hamada, T.; Kubota, S.; Shibamura, H.; Doke, T., Average energy expended per ion pair in liquid argon, PHYS. REV. PT. A 9(3): 1438-43 (1974).
- [740092] Kroh, J.; Piekarska, J., Non-homogeneous kinetics of ionic reactions in liquid n-hexane and 2,2,4-trimethylpentane, BULL. ACAD. POL. SCI. SER. SCI. CHIM. 22(2): 167-72 (1974).
- [740153] Takahashi, T.; Konno, S.; Doke, T., The average energies, W, required to form an ion pair in liquefied rare gases, J. PHYS. C: SOLID STATE PHYS. 7(1): 230-40 (1974).
- [740159] Belloni, J.; Cordier, P.; Delaire, J., Transient spectrum from nanosecond pulse radiolysis of liquid ammonia, CHEM. PHYS. LETT. 27(2): 241-4 (1974).
- [741012] Robinson, M.G.; Freeman, G.R., Electron mobilities and ranges in liquid C₁-C₃ hydrocarbons and in xenon: Effects of temperature and field strength, CAN. J. CHEM. 52(3): 440-6 (1974).
- [741030] Shaede, E.A.; Kurihara, H.; Dorfman, L.M., Observation of initial ion-pairs in pulse-irradiated weakly polar liquids, INT. J. RADIAT. PHYS. CHEM. 6: 47-54 (1974).
- [741035] Brandon, J.R.; Firestone, R.F., Solvated electrons in alcohol-alkane and alcohol-amine solutions, J. PHYS. CHEM. 78(8): 792-6 (1974).
- [741062] Farhataziz; Perkey, L.M.; Hentz, R.R., Pulse radiolysis of liquids at high pressures. V. Absorption spectrum and yield of the solvated electron in liquid ammonia at 23°C and pressures up to 6.7 kbar, J. CHEM. PHYS. 60(11): 4383-9 (1974).
- [741084] Brown, B.J.; Barker, N.T.; Sangster, D.F., Pulse-radiolysis of binary liquid alcohol systems containing benzene, AUST. J. CHEM. 27(5): 1129-32 (1974).
- [741094] Seddon, W.A.; Fletcher, J.W.; Sopchyshyn, F.C., Jevcak, J., Pulse radiolysis of liquid methylamine, ethylamine, and ammonia: the yield of solvated electrons, CAN. J. CHEM. 52: 3269-73 (1974).
- [741100] Kira, A.; Thomas, J.K., Formation of ions and excited states in the pulse radiolysis of benzonitrile, J. PHYS. CHEM. 78(21): 2094-8 (1974).
- [741110] Shinsaka, K.; Freeman, G.R., Epithermal electron ranges and thermal electron mobilities in liquid aromatic hydrocarbons, CAN. J. CHEM. 52(20): 3495-506 (1974).
- [748021] Mozumder, A., Effect of an external electric field on the yield of free ions. II. The initial distribution of ion pairs in liquid hydrocarbons, J. CHEM. PHYS. 60(11): 4305-10 (1974).
- [749139] Ertl, G.; Kueppers, J., Low energy electrons and surface chemistry, Verlag Chemie G.m.b.H., Weinheim, 1974, Chapter 1.
- [751047] Dodelet, J.-P.; Freeman, G.R., Electron mobilities and ranges in liquid ethers: ion-like and conduction band mobilities, CAN. J. CHEM. 53(9): 1263-74 (1975).
- [751054] Dixon, R.S.; Lopata, V.J., Solvated electron yields in pulse-irradiated liquid 1-propanol, J. CHEM. PHYS. 62(11): 4573-4 (1975).
- [758020] Dodelet, J.-P.; Freeman, G.R., Electric field effect on free ion yields in x-irradiated liquids: a multipair spur model, INT. J. RADIATION PHYS. CHEM. 7(2/3): 183-92 (1975).

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET		1. PUBLICATION OR REPORT NO. NBS-NSRDS 57	2. Gov't Accession No.	3. Recipient's Accession No.
4. TITLE AND SUBTITLE Yields of Free Ions Formed in Liquids by Radiation			5. Publication Date April 1976	
			6. Performing Organization Code	
7. AUTHOR(S) A. O. Allen			8. Performing Organ. Report No.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, D.C. 20234			10. Project/Task/Work Unit No.	
			11. Contract/Grant No.	
12. Sponsoring Organization Name and Complete Address (Street, City, State, ZIP) Same as Item 9			13. Type of Report & Period Covered N/A	
			14. Sponsoring Agency Code	
15. SUPPLEMENTARY NOTES Library of Congress Catalog Card Number: 75-619361				
16. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here.) Free ions are those produced by ionizing radiation which escape initial recombination. Yields of free-ion pairs are tabulated for liquid alkanes, alkenes, alkynes, dienes and aromatic hydrocarbons, and other non-polar and polar compounds, including alcohols, ethers, esters, halides, amines, nitriles, etc. Total ion yields for a few liquids are compared with gas phase ion yields. Theoretical treatments of these data are discussed.				
17. KEY WORDS (six to twelve entries; alphabetical order; capitalize only the first letter of the first key word unless a proper name; separated by semicolons) Electrons; free ions; ion yields; liquid; non-polar systems; polar systems; radiation chemistry; radiation physics.				
18. AVAILABILITY <input checked="" type="checkbox"/> Unlimited <input type="checkbox"/> For Official Distribution. Do Not Release to NTIS <input checked="" type="checkbox"/> Order From Sup. of Doc., U.S. Government Printing Office Washington, D.C. 20402, SD Cat. No. C13. 48:57 <input type="checkbox"/> Order From National Technical Information Service (NTIS) Springfield, Virginia 22151		19. SECURITY CLASS (THIS REPORT) UNCLASSIFIED		21. NO. OF PAGES 25
		20. SECURITY CLASS (THIS PAGE) UNCLASSIFIED		22. Price .55

get
a line on
science and
technology.
subscribe to

DIMENSIONS



Whether you're in business, or a teacher, scientist, or consumer, you'll want to keep up with the latest developments in science and technology. DIMENSIONS/NBS, the monthly magazine from the Commerce Department's National Bureau of Standards, can help keep you informed. Every day at NBS, one of the nation's largest research laboratories, scientists seek new answers to a host of national problems, including energy conservation, product safety, metric conversion, and pollution abatement. Their findings, reported each month in DIMENSIONS/NBS, have a direct impact on our daily lives.

Subscription price: \$9.45 per year.
Order prepaid from the
Superintendent of Documents,
U.S. Government Printing Office,
Washington, D.C. 20402
SD Catalog No. C13.13

Announcement of New Publications on Standard Reference Data

Superintendent of Documents,
Government Printing Office,
Washington, D.C. 20402

Dear Sir:

Please add my name to the announcement list of new publications to be issued in the series:
National Standard Reference Data Series—National Bureau of Standards.


Name _____

Company _____

Address _____

City _____ State _____ Zip Code _____

(Notification Key N337)



Important Additions

TO THE JOURNAL OF PHYSICAL AND CHEMICAL REFERENCE DATA

Two comprehensive reference volumes, each, like the Journal itself, published by The American Institute of Physics, The National Bureau of Standards and The American Chemical Society . . . your triple assurance of their accuracy, immediacy, and usefulness.

SUPPLEMENT NO. 1 TO VOLUME 2

"Physical and Thermodynamic Properties of Aliphatic Alcohols"

by R. C. Wilhoit and B. J. Zwolinski, *Thermodynamics Research Center, Department of Chemistry, Texas A&M University.*

The most exhaustive review and critical analysis of selected physical and thermodynamic properties of aliphatic alcohols that has been published. Coverage of the important properties of the liquid, vapor, and ideal gaseous states as well as pertinent solid state data necessary for equilibrium calculations ordinarily encountered in chemical and chemical engineering applications. All available quantitative data on each property for each aliphatic alcohol are fully documented and critically analyzed providing a "data bank" for the 640 monohydroxy alcohols in the carbon range of C₁ to C₅₀. Internally consistent tables of critical, standard, or selected "best" values are tabulated for each compound. Index included.

SUPPLEMENT NO. 1 TO VOLUME 3

"Thermal Conductivity of the Elements: A Comprehensive Review"

by C. Y. Ho, R. W. Powell, and P. E. Lilly, *1974 Thermophysical Properties Research Center, Purdue University.*

Comprehensive review of the world's thermal conductivity data presents recommended or estimated values for all 105 elements. Reliable thermal

conductivity data for those elements which can be used as standard reference materials to calibrate or check apparatus. Original data, specimen characterization and measurement information for 5200 sets of raw data. Detailed discussions for each element, review of the available experimental data and considerations by which the authors arrived at final assessments. Complete bibliographic citations for 1658 references. Only original sources have been used. All cited documents are available at TPRC in standard microfiche.

Special Issue Sales

American Chemical Society
1155 16th St., N.W., Washington, D. C. 20036

Please send _____ copies of the 420 page supplement to the Journal of Physical and Chemical Reference Data, Volume 2, "Physical and Thermodynamic Properties of Aliphatic Alcohols," at the prices checked below.

Members: Hard cover edition: ☐ \$33.00
Nonmembers: Hard cover edition: ☐ \$33.00
Members: Soft cover edition: ☐ \$10.00
Nonmembers: Soft cover edition: ☐ \$30.00

Check or money order must accompany order. \$1.00 extra for foreign postage and handling.

Name _____

Address _____

City _____ State _____ Zip _____

Please send _____ copies of the 796 page supplement to the Journal of Physical and Chemical Reference Data, Volume 3, "Thermal Conductivity of the Elements: A Comprehensive Review," at the prices checked below.

Members: Hard cover edition: ☐ \$60.00
Nonmembers: Hard cover edition: ☐ \$60.00
Members: Soft cover edition: ☐ \$25.00
Nonmembers: Soft cover edition: ☐ \$55.00



American Chemical Society

NBS TECHNICAL PUBLICATIONS

PERIODICALS

JOURNAL OF RESEARCH reports National Bureau of Standards research and development in physics, mathematics, and chemistry. It is published in two sections, available separately:

- **Physics and Chemistry (Section A)**

Papers of interest primarily to scientists working in these fields. This section covers a broad range of physical and chemical research, with major emphasis on standards of physical measurement, fundamental constants, and properties of matter. Issued six times a year. Annual subscription: Domestic, \$17.00; Foreign, \$21.25.

- **Mathematical Sciences (Section B)**

Studies and compilations designed mainly for the mathematician and theoretical physicist. Topics in mathematical statistics, theory of experiment design, numerical analysis, theoretical physics and chemistry, logical design and programming of computers and computer systems. Short numerical tables. Issued quarterly. Annual subscription: Domestic, \$9.00; Foreign, \$11.25.

DIMENSIONS/NBS (formerly *Technical News Bulletin*)—This monthly magazine is published to inform scientists, engineers, businessmen, industry, teachers, students, and consumers of the latest advances in science and technology, with primary emphasis on the work at NBS. The magazine highlights and reviews such issues as energy research, fire protection, building technology, metric conversion, pollution abatement, health and safety, and consumer product performance. In addition, it reports the results of Bureau programs in measurement standards and techniques, properties of matter and materials, engineering standards and services, instrumentation, and automatic data processing.

Annual subscription: Domestic, \$9.45; Foreign, \$11.85.

NONPERIODICALS

Monographs—Major contributions to the technical literature on various subjects related to the Bureau's scientific and technical activities.

Handbooks—Recommended codes of engineering and industrial practice (including safety codes) developed in cooperation with interested industries, professional organizations, and regulatory bodies.

Special Publications—Include proceedings of conferences sponsored by NBS, NBS annual reports, and other special publications appropriate to this grouping such as wall charts, pocket cards, and bibliographies.

Applied Mathematics Series—Mathematical tables, manuals, and studies of special interest to physicists, engineers, chemists, biologists, mathematicians, computer programmers, and others engaged in scientific and technical work.

National Standard Reference Data Series—Provides quantitative data on the physical and chemical properties of materials, compiled from the world's literature and critically evaluated. Developed under a world-wide

program coordinated by NBS. Program under authority of National Standard Data Act (Public Law 90-396).

NOTE: At present the principal publication outlet for these data is the *Journal of Physical and Chemical Reference Data* (JPCRD) published quarterly for NBS by the American Chemical Society (ACS) and the American Institute of Physics (AIP). Subscriptions, reprints, and supplements available from ACS, 1155 Sixteenth St. N. W., Wash. D. C. 20056.

Building Science Series—Disseminates technical information developed at the Bureau on building materials, components, systems, and whole structures. The series presents research results, test methods, and performance criteria related to the structural and environmental functions and the durability and safety characteristics of building elements and systems.

Technical Notes—Studies or reports which are complete in themselves but restrictive in their treatment of a subject. Analogous to monographs but not so comprehensive in scope or definitive in treatment of the subject area. Often serve as a vehicle for final reports of work performed at NBS under the sponsorship of other government agencies.

Voluntary Product Standards—Developed under procedures published by the Department of Commerce in Part 10, Title 15, of the Code of Federal Regulations. The purpose of the standards is to establish nationally recognized requirements for products, and to provide all concerned interests with a basis for common understanding of the characteristics of the products. NBS administers this program as a supplement to the activities of the private sector standardizing organizations.

Federal Information Processing Standards Publications (FIPS PUBS)—Publications in this series collectively constitute the Federal Information Processing Standards Register. Register serves as the official source of information in the Federal Government regarding standards issued by NBS pursuant to the Federal Property and Administrative Services Act of 1949 as amended, Public Law 89-306 (79 Stat. 1127), and as implemented by Executive Order 11717 (38 FR 12315, dated May 11, 1973) and Part 6 of Title 15 CFR (Code of Federal Regulations).

Consumer Information Series—Practical information, based on NBS research and experience, covering areas of interest to the consumer. Easily understandable language and illustrations provide useful background knowledge for shopping in today's technological marketplace.

NBS Interagency Reports (NBSIR)—A special series of interim or final reports on work performed by NBS for outside sponsors (both government and non-government). In general, initial distribution is handled by the sponsor; public distribution is by the National Technical Information Service (Springfield, Va. 22161) in paper copy or microfiche form.

Order NBS publications (except NBSIR's and Bibliographic Subscription Services) from: Superintendent of Documents, Government Printing Office, Washington, D.C. 20402.

BIBLIOGRAPHIC SUBSCRIPTION SERVICES

The following current-awareness and literature-survey bibliographies are issued periodically by the Bureau:

Cryogenic Data Center Current Awareness Service

A literature survey issued biweekly. Annual subscription: Domestic, \$20.00; foreign, \$25.00.

Liquefied Natural Gas. A literature survey issued quarterly. Annual subscription: \$20.00.

Superconducting Devices and Materials. A literature

survey issued quarterly. Annual subscription: \$20.00. Send subscription orders and remittances for the preceding bibliographic services to National Bureau of Standards, Cryogenic Data Center (275.02) Boulder, Colorado 80302.

Electromagnetic Metrology Current Awareness Service Issued monthly. Annual subscription: \$24.00. Send subscription order and remittance to Electromagnetics Division, National Bureau of Standards, Boulder, Colo. 80302.

U.S. DEPARTMENT OF COMMERCE
National Bureau of Standards
Washington, D.C. 20234

OFFICIAL BUSINESS

Penalty for Private Use, \$300

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-215



SPECIAL FOURTH-CLASS RATE
BOOK



75 YEARS
NBS
1901-1976