

A11102 146420

NAT'L INST. OF STANDARDS & TECH R.I.C.



A11102146420

Sleck, L. Wayne/Rate coefficients for Io
QC100 .U573 V64:1979 C.2 NBS-PUB-C 1979



NSRDS-NBS 64

U.S. DEPARTMENT OF COMMERCE / National Bureau of Standards



Rate Coefficients for Ion-Molecule Reactions

Organic Ions Other Than Those Containing Only C and H

QC

100

.U573

No. 64

1979

C.2

NATIONAL BUREAU OF STANDARDS

The National Bureau of Standards¹ was established by an act of Congress March 3, 1901. The Bureau's overall goal is to strengthen and advance the Nation's science and technology and facilitate their effective application for public benefit. To this end, the Bureau conducts research and provides: (1) a basis for the Nation's physical measurement system, (2) scientific and technological services for industry and government, (3) a technical basis for equity in trade, and (4) technical services to promote public safety. The Bureau's technical work is performed by the National Measurement Laboratory, the National Engineering Laboratory, and the Institute for Computer Sciences and Technology.

THE NATIONAL MEASUREMENT LABORATORY provides the national system of physical and chemical and materials measurement; coordinates the system with measurement systems of other nations and furnishes essential services leading to accurate and uniform physical and chemical measurement throughout the Nation's scientific community, industry, and commerce; conducts materials research leading to improved methods of measurement, standards, and data on the properties of materials needed by industry, commerce, educational institutions, and Government; provides advisory and research services to other Government Agencies; develops, produces, and distributes Standard Reference Materials; and provides calibration services. The Laboratory consists of the following centers:

Absolute Physical Quantities² — Radiation Research — Thermodynamics and Molecular Science — Analytical Chemistry — Materials Science.

THE NATIONAL ENGINEERING LABORATORY provides technology and technical services to users in the public and private sectors to address national needs and to solve national problems in the public interest; conducts research in engineering and applied science in support of objectives in these efforts; builds and maintains competence in the necessary disciplines required to carry out this research and technical service; develops engineering data and measurement capabilities; provides engineering measurement traceability services; develops test methods and proposes engineering standards and code changes; develops and proposes new engineering practices; and develops and improves mechanisms to transfer results of its research to the ultimate user. The Laboratory consists of the following centers:

Applied Mathematics — Electronics and Electrical Engineering² — Mechanical Engineering and Process Technology² — Building Technology — Fire Research — Consumer Product Technology — Field Methods.

THE INSTITUTE FOR COMPUTER SCIENCES AND TECHNOLOGY conducts research and provides scientific and technical services to aid Federal Agencies in the selection, acquisition, application, and use of computer technology to improve effectiveness and economy in Government operations in accordance with Public Law 89-306 (40 U.S.C. 759), relevant Executive Orders, and other directives; carries out this mission by managing the Federal Information Processing Standards Program, developing Federal ADP standards guidelines, and managing Federal participation in ADP voluntary standardization activities; provides scientific and technological advisory services and assistance to Federal Agencies; and provides the technical foundation for computer-related policies of the Federal Government. The Institute consists of the following divisions:

Systems and Software — Computer Systems Engineering — Information Technology.

¹Headquarters and Laboratories at Gaithersburg, Maryland, unless otherwise noted; mailing address Washington, D.C. 20234.

²Some divisions within the center are located at Boulder, Colorado, 80303.

National Bureau of Standards
APR 12 1979
NOT ACC-Circ
GC100
.US\$2
no. 64
1979
c 2

Rate Coefficients for Ion-Molecule Reactions

Organic Ions Other than Those Containing Only C and H

L. Wayne Sieck

National Measurement Laboratory
National Bureau of Standards
Washington, DC 20234



T
NSRDS-NBS #64

U.S. DEPARTMENT OF COMMERCE, Juanita M. Kreps, Secretary

Jordan J. Baruch, Assistant Secretary for Science and Technology

US NATIONAL BUREAU OF STANDARDS, Ernest Ambler, Director

Issued February 1979

Library of Congress Cataloging in Publication Data

Sieck, L Wayne.

Rate coefficients for ion-molecule reactions.

(Nat. stand. ref. data ser.; NSRDS-NBS 64)

"National Standard Reference Data System."

Supt. of Docs. no.: C 13.48:64

1. Chemical reaction, Rate of. 2. Ions. 3. Molecules.

4. Chemistry, Physical organic. I. Title. II. Series: United States.

National Bureau of Standards. National standard reference data series

; NSRDS-NBS 64.

QD502.S52 547'.139 78-10710

NSRDS-NBS 64

Nat. Stand. Ref. Data Ser., Nat. Bur. Stand. (U.S.), 64, 27 pages (Feb. 1979)

© 1979 by the Secretary of Commerce on Behalf of the United States Government

U.S. GOVERNMENT PRINTING OFFICE
WASHINGTON: 1979

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402

Stock No. 003-003-02027-3 - Price \$1.30

(Add 25 percent additional for other than U.S. mailing).

Foreword

The National Standard Reference Data System provides access to the quantitative data of physical science, critically evaluated and compiled for convenience and readily accessible through a variety of distribution channels. The System was established in 1963 by action of the President's Office of Science and Technology and the Federal Council for Science and Technology, and responsibility to administer it was assigned to the National Bureau of Standards.

NSRDS receives advice and planning assistance from a Review Committee of the National Research Council of the National Academy of Sciences-National Academy of Engineering. A number of Advisory Panels, each concerned with a single technical area, meet regularly to examine major portions of the program, assign relative priorities, and identify specific key problems in need of further attention. For selected specific topics, the Advisory Panels sponsor subpanels which make detailed studies of users' needs, the present state of knowledge, and existing data resources as a basis for recommending one or more data compilation activities. This assembly of advisory services contributes greatly to the guidance of NSRDS activities.

The System now includes a complex of data centers and other activities in academic institutions and other laboratories. Components of the NSRDS produce compilations of critically evaluated data, reviews of the state of quantitative knowledge in specialized areas, and computations of useful functions derived from standard reference data. The centers and projects also establish criteria for evaluation and compilation of data and recommend improvements in experimental techniques. They are normally associated with research in the relevant field.

The technical scope of NSRDS is indicated by the categories of projects active or being planned: nuclear properties, atomic and molecular properties, solid state properties, thermodynamic and transport properties, chemical kinetics, and colloid and surface properties.

Reliable data on the properties of matter and materials are a major foundation of scientific and technical progress. Such important activities as basic scientific research, industrial quality control, development of new materials for building and other technologies, measuring and correcting environmental pollution depend on quality reference data. In NSRDS, the Bureau's responsibility to support American science, industry, and commerce is vitally fulfilled.

A handwritten signature in black ink, appearing to read "Ernest Ambler". The signature is fluid and cursive, with "Ernest" on the left and "Ambler" on the right, separated by a small gap.

ERNEST AMBLER, Director

Contents

	Page
Foreword	III
1. Introduction	1
2. Reaction Rate Tables	2
Table 1. Rate Coefficients for Reactions of Organic Cations	4
Table 2. Rate Coefficients for Reactions of Organic Anions	17
Acknowledgements	19
References	19

Rate Coefficients for Ion-Molecule Reactions.

Organic Ions Other Than Those Containing Only C and H

L. Wayne Sieck

National Measurement Laboratory, National Bureau of Standards, Washington, DC 20234

A compilation is presented of all experimentally determined bimolecular and termolecular rate coefficients for the reactions of organic ions (other than those containing only C and H) with neutral molecules in the vapor phase. The literature covered is from 1960 to the present, and both positive and negative ions are considered. Five hundred and seventy-seven separate reaction-partners are tabulated, and experimental conditions are specified wherever possible. Preferred values are suggested for a number of these processes.

Key words: Chemical kinetics; data evaluation; gas phase ion-molecule reactions; mass spectrometry; organic molecules; rate coefficients.

1. Introduction

During the past decade, the detailed investigation of the dynamics, rates, and mechanisms associated with the interactions of ions with molecules in the vapor phase has been characterized by an almost exponential growth. The current widespread interest in these processes can be attributed to their recognized role in the upper atmosphere, combustion systems, in materials exposed to high energy radiation, and to their application in industrial chemical synthesis and trace analysis, especially in air and water pollution. This is the second in a series of compilations and evaluations of rate coefficients for ion-molecule reactions occurring at thermal or near thermal ion kinetic energies. The first segment, which dealt exclusively with the reactions of organic ions containing only C and H, was published in 1976 [1]¹. This portion deals with the reactivities of all other organic ions (other than those containing C and H only). Ions which constitute "borderline" organics, such as HCN^+ , CS_2^+ , CS_2^- , and CO_3^- , CO_4^- , and CN^- , as well as protonated CO and CO_2 , will be considered in a subsequent article.

Our goal in preparing these articles is twofold. First, we wish to provide a comprehensive, up-to-date compilation of those bimolecular and termolecular ion-molecule reactions and the associated rate constants (coefficients) recorded under reasonably well-specified experimental conditions at ion kinetic energies ≤ 0.5 eV. We have restricted our entries to those data expressed in units of $\text{cm}^3/\text{molecule}\cdot\text{s}$ or $\text{cm}^6/\text{molecule}^2\cdot\text{s}$, which encompass more than 95 percent of the

recent measurements carried out at thermal or nearly thermal energies. Second, since rate coefficients have now been determined under a variety of experimental conditions it is possible, in individual cases, to subject the combined literature to critical evaluation and to suggest a preferred value. We feel that this objective is of prime importance since many investigators use reference reactions for instrumental calibration purposes. Unfortunately, depending upon the particular laboratory, different values have been assumed in many cases for the same calibration reaction, and in several instances the value chosen by a given laboratory has changed with time. One would hope that the assignment of a preferred value with appropriate error limits would provide a common reference base for use by the various research groups.

The following journals were searched for entries from 1960 to the cut-off date, approximately 1 January 1978.

- Journal of Chemical Physics
- Journal of Physical Chemistry
- Journal of the American Chemical Society
- Journal of Research of the National Bureau of Standards
- The International Journal of Mass Spectrometry and Ion Physics
- International Journal of Chemical Kinetics
- Canadian Journal of Chemistry
- Chemical Physics Letters
- Transactions of the Faraday Society

Other pertinent entries were found by scanning Current Contents, Advances in Mass Spectrometry, and the published proceedings of various symposia and meet-

¹Figures in brackets indicate literature references at the end of this paper.

ings dealing with this subject. Unpublished results, as well as any which were not subjected to prior critical review via the normal scientific refereeing channels, are not included. In spite of the fact that every effort has been made to extract all of the relevant kinetic data from the literature it is possible, in an undertaking of this magnitude, that pertinent references may have been inadvertently overlooked. Any omissions which may have accidentally occurred are not intended to reflect the authors' judgments concerning the validity of individual rate measurements or the potential accuracy of the experimental technique used by any particular laboratory.

1.1. Evaluation of Data

With respect to error limits and the reliability of data, it is often difficult to make an objective judgment concerning the validity of a particular rate coefficient. However, it is our feeling that greater than 98 percent of the tabulated values for fast reactions are most certainly correct to within a factor of 2 of the true value at the stated temperature and/or kinetic energy. Difficulties often arise in assigning more stringent limits. For example, one of the perplexing aspects of the chemistry associated with complex ions is the fact that the reactivity of these species may depend critically upon the internal energy content of the reactant ion. Whenever specific reactions have been shown to exhibit such effects it is so stated in the compilation. Second, many of the older values reflect the reactivities of ions which were translationally excited due to the presence of electric fields within the reaction zone. Although this condition would not necessarily affect the total reactivity of any given ion, it is well established that resultant product distributions are often seriously affected. Moreover, in many cases the original authors did not properly state their experimental conditions, particularly with respect to the temperature of the bulk gas. Even when the temperature was stated it is unlikely that the neutral flow component had achieved equilibrium, especially in measurements carried out at reduced pressures. Taken together, these complications, as well as others discussed elsewhere [2,3], introduce a large and often undefinable uncertainty in a number of the tabulated values.

When the accuracy limits given by the original authors seem appropriate within the framework of the possible sources of error, the original error limits are cited. Otherwise, either a reasonable estimate is applied or limits are simply not stated. In general, the most

recent values tend to be the most accurate for any given reaction-pair.

2. Reaction Rate Tables

Two tables are presented, table 1 covering positive ion reactions and table 2, negative ion reactions. The entries in the various columns are described as follows:

2.1. Reaction

The reactions are listed sequentially according to the following format:

(a) Carbon number of the reactant ion; i.e., all of the C₁'s appear first.

(b) Increasing atomic number (Z) of the highest Z element. A C₁ ion containing only H and B (regardless of the number of boron atoms) would therefore appear before CHNH⁺, C₂F₆⁺ would be listed before C₂H₄FBr⁺, and so forth.

(c) When the reactivity of any single ion towards two or more neutrals has been reported, the ordering sequence reflects the hierarchy of the neutral within the procedure outlined in (a) and (b).

Only the empirical formula of the reactant ion is given. The neutral from which the ion was derived, or the structure suggested by the original author, is given under the column entitled Comments (see below). The name or the structural formula of the neutral reactant is always given under the Reaction column. In order to conserve space, the abbreviations *n* (normal), *i* (iso), *t* (tertiary), and *c* (cyclo) are also sometimes used to define structural isomers. Due to the complexity of the product distributions associated with many of the reactions which are listed, as well as the fact that the branching ratios are often sensitive to changes in temperature, pressure, and internal energy, the reaction products are not specified. Moreover, in many cases only the total reactivity of a given ion towards a given neutral was reported, without specifying reaction products.

2.2. Rate Coefficient (*k*)

Rate coefficients are given in units of cm³ molecule⁻¹ s⁻¹ for bimolecular reactions and cm⁶ molecule⁻² s⁻¹ for third order processes. Unless otherwise stated, the error limits given are those imposed by the original authors (see Introduction for discussion). The most recent experimental determination is listed first, the other following in reverse chronological order. When a given laboratory has reported two or more values, or a corrected value, for a given reaction using the same

technique, then only the more recent or corrected value is reported.

2.3. Temperature (*T*)

When the system temperature was specified, then that value is given in degrees K. All ion cyclotron resonance measurements may be assumed to have been carried out at, or slightly above, ambient laboratory temperatures (295 + 15, -5 K).

2.4. Experimental Methods

The nomenclature used to describe the various measurements techniques is similar to that used by Ferguson [4].

Method	See below for short description
DT	Drift Tube
FA	Flowing Afterglow
ICR	Ion Cyclotron Resonance
MS	Mass Spectrometer Ion Source
B	Beam Apparatus
TI	Ion Trap

a. Drift Tube (DT)

This is a relatively new technique which yields rate coefficients from thermal to several electron volts kinetic energy. It has been mainly applied to the reactions of inorganic ions of aeronomic interest.

b. Flowing Afterglow (FA)

This versatile method, which utilizes a buffer gas at relatively high pressures, assures that the reactant ions are essentially always in thermodynamic equilibrium prior to reaction. The gas temperature is variable over a considerable range (up to 900 K). FA has been especially useful for the determination of equilibrium constants and other thermodynamic quantities pertinent to ionic association and particle transfer reactions as well as reactions of aeronomic importance.

c. Ion Cyclotron Resonance (ICR)

Ion Cyclotron Resonance mass spectrometry is a low pressure technique (10^{-7} to 10^{-3} Nm $^{-2}$) in which the reaction time is usually the variable experimental parameter. Most of the rate data have been obtained at nearly thermal kinetic energies and at ambient room temperature.

d. Mass Spectrometer Ion Source (MS)

This is a catchall category used to denote measurements involving a single reaction chamber associated with a mass analysis system. The application of this technique varies considerably from laboratory to laboratory. Some groups work in the pressure range 10^{-3} to 10^{-1} Nm $^{-2}$, with or without pulsed chambers, while

others have extended the pressure range up to nearly one atmosphere. Both photoionization and electron impact have been utilized to produce reactant ions. Many of the earlier low pressure studies involved the application of an electric field across the reaction zone for the purposes of ion extraction. In these cases the reactant ions were continuously accelerated, giving an epithermal kinetic energy distribution.

e. Beam (B)

Beam experiments involve the generation of a mass and energy selected ion beam which is impacted on a neutral target in a collision chamber coupled with a second mass analyzer. The kinetic energy may be varied from nearly thermal values up to tens of kilovolts.

f. Ion Trap (TI)

This relatively new technique involves the trapping of ions for a variable period of time in the negative space charge of a continuous low energy electron beam. The reactant ions may have kinetic energies in excess of thermal values, in some cases up to several tenths of an electron volt.

In many cases characterization of an experiment as being of the Beam type, Mass Spectrometer ion source type, etc., is not sufficient for defining the exact measurement technique. The reader should always refer to the specific literature citation associated with the quoted rate coefficient for details of the variation used to generate any particular value.

2.5. Comments

This column is used to provide descriptive information pertinent to the measurement such as "*k* varies with temperature, photoionization," etc. When the quoted value of *k* for any particular reaction was determined relative to an assumed value for a calibration reaction, then the assumed value for the calibration reaction is given. For ions containing two or more C atoms, one or more structural isomers may be present in any given experiment. In every case where the identity of the reactant ion is not obvious, we have listed either the neutral from which the ion was derived or the probable structural formula. The following abbreviations are also used: KE = ion kinetic energy, IEE = ion exit energy, and V cm $^{-1}$ to denote the magnitude of the imposed potential within the reaction chamber. Several of the ICR values were determined by pressure variation. When such measurements are reported, the entry "Pressure varied" is included. Comments pertinent to our choice of a preferred value are also given in this column.

TABLE I. Rate coefficients^a for reactions of organic cations

	Reaction	T	k	Method	Comments	Ref.
CH_2N^+	+ $\text{CH}_3\text{NH}_2 \rightarrow$	370	1.8(-9) ^b	MS	13eV e^- , 3.4eV IEE	5
		370	2.4(-9)	MS	15eV e^- , 3.4eV IEE	5
		370	1.6(-9)	MS	50eV e^- , 3.4eV IEE	5
			9±1(-10)	MS		6
	+ $(\text{CH}_3)_2\text{NH} \rightarrow$		5(-10)	MS		6
	+ $c\text{-C}_3\text{H}_5\text{NH}_2 \rightarrow$		8.0±0.2(-10)	ICR		7
	+ $(\text{CH}_3)_3\text{N} \rightarrow$		5(-10)	MS		6
	CH_3N^+ + $\text{CH}_3\text{NH}_2 \rightarrow$	370	2.6(-9)	MS	13eV e^- , 3.4eV IEE	5
		370	2.6(-9)	MS	15eV e^- , 3.4eV IEE	5
		370	1.9(-9)	MS	50eV e^- , 3.4eV IEE	5
			9±1(-10)	MS		6
CH_4N^+	+ $\text{CH}_3\text{NH}_2 \rightarrow$	295	1.1±0.1(-9)	MS	Photoionization	8
		370	1.8(-9)	MS	13eV e^- , 3.4eV IEE	5
	+ $\text{CH}_3\text{NH}_2 \rightarrow$	370	1.8(-9)	MS	15eV e^- , 3.4eV IEE	5
		370	1.2(-9)	MS	50eV e^- , 3.4eV IEE	5
			7±1(-10)	MS		6
	+ $(\text{CH}_3)_2\text{NH} \rightarrow$		6±2(-10)	MS		6
	+ $\text{C}_2\text{H}_5\text{NH}_2 \rightarrow$	298	1.97±0.2(-9)	MS	Photoionization	9
	+ $n\text{-C}_8\text{H}_7\text{NH}_2 \rightarrow$	298	1.56±0.2(-9)	MS	Photoionization	9
	+ $(\text{CH}_3)_3\text{N} \rightarrow$		4(-10)	MS		6
	CH_5N^+ + $\text{CH}_3\text{NH}_2 \rightarrow$		1.64(-9)	ICR	Time varied	10
			2.06(-9)	ICR	Pressure varied	10
		295	1.21±0.1(-9)	MS	Photoionization	8
		370	2.7(-9)	MS	13eV e^- , 3.4eV IEE	5
		370	2.6(-9)	MS	15eV e^- , 3.4eV IEE	5
		370	1.9(-9)	MS	50eV e^- , 3.4eV IEE	5
			9±1(-10)	MS		6
$\text{CH}_2\text{D}_3\text{N}^+$	+ $\text{CD}_3\text{NH}_2 \rightarrow$		1.85(-9)	ICR	From CD_3NH_2	10
CH_6N^+	+ $2\text{CH}_3\text{NH}_2 \rightarrow$	250	3.0(-26)	MS	CH_3NH_3^+	11
		350	1.6(-26)	MS	CH_3NH_3^+	11
CH_5DN^+	+ $\text{CH}_3\text{NH}_2 \rightarrow$		6.4±0.2(-10)	TI	$\text{CH}_5\text{NH}_2\text{D}^+$	12
$\text{CH}_2\text{D}_4\text{N}^+$	+ $\text{CD}_3\text{NH}_2 \rightarrow$		4.8(-10)	ICR	$\text{CD}_3\text{NH}_2\text{D}^+$	10
			3.1(-10)	ICR	$\text{CD}_3\text{NH}_2\text{D}^+$	10
CH_3N_2^+	+ $\text{CH}_3\text{N}_2\text{CH}_3 \rightarrow$		7.8±1.0(-10)	ICR		13
CH_2O^+	+ $\text{CH}_2\text{O} \rightarrow$		1.3(-10)	ICR		14
			1.77(-9)	ICR		15
		373	1.93±0.02(-9)	MS	3.4eV IEE	16
		373	2.00±0.01(-9)	MS	2.4eV IEE	16
	+ $\text{CH}_3\text{OH} \rightarrow$	373	2.03±0.1(-9)	MS	Thermal ions	17
		373	1.40±0.2(-9)	MS	3.4eV IEE	17
CH_3O^+	+ $(\text{CH}_3)_3\text{B} \rightarrow$		2.2±0.4(-10)	ICR		18
	+ $\text{CH}_2\text{O} \rightarrow$		6.7(-11)	ICR		14
	+ $\text{CH}_3\text{OH} \rightarrow$	373	2.11±0.3(-9)	MS	11eV e^- , Thermal	17
		373	1.68±0.2(-9)	MS	50eV e^- , Thermal	17
		373	1.58±0.2(-9)	MS	11eV e^- , 3.4eV IEE	17
		373	8.0±2.0(-10)	MS	50eV e^- , 3.4eV IEE	17
			6.8(-10)	TI		19
	+ $\text{CH}_3\text{CHO} \rightarrow$		1.80(-9)	ICR		15
	+ Ethylene oxide →		1.16±0.15(-9)	MS		20
	+ $\text{HCO}_2\text{CH}_3 \rightarrow$		2.0±0.2(-9)	MS		21
	+ Dimethylsulfoxide →	408	4.5±0.5(-9)	MS	0.71eV IEE	22
		408	4.5±0.5(-9)	MS	1.07eV IEE	22
CH_2DO^+	+ $\text{DCO}_2\text{CH}_3 \rightarrow$		1.9±0.2(-9)	MS	2.4eV IEE	21
CHD_2O^+	+ $\text{CD}_3\text{OH} \rightarrow$	373	1.44(-9)	MS	From CD_3OH , thermal	17
CH_3O^+	+ $\text{CH}_3\text{OH} \rightarrow$		2.17(-9)	ICR	Pressure varied, CH_3OH^+	10
			2.08(-9)	ICR	Time varied, CH_3OH^+	10
			1.86(-9)	ICR	CH_3OH^+	15
			1.90(-9)	ICR	CH_3OH^+	23

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction		T	k	Method	Comments	Ref.
CH_4O^+	$+ \text{CH}_3\text{OH} \rightarrow$	373	$2.53 \pm 0.2(-9)$	MS	11eV e^- , thermal, CH_3OH^+	17
		373	$2.47 \pm 0.2(-9)$	MS	50eV e^- , thermal, CH_3OH^+	17
		373	$2.56 \pm 0.4(-9)$	MS	11eV e^- , 3.4eV IEE , CH_3OH^+	17
		373	$2.5 \pm 0.2(-9)$	MS	50eV e^- , 3.4eV IEE , CH_3OH^+	17
		450	$2.0(-9)$	MS	9.1 V cm^{-1} , CH_3OH^+	24
		420	$2.45(-9)$	MS	0.86 V cm^{-1} , CH_3OH^+	19
					See footnote c for further comment.	
	$+ \text{HCO}_2\text{CH}_3 \rightarrow$	370	$2.54 \pm 0.1(-9)$	MS	2.4eV IEE , CH_3OH^+	21
	$+ 2\text{-pentanone} \rightarrow$		$7.5(-10)$	B	0.3eV KE , CH_3OH^+	25
	CH_3DO^+	$+ \text{CH}_3\text{OH} \rightarrow$	450	$2.0(-9)$	MS	9.1 V cm^{-1} , CH_3OD^+
CHD_3O^+	$+ \text{DCO}_2\text{CH}_3 \rightarrow$	370	$2.41 \pm 0.1(-9)$	MS	2.4eV IEE , CH_3OD^+	21
	$+ \text{CD}_3\text{OH} \rightarrow$		$2.56(-9)$	ICR	CD_3OH^+	10
		373	$2.19(-9)$	MS	Thermal, CD_3OH^+	17
		450	$2.0(-9)$	MS	0.9eV IEE , CD_3OH^+	24
CH_3O^+	$+ \text{CH}_3\text{OH} \rightarrow$		$1.1(-10)$	ICR	Time varied, CH_3OH_2^+	10
			$1.0(-10)$	ICR	Pressure varied, CH_3OH_2^+	10
			$8(-11)$	ICR	CH_3OH_2^+	15
	$+ \text{CD}_3\text{OH} \rightarrow$		$1.5(-10)$	ICR	Pressure varied, CD_3OH_2^+	10
$\text{CH}_2\text{D}_3\text{O}^+$	$+ \text{CD}_3\text{OH} \rightarrow$		$1.3(-10)$	ICR	Time varied, CD_3OH_2^+	10
	$+ \text{CD}_3\text{OH} \rightarrow$		$4.4(-10)$	ICR	Pressure varied, CD_3OHD^+	10
			$4.2(-10)$	ICR	Time varied, CD_3OHD^+	10
			$4.0 \pm 0.5(-10)$	TI	CD_3OHD^+	12
CDO_2^+	$+ \text{DCO}_2\text{H} \rightarrow$	370	$1.72 \pm 0.1(-9)$	MS	2.4eV IEE	21
CH_2O_2^+	$+ \text{H}_2\text{O} \rightarrow$	174	$3(-10)$	MS		26
		200	$3(-10)$	MS		26
		280	$3(-10)$	MS		26
CHDO_2^+	$+ \text{DCO}_2\text{H} \rightarrow$	370	$1.59 \pm 0.1(-9)$	MS	2.4eV IEE	21
CH_4NO_2^+	$+ 2 \text{CH}_3\text{OH} \rightarrow$		$1.9 \pm 0.4(-27)$	MS	$(\text{CH}_3\text{OH})\text{NO}^+$	27
CF^+	$+ \text{CH}_3\text{F} \rightarrow$	373	$4.7(-10)$	MS	0.24eV IEE	28
		373	$4.8(-10)$	MS	0.65eV IEE	28
	$+ \text{HF}\text{CO} \rightarrow$		$5(-11)$	ICR		29
	$+ \text{CH}_2\text{F}_2 \rightarrow$	320	$1.12(-9)$	MS		30
	$+ \text{CF}_4 \rightarrow$	320	$3.3(-10)$	MS		30
	$+ \text{CHCl}_2\text{F} \rightarrow$		$4(-10)$	MS	0.2eV KE	31
	$+ \text{CCl}_4 \rightarrow$		$5.9(-10)$	MS	0.2eV KE	31
	$+ \text{C}_2\text{H}_2\text{F}_2 \rightarrow$	320	$1.12(-9)$	MS		30
	$+ \text{C}_2\text{F}_6 \rightarrow$	295	$1.9 \pm 0.4(-10)$	MS	Photoionization	32
			$3.9(-11)$	B	0.3eV KE	33
CHF^+	$+ \text{C}_3\text{F}_8 \rightarrow$		$7.1(-11)$	B	0.3eV KE	34
	$+ 1,1,2,2,\text{-}c\text{-C}_4\text{H}_4\text{F}_4 \rightarrow$		$1.78(-10)$	MS	Average of two methods	35
	$+ \text{SiD}_4 \rightarrow$		$1.1 \pm 0.1(-10)$	B	1.0eV KE	36
	$+ \text{CH}_3\text{F} \rightarrow$	373	$1.21(-9)$	MS	0.24eV KE	28
		373	$1.10(-9)$	MS	0.65eV KE	28
			$1.6(-9)$	MS	0.2eV IEE	37
	$+ \text{CH}_2\text{F}_2 \rightarrow$	320	$1.32(-9)$	MS		30
		317	$1.77(-9)$	MS		38
	$+ \text{CHCl}_2\text{F} \rightarrow$		$1.53(-9)$	MS	0.2eV KE	31
	$+ \text{CHClF}_2 \rightarrow$		$3(-10)$	MS	0.2eV KE	31
CH_2F^+	$+ \text{CHF}_3 \rightarrow$		$1(-10)$	MS	0.2eV KE	31
	$+ 2 \text{CH}_3\text{F} \rightarrow$	320	$2.8(-26)$	MS	200eV e^-	30
		320	$3.45(-26)$	MS	14.5eV e^-	30
	$+ \text{CH}_2\text{F}_2 \rightarrow$		$1.30(-9)$	ICR		39
			$2.2(-10)$	ICR		40
	$+ 2 \text{CH}_2\text{F}_2 \rightarrow$	320	$1.1(-26)$	MS		30
	$+ \text{CH}_2\text{FCl} \rightarrow$		$1.04 \pm 0.2(-9)$	ICR		41
$\text{C}_2\text{H}_5\text{F}$	$+ \text{C}_2\text{H}_5\text{F} \rightarrow$		$1.9 \pm 0.1(-9)$	MS		42
	$+ 2 \text{CH}_2\text{CF}_2 \rightarrow$	317	$7.8(-25)$	MS		38

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction		T	k	Method	Comments	Ref.
CH_3F^+	+ $\text{CH}_3\text{F} \rightarrow$	320	1.86(-9)	MS	14.5eV e ⁻	30
		320	2.30(-9)	MS	200eV e ⁻	30
			1.73(-9)	ICR		39
			1.37(-9)	ICR		43
			1.38(-9)	ICR		44
		373	1.53(-9)	MS	0.24eV IEE	28
		373	2.27(-9)	MS	1.1eV IEE, Increases with increasing IEE	28
			2.2(-9)	MS	0.2eV KE See footnote d for further comment.	37
	+ $\text{CH}_3\text{F} \rightarrow$	320	1.36(-9)	MS	200eV e ⁻	30
		320	9.5(-10)	MS	14.5eV e ⁻	30
CH_4F^+	+ $\text{CH}_3\text{F} \rightarrow$		9.9(-10)	ICR		39
			7.1(-10)	ICR		44
			9.0(-10)	MS		28
			8.0(-10)	ICR		43
	+ $\text{HFCO} \rightarrow$		2.5(-9)	ICR	HFCO^+	29
	+ $\text{CH}_2\text{FO}^+ \rightarrow$		1.2(-10)	ICR	$(\text{HFCO})\text{H}^+$	29
	+ $\text{CF}_3^- \rightarrow$	320	9.3(-10)	MS		30
			9.2(-10)	MS	0.2eV KE	31
	+ $\text{CF}_4 \rightarrow$	320	2.7(-10)	MS		30
			1.7(-10)	ICR		39
CHO^+	+ $\text{CHClF}_3 \rightarrow$	390	4.0(-10)	MS	0.2eV KE	31
	+ $\text{CCl}_2\text{F}_2 \rightarrow$	390	4.0(-10)	MS	0.2eV KE	31
	+ $\text{CH}_2\text{CF}_3 \rightarrow$	317	2.15(-9)	MS		38
	+ $\text{C}_2\text{F}_6 \rightarrow$	295	$1.45 \pm 0.3 (-10)$	MS	Photoionization	32
			6.7(-11)	B	0.3eV KE	33
	+ $\text{C}_3\text{F}_8 \rightarrow$		2.8(-10)	B	0.3eV KE	34
	+ $\text{SiH}_4 \rightarrow$		$1.1 \pm 0.1 (-9)$	B	1.0eV KE	36
	+ $i\text{-C}_5\text{H}_{10} \rightarrow$		$1.24 \pm 0.08 (-9)$	ICR		45
	+ $neo\text{-C}_5\text{H}_{12} \rightarrow$		$1.22 \pm 0.06 (-9)$	ICR		45
	+ $\text{CH}_3\text{OH} \rightarrow$		1.41(-9)	ICR		46
CH_2F_2^+	+ $\text{HCOOH} \rightarrow$		1.16(-9)	ICR		46
	+ $\text{CH}_3\text{CHO} \rightarrow$		9.6(-10)	ICR		46
	+ $\text{C}_2\text{H}_5\text{OH} \rightarrow$		1.11(-9)	ICR		46
	+ $\text{CH}_3\text{OCH}_3 \rightarrow$		1.0(-9)	ICR		46
	+ $\text{CH}_3\text{COOH} \rightarrow$		1.9(-9)	ICR		46
	+ $\text{HCOOCH}_3 \rightarrow$		1.53(-9)	ICR		46
	+ $\text{CH}_3\text{COCH}_3 \rightarrow$		1.84(-9)	ICR		46
	+ $\text{HCOOC}_2\text{H}_5 \rightarrow$		1.86(-9)	ICR		46
	+ $\text{CH}_3\text{COOCH}_3 \rightarrow$		1.51(-9)	ICR		46
	+ $\text{CH}_3\text{COC}_2\text{H}_5 \rightarrow$		1.27(-9)	ICR		46
CH_2F_2^+	+ $(\text{CH}_3)_2\text{CHCHO} \rightarrow$		2.05(-9)	ICR		46
	+ $\text{HCOOC}_2\text{H}_5 \rightarrow$		1.56(-9)	ICR		46
	+ $\text{CH}_2\text{F}_2 \rightarrow$	320	6.0(-10)	MS		30
			1.9(-10)	ICR		39
			2.2(-10)	ICR		39
			1.4(-10)	ICR		40
	+ $\text{CHF}_3 \rightarrow$		1.4(-10)	ICR		39
	+ $\text{CF}_2\text{Cl}_2 \rightarrow$		$9.7 \pm 1.4 (-10)$	ICR		41
	+ $\text{CCl}_3\text{H} \rightarrow$		$1.35 \pm 0.20 (-9)$	ICR		41
	+ $\text{CFCl}_3 \rightarrow$		$1.21 \pm 0.18 (-9)$	ICR		41
CH_2F_2^+	+ $\text{CCl}_4 \rightarrow$		$1.28 \pm 0.19 (-9)$	ICR		41
	+ $\text{NH}_3 \rightarrow$		1.0(-9)	ICR		46
	+ $\text{CH}_2\text{F}_2 \rightarrow$	320	1.77(-9)	MS	200eV e ⁻	30
		320	1.13(-9)	MS	14.5eV e ⁻	30

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

	Reaction	T	k	Method	Comments	Ref.
CH_2F_2^+	+ $\text{CH}_2\text{F}_2 \rightarrow$		1.30(-9)	ICR		39
CH_2F_2^+	+ $\text{CH}_2\text{F}_2 \rightarrow$		1.40(-9)	ICR		39
CF_3^+	+ $\text{C}_2\text{H}_6 \rightarrow$	323	$3.6 \pm 0.4 (-10)$	MS	15 V cm^{-1}	47
	+ $i\text{-C}_4\text{H}_{10} \rightarrow$		$5.5 \pm 0.5 (-10)$	ICR		45
	+ $i\text{-C}_5\text{H}_{12} \rightarrow$		$7.3 \pm 0.5 (-10)$	ICR		45
	+ $neo\text{-C}_5\text{H}_{12} \rightarrow$		$4.9 \pm 0.4 (-10)$	ICR		45
	+ $\text{CH}_3\text{F} \rightarrow$		$2.7 \pm 0.7 (-10)$	ICR		41
	+ $\text{CHF}_3 \rightarrow$	320	$6.4 (-10)$	MS		30
			$2.1 (-10)$	ICR		39
			5.9(-10)	MS	0.2eV KE	31
	+ $\text{CH}_2\text{FCl} \rightarrow$		$7.3 \pm 1.1 (-10)$	ICR		41
	+ $\text{CF}_2\text{Cl}_2 \rightarrow$		$5.8 \pm 0.9 (-10)$	ICR		41
	+ $\text{CHCl}_3 \rightarrow$		$9.1 \pm 1.4 (-10)$	ICR		41
	+ $\text{CFCl}_3 \rightarrow$		$7.7 \pm 1.2 (-10)$	ICR		41
	+ $\text{CCl}_4 \rightarrow$		$9.1 \pm 1.4 (-10)$	ICR		41
	+ $\text{C}_2\text{F}_6 \rightarrow$	295	$4 \pm 1 (-11)$	MS	Photoionization	32
			3.4(-11)	B	0.3eV KE	33
	+ $\text{C}_3\text{F}_8 \rightarrow$		2.9(-11)	B	0.3eV KE	34
	+ $\text{SiH}_4 \rightarrow$		$6.5 \pm 0.7 (-10)$	B	1.0eV KE	36
CH_2F_3^+	+ $\text{CHF}_3 \rightarrow$		8.2(-11)	ICR	$(\text{CHF}_3)\text{H}^+$	39
CHSi^+	+ $\text{CH}_3\text{SiH}_3 \rightarrow$		$1.9 \pm 0.3 (-10)$	MS		48
CH_3Si^+	+ $\text{CH}_3\text{SiH}_3 \rightarrow$		$1.7 \pm 0.4 (-10)$	MS		48
CH_3Si^+	+ $\text{CH}_3\text{SiH}_3 \rightarrow$		$1.26 \pm 0.33 (-9)$	MS		48
	+ $(\text{CH}_3)_4\text{Si} \rightarrow$		$1.6 \pm 0.4 (-9)$	MS	11.7 V cm^{-1}	49
CH_3Si^+	+ $\text{CH}_3\text{SiH}_3 \rightarrow$		$2.45 \pm 0.6 (-9)$	MS	$(\text{CH}_3\text{SiH})^+$	48
	+ $(\text{CH}_3)_2\text{SiH}_2 \rightarrow$		1.3(-9)	MS	$(\text{CH}_3\text{SiH})^+$	50
CH_3Si^+	+ $\text{CH}_3\text{SiH}_3 \rightarrow$		$3.2 \pm 0.8 (-10)$	MS	$(\text{CH}_3\text{SiH}_2)^+$	48
	+ $(\text{CH}_3)_2\text{SiH}_2 \rightarrow$		7.5(-10)	MS	$(\text{CH}_3\text{SiH}_2)^+$	50
$\text{CH}_3\text{SiF}_2^+$	+ $\text{CH}_3\text{SiF}_3 \rightarrow$		6.1(-11)	ICR		51
$\text{CH}_3\text{SiF}_2^+$	+ $\text{CH}_3\text{SiF}_3 \rightarrow$		1.3(-11)	ICR		51
	+ $(\text{CH}_3)_2\text{SiF}_2 \rightarrow$		3.2(-10)	ICR		51
$\text{CH}_3\text{SiF}_3^+$	+ $\text{CH}_3\text{SiF}_3 \rightarrow$		6.7(-11)	ICR		51
CH_3P^+	+ $\text{CH}_3\text{PH}_2 \rightarrow$		$9.5 \pm 2.0 (-10)$	ICR	CH_3PH_2^+	52
CS^+	+ $\text{H}_2 \rightarrow$		$7.3 \pm 1.1 (-10)$	ICR		53
	+ $\text{CH}_4 \rightarrow$		$9.5 \pm 1.4 (-10)$	ICR		53
CHS^+	+ $\text{CH}_3\text{SH} \rightarrow$		$8.0 \pm 1.0 (-10)$	MS	10.7 V cm^{-1}	54
			7(-10)	MS	3.4eV IEE	55
	+ $(\text{CH}_3)_2\text{SH} \rightarrow$		$1.0 \pm 0.2 (-9)$	MS	10.7 V cm^{-1}	54
CDS^+	+ $\text{CD}_3\text{SH} \rightarrow$		$3.5 \pm 0.2 (-10)$	TI	0.4eV KE	56
CH_2S^+	+ $\text{CH}_3\text{SH} \rightarrow$		$8 \pm 1 (-10)$	TI	0.4eV KE	56
	+ $(\text{CH}_3)_2\text{S} \rightarrow$		$2.0 \pm 0.3 (-9)$	MS	10 V cm^{-1}	54
	+ dimethylsulfoxide →	408	$3.2 \pm 0.3 (-9)$	MS	$\geq 0.7 \text{ eV KE}$	22
CHDS^+	+ $\text{CD}_3\text{SH} \rightarrow$		$7.9 \pm 1.5 (-10)$	TI	0.4eV KE	56
CD_2S^+	+ $\text{CD}_3\text{SH} \rightarrow$		$4.7 \pm 0.3 (-10)$	TI	0.4eV KE	56
CH_3S^+	+ $\text{CH}_3\text{SH} \rightarrow$		$6.3 \pm 1.6 (-10)$	TI	0.4eV KE	56
			1.6±0.2(-9)	MS	10.7 V cm^{-1}	54
			1.04±0.03(-9)	MS	3.4eV IEE	55
	+ $(\text{CH}_3)_2\text{S} \rightarrow$		1.5±0.2(-9)	MS	10.7 V cm^{-1}	54
	+ dimethylsulfoxide →	408	$2.7 \pm 0.3 (-9)$	MS	0.7eV IEE	22
			2.6±0.3(-9)	MS	1.1eV IEE	22
CHD_2S^+	+ $\text{CD}_3\text{SH} \rightarrow$		$6.7 \pm 0.2 (-10)$	TI	0.4eV KE	56
			8.0±0.6(-10)	MS	3.4eV IEE	55
CD_3S^+	+ $\text{CD}_3\text{SH} \rightarrow$		$7.2 \pm 0.6 (-10)$	TI	0.4eV KE	56
			1.00±0.04(-9)	MS	3.4eV IEE	55
CH_3S^+	+ $\text{CH}_3\text{SH} \rightarrow$		$1.41 \pm 0.07 (-9)$	MS	Photoionization	57
			7.7±0.2(-10)	TI	0.41V KE	56
			1.19±0.06(-10)	MS	3.4eV IEE	55
	+ $\text{C}_2\text{H}_6\text{CHO} \rightarrow$		1.8±0.1(-9)	MS		58

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

	Reaction	T	k	Method	Comments	Ref.
CHD ₃ S ⁺	.+ CH ₃ CHO →		1.2±0.2(-9)	MS	(CD ₃ SH) ⁺	58
	+ CD ₃ SH →		8.2±0.2(-10)	TI	0.4eV KE, (CD ₃ SH) ⁺	56
			1.21±0.03(-9)	MS	(CD ₃ SH) ⁺	55
	+ CH ₃ OCH ₃ →		1.51±0.36(-9)	MS	(CD ₃ SH) ⁺	58
CH ₅ S ⁺	+ CH ₃ SH →		7.6(-12)	TI	0.4eV KE, CH ₃ SH ₂ ⁺	56
			5.3±0.8(-10)	MS	10 V cm ⁻¹ , CH ₃ SH ₂ ⁺	54
	+ C ₂ H ₅ CHO →		5.6±0.3(-10)	MS	CH ₃ SH ₂ ⁺	58
	+ (CH ₃) ₂ S →		1.74±0.05(-9)	MS	CH ₃ SH ₂ ⁺	58
	+ CH ₃ COCH ₃ →		2.30±0.21(-9)	MS	CH ₃ SH ₂ ⁺	58
CH ₄ DS ⁺	+ CH ₄ S →		4.4±0.6(-10)	TI	CH ₃ SHD ⁺	12
CH ₂ D ₃ S ⁺	+ CH ₃ OCH ₃ →		1.24±0.10(-9)	MS	CD ₃ SH ₂ ⁺	58
CHD ₄ S ⁺	+ CHD ₃ S →		1.3±0.2(-10)	TI	<i>k</i> reflects sources of CD ₃ SHD ⁺	56
			2.6±0.3(-10)	TI	<i>k</i> reflects source CD ₃ SHD ⁺	56
	+ CH ₃ CHO →		7.3±0.9(-10)	MS	CD ₃ SHD ⁺	58
CH ₃ SO ⁺	+ dimethylsulfoxide →	408	3.4±0.3(-9)	MS	1.07eV IEE	22
		408	3.4±0.3(-9)	MS	0.71 IEE	22
		408	3.2±0.3(-9)	MS	Thermal	22
CCl ⁺	+ CH ₃ Cl →	373	1.2(-10)	MS	0.21eV IEE	28
		373	1.7(-10)	MS	0.46eV IEE	28
	+ CH ₃ Cl →	373	2.1(-10)	MS	0.65eV IEE	28
		373	1.4(-10)	MS	1.1eV IEE	28
	+ CHClF ₂ →		2.0(-10)	MS	0.2eV IEE	31
	+ CHFCl ₂ →		4.0(-10)	MS	0.2eV IEE	31
	+ CHCl ₃ →		7.0(-10)	MS	0.2eV IEE	31
	+ CCl ₄ →		4.0(-10)	MS	0.2eV IEE	31
CHCl ⁺	+ CH ₃ Cl →	373	6.1(-10)	MS	0.2eV IEE	28
		373	4.4(-10)	MS	0.46eV IEE	28
		373	3.3(-10)	MS	0.65eV IEE	28
		373	3.2(-10)	MS	1.1eV IEE	28
	+ CHFCl ₂ →		7.6(-10)	MS	0.2eV KE	31
	+ CHCl ₃ →		7(-10)	MS	0.2eV KE	31
CH ₂ Cl ⁺	+ CH ₃ Cl →	373	4.6(-10)	MS		17
	+ CHF ₂ Cl →		1.5±0.5(-10)	ICR		41
	+ CH ₂ Cl ₂ →		<1.0(-11)	ICR		41
	+ C ₂ H ₅ Cl →	320	1.15±0.08(-9)	MS		42
CH ₃ Cl ⁺	+ CH ₃ Cl →		1.25(-9)	ICR		44
		373	1.54±0.08(-9)	MS	Value over range 0.21 to 1.1eV IEE	28
			1.66(-9)	MS	0.2eV KE	59
		373	1.70(-9)	MS		17
CH ₄ Cl ⁺	+ CH ₃ Cl →		1.4(-10)	ICR		44
		373	1.0(-10)	MS	0.45eV IEE	28
CNCl ⁺	+ CNCl →		1.2(-9)	MS		60
CFCl ⁺	+ CHClF ₂ →		4.2(-10)	MS	0.2eV KE	31
	+ CHFCl ₂ →		5.0(-10)	MS	0.2eV KE	31
	+ CF ₂ Cl ₂ →		6.1(-10)	MS	0.2eV KE	31
CHFCl ⁺	+ CHFCl ₂ →		3.0±0.5(-11)	ICR		41
CF ₂ Cl ⁺	+ CHFCl ₂ →		2.2±0.3(-10)	ICR		41
	+ CFCl ₃ →		4.0±0.6(-10)	ICR		41
	+ CCl ₄ →		5.0±0.8(-10)	ICR		41
	+ i-C ₄ H ₁₀ →		3.3(-10)	MS	0.2eV KE	31
CCl ₂ ⁺	+ CHFCl ₂ →		2.4±0.1(-10)	ICR		45
	+ CHCl ₃ →		3.6(-10)	MS	0.2eV KE	31
	+ CCl ₄ →		5.9(-10)	MS	0.2eV KE	31
CHCl ₂ ⁺	+ CH ₃ OH →		7.5(-11)	ICR		46
	+ HCOOH →		5.7(-11)	ICR		46

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

	Reaction	T	k	Method	Comments	Ref.
CHCl ₂ ⁺	+ CH ₃ CHO →		5.6(-10)	ICR		46
	+ C ₂ H ₅ OH →		5.2(-10)	ICR		46
	+ CH ₃ OCH ₃ →		4.0(-10)	ICR		46
	+ HCOOCH ₃ →		3.9(-10)	ICR		46
	+ CH ₃ COCH ₃ →		5.7(-10)	ICR		46
	+ HCOOC ₂ H ₅ →		8.3(-10)	ICR		46
	+ CH ₃ COOCH ₃ →		9.8(-10)	ICR		46
	+ i-C ₄ H ₁₀ →		6.7±0.3(-10)	ICR		45
	+ CH ₃ COC ₂ H ₅ →		7.2(-10)	ICR		46
	+ (CH ₃) ₂ CHCHO →		1.33(-9)	ICR		46
	+ C ₂ H ₅ OC ₂ H ₅ →		9.0(-10)	ICR		46
	+ i-C ₅ H ₁₂ →		8.4±0.3(-10)	ICR		45
	+ neo-C ₅ H ₁₂ →		3.5±0.3(-10)	ICR		45
	+ HCOOC ₄ H ₉ →		1.01(-9)	ICR		46
	+ CCl ₄ →		8.0±1.2(-11)	ICR		41
CDCl ₂ ⁺	+ NH ₃ →		7.8(-10)	ICR		46
	+ (CH ₃) ₃ N →		1.37(-9)	ICR		46
CFCl ₂ ⁺	+ CHCl ₃ →		<4(-12)	ICR		41
	+ CCl ₃ F →		1.3±0.2(-11)	ICR		41
	+ CCl ₄ →		<4(-12)	ICR		41
	+ i-C ₄ H ₁₀ →		7.4±1.0(-11)	ICR		45
	+ 2,2-dimethylpentane →		1.2±0.2(-10)	ICR		45
	+ 3,3-dimethylpentane →		1.8±0.1(-10)	ICR		45
	+ 2-methylheptane →		7.3±0.4(-10)	ICR		45
	+ 2,3-dimethylhexane →		5.8±0.3(-10)	ICR		45
	+ 2,4-dimethylhexane →		3.7±0.3(-10)	ICR		45
	+ 2,5-dimethylhexane →		4.5±0.1(-10)	ICR		45
	+ 2,3,4-trimethylpentane →		3.4±0.2(-10)	ICR		45
CCl ₃ ⁺	+ 2-methylheptane →		6.0±0.4(-11)	ICR		45
	+ 3-methylheptane →		6.8±0.4(-11)	ICR		45
	+ 2,3-dimethylhexane →		6.2±0.5(-11)	ICR		45
	+ 2,4-dimethylhexane →		4.6±0.6(-11)	ICR		45
	+ 2,5-dimethylhexane →		1.1±0.1(-10)	ICR		45
	+ 2,3,4-trimethylpentane →		4.9±0.7(-11)	ICR		45
CH ₃ Br ⁺	+ CH ₃ Br →	295	5.8±0.4(-11)	MS	Photoionization at 10.0eV	61
		295	3.8±0.3(-11)	MS	Photoionization at 11.7eV	61
			5.5(-11)	ICR		44
CH ₃ I ⁺	+ CH ₃ I →	220	2.46±0.25(-11)	MS	Photoionization	62
		295	2.37±0.12(-11)	MS	Photoionization	62
		320	2.37±0.05(-11)	MS	Photoionization	62
	+ CH ₃ I →	295	1.2±0.5(-11)	MS	Photoionization	61
			5.5(-12)	ICR		44
CD ₃ I ⁺	+ 2CH ₃ I →	295	3.3±0.1(-25)	MS	Photoionization	62
	+ CD ₃ I →	220	2.04±0.12(-11)	MS	Photoionization	62
		295	2.23±0.05(-11)	MS	Photoionization	62
		320	2.16±0.12(-11)	MS	Photoionization	62
CH ₃ Hg ⁺	+ 2CD ₃ I →	295	9.4±3.0(-25)	MS	Photoionization	62
	+ C ₂ H ₄ →		2.7±0.4(-10)	ICR		63
	+ C ₃ H ₆ →		2.9±0.3(-10)	ICR		63
	+ Allene →		1.5±0.5(-10)	ICR		63
	+ 1,3-butadiene →		3.9±1.3(-10)	ICR		63
	+ 1-C ₄ H ₈ →		1.6±0.1(-10)	ICR		63
	+ iso-C ₄ H ₈ →		3.7±0.9(-10)	ICR		63
	+ trans-2-C ₄ H ₈ →		5.2(-10)	ICR		63
	+2-methyl-2-butene →		5.2(-10)	ICR		63
C ₂ H ₆ B ⁺	+ (CH ₃) ₃ B →		2.6±0.4(-10)	ICR	(CH ₃) ₂ B ⁺	18
C ₂ N ⁺	+ CH ₃ CN →	300	2.6(-9)	MS	Pressure varied	64
	+ C ₂ H ₅ CN →		2.1(-9)	MS	Pressure varied	64
			2.0(-9)	MS	Time varied	64

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction		T	k	Method	Comments	Ref.
C ₂ HN ⁺	+ CH ₃ CN →		3.54(-9)	ICR	CHCN ⁺	15
		300	3.0(-9)	MS	Pressure varied, CHCN ⁺	64
C ₂ DN ⁺	+ CD ₃ CN →	300	2.5(-9)	MS	Pressure varied, CDCN ⁺	64
		300	6.7(-9)	MS	Time varied, CDCN ⁺	64
C ₂ H ₂ N ⁺	+ CH ₃ CN →	300	1.78(-9)	MS	Pressure varied, CH ₂ CN ⁺	64
			2.09(-9)	ICR	CH ₂ CN ⁺	15
	+ C ₂ H ₅ CN →	300	9(-10)	MS	Pressure varied, CH ₂ CN ⁺	64
		300	1(-9)	MS	Time varied, CH ₂ CN ⁺	64
C ₂ D ₂ N ⁺	+ CD ₃ CN →	300	1.15(-9)	MS	Pressure varied, CD ₂ CN ⁺	64
		300	1.08(-9)	MS	Time varied, CD ₂ CN ⁺	64
C ₂ H ₃ N ⁺	+ D ₂ →		1.0(-10)	MS	3.4eV IEE, CH ₃ CN ⁺	65
	+ CH ₃ CN →		2.09(-9)	ICR	CH ₃ CN ⁺	15
		373	1.96(-9)	MS	Thermal, CH ₃ CN ⁺	17
		300	2.01(-9)	MS	Pressure varied, CH ₃ CN ⁺	64
		300	2.38(-9)	MS	Time varied, CH ₃ CN ⁺	64
			2.15±0.21(-9)	MS	3.7eV IEE, CH ₃ CN ⁺ See footnote e for further information	65
C ₂ D ₃ N ⁺	+ CD ₃ CN →	300	2.45(-9)	MS	Pressure varied, CD ₃ CN ⁺	64
		300	1.98(-9)	MS	Time varied, CD ₃ CN ⁺	64
C ₂ H ₄ N ⁺	+ (CH ₃) ₂ NH →		4(-10)	MS	From (CH ₃) ₂ NH	6
C ₂ H ₅ N ⁺	+ (CH ₃) ₂ NH →		5(-10)	MS	From (CH ₃) ₂ NH	6
C ₂ H ₆ N ⁺	+ (CH ₃) ₂ NH →		1.00±0.13(-9)	MS	CH ₃ NHCH ₂ ⁺	8
		373	8(-10)	MS	3.4eV IEE, (CH ₃) ₂ NH ⁺	5
			4±1(-10)	MS	CH ₃ NHCH ₂ ⁺	6
C ₂ H ₇ N ⁺	+ (CH ₃) ₂ NH →		1.28(-9)	ICR	Pressure varied, (CH ₃) ₂ NH ⁺	10
			1.49(-9)	ICR	Time varied, (CH ₃) ₂ NH ⁺	10
		295	1.16±0.08(-9)	MS	Photoionization at 8.4eV	8
		295	1.02±0.08(-9)	MS	Photoionization at 10.0, 11.7eV	8
		373	1.2(-9)	MS	3.4 eV IEE, (CH ₃) ₂ NH ⁺	5
			6±1(-10)	MS	(CH ₃) ₂ NH ⁺	6
	+ C ₂ H ₅ NH ₂ →	298	1.83±0.09(-9)	MS	Photoionization, C ₂ H ₅ NH ₂ ⁺	9
C ₂ HD ₆ N ⁺	+ (CD ₃) ₂ NH →		1.4(-9)	ICR	(CD ₃) ₂ NH ⁺	10
C ₂ H ₈ N ⁺	+ 2(CH ₃) ₂ NH →	350	2.6(-26)	MS	(CH ₃) ₂ NH ₂ ⁺	11
	+ (CH ₃) ₃ N →		3.5(-10)	MS	From (CH ₃) ₃ N	6
C ₂ H ₇ DN ⁺	+ (CH ₃) ₂ NH →		3.1±0.2(-10)	TI	(CH ₃) ₂ NHD ⁺	12
C ₂ HD ₇ N ⁺	+ (CD ₃) ₂ NH →		3.4(-10)	ICR	Pressure varied, (CD ₃) ₂ NHD ⁺	10
			2.8(-10)	ICR	Time varied, (CD ₃) ₂ NHD ⁺	10
C ₂ H ₃ O ⁺	+ CH ₃ CHO →	373	1.3±1.0(-10)	MS	2.4eV IEE, CH ₃ CO ⁺	16
	+ CH ₃ COCH ₃ →		2.0(-10)	ICR	CH ₃ CO ⁺	66
			2.33±0.10(-10)	ICR	CH ₃ CO ⁺ from acetone	67
			2.30±0.10(-10)	ICR	CH ₃ CO ⁺ from protonated ketene	67
			2.2±0.2(-10)	TI	CH ₃ CO ⁺	20
		295	4.6±0.6(-10)	MS	CH ₃ CO ⁺ , Photoionization	68
			4.3±0.3(-10)	ICR	CH ₃ CO ⁺	69
		473	3.0(-10)	MS	CH ₃ CO ⁺	69
	+ CH ₃ COC ₂ H ₅ →		7.0(-10)	ICR	CH ₃ CO ⁺	66
	+ vinyl formate →		1.4(-9)	MS	2.4eV IEE	16
	+ C ₃ H ₅ COCH ₃ →		7.1±2.1(-10)	ICR	CH ₃ CO ⁺	7
	+ C ₆ H ₅ CHO →		1.2(-10)	ICR	CH ₃ CO ⁺	70
	+ C ₆ H ₅ OC ₂ H ₅ →		2.3(-10)	ICR	CH ₃ CO ⁺	70
	+ o-methylanisole →		2(-10)	ICR	CH ₃ CO ⁺ from acetone	71
			1.9±0.4(-9)	ICR	CH ₃ CO ⁺ from biacetyl	71
	+ m-methylanisole →		1.8±0.4(-10)	ICR	CH ₃ CO ⁺ from biacetyl	71
	+ p-methylanisole →		3.5±0.7(-10)	ICR	CH ₃ CO ⁺ from biacetyl	71
	+ o-cresol →		5.6±0.6(-10)	ICR	CH ₃ CO ⁺ from biacetyl	72

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction		T	k	Method	Comments	Ref.	
C ₂ D ₃ O ⁺	+ CD ₃ COCD ₃ →	295	4.6±0.6(-10)	MS	Photoionization, CD ₃ CO ⁺	69	
C ₂ H ₄ O ⁺	+ CH ₃ CHO →		1.99(-9)	ICR	CH ₃ CHO ⁺	15	
			2.49±0.10(-9)	TI	CH ₃ CHO ⁺	20	
	+ ethylene oxide →	373	1.91±0.20(-9)	MS	Thermal, CH ₃ CHO ⁺	16	
			1.9±0.2(-10)	ICR	From ethylene oxide	73	
			1.5±0.3(-10)	TI	From ethylene oxide	20	
	+ ethylene oxide (d ₄) →	373	2±1(-10)	MS	2.4eV IEE	16	
C ₂ D ₄ O ⁺	+ CD ₃ CDO →		2.46(-9)	TI	CD ₃ CDO ⁺	20	
	+ ethylene oxide (d ₄) →		1.9±0.2(-10)	TI	From deuterated ethylene oxide	20	
C ₂ H ₅ O ⁺	+ 2CH ₃ CHO →		1.25(-25)	TI	CH ₃ CHOH ⁺	20	
	+ C ₂ H ₅ OH →		1.12(-9)	ICR	From C ₂ H ₅ OH	10	
	+ (CH ₃) ₂ O →	473	1.2±0.6(-9)	MS	From C ₂ H ₅ OH	74	
	+ (CD ₃) ₂ O →		1.3±0.9(-11)	TI	CH ₃ COCH ₂ ⁺	20	
	+ (CH ₃) ₂ CO →		1.88±0.05(-9)	TI	CH ₃ CHOH ⁺	20	
			3.3±1.0(-11)	TI	CH ₃ OCH ₂ ⁺	20	
			2.25(-9)	TI	CH ₃ CHOH ⁺	20	
	+ i-C ₃ H ₇ OH →	295	1.1±0.1(-9)	MS	Photoionization, CH ₃ CHOH ⁺	75	
			1.5±0.3(-9)	ICR	CH ₃ CHOH ⁺	76	
C ₂ H ₂ D ₃ O ⁺	+ CD ₃ OC ₂ H ₅ →		1.75±0.05(-10)	TI	CD ₃ OCH ₂ ⁺	20	
C ₂ HD ₄ O ⁺	+ C ₂ D ₅ OH →		1.04(-9)	ICR	CD ₃ CDOH ⁺	10	
C ₂ H ₆ O ⁺	+ (CH ₃) ₂ O ⁺ →		1.58±0.15(-9)	TI	(CH ₃) ₂ O ⁺	20	
	+ (CH ₃) ₂ O ⁺ →	373	1.45±0.1(-9)	MS	2.4eV IEE, (CH ₃) ₂ O ⁺	16	
	+ C ₂ H ₅ OH →	373	1.93(-9)	MS	Thermal, (CH ₃) ₂ O ⁺	17	
			1.78(-10)	ICR	C ₂ H ₅ OH ⁺	10	
			3.03(-9)	MS	10 V cm ⁻¹ , C ₂ H ₅ OH ⁺	77	
C ₂ H ₃ D ₃ O ⁺	+ CH ₃ OCD ₃ →	373	2.09(-9)	MS	Thermal, CH ₃ OCD ₃ ⁺	17	
C ₂ D ₆ O ⁺	+ CD ₃ OCD ₃ →		1.58(-9)	TI	(CD ₃) ₂ O ⁺	20	
C ₂ H ₇ O ⁺	+ C ₂ H ₅ OH →		2.2(-9)	ICR	Pressure varied, C ₂ H ₅ OH ₂ ⁺	10	
			2.4(-9)	ICR	Time varied, C ₂ H ₅ OH ₂ ⁺	10	
	+ 2(CH ₃) ₂ O →		1.0±0.2(-25)	TI	(CH ₃) ₂ O ⁺	20	
	+ (CH ₃) ₂ CO →		1.0±0.1(-9)	TI	(CH ₃) ₂ O ⁺	20	
C ₂ H ₂ D ₅ O ⁺	+ C ₂ D ₅ OH →		1.7(-10)	ICR	Pressure varied, C ₂ D ₅ OH ₂ ⁺	10	
			1.4(-10)	ICR	Time varied, C ₂ D ₅ OH ₂ ⁺	10	
C ₂ HD ₆ O ⁺	+ C ₂ D ₅ OH →		3.7(-10)	ICR	Pressure varied, C ₂ D ₅ OHD ⁺	10	
			2.8(-10)	ICR	Time varied, C ₂ D ₅ OHD ⁺	10	
C ₂ H ₄ O ₂ ⁺	+ HCO ₂ CH ₃ →	370	1.57±0.10(-9)	MS	2.4eV IEE, HCO ₂ CH ₃ ⁺	21	
C ₂ H ₃ DO ₂ ⁺	+ DCO ₂ CH ₃ →	370	1.65±0.10(-9)	MS	2.4eV IEE, DCO ₂ CH ₃ ⁺	21	
C ₂ H ₈ NO ₃ ⁺	+ 2CH ₃ OH →		2(-27)	MS	(CH ₃ OH) ₂ NO ⁺	27	
C ₂ F ⁺	+ CH ₂ CF ₂ →	317	1.91(-9)	MS		38	
C ₂ HF ⁺	+ C ₂ H ₃ F →		1.93±0.12(-9)	MS		78	
			370	1.95(-9)	MS		79
	+ CH ₂ CF ₂ →	317	1.84(-9)	MS		38	
C ₂ H ₂ F ⁺	+ C ₂ H ₃ F →		1.03±0.06(-9)	MS		78	
			370	7.5±1.0(-10)	MS		79
	+ C ₂ H ₅ F →		1.75±0.19(-9)	MS		42	
C ₂ H ₃ F ⁺	+ C ₂ H ₄ →		5.9±1.8(-10)	ICR		80	
	+ C ₂ D ₄ →		5.1(-10)	ICR		81	
	+ C ₂ H ₃ F →	300	2.9(-10)	ICR	k varies with internal energy	82	
	+ C ₂ H ₃ F →		2.1(-10)	ICR		81	
	+ 2C ₂ H ₃ F →	295	3.2±0.2(-10)	MS	Photoionization	83	
			2.1(-10)	ICR		84	
			1.34(-10)	ICR		85	
			2.2±0.4(-10)	TI		86	
	+ 2C ₂ H ₃ F →	370	3.0(-11)	MS	19eV e ⁻	79	
	+ C ₂ H ₅ F →	370	2.3(-11)	MS	11eV e ⁻	79	
	+ CH ₂ CF ₂ →		1.61(-25)	MS		78	
	+ 2C ₂ H ₃ F →	320	5.9(-10)	MS	From C ₂ H ₅ F	42	
	+ C ₂ H ₅ F →		1.02(-9)	ICR		81	

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

	Reaction	T	k	Method	Comments	Ref.
C ₂ H ₃ F ⁺	+ <i>cis</i> -CHFCHF →		1.4(-9)	ICR		81
	+ <i>trans</i> -CHFCHF →		1.3(-9)	ICR		81
	+ C ₂ HF ₃ →		1.3(-9)	ICR		81
	+ C ₂ F ₄ →		8.8(-10)	ICR		81
	+ 2-fluoropropane →		1.3±0.2(-9)	ICR	From 2-fluoropropane	87
C ₂ H ₄ F ⁺	+ C ₂ H ₃ F →		4.2±0.7(-10)	MS		78
	+ C ₂ H ₅ F →	320	1.06±0.10(-9)	MS		42
	+ 2-fluoropropane →		1.4±0.2(-9)	ICR	From 2-fluoropropane	87
C ₂ H ₅ F ⁺	+ C ₂ H ₅ F →	320	1.53±0.12(-9)	MS	200eV e ⁻	42
		320	1.18±0.20(-9)	MS	14eV e ⁻	42
			1.50(-9)	ICR		44
C ₂ H ₆ F ⁺	+ 2CH ₃ F →	320	1.5(-27)	MS	200eV e ⁻ , (CH ₃) ₂ F ⁺	30
		320	2.3(-27)	MS	14.5eV e ⁻ , (CH ₃) ₂ F ⁺	30
	+ 2CH ₂ F ₂ →	320	5.88(-28)	MS		30
	+ C ₂ H ₅ F →	320	1.33±0.14(-9)	MS		42
			1.4(-9)	ICR		44
C ₂ F ₂ ⁺	+ C ₃ F ₈ →		1.2(-10)	B	0.3eV KE	34
C ₂ H ₂ F ₂ ⁺	+ C ₂ H ₄ →		3.4±0.5(-10)	ICR	CHFCHF ⁺	80
			2.5±0.5(-10)	ICR	CH ₂ CF ₂ ⁺	80
			1.6(-10)	ICR	From <i>cis</i> -CHFCHF	81
			8.5(-11)	ICR	From <i>trans</i> -CHFCHF	81
			1.7(-10)	ICR	CH ₂ CF ₂ ⁺	81
	+ C ₂ D ₄ →		2.9(-10)	ICR	CH ₂ CF ₂ ⁺	81
	+ C ₂ H ₃ F →		5.1(-10)	ICR	CH ₂ CF ₂ ⁺	81
			1.9(-10)	ICR	From <i>trans</i> -CHFCHF	81
	+ CH ₂ CF ₂ →		3.5(-10)	ICR	From <i>cis</i> -CHFCHF	81
			2.2(-10)	ICR	From <i>trans</i> -CHFCHF	81
	+ <i>cis</i> -CHFCHF →		8.9(-10)	ICR	CH ₂ CF ₂ ⁺	81
			6.3(-12)	ICR	From <i>cis</i> -CHFCHF, charge transfer only	88
			8.3(-11)	ICR	From <i>trans</i> -CHFCHF, charge transfer only	88
			6(-12)	ICR	From <i>cis</i> -CHFCHF, condensation only	81
			2.6(-10)	ICR	From <i>trans</i> -CHFCHF	81
	+ <i>trans</i> -CHFCHF →		5.7(-10)	ICR	From <i>cis</i> -CHFCHF	81
			3(-12)	ICR	From <i>trans</i> -CHFCHF, condensation only	81
			2.4(-10)	ICR	From <i>cis</i> -CHFCHF, charge transfer only	88
			3(-12)	ICR	From <i>trans</i> -CHFCHF, charge transfer only	88
	+ 2CH ₂ CF ₂ →	320	5.98(-26)	MS	CH ₂ CF ₂ ⁺	38
			1.1±0.2(-24)	ICR	CH ₂ CF ₂ ⁺	89
	+ C ₂ HF ₃ →		8.9(-10)	ICR	CH ₂ CF ₂ ⁺	81
			7.8(-10)	ICR	From <i>cis</i> -CHFCHF	81
			5.4(-10)	ICR	From <i>trans</i> -CHFCHF	81
	+ C ₂ F ₄ →		6.4(-10)	ICR	CH ₂ CF ₂ ⁺	81
			5.7(-10)	ICR	From <i>cis</i> -CHFCHF	81
			3.9(-10)	ICR	From <i>trans</i> -CHFCHF	81
C ₂ D ₂ F ₂ ⁺	+ C ₂ H ₃ F →		3.5(-10)	ICR	From <i>cis</i> -CDFCDF	81
			1.9±0.4(-10)	MS		78
C ₂ F ₃ ⁺	+ C ₃ F ₈ →		8.3(-11)	B	0.3eV KE	34
C ₂ HF ₃ ⁺	+ C ₂ H ₄ →		1.7(-10)	ICR		81
	+ CH ₂ CF ₂ →		6.7±0.7(-10)	ICR		80
	+ <i>cis</i> -CHFCHF →		1.8(-10)	ICR	CHFCF ₂ ⁺	81
	+ <i>trans</i> -CHFCHF →		2.7(-10)	ICR	CHFCF ₂ ⁺	81
	+ C ₂ HF ₃ →		2.8(-10)	ICR	CHFCF ₂ ⁺	81
			5(-11)	ICR	Condensation only	81

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction	T	k	Method	Comments	Ref.		
C ₂ HF ₃ ⁺ + C ₂ HF ₃ →		5.6(-11)	ICR	CHFCF ₂ ⁺	90		
		4.8(-11)	ICR	CHFCF ₂ ⁺	85		
+ C ₂ F ₄ →		2.2(-10)	ICR	CHFCF ₂ ⁺	81		
		5.7(-11)	ICR	CHFCF ₂ ⁺	90		
C ₂ DF ₃ ⁺ + C ₂ H ₃ F →		1.4(-10)	ICR	CDFCF ₂ ⁺	81		
C ₂ H ₄ F ₃ ⁺ + 2CH ₂ F ₂ →		5.16(-28)	MS		30		
C ₂ F ₄ ⁺ + C ₂ H ₄ →		1.7(-10)	ICR		81		
		1.07±0.30(-10)	ICR		80		
+ C ₂ H ₃ F →		1.6(-10)	ICR		81		
+ C ₂ F ₂ →		1.0(-11)	ICR		90		
+ CH ₂ CF ₂ →		1.5(-10)	ICR		81		
+ cis-CHFCHF →		1.9(-10)	ICR		81		
+ trans-CHFCHF →		2.0(-10)	ICR		81		
+ C ₂ HF ₃ →		2.1(-10)	ICR		81		
		1.06(-10)	ICR		90		
+ C ₂ F ₄ →		1.5(-11)	ICR	Condensation only	81		
		1.0(-11)	ICR	Condensation only	85		
C ₂ F ₅ ⁺ + C ₂ F ₆ →		7.1(-11)	B	0.3eV KE	34		
+ C ₃ F ₈ →		8.6(-12)	B	0.3eV KE	34		
C ₂ H ₆ Si ⁺ + (CH ₃) ₂ SiH ₂ →		7.0(-10)	MS	(CH ₃) ₂ Si ⁺	50		
C ₂ H ₆ FSi ⁺ + (CH ₃) ₂ SiF ₂ →		5(-10)	ICR	(CH ₃) ₂ SiF ⁺	51		
		7.4(-10)	ICR	(CH ₃) ₂ SiF ⁺	51		
C ₂ H ₆ F ₂ Si ⁺ + (CH ₃) ₂ SiF ₂ →		2.0(-10)	ICR	(CH ₃) ₂ SiF ⁺	51		
C ₂ H ₇ P ⁺ + (CH ₃) ₂ PH →		6.3±1.2(-10)	ICR	(CH ₃) ₂ PH ⁺	52		
C ₂ H ₅ S ⁺ + CH ₃ SH →		1.5±0.2(-10)	MS	10.7 V cm ⁻¹	54		
		3.5±0.5(-10)	MS	10.7 V cm ⁻¹	54		
C ₂ H ₆ S ⁺ + C ₂ H ₅ SH →		1.25±0.06(-9)	MS	Photoionization, C ₂ H ₅ SH ⁺	57		
		3.2(-12)	MS	(CH ₃) ₂ S ⁺	58		
C ₂ H ₇ S ⁺ + (CH ₃) ₂ S →		2.8±0.4(-10)	MS		54		
C ₂ H ₆ OS ⁺ + dimethylsulfoxide →	408	1.9±0.2(-9)	MS	From dimethylsulfoxide	22		
C ₂ H ₅ Cl ⁺ + C ₂ H ₅ Cl →	295	1.78±0.09(-10)	MS	Photoionization	83		
		9.6(-11)	ICR		84		
		5.2(-11)	ICR		85		
	370	1.81±0.06(-11)	MS	20eV e ⁻	91		
	370	1.35(-11)	MS	15eV e ⁻	91		
	370	1.03(-11)	MS	10.5eV e ⁻	91		
C ₂ H ₅ Cl ⁺ + C ₂ H ₅ Cl →	320	1.78±0.09(-9)	MS		42		
C ₂ H ₅ Cl ⁺ + C ₂ H ₅ Cl →		1.50(-9)	ICR		44		
C ₂ H ₆ Cl ⁺ + C ₂ H ₅ Cl →	320	1.36(-9)	MS	C ₂ H ₅ ClH ⁺	42		
		5.5(-10)	ICR	C ₂ H ₅ ClH ⁺	44		
C ₂ H ₃ Br ⁺ + C ₂ H ₃ Br →	295	1.25(-10)	MS	Photoionization	83		
	370	6.5±0.2(-12)	MS	2.4eV IEE	91		
+ 2C ₂ H ₃ Br →	370	5.5±0.2(-26)	MS		91		
C ₂ H ₅ Br ⁺ + C ₂ H ₅ Br →	295	6.7±0.5(-11)	MS	Photoionization	61		
		5.6(-11)	ICR		44		
C ₂ H ₆ Br ⁺ + C ₂ H ₅ Br →		5(-11)	ICR	C ₂ H ₅ BrH ⁺	44		
C ₂ H ₅ I ⁺ + C ₂ H ₅ I →	295	2.0±0.1(-11)	MS	Photoionization	62		
		1.6(-11)	ICR		44		
+ 2C ₂ H ₅ I →	295	4.1(-24)	MS	Photoionization	62		
C ₂ H ₃ D ₂ I ⁺ + CH ₃ CD ₂ I →	295	2.0±0.1(-11)	MS	Photoionization, CH ₃ CD ₂ I ⁺	62		
C ₂ H ₃ D ₂ I ⁺ + 2CH ₃ CD ₂ I →	295	5.3(-24)	MS	Photoionization, CH ₃ CD ₂ I ⁺	62		
C ₂ H ₂ D ₃ I ⁺ + CD ₃ CH ₂ I →	295	2.0±0.1(-11)	MS	Photoionization, CD ₃ CH ₂ I ⁺	62		
		2CD ₃ CH ₂ I →	295	7.4(-24)	MS	Photoionization, CD ₃ CH ₂ I ⁺	62
C ₂ D ₅ I ⁺ + C ₂ D ₅ I →	295	2.0±0.1(-11)	MS	Photoionization	62		
		+ 2C ₂ D ₅ I →	295	1.01(-23)	MS	Photoionization	62
C ₃ HN ⁺ + C ₂ H ₅ CN →	300	3.7(-9)	MS	From C ₂ H ₅ CN	64		
C ₃ H ₂ N ⁺ + C ₂ H ₅ CN →	300	1.9(-9)	MS	From C ₂ H ₅ CN	64		
C ₃ H ₄ N ⁺ + C ₂ H ₅ CN →	300	1.2(-9)	MS	Pressure varied, C ₂ H ₄ CN ⁺	64		

TABLE 1. Rate coefficients* for reactions of organic cations—Continued

	Reaction	T	k	Method	Comments	Ref.
C ₃ H ₅ N ⁺	+ C ₂ H ₅ CN →		0.6(-9)	MS	Time varied, C ₂ H ₄ CN ⁺	64
C ₃ D ₄ N ⁺	+ C ₂ D ₅ CN →	300	1.7(-9)	MS	Pressure varied, C ₂ D ₄ CN ⁺	64
		300	0.5(-9)	MS	Time varied, C ₂ D ₄ CN ⁺	64
C ₃ H ₆ N ⁺	+ cyclo-C ₃ H ₅ NH ₂ →		1.3±0.4(-10)	ICR	From cyclo-C ₃ H ₅ NH ₂	7
C ₃ H ₇ N ⁺	+ cyclo-C ₃ H ₅ NH ₂ →		1.9(-10)	ICR	From cyclo-C ₃ H ₅ NH ₂	7
	+ (CH ₃) ₃ N →		1.0(-10)	MS	From (CH ₃) ₃ N	6
C ₃ H ₈ N ⁺	+ diethylamine →	298	1.36±0.14(-9)	MS	From diethylamine	9
	+ pyridine →		5.3(-10)	ICR	(CH ₃) ₂ NH ₂ ⁺	92
C ₃ H ₉ N ⁺	+ n-C ₃ H ₇ NH ₂ →	298	1.56±0.08(-9)	MS	From n-C ₃ H ₇ NH ₂	9
	+ (CH ₃) ₃ N →	295	8.2±0.3(-10)	MS	Photoionization at 8.4eV, (CH ₃) ₃ N ⁺	8
	+ (CH ₃) ₃ N →	295	5.5±0.5(-10)	MS	Photoionization at 10.0eV, (CH ₃) ₃ N ⁺	8
		295	4.9±0.3(-10)	MS	Photoionization at 11.7eV, (CH ₃) ₃ N ⁺	8
C ₃ H ₁₀ N ⁺	+ 4-methylpyridine →		3.3(-10)	MS	(CH ₃) ₃ N ⁺	6
C ₆ H ₁₂ B ₃ N ₃ ⁺	+ (CH ₃) ₃ B ₃ N ₃ H ₃ →		2.6(-10)	ICR	(CH ₃) ₃ NH ⁺	92
C ₃ H ₈ O ⁺	+ cyclo-C ₃ H ₅ OH →		7.0±0.4(-10)	MS	(CH ₃) ₃ B ₃ N ₃ H ₃ ⁺	93
	+ 2-butanone →		2.0±0.6(-10)	ICR	From cyclo-C ₃ H ₅ OH	7
C ₃ H ₈ O ⁺	+ cyclo-C ₃ H ₅ OH →		6.0(-11)	ICR	C ₂ H ₅ CO ⁺	66
	+ CH ₃ COCH ₃ →		1.5±0.5(-10)	ICR	From cyclo-C ₃ H ₅ OH	7
			6.0(-10)	ICR	CH ₃ COCH ₃ ⁺	66
			2.0(-10)	ICR	Condensation only	72
			6.5±0.3(-10)	TI	CH ₃ COCH ₃ ⁺	20
			5.4±0.4(-10)	ICR	CH ₃ COCH ₃ ⁺	69
	+ CH ₃ COCH ₃ →	473	5.1(-10)	MS	CH ₃ COCH ₃ ⁺	69
		300	8.5±0.3(-10)	MS	Photoionization at 10.0eV, CH ₃ COCH ₃ ⁺	68
		300	7.6±0.3(-10)	MS	Photoionization at 11.7eV, CH ₃ COCH ₃ ⁺	68
C ₃ D ₆ O ⁺	+ CD ₃ COCD ₃ →	300	7.0±0.3(-10)	MS	Photoionization at 10.0eV, CD ₃ COCD ₃ ⁺	68
		300	6.8±0.3(-10)	MS	Photoionization at 11.7eV, CD ₃ COCD ₃ ⁺	68
C ₃ H ₇ O ⁺	+ CH ₂ CO →		2.98±0.10(-10)	ICR	(CH ₃) ₂ COH ⁺	67
	+ CH ₃ COCH ₃ →		4.0(-11)	ICR	(CH ₃) ₂ COH ⁺	66
		473	4.3(-10)	MS	12 V cm ⁻¹ , (CH ₃) ₂ COH ⁺	69
	+ 2CH ₃ COCH ₃ →		6.5±0.5(-25)	TI	(CH ₃) ₂ COH ⁺	20
	+ iso-C ₄ H ₉ →		7.1(-11)	ICR	(CH ₃) ₂ COH ⁺	92
	+ t-C ₄ H ₉ OH →		1.1(-9)	ICR	(CH ₃) ₂ COH ⁺	94
			3.7±0.5(-10)	ICR	(CH ₃) ₂ COH ⁺	95
C ₃ H ₈ D ₃ O ⁺	+ CD ₃ OC ₂ H ₅ →		1.43±0.05(-9)	TI	CD ₃ OC ₂ H ₅ ⁺	20
C ₃ H ₈ O ₂ ⁺	+ vinyl formate →	373	1.4(-9)	MS	From vinyl formate	16
C ₃ H ₈ O ₂ ⁺	+ t-C ₄ H ₉ OH →		6.8(-10)	ICR	(CH ₃) ₂ COHOH ₂ ⁺	94
C ₃ H ₈ F ⁺	+ C ₂ H ₃ F →		2.2±0.7(-10)	MS		78
	+ CH ₂ CF ₂ →	320	3.49(-10)	MS		38
C ₃ H ₈ F ⁺	+ 2-fluoropropane →		4.4±0.6(-10)	ICR	(CH ₃) ₂ CF ⁺	87
C ₃ H ₈ F ₂ ⁺	+ C ₂ H ₃ F →		5(-11)	MS		78
	+ CH ₂ CF ₂ →	320	1.24(-10)	MS		38
C ₃ H ₈ F ₃ ⁺	+ CH ₂ CF ₂ →	320	6.6(-11)	MS		38
	+ C ₂ HF ₃ →		3.0(-11)	ICR		90
C ₃ H ₁₀ Si ⁺	+ (CH ₃) ₃ SiH →		7(-11)	MS	(CH ₃) ₃ SiH ⁺	50
C ₃ H ₈ FSi ⁺	+ (CH ₃) ₃ SiF →		6.5(-10)	ICR	(CH ₃) ₃ SiF ⁺	51
C ₃ H ₈ P ⁺	+ (CH ₃) ₃ P →		5.9±1.2(-10)	ICR	(CH ₃) ₃ P ⁺	52
C ₃ H ₈ S ⁺	+ C ₃ H ₇ SH →		1.13±0.56(-9)	MS	Photoionization, C ₃ H ₇ SH ⁺	57
C ₃ H ₈ S ₂ ⁺	+ (CH ₃) ₂ S →		1.6±0.2(-11)	MS	10.7 V cm ⁻¹	54
C ₄ H ₉ N ⁺	+ C ₃ H ₅ CN →		1.78±0.54(-9)	ICR	C ₃ H ₅ CN ⁺	7
C ₄ H ₁₁ N ⁺	+ diethylamine →	298	1.25±0.06(-9)	MS	From diethylamine	9

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction	T	k	Method	Comments	Ref.
C ₄ H ₄ O ⁺ + toluene →		1.28±0.19(−9)	ICR	From furan	96
C ₄ H ₇ O ⁺ + diisopropyl ketone →		1.1(−10)	ICR	(CH ₃) ₂ CHCO ⁺	66
C ₄ H ₈ O ⁺ + 2-butanone →		9(−11)	ICR	CH ₃ COC ₂ H ₅ ⁺	66
C ₄ H ₉ O ⁺ + 2-butanone →		9.0(−11)	ICR	(CH ₃ COC ₂ H ₅)H ⁺	66
C ₄ H ₁₁ O ⁺ + <i>t</i> -butanol →		7.2(−10)	ICR	(CH ₃) ₃ COH ₂ ⁺	94
C ₄ H ₈ O ₂ ⁺ + biacetyl →		7.5±1.0(−10)	ICR	(CH ₃ CO) ₂ ⁺	97
+ pyrrole →		4.3(−10)	ICR	(CH ₃ CO) ₂ ⁺	97
C ₄ H ₁₃ O ₂ ⁺ + <i>t</i> -butanol →		2.3(−10)	ICR	(CH ₃) ₃ COH ₂ OH ₂ ⁺	94
C ₄ H ₁₀ S ⁺ + (CH ₃) ₃ CSH →		5.7(−10)	MS	(CH ₃) ₃ CSH ⁺	57
C ₅ H ₆ N ⁺ + (CH ₃) ₂ NH →		3.3(−10)	ICR	Protonated pyridine	92
C ₅ H ₁₂ N ⁺ + triethylamine →	298	1.28±0.13(−10)	MS	(C ₂ H ₅) ₂ NCH ₂ ⁺	9
C ₅ H ₈ O ⁺ + cyclo-C ₃ H ₅ COCH ₃ →		2.0±0.6(−10)	ICR	From cyclo-C ₃ H ₅ COCH ₃	7
C ₅ H ₁₀ O ⁺ + 3-pentanone →		8(−11)	ICR	From 3-pentanone	66
C ₅ H ₁₁ O ⁺ + 3-pentanone →		5(−11)	ICR	Protonated 3-pentanone	66
C ₅ H ₈ O ₂ ⁺ + CH ₃ COCH ₃ →	473	8(−11)	ICR	(CH ₃ COCH ₃)CH ₃ CO ⁺	66
+ C ₆ H ₅ NH ₂ →		5.4±0.4(−10)	ICR	(CH ₃ COCH ₃)CH ₃ CO ⁺	69
+ C ₆ H ₅ NO ₂ →		5.1(−10)	MS	(CH ₃ COCH ₃)CH ₃ CO ⁺	69
+ anisole (H-substituted) →		1.5(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	98
+ anisole (<i>o</i> -methyl substituted) →		4(−11)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>m</i> -methyl substituted) →		2.9±0.6(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>p</i> -methyl substituted) →		4.0±0.8(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>p</i> -ethyl substituted) →		5.4±1.1(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>p</i> -ethyl substituted) →		3.0±0.6(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>p</i> -ethyl substituted) →		3.7±0.7(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	71
+ <i>o</i> -cresol →		5.8±0.6(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	72
+ <i>m</i> -cresol →		6.1±0.6(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	72
+ <i>p</i> -cresol →		4.6±0.5(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	72
+ C ₆ H ₅ OH →		4.3(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	70
+ C ₆ H ₅ CHO →		8.5(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	70
+ C ₆ H ₅ OCH ₃ →		2.9(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	70
+ C ₆ H ₅ OCD ₃ →		2.9(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	70
+ C ₆ H ₅ COCH ₃ →		6.3(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	70
+ C ₆ H ₅ OC ₂ H ₅ →		4.8(−10)	ICR	CH ₃ CO(COCH ₃)CH ₃ ⁺	70
C ₅ H ₅ Fe ⁺ + Fe(C ₅ H ₅) ₂ →		2.4±0.5(−9)	MS	From cyclopentyl iron	99
+ C ₆ H ₅ Ni ⁺ + CH ₃ CHO →		7.9±0.7(−10)	ICR	From cyclopentyl iron	100
C ₆ H ₈ N ⁺ + (CH ₃) ₃ N →		8.0±1.6(−10)	ICR	from cyclopentyl nickel	101
+ (CH ₃) ₃ N →		6.5(−10)	ICR	Protonated 4-methyl	
C ₆ H ₁₂ N ⁺ + (C ₂ H ₅) ₃ N →	298	pyridine			92
C ₆ H ₁₄ N ⁺ + quinuclidine →		7.05±0.35(−10)	MS	(C ₂ H ₅) ₃ N ⁺	9
C ₆ H ₁₂ O ₂ ⁺ + 2-butanone →		5.6(−10)	ICR	(C ₂ H ₅) ₃ NH ⁺	92
C ₆ H ₉ O ₃ ⁺ + C ₆ H ₅ NH ₂ →		9.0(−11)	ICR	(CH ₃ COC ₂ H ₅)CH ₃ CO ⁺	66
+ <i>o</i> -cresol →		1.8(−9)	ICR	CH ₃ CO(COCH ₃)COCH ₃ ⁺	98
C ₆ H ₉ O ₃ ⁺ + <i>m</i> -cresol →		4.8±0.5(−10)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	72
+ <i>p</i> -cresol →		4.9±0.5(−10)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	72
+ anisole (H-substituted) →		4.9±0.5(−10)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	72
+ anisole (<i>o</i> -methyl substituted) →		1.4±0.2(−9)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>m</i> -methyl substituted) →		9.5±1.9(−10)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>p</i> -methyl substituted) →		1.6±0.3(−9)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	71
+ anisole (<i>p</i> -ethyl substituted) →		1.6±0.3(−9)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	71
+ pyrrole →		2.5±0.5(−9)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	71
+ C ₆ H ₅ OH →		1.1(−9)	ICR	CH ₃ COCO(COCH ₃)CH ₃ ⁺	71
+ C ₆ H ₅ NO ₂ →		4.2(−9)	ICR	CH ₃ COCO(COCH ₃)COCH ₃ ⁺	98
+ C ₆ H ₅ CN →		1.7(−9)	ICR	CH ₃ COCO(COCH ₃)COCH ₃ ⁺	98
+ C ₆ H ₅ CHO →		3.0(−9)	ICR	CH ₃ COCO(COCH ₃)COCH ₃ ⁺	98
+ C ₆ H ₅ OCH ₃ →		1.5(−9)	ICR	(CH ₃ CO) ₃ ⁺	70
+ C ₆ H ₅ OCD ₃ →		1.4(−9)	ICR	(CH ₃ CO) ₃ ⁺	70
+ C ₆ H ₅ COCH ₃ →		1.4(−9)	ICR	(CH ₃ CO) ₃ ⁺	70
+ C ₆ H ₅ OC ₂ H ₅ →		6.8(−9)	ICR	(CH ₃ CO) ₃ ⁺	70
+ C ₆ H ₅ OC ₂ H ₅ →		1.2(−9)	ICR	(CH ₃ CO) ₃ ⁺	70

TABLE 1. Rate coefficients^a for reactions of organic cations—Continued

Reaction	T	k	Method	Comments	Ref.	
$C_6H_5O_3^+$		9.0(−10)	ICR	$CH_3CO(COCH_3)COCH_3^+$	98	
		<3.0(−11)		$CH_3CO(COCH_3)COCH_3^+$	98	
$C_6H_5F^+$	295	1.02±0.15(−9)	ICR	From fluorobenzene	96	
		4±1(−26)		From fluorobenzene	102	
$C_6H_4F_2^+$	+ $C_6H_5F \rightarrow$	2.4±0.4(−10)	ICR	From <i>p</i> -difluorobenzene	96	
$C_6H_5Cl^+$	295	5.3±1.0(−26)	MS	From chlorobenzene	102	
		530		From chlorobenzene	103	
	570	1.1(−9)	MS	From chlorobenzene	103	
		1.4(−9)		From chlorobenzene	103	
$C_6H_5N^+$	+ $C_6H_6 \rightarrow$	500	1.3(−9)	MS	$C_6H_5CN^+$	103
$C_7H_{14}N^+$	+ $(C_2H_5)_3N \rightarrow$	3.3(−10)	ICR	Protonated quinulidine	92	
$C_7H_{14}O^+$	+ $(i-C_3H_7)_2CO \rightarrow$	2(−11)	ICR	$(i-C_3H_7)_2CO^+$	66	
$C_7H_{15}O^+$	+ $(i-C_3H_7)_2CO \rightarrow$	2(−11)	ICR	$(i-C_3H_7)_2COH^+$	66	
$C_7H_7F^+$	+ toluene →	2.8±0.4(−10)	ICR	From <i>p</i> -fluoro toluene	96	
$C_7H_5F_3^+$	+ $C_6F_5H \rightarrow$	1.16±0.17(−9)	ICR	$C_6H_5CF_3^+$	96	
$C_8H_{15}O_2^+$	+ $(C_2H_5)_2CO \rightarrow$	3(−11)	ICR	$(C_2H_5)_2COC_2H_5CO^+$	66	
$C_9H_{11}O_2^+$	360	2.8(−9)	MS	$(CH_3COOC_7H_7)H^+$	104	
		2(−11)		$(i-C_3H_7)_2COC_3H_7CO^+$	66	

^aRate coefficients are in units of $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ for bimolecular reactions and $\text{cm}^6 \text{molecule}^{-2} \text{s}^{-1}$ for termolecular reactions.

^bThe numbers in parenthesis should be read as the power 10 of the number. Thus (−9) is 10^{-9} .

^cThe reaction $CH_3OH^+ + CH_3OH \rightarrow$ products has been investigated in five different laboratories. The most recent determinations (refs. 10, 15, and 23) of the overall conversion coefficient were obtained by the ICR technique, and suggest an average value of 2.00 ± 0.17 (−9) for this process. All earlier values were obtained by single source mass spectrometry (refs. 17, 19, and 24). In each of those studies the rate coefficient for the reaction $CH_4^+ + CH_4 \rightarrow$ products was independently determined and reported. Taking the reference value which we suggested earlier [1.14 (−9), ref. 1] for the methane reaction together with the methanol values given in refs. 17, 19, and 24, and appropriately correcting the individual values given for methanol, an average value of 2.47 ± 0.16 (−9) is obtained for the MS studies. Although one is tempted to suggest that the recent ICR values are more accurate, it is disconcerting to note that ref. 10 (the most recent ICR study) also reports a value of 2.56 ± 0.17 (−9) for the reaction $CD_3OH^+ + CD_3OH \rightarrow$ products, which is essentially equivalent to the MS value for the unlabeled analogue. Taking all factors into consideration, and recognizing that the methanol reaction will exhibit a negative temperature and ion kinetic dependence due to ion-dipole interactions (ref. 23), it would seem that the actual value of the thermal rate coefficient (300 K) for the reaction $CH_3OH^+ + CH_3OH \rightarrow$ products other than CH_3OH^+ (resonance electron or hydrogen exchange also gives CH_3OH^+ , and would be undetected) should fall within the limits 2.2 ± 0.3 (−9). Additional careful measurements involving this reaction pair are encouraged since it appears to represent a model highly reactive oxygenated organic system.

^dTaking the most recent value from Beauchamp's laboratory (ref. 39), the lowest kinetic energy value from ref. 28, the lowest electron impact energy value of ref. 30, and the earlier ICR value of ref. 43, the current thermal value for this fast reaction falls within the limits 1.64 ± 0.22 (−9), which is essentially equivalent to a collision efficiency of unity for conversion to products. Contributions from resonant electron or atom exchange cannot be evaluated.

^eThe rate coefficients reported for this very fast reaction have remained remarkably consistent over the years, and a value of 2.12 ± 0.20 (−9) is suggested at moderate temperatures (~ 300 K) for thermal ions.

TABLE 2. Rate coefficients^a for reactions of organic anions

	Reaction	T	k	Method	Comments	Ref.
CH_4N^-	+ $\text{CH}_3\text{F} \rightarrow$	297	5(-11) ^b	FA	CH_3NH^-	105
	+ $\text{CH}_3\text{Cl} \rightarrow$	297	$1.7 \pm 0.1 (-9)$	FA	CH_3NH^-	105
	+ $\text{C}_2\text{H}_2 \rightarrow$	297	$1.33 \pm 0.33 (-9)$	FA	CH_3NH^-	106
	+ $\text{CH}_3\text{OH} \rightarrow$		$1.2 \pm 0.3 (-9)$	ICR		107
CH_3O^-	+ $\text{CH}_3\text{F} \rightarrow$	297	$1.4 \pm 0.2 (-11)$	FA		105
	+ $\text{CH}_3\text{Cl} \rightarrow$	297	$1.3 \pm 0.1 (-9)$	FA		105
	+ $\text{C}_2\text{H}_2 \rightarrow$	297	$1.64 \pm 0.41 (-9)$	FA		106
		300	$1.3 (-9)$	FA		108
	+ $\text{CH}_3\text{CN} \rightarrow$	297	$3.5 \pm 0.3 (-9)$	FA		109
	+ $\text{C}_2\text{H}_5\text{OH} \rightarrow$	300	$\geq 5 (-11)$	FA		110
	+ allene \rightarrow	300	$\sim 5 (-12)$	FA		110
	+ propylene \rightarrow	300	$< 5 (-12)$	FA		110
	+ $t\text{-C}_4\text{H}_9\text{OH} \rightarrow$	300	$\geq 5 (-11)$	FA		110
	+ $\text{C}_6\text{H}_5\text{CH}_3 \rightarrow$		$2.0 \pm 0.2 (-10)$	ICR		107
CD_3O^-		300	$\geq 5 (-12)$	FA		110
	+ $\text{C}_6\text{H}_5\text{CH}(\text{CH}_3)_2 \rightarrow$	300	$\leq 5 (-12)$	FA		110
	+ $\text{CH}_3\text{CH}_2\text{F} \rightarrow$		$1.3 (-10)$	ICR		111
	+ $\text{CH}_3\text{CHF}_2 \rightarrow$		$1.26 (-9)$	ICR		111
	+ $\text{CH}_2\text{FCHF}_2 \rightarrow$		$1.68 (-9)$	ICR		111
	+ $\text{CH}_3\text{CF}_3 \rightarrow$		$1.55 (-10)$	ICR		111
	+ $\text{CHF}_2\text{CHF}_2 \rightarrow$		$1.69 (-9)$	ICR		111
	+ $\text{CH}_2\text{FCF}_3 \rightarrow$		$8.8 (-10)$	ICR		111
	+ $\text{CHF}_2\text{CF}_3 \rightarrow$		$1.9 (-9)$	ICR		111
	+ $(\text{CH}_3)_3\text{B} \rightarrow$		$3.85 \pm 0.77 (-10)$	ICR		18
CH_2NO_2^-	+ $\text{CHCl}_3 \rightarrow$	300	$\leq 5 (-12)$	FA		110
	+ $n\text{-C}_4\text{H}_9\text{SH} \rightarrow$	300	$> 1.6 (-10)$	FA		110
	+ $\text{C}_5\text{H}_6 \rightarrow$	300	$< 1 (-11)$	FA		110
CHF^-	+ $\text{CH}_3\text{F} \rightarrow$	297	$\geq 4.1 (-10)$	FA		105
CF_3^-	+ $\text{NO}_2 \rightarrow$		$3.7 (-10)$	B	0.3 eV KE	112
	+ $\text{Xe} \rightarrow$		$3 (-12)$	B	0.3 eV KE	113
CH_3S^-	+ $\text{CH}_3\text{F} \rightarrow$	297	$\leq 1 (-12)$	FA		105
	+ $\text{CH}_3\text{Cl} \rightarrow$	297	$1.1 \pm 0.1 (-10)$	FA		105
CHCl_2^-			$7.8 \pm 1.2 (-11)$	ICR		114
	+ $\text{CH}_3\text{Br} \rightarrow$		$1.4 \pm 0.2 (-10)$	ICR		114
	+ $\text{CH}_3\text{Cl} \rightarrow$	297	$5.1 \pm 0.4 (-10)$	FA		105
	+ $\text{C}_2\text{H}_2 \rightarrow$	300	$\sim 3 (-11)$	FA		110
	+ $\text{CH}_3\text{CN} \rightarrow$	300	$\geq 2 (-10)$	FA		110
CCl_3^-	+ $\text{C}_2\text{H}_5\text{OH} \rightarrow$	300	$\leq 1 (-11)$	FA		110
	+ $\text{CH}_3\text{SOCH}_3 \rightarrow$	300	$\geq 1 (-11)$	FA		110
	+ $\text{CH}_3\text{NO}_2 \rightarrow$	300	$\geq 1 (-10)$	FA		110
	+ $\text{CH}_3\text{CN} \rightarrow$	300	$< 5 (-12)$	FA		110
	+ $\text{CH}_3\text{COCH}_3 \rightarrow$	300	$< 1 (-12)$	FA		110
$\text{C}_2\text{H}_2\text{N}^-$	+ $\text{C}_6\text{H}_6 \rightarrow$	300	$\sim 2 (-11)$	FA		110
	+ $\text{CHCl}_3 \rightarrow$	300	$\geq 1 (-10)$	FA	CH_2CN^-	110
	+ $\text{CH}_3\text{SOCH}_3 \rightarrow$	300	$< 1 (-11)$	FA	CH_2CN^-	110
$\text{C}_2\text{H}_5\text{N}^-$	+ $\text{CH}_3\text{COCH}_3 \rightarrow$	300	$\geq 1 (-10)$	FA	CH_2CN^-	110
	+ $\text{C}_2\text{H}_2 \rightarrow$	297	$1.04 \pm 0.26 (-9)$	FA	$(\text{CH}_3)_2\text{N}^-$	106
		297	$1.18 \pm 0.30 (-9)$	FA	$\text{C}_2\text{H}_5\text{NH}^-$	106
$\text{C}_2\text{H}_7\text{N}^-$	+ $\text{H}_2\text{O} \rightarrow$		$2.4 (-9)$	FA	$\text{C}_2\text{H}_5\text{NH}_2^-$	115
$\text{C}_2\text{H}_5\text{O}^-$	+ $\text{CH}_3\text{OH} \rightarrow$	300	$\geq 5 (-11)$	FA		110
	+ $\text{CH}_2\text{Cl}_2 \rightarrow$	300	$\geq 5 (-11)$	FA		110
$\text{C}_2\text{H}_6\text{O}^-$	+ $\text{C}_2\text{H}_2 \rightarrow$	297	$1.41 \pm 0.35 (-9)$	FA		106
		300	$1.1 (-9)$	FA		108
	+ $(\text{CH}_3)_3\text{B} \rightarrow$		$4.6 \pm 0.9 (-10)$	ICR		18
	+ $i\text{-C}_3\text{H}_7\text{OH} \rightarrow$	300	$\geq 5 (-11)$	FA		110
	+ $t\text{-C}_4\text{H}_9\text{OH} \rightarrow$	300	$\geq 5 (-11)$	FA		110
	+ $\text{C}_6\text{H}_5\text{CH}_3 \rightarrow$	300	$< 5 (-11)$	FA		110
	+ $\text{C}_6\text{H}_5\text{CH}(\text{CH}_3)_2 \rightarrow$	300	$\leq 1 (-11)$	FA		110

TABLE 2. Rate coefficients^a for reactions of organic cations—Continued

	Reaction	T	k	Method	Comments	Ref.
C ₂ H ₃ F ₂ ⁻	+ CH ₃ CHF ₂ →		1.5±0.4(-10)	ICR	CHF ₂ CH ₂ ⁻	111
C ₂ H ₂ F ₃ ⁻	+ CHF ₂ CH ₂ F →		1.5±0.2(-10)	ICR	CHF ₂ CHF ⁻	111
	+ CH ₃ CF ₃ →		7.9±0.9(-10)	ICR	CF ₃ CH ₃ ⁻	111
C ₂ HF ₄ ⁻	+ CHF ₂ CHF ₂ →		1.07±0.10(-9)	ICR	CHF ₂ CF ₂ ⁻	111
	+ CH ₂ FCF ₃ →		1.5±0.3(-10)	ICR	CF ₃ CFH ⁻	111
C ₂ F ₅ ⁻	+ NO ₃ →		3.2(-10)	B	0.3 eV KE	112
	+ Xe →		8(-12)	B	0.3 eV KE	113
C ₂ H ₅ OS ⁻	+ CH ₂ Cl ₂ →	300	≥2(-10)	FA	CH ₃ SOCH ₂ ⁻	110
	+ C ₂ H ₂ →	300	≤1(-11)	FA	CH ₃ SOCH ₂ ⁻	110
	+ CH ₃ CN →	300	≥2(-10)	FA	CH ₃ SOCH ₂ ⁻	110
	+ C ₆ H ₅ CH ₃ →	300	≤1(-12)	FA	CH ₃ SOCH ₂ ⁻	110
C ₃ H ₅ O ⁻	+ CHCl ₃ →	300	>2(-10)	FA	CH ₃ COCH ₂ ⁻	110
	+ CH ₃ CN →	300	≤3(-11)	FA	CH ₃ COCH ₂ ⁻	110
	+ CD ₃ COCD ₃ →		3.2±1.9(-12)	ICR	CH ₃ COCH ₂ ⁻	116
	+ (C ₂ H ₅) ₂ CO →		8±4(-12)	ICR	CH ₃ COCH ₂ ⁻	116
	+ CH ₃ COCD ₂ CODCH ₃ →		3.6±0.3(-10)	ICR	CH ₃ COCH ₂ ⁻	116
C ₃ D ₆ O ⁻	+ CH ₃ COC ₂ H ₅ →		6.4±4.0(-12)	ICR	CD ₃ COCD ₂ ⁻	116
C ₃ H ₇ O ⁻	+ C ₂ H ₂ →	297	≥8.9±2.3(-10)	FA	(CH ₃) ₂ CHO ⁻	106
	+ C ₂ H ₅ OH →	300	5.2(-10)	FA	i-C ₃ H ₇ O ⁻	108
	+ t-C ₄ H ₉ OH →	300	≥5(-11)	FA	i-C ₃ H ₇ O ⁻	110
	+ C ₄ H ₉ DO ⁻	300	≥5(-11)	FA	i-C ₃ H ₇ O ⁻	110
C ₄ H ₆ DO ⁻	+ CH ₃ COCD ₂ CH ₃ →		1.8±1.5(-12)	ICR	CH ₃ COCDCH ₃ ⁻	116
C ₄ H ₅ D ₂ O ⁻	+ CH ₃ COCD ₂ CH ₃ →		2.7±2.2(-12)	ICR	CH ₂ COCD ₂ CH ₃ ⁻	116
C ₄ H ₉ O ⁻	+ CH ₃ OH →	300	≥5(-11)	FA	t-C ₄ H ₉ O ⁻	110
	+ CH ₂ Cl ₂ →	300	≥1(-10)	FA	t-C ₄ H ₉ O ⁻	110
	+ C ₂ H ₂ →	300	1.6(-10)	FA	t-C ₄ H ₉ O ⁻	108
	+ C ₂ H ₅ OH →	300	≥5(-11)	FA	t-C ₄ H ₉ O ⁻	110
	+ i-C ₃ H ₇ OH →	300	≥5(-11)	FA	t-C ₄ H ₉ O ⁻	110
C ₄ F ₈ ⁻	+ NO ₂ →		1.0(-10)	B	0.3 eV KE	117
	+ SF ₆ →	300	<1(-10)	FA		118
C ₄ H ₉ S ⁻	+ CH ₃ NO ₂ →	300	≤2.6(-11)	FA	n-C ₄ H ₉ S ⁻	110
C ₅ H ₉ O ⁻	+ CH ₃ CD ₂ COCD ₂ CH ₃ →		3.8±3.0(-12)	ICR	CH ₃ CH ₂ COCHCH ₃ ⁻	116
C ₅ H ₇ O ₂ ⁻	+ C ₆ H ₅ SH →		3.0±0.3(-10)	ICR	From CH ₃ COCHCOHCH ₃	116
	+ parachlorophenol →		1.5±0.4(-10)	ICR	From CH ₃ COCHCOHCH ₃	116
C ₅ H ₆ DO ₂ ⁻	+ CD ₃ COCDCODCD ₃ →		1.2±0.3(-11)	ICR	CH ₃ COCDCOCH ₃ ⁻	116
C ₅ H ₅ D ₂ O ₂ ⁻	+ CH ₃ COCDCODCH ₃ →		3.0±0.5(-11)	ICR	CH ₂ COCDCODCH ₃ ⁻	116
C ₆ H ₅ O ⁻	+ parahydroxytoluene →		6.2±2.0(-11)	ICR	Phenoxy anion	116
C ₆ H ₅ NO ₂ ⁻	+ NO ₂ →		3.6(-10)	B	0.3 eV KE	117
C ₆ F ₆ ⁻	+ NO ₂ →		2.0(-10)	B	0.3 eV KE	117
C ₆ F ₁₀ ⁻	+ NO ₂ →		1.3(-11)	B	0.3 eV KE	117
C ₇ H ₇ O ⁻	+ phenol →		1.8±0.3(-10)	ICR	From parahydroxytoluene	116
C ₇ F ₈ ⁻	+ NO ₂ →		7.0(-11)	B	0.3 eV KE	117

^aRate coefficients are in units of cm³ molecule⁻¹ s⁻¹ for bimolecular reactions and cm⁶ molecule⁻² s⁻¹ for termolecular reactions.^bThe numbers in parenthesis should be read as the power 10 of the number. This (-9) is 10⁻⁹.

Acknowledgements

The author would like to express his sincere gratitude to Dr. Sharon Lias for her assistance in the literature search and indexing, to Dr. Pierre Ausloos for his valuable critique and encouragement, and to Ms. Jeanne Miller, of Yale University, for her help in structuring the tabular material. This work was supported by the Office of Standard Reference Data, National Bureau of Standards.

References

- [1] Sieck, L. W., and Lias, S. G., *J. Phys. Chem. Ref. Data*, **5**, 1123 (1976).
- [2] Meisels, G. G., 'Determination of Rate Coefficients of Ion-Molecule Reactions,' in *NATO Advanced Study Institute Series B6*, P. Ausloos, ed., Plenum Press, New York (1975).
- [3] Henchman, M. J., *Ion-Molecule Reactions*, J. L. Franklin, ed., **1**, 101 Plenum Press, New York (1972).
- [4] Ferguson, E. E., *Atomic and Nuclear Data Tables* **12**, 159 (1973).
- [5] Jones, E. G., and Harrison, A. G., *Can. J. Chem.* **45**, 3119 (1967).
- [6] Munson, M. S. B., *J. Phys. Chem.* **70**, 2034 (1966).
- [7] Luippold, D. A., and Beauchamp, J. L., *J. Phys. Chem.* **80**, 795 (1976).
- [8] Hellner, L., and Sieck, L. W., *Int. J. Chem. Kinet.* **5**, 177 (1973).
- [9] Brupbacher, J. M., Eagle, C. J., and Tschukow-Roux, E., *J. Phys. Chem.* **79**, 671 (1975).
- [10] McMahon, T. B., and Beauchamp, J. L., *J. Phys. Chem.* **81**, 593 (1977).
- [11] Meot-Ner, M., and Field, F. H., *J. Am. Chem. Soc.* **97**, 5339 (1975).
- [12] Harrison, A. G., Lin, P. H., and Tsang, C. W., *Int. J. Mass Spectrom. Ion Phys.* **19**, 23 (1976).
- [13] Foster, M. S., and Beauchamp, J. L., *J. Am. Chem. Soc.* **94**, 2425 (1972).
- [14] Karpas, Z., and Klein, F. S., *Int. J. Mass Spectrom. Ion Phys.* **16**, 289 (1975).
- [15] Vogt, J., and Beauchamp, J. L., *J. Am. Chem. Soc.* **97**, 6682 (1975).
- [16] Pritchard, H., and Harrison, A. G., *J. Chem. Phys.* **48**, 5623 (1968).
- [17] Gupta, S. K., Jones, E. G., Harrison, A. G., and Myher, J. J., *Can. J. Chem.* **45**, 3107 (1967).
- [18] Murphy, M. K., and Beauchamp, J. L., *J. Am. Chem. Soc.* **98**, 1433 (1976).
- [19] Ryan, K. R., Sieck, L. W., and Futrell, J. H., *J. Chem. Phys.* **41**, 111 (1964).
- [20] Blair, A. S., and Harrison, A. G., *Can. J. Chem.* **51**, 703 (1973).
- [21] Pritchard, H., Thynne, J. C. J., and Harrison, A. G., *Can. J. Chem.* **46**, 2141 (1968).
- [22] Peers, A. M., and Muller, J. C., *Int. J. Mass Spectrom. Ion Phys.* **16**, 321 (1975).
- [23] Bowers, M. T., Su, T., and Anicich, V. G., *J. Chem. Phys.* **58**, 5175 (1973).
- [24] Hyatt, D. J. Dodman, E. A., and Henchman, M. J., *Advan. Chem. Ser.*, No. 58, 131 (1966).
- [25] Lifshitz, C., and Tiernan, T. O., *J. Chem. Phys.* **57**, 1515 (1972).
- [26] Horton, R. L., Franklin, J. L., and Mazzeo, B., *J. Chem. Phys.* **62**, 1739 (1975).
- [27] Turner, D. L., and Bone, L. I., *J. Phys. Chem.* **78**, 501 (1974).
- [28] Herod, A. A., Harrison, A. G., and McAskill, N. A., *Can. J. Chem.* **49**, 2217 (1971).
- [29] Karpas, Z., and Klein, F. S., *Int. J. Mass Spectrom. Ion Phys.* **24**, 137 (1977).
- [30] Pabst, M. J. K., Tan, H. S., and Franklin, J. L., *Int. J. Mass Spectrom. Ion Phys.* **20**, 191 (1976).
- [31] McAskill, N. A., *Aust. J. Chem.* **23**, 893 (1970).
- [32] Sieck, L. W., Gorden, R., and Ausloos, P., *J. Res. Natl. Bur. Stand.* **78A**, 151 (1974).
- [33] Marcotte, R. E., and Tiernan, T. O., *J. Chem. Phys.* **54**, 4871 (1971).
- [34] Su, T., Kevan, L., and Tiernan, T. O., *J. Chem. Phys.* **54**, 4871 (1971).
- [35] Ravishankara, A. R., Eyler, J. R., and Hanrahan, R. J., *Int. J. Mass Spectrom. Ion Phys.* **22**, 315 (1976).
- [36] Krause, J. R., and Lampe, F. W., *J. Phys. Chem.* **81**, 281 (1977).
- [37] McAskill, N. A., *Aust. J. Chem.* **23**, 2301 (1970).
- [38] Tan, H. S., Pabst, M. J. K., and Franklin, J. L., *Int. J. Mass Spectrom. Ion Phys.* **20**, 247 (1976).
- [39] Blint, R. J., McMahon, T. B., and Beauchamp, J. L., *J. Am. Chem. Soc.* **96**, 1269 (1974).
- [40] McMahon, T. B., Blint, R. J., Ridge, D. P., and Beauchamp, J. L., *J. Am. Chem. Soc.* **94**, 8934 (1972).
- [41] Lias, S. G., and Ausloos, P., *Int. J. Mass Spectrom. Ion Phys.* **23**, 273 (1977).
- [42] Tan, H. S., Pabst, M. J. K., and Franklin, J. L., *Int. J. Mass Spectrom. Ion Phys.* **21**, 297 (1976).
- [43] Marshall, A. G., and Butrill, S. E., *J. Chem. Phys.* **52**, 2752 (1970).
- [44] Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., *J. Am. Chem. Soc.* **94**, 2798 (1972).
- [45] Lias, S. G., Eyler, J. R., and Ausloos, P., *Int. J. Mass Spectrom. Ion Phys.* **19**, 219 (1976).
- [46] Lias, S. G., and Ausloos, P., *Int. J. Mass Spectrom. Ion Phys.* **22**, 135 (1976).
- [47] Heckel, E., and Hanrahan, F. J., *J. Chem. Phys.* **62**, 1027 (1975).
- [48] Mayer, T. M., and Lampe, F. W., *J. Phys. Chem.* **78**, 2424 (1974).
- [49] Klevan, L., and Munson, M. S. B., *Int. J. Mass Spectrom. Ion Phys.* **13**, 261 (1974).
- [50] Potzinger, P., and Lampe, F. W., *J. Phys. Chem.* **75**, 13 (1971).
- [51] Murphy, M. K., and Beauchamp, J. L., *J. Am. Chem. Soc.* **98**, 5781 (1976).
- [52] Staley, R. H., and Beauchamp, J. L., *J. Am. Chem. Soc.* **96**, 6252 (1974).
- [53] McAllister, T., *Int. J. Mass Spectrom. Ion Phys.* **13**, 63 (1974).
- [54] Ruska, W. E. W., and Franklin, J. L., *Int. J. Mass Spectrom. Ion Phys.* **3**, 221 (1969).
- [55] Nagy, G. P., Thynne, J. C. J., and Harrison, A. G., *Can. J. Chem.* **46**, 3609 (1968).

- [56] Solka, B. H., and Harrison, A. G., Int. J. Mass Spectrom. Ion Phys. **14**, 295 (1974).
- [57] Bruppacher, J. M., Eagle, C. J., Koprio, J. A., and Tschiukow-Roux, E., J. Phys. Chem. **81**, 1125 (1977).
- [58] Solka, B. H., and Harrison, A. G., Int. J. Mass Spectrom. Ion Phys. **17**, 379 (1975).
- [59] McAskill, N. A., Aust. J. Chem. **22**, 2275 (1969).
- [60] Yerger, A. J., and Franklin, J. L., Int. J. Mass Spectrom. Ion Phys. **9**, 354 (1972).
- [61] Sieck, L. W., and Gorden, R., Int. J. Chem. Kinet. **5**, 445 (1973).
- [62] Sieck, L. W., J. Res. Nat. Bur. Stand. **81A**, 267 (1977).
- [63] Bach, R. D., Weibel, A. T., Patane, J., and Kevan, L., J. Am. Chem. Soc. **98**, 6237 (1976).
- [64] Franklin, J. L., Wada, Y., Natalis, P., and Hierl, P. M., J. Phys. Chem. **70**, 2353 (1966).
- [65] Shannon, T. W., and Harrison, A. G., J. Chem. Phys. **43**, 4206 (1965).
- [66] Van der Hart, W. J., and van Sprang, H. A., J. Am. Chem. Soc. **99**, 32 (1977).
- [67] Ausloos, P., and Lias, S. G., Chem. Phys. Lett. **51**, 53 (1977).
- [68] Sieck, L. W., and Ausloos, P., Rad. Res. **52**, 47 (1972).
- [69] MacNeil, K. A. G., and Futrell, J. H., J. Phys. Chem. **76**, 409 (1972).
- [70] Chatfield, D. A., and Bursey, M. M., Int. J. Mass Spectrom. Ion Phys. **18**, 239 (1975).
- [71] Chatfield, D. A., and Bursey, M. M., J. Am. Chem. Soc. **98**, 6492 (1976).
- [72] Chatfield, D. A., and Bursey, M. M., J. Am. Chem. Soc. **97**, 3600 (1975).
- [73] Corderman, R. R., LeBreton, P. R., Buttrill, S. E., Williamson, A. D., and Beauchamp, J. L., J. Chem. Phys. **65**, 4929 (1976).
- [74] Sieck, L. W., Abramson, F. P., and Futrell, J. H., J. Chem. Phys. **45**, 2859 (1966).
- [75] Gorden, R., and Sieck, L. W., J. Res. Nat. Bur. Stand. **76A**, 655 (1972).
- [76] Beauchamp, J. L., and Dunbar, R. C., J. Am. Chem. Soc. **92**, 1477 (1970).
- [77] Ryan, K. R., and Futrell, J. H., J. Chem. Phys. **42**, 824 (1965).
- [78] Tan, H. S., Pabst, M. J. K., and Franklin, J. L., Int. J. Mass Spectrom. Ion Phys. **19**, 163 (1976).
- [79] Herman, J. A., and Harrison, A. G., Can. J. Chem. **47**, 957 (1969).
- [80] Ferrer-Correia, A. J., and Jennings, K. R., Int. J. Mass Spectrom. Ion Phys. **11**, 111 (1973).
- [81] Anicich, V. G., and Bowers, M. T., Int. J. Mass Spectrom. Ion Phys. **13**, 359 (1974).
- [82] Williamson, A. D., and Beauchamp, J. L., J. Chem. Phys. **65**, 3196 (1976).
- [83] Sieck, L. W., Gorden, R., Lias, S. G., and Ausloos, P., Int. J. Mass Spectrom. Ion Phys. **15**, 181 (1974).
- [84] Anicich, V. G., and Bowers, M. T., Int. J. Mass Spectrom. Ion Phys. **12**, 231 (1973).
- [85] Anicich, V. G., and Bowers, M. T., Int. J. Mass Spectrom. Ion Phys. **11**, 329 (1973).
- [86] McAskill, N. A., and Harrison, A. G., Int. J. Mass Spectrom. Ion Phys. **5**, 193 (1970).
- [87] Beauchamp, J. L., and Park, J. Y., J. Phys. Chem. **80**, 575 (1976).
- [88] Anicich, V. G., and Bowers, M. T., Int. J. Mass Spectrom. Ion Phys. **13**, 351 (1974).
- [89] O'Malley, R. M., Jennings, K. R., Bowers, M. T., and Anicich, V. G., Int. J. Mass Spectrom. Ion Phys. **11**, 89 (1973).
- [90] Anicich, V. G., Bowers, M. T., O'Malley, R. M., and Jennings, K. R., Int. J. Mass Spectrom. Ion Phys. **11**, 99 (1973).
- [91] Herman, J. A., Myher, J. J., and Harrison, A. G., Can. J. Chem. **47**, 647 (1969).
- [92] Davidson, W. R., Bowers, M. T., Su, T., and Aue, D. H., Int. J. Mass Spectrom. Ion Phys. **24**, 83 (1977).
- [93] DeStefano, A. J., and Porter, R. F., J. Phys. Chem. **80**, 2918 (1976).
- [94] Beauchamp, J. L., Caserio, M. C., and McMahon, T. B., J. Am. Chem. Soc. **96**, 6243 (1974).
- [95] Beauchamp, J. L., J. Am. Chem. Soc. **91**, 5925 (1969).
- [96] Lias, S. G., and Ausloos, P., J. Am. Chem. Soc. **99**, 4831 (1977).
- [97] Dunbar, R. C., Bursey, M. M., and Chatfield, D. A., Int. J. Mass Spectrom. Ion Phys. **13**, 195 (1974).
- [98] Chatfield, D. A., and Bursey, M. M., Int. J. Mass Spectrom. Ion Phys. **20**, 101 (1976).
- [99] Schildcrout, S. M., J. Am. Chem. Soc. **95**, 3846 (1973).
- [100] Foster, M. S., and Beauchamp, J. L., J. Am. Chem. Soc. **97**, 4814 (1975).
- [101] Corderman, R. R., and Beauchamp, J. L., J. Am. Chem. Soc. **98**, 5700 (1976).
- [102] Sieck, L. W., and Gorden, R., Int. J. Mass Spectrom. Ion Phys. **19**, 269 (1976).
- [103] Meot-Ner, M., and Field, F. H., Chem. Phys. Lett. **44**, 484 (1976).
- [104] Meot-Ner, M., Hunter, E. P., and Field, F. H., J. Am. Chem. Soc. **99**, 5576 (1977).
- [105] Tanaka, K., Mackay, G. I., Payzant, J. D., and Bohme, D. K., Can. J. Chem. **54**, 1643 (1976).
- [106] Mackay, G. I., Tanaka, K., and Bohme, D. K., Int. J. Mass Spectrom. Ion Phys. **24**, 125 (1977).
- [107] Brauman, J. I., Lieder, C. A., and White, M. J., J. Am. Chem. Soc. **95**, 927 (1973).
- [108] Bohme, D. K., Lee-Ruff, E., and Young, L. B., J. Am. Chem. Soc. **93**, 4608 (1971).
- [109] Mackay, G. I., Betowski, L. D., Payzant, J. D., Schiff, H. I., and Bohme, D. K., J. Phys. Chem. **80**, 2919 (1976).
- [110] Bohme, D. K., Lee-Ruff, E., and Young, L. B., J. Am. Chem. Soc. **94**, 5153 (1972).
- [111] Ridge, D. P., and Beauchamp, J. L., J. Am. Chem. Soc. **96**, 3595 (1974).
- [112] Hughes, B. M., Lifshitz, C., and Tiernan, T. O., J. Chem. Phys. **59**, 3162 (1973).
- [113] Su, T., Kevan, L., and Tiernan, T. O., J. Phys. Chem. **75**, 2534 (1971).
- [114] Brauman, J. I., and Olmstead, W. N., J. Am. Chem. Soc. **96**, 4030 (1974).
- [115] Betowski, D., Payzant, J. D., Mackay, G. I., and Bohme, D. K., Chem. Phys. Lett. **31**, 321 (1975).
- [116] Farneth, W. E., and Brauman, J. I., J. Am. Chem. Soc. **98**, 7891 (1976).
- [117] Lifshitz, C., Tiernan, T. O., and Hughes, B. M., J. Chem. Phys. **59**, 3182 (1973).
- [118] Fehsenfeld, F. C., J. Chem. Phys. **54**, 438 (1971).

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET		1. PUBLICATION OR REPORT NO. NSRDS-NBS 64	2. Gov't Accession No.	3. Recipient's Accession No.
4. TITLE AND SUBTITLE Rate Coefficients for Ion-Molecule Reactions. Organic Ions Other Than Those Containing Only C and H		5. Publication Date February 1979		
7. AUTHOR(S) L. Wayne Sieck		8. Performing Organ. Report No.		
9. PERFORMING ORGANIZATION NAME AND ADDRESS NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, D.C. 20234		10. Project/Task/Work Unit No.		
12. Sponsoring Organization Name and Complete Address (Street, City, State, ZIP) Same as Item 9		13. Type of Report & Period Covered N/A		
15. SUPPLEMENTARY NOTES Library of Congress Catalog Card Number: 78-10710				
16. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here.) A compilation is presented of all experimentally determined bimolecular and termolecular rate coefficients for the reactions of organic ions (other than those containing only C and H) with neutral molecules in the vapor phase. The literature covered is from 1960 to the present, and both positive and negative ions are considered. Five hundred and seventy-seven separate reaction-partners are tabulated, and experimental conditions are specified wherever possible. Preferred values are suggested for a number of these processes.				
17. KEY WORDS (six to twelve entries; alphabetical order; capitalize only the first letter of the first key word unless a proper name; separated by semicolons) Chemical kinetics; data evaluation; gas phase ion-molecule reactions; mass spectrometry; organic molecules; rate coefficients.				
18. AVAILABILITY <input checked="" type="checkbox"/> Unlimited <input type="checkbox"/> For Official Distribution. Do Not Release to NTIS <input checked="" type="checkbox"/> Order From Sup. of Doc., U.S. Government Printing Office Washington, D.C. 20402, Stock No. 003-003-02027-3 <input type="checkbox"/> Order From National Technical Information Service (NTIS) Springfield, Virginia 22151		19. SECURITY CLASS (THIS REPORT) UNCLASSIFIED	21. NO. OF PAGES 27	
		20. SECURITY CLASS (THIS PAGE) UNCLASSIFIED	22. Price \$1.30	

**Where can you find
all the reference data
you need?**


**Right in the
Journal of Physical
and Chemical
Reference Data!**

Now in its sixth year, this valuable publication has proved that it fills the important gaps for you in the literature of the physical sciences.

Published by the American Institute of Physics and the American Chemical Society for the National Bureau of Standards, this quarterly gives you quantitative numerical data, with recommended values and uncertainty limits chosen by experts in the field.

Critical commentary on methods of measurement and sources of error, as well as full references to the original literature, is an integral part of each of your four issues a year.

Can you afford to be without this prime source of reliable data on physical and chemical properties? To start receiving your copies, just fill in the order form and drop into the mail. If you do use a purchase order, please attach the printed form as this will help us to expedite your order. Send for complete list of reprints!



Journal of Physical and Chemical Reference Data
American Chemical Society
1155 Sixteenth Street, N.W., Washington, D.C. 20036

Yes, I would like to receive the JOURNAL OF PHYSICAL AND CHEMICAL REFERENCE DATA at the one-year rate checked below:

Name _____

Home

Business

Street _____

Members Nonmembers

City _____ State _____ Zip _____

*U.S., Canada,
Mexico* \$25.00 \$100.00

Other Countries \$29.00 \$104.00

Bill me

Bill company or school

Payment enclosed

Please Attach This Order Form To Purchase Order.

**Announcement of New Publications in
National Standard Reference Data Series**

Superintendent of Documents,
Government Printing Office,
Washington, D.C. 20402

Dear Sir:

Please add my name to the announcement list of new publications to be issued in the series: National Standard Reference Data Series—National Bureau of Standards.


Name_____

Company_____

Address_____

City_____ State_____ Zip Code_____

(Notification Key N-519)



NBS TECHNICAL PUBLICATIONS

PERIODICALS

JOURNAL OF RESEARCH—The Journal of Research of the National Bureau of Standards reports NBS research and development in those disciplines of the physical and engineering sciences in which the Bureau is active. These include physics, chemistry, engineering, mathematics, and computer sciences. Papers cover a broad range of subjects, with major emphasis on measurement methodology, and the basic technology underlying standardization. Also included from time to time are survey articles on topics closely related to the Bureau's technical and scientific programs. As a special service to subscribers each issue contains complete citations to all recent NBS publications in NBS and non-NBS media. Issued six times a year. Annual subscription: domestic \$17.00; foreign \$21.25. Single copy, \$3.00 domestic; \$3.75 foreign.

Note: The Journal was formerly published in two sections: Section A "Physics and Chemistry" and Section B "Mathematical Sciences."

DIMENSIONS/NBS

This monthly magazine is published to inform scientists, engineers, businessmen, industry, teachers, students, and consumers of the latest advances in science and technology, with primary emphasis on the work at NBS. The magazine highlights and reviews such issues as energy research, fire protection, building technology, metric conversion, pollution abatement, health and safety, and consumer product performance. In addition, it reports the results of Bureau programs in measurement standards and techniques, properties of matter and materials, engineering standards and services, instrumentation, and automatic data processing.

Annual subscription: Domestic, \$11.00; Foreign \$13.75

NONPERIODICALS

Monographs—Major contributions to the technical literature on various subjects related to the Bureau's scientific and technical activities.

Handbooks—Recommended codes of engineering and industrial practice (including safety codes) developed in cooperation with interested industries, professional organizations, and regulatory bodies.

Special Publications—Include proceedings of conferences sponsored by NBS, NBS annual reports, and other special publications appropriate to this grouping such as wall charts, pocket cards, and bibliographies.

Applied Mathematics Series—Mathematical tables, manuals, and studies of special interest to physicists, engineers, chemists, biologists, mathematicians, computer programmers, and others engaged in scientific and technical work.

National Standard Reference Data Series—Provides quantitative data on the physical and chemical properties of materials, compiled from the world's literature and critically evaluated. Developed under a world-wide program co-ordinated by NBS. Program under authority of National Standard Data Act (Public Law 90-396).

NOTE: At present the principal publication outlet for these data is the Journal of Physical and Chemical Reference Data (JPCRD) published quarterly for NBS by the American Chemical Society (ACS) and the American Institute of Physics (AIP). Subscriptions, reprints, and supplements available from ACS, 1155 Sixteenth St. N.W., Wash., D.C. 20056.

Building Science Series—Disseminates technical information developed at the Bureau on building materials, components, systems, and whole structures. The series presents research results, test methods, and performance criteria related to the structural and environmental functions and the durability and safety characteristics of building elements and systems.

Technical Notes—Studies or reports which are complete in themselves but restrictive in their treatment of a subject. Analogous to monographs but not so comprehensive in scope or definitive in treatment of the subject area. Often serve as a vehicle for final reports of work performed at NBS under the sponsorship of other government agencies.

Voluntary Product Standards—Developed under procedures published by the Department of Commerce in Part 10, Title 15, of the Code of Federal Regulations. The purpose of the standards is to establish nationally recognized requirements for products, and to provide all concerned interests with a basis for common understanding of the characteristics of the products. NBS administers this program as a supplement to the activities of the private sector standardizing organizations.

Consumer Information Series—Practical information, based on NBS research and experience, covering areas of interest to the consumer. Easily understandable language and illustrations provide useful background knowledge for shopping in today's technological marketplace.

Order above NBS publications from: Superintendent of Documents, Government Printing Office, Washington, D.C. 20402.

Order following NBS publications—NBSIR's and FIPS from the National Technical Information Services, Springfield, Va. 22161.

Federal Information Processing Standards Publications (FIPS PUB)—Publications in this series collectively constitute the Federal Information Processing Standards Register. Register serves as the official source of information in the Federal Government regarding standards issued by NBS pursuant to the Federal Property and Administrative Services Act of 1949 as amended, Public Law 89-306 (79 Stat. 1127), and as implemented by Executive Order 11717 (38 FR 12315, dated May 11, 1973) and Part 6 of Title 15 CFR (Code of Federal Regulations).

NBS Interagency Reports (NBSIR)—A special series of interim or final reports on work performed by NBS for outside sponsors (both government and non-government). In general, initial distribution is handled by the sponsor; public distribution is by the National Technical Information Services (Springfield, Va. 22161) in paper copy or microfiche form.

BIBLIOGRAPHIC SUBSCRIPTION SERVICES

The following current-awareness and literature-survey bibliographies are issued periodically by the Bureau:

Cryogenic Data Center Current Awareness Service. A literature survey issued biweekly. Annual subscription: Domestic, \$25.00; Foreign, \$30.00.

Liquified Natural Gas. A literature survey issued quarterly. Annual subscription: \$20.00.

Superconducting Devices and Materials. A literature survey issued quarterly. Annual subscription: \$30.00. Send subscription orders and remittances for the preceding bibliographic services to National Bureau of Standards, Cryogenic Data Center (275.02) Boulder, Colorado 80302.

U.S. DEPARTMENT OF COMMERCE
National Bureau of Standards
Washington, D.C. 20234

OFFICIAL BUSINESS

Penalty for Private Use, \$300

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-215



SPECIAL FOURTH-CLASS RATE
BOOK
