

NBS  
PUBLICATIONS

A11102 466773

NAT'L INST OF STANDARDS & TECH R.I.C.



A1102466773

Ross, Alberta B/Rate constants for react  
QC100 .U573 NO.70;1982 C.1 NBS-PUB-C 198



NSRDS-NBS 70

U.S. DEPARTMENT OF COMMERCE / National Bureau of Standards



# Rate Constants for Reactions of Aliphatic Carbon-Centered Radicals in Aqueous Solution

QC  
100  
.U573  
No. 70  
1982

## NATIONAL BUREAU OF STANDARDS

The National Bureau of Standards<sup>1</sup> was established by an act of Congress on March 3, 1901. The Bureau's overall goal is to strengthen and advance the Nation's science and technology and facilitate their effective application for public benefit. To this end, the Bureau conducts research and provides: (1) a basis for the Nation's physical measurement system, (2) scientific and technological services for industry and government, (3) a technical basis for equity in trade, and (4) technical services to promote public safety. The Bureau's technical work is performed by the National Measurement Laboratory, the National Engineering Laboratory, and the Institute for Computer Sciences and Technology.

**THE NATIONAL MEASUREMENT LABORATORY** provides the national system of physical and chemical and materials measurement; coordinates the system with measurement systems of other nations and furnishes essential services leading to accurate and uniform physical and chemical measurement throughout the Nation's scientific community, industry, and commerce; conducts materials research leading to improved methods of measurement, standards, and data on the properties of materials needed by industry, commerce, educational institutions, and Government; provides advisory and research services to other Government agencies; develops, produces, and distributes Standard Reference Materials; and provides calibration services. The Laboratory consists of the following centers:

Absolute Physical Quantities<sup>2</sup> — Radiation Research — Chemical Physics —  
Analytical Chemistry — Materials Science

**THE NATIONAL ENGINEERING LABORATORY** provides technology and technical services to the public and private sectors to address national needs and to solve national problems; conducts research in engineering and applied science in support of these efforts; builds and maintains competence in the necessary disciplines required to carry out this research and technical service; develops engineering data and measurement capabilities; provides engineering measurement traceability services; develops test methods and proposes engineering standards and code changes; develops and proposes new engineering practices; and develops and improves mechanisms to transfer results of its research to the ultimate user. The Laboratory consists of the following centers:

Applied Mathematics — Electronics and Electrical Engineering<sup>2</sup> — Manufacturing Engineering — Building Technology — Fire Research — Chemical Engineering<sup>2</sup>

**THE INSTITUTE FOR COMPUTER SCIENCES AND TECHNOLOGY** conducts research and provides scientific and technical services to aid Federal agencies in the selection, acquisition, application, and use of computer technology to improve effectiveness and economy in Government operations in accordance with Public Law 89-306 (40 U.S.C. 759), relevant Executive Orders, and other directives; carries out this mission by managing the Federal Information Processing Standards Program, developing Federal ADP standards guidelines, and managing Federal participation in ADP voluntary standardization activities; provides scientific and technological advisory services and assistance to Federal agencies; and provides the technical foundation for computer-related policies of the Federal Government. The Institute consists of the following centers:

Programming Science and Technology — Computer Systems Engineering.

<sup>1</sup>Headquarters and Laboratories at Gaithersburg, MD, unless otherwise noted; mailing address Washington, DC 20234.

<sup>2</sup>Some divisions within the center are located at Boulder, CO 80303.

QC100  
.U573  
#70  
1982

# Rate Constants for Reactions of Aliphatic Carbon-Centered Radicals in Aqueous Solution

---

Alberta B. Ross and Pedatsur Neta

Radiation Chemistry Data Center  
Radiation Laboratory  
University of Notre Dame  
Notre Dame, IN 46556



---

U.S. DEPARTMENT OF COMMERCE, Malcolm Baldrige, Secretary  
NATIONAL BUREAU OF STANDARDS, Ernest Ambler, Director

Issued October 1982

**Library of Congress Cataloging in Publication Data**

Ross, Alberta B.

Rate constants for reactions of aliphatic carbon-centered radicals in aqueous solution.

(NSRDS-NBS ; 70)

"Issued October 1982."

1. Chemical reaction, Rate of—Tables. 2. Radicals (Chemistry)—  
Tables. I. Neta, Pedatsur. II. Title. III. Series.

QC100.U573 no. 70 [QD502]	602'.18s	82-2096
	[547.1'394]	AACR2

**NSRDS-NBS 70**

Natl. Stand. Ref. Data Ser., Natl. Bur. Stand. (U.S.), 70, 103 pages (Oct. 1982)

CODEN: NSRDAP

© 1982 by the Secretary of Commerce on Behalf of the United States Government

U.S. GOVERNMENT PRINTING OFFICE  
WASHINGTON: 1982

---

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402

Price \$5.50

(Add 25 percent for other than U.S. mailing).



## Foreword


The National Standard Reference Data System provides access to the quantitative data of physical science, critically evaluated and compiled for convenience and readily accessible through a variety of distribution channels. The System was established in 1963 by action of the President's Office of Science and Technology and the Federal Council for Science and Technology, and responsibility to administer it was assigned to the National Bureau of Standards.

NSRDS receives advice and planning assistance from a Review Committee of the National Research Council of the National Academy of Sciences-National Academy of Engineering. A number of Advisory Panels, each concerned with a single technical area, meet regularly to examine major portions of the program, assign relative priorities, and identify specific key problems in need of further attention. For selected specific topics, the Advisory Panels sponsor subpanels which make detailed studies of users' needs, the present state of knowledge, and existing data resources as a basis for recommending one or more data compilation activities. This assembly of advisory services contributes greatly to the guidance of NSRDS activities.

The System now includes a complex of data centers and other activities in academic institutions and other laboratories. Components of the NSRDS produce compilations of critically evaluated data, reviews of the state of quantitative knowledge in specialized areas, and computations of useful functions derived from standard reference data. The centers and projects also establish criteria for evaluation and compilation of data and recommend improvements in experimental techniques. They are normally associated with research in the relevant field.

The technical scope of NSRDS is indicated by the categories of projects active or being planned: nuclear properties, atomic and molecular properties, solid state properties, thermodynamic and transport properties, chemical kinetics, and colloid and surface properties.

Reliable data on the properties of matter and materials are a major foundation of scientific and technical progress. Such important activities as basic scientific research, industrial quality control, development of new materials for building and other technologies, measuring and correcting environmental pollution depend on quality reference data. In NSRDS, the Bureau's responsibility to support American science, industry, and commerce is vitally fulfilled.



ERNEST AMBLER, *Director*

## Preface

This report is one of a series of data publications on radiation chemistry; the aim of the series is to compile, evaluate, and present the numerical results on processes occurring in systems which have been subjected to ionizing radiation. Various kinds of data are important in radiation chemistry. The quantities which were measured first were the observed radiation yields or  $G$  values (molecules formed or destroyed per 100 eV). Various indirect methods based on  $G$  values have been used to determine yields of transient species and relative rates of reactions. The spectral properties (optical, electron spin resonance) of transients have provided a direct method for their identification, and rates of the very fast reactions of transients which occur in irradiated systems have been measured directly by spectroscopic methods. Conductivity and luminescence methods have also provided a means of measuring properties of transients and their kinetics. Some reactions which occur in irradiated systems have also been studied by other methods, such as photochemistry, electric discharge, ultrasonics, chemical initiation, electron impact, etc. The emphasis in these publications is on the data of radiation chemistry, but where other pertinent data exist, they are included.

The data of radiation chemistry are voluminous; thousands of systems have been investigated. As a result there are certain collections, *e.g.* rate constants of particular types of reactions or certain properties of transients, for which tabulations of the data are considered essential, but for which critical assessment of each value is impossible. On the other hand, certain systems and properties have been studied so extensively that critical examination of these data is desirable and timely. Authors of this series of data publications have been asked to evaluate the extent to which the data can be critically assessed, to describe their criteria for evaluation, and to designate preferred values whenever possible.

## Contents

	Page
Introduction .....	1
List of Abbreviations and Symbols .....	3
List of Aliphatic Radicals .....	4
Table 1.    Rate Constants for Reactions of Aliphatic Radicals with Inorganic Solutes in Aqueous Solution .....	7
Table 2.    Rate Constants for Reactions of Aliphatic Radicals with Organic Solutes in Aqueous Solution .....	33
Table 3.    Rate Constants for Acid- and Base-catalyzed Proton Transfer Reactions of Aliphatic Radicals in Aqueous Solution .....	73
Table 4.    Rate Constants for Second Order Decay of Selected Aliphatic Radicals in Aqueous Solution .....	77
Index to Radical Reactions .....	80
Radical Formula Index .....	83
Substrate Formula Index .....	85
References .....	89



e.g. the OH radical abstracts hydrogen from alcohols mostly from the alpha position and to some extent from other sites. As a result the system contains a mixture of the reducing  $\alpha$ -hydroxyalkyl radicals and the much less reactive  $\beta$ -hydroxy- and  $\gamma$ -hydroxy-, etc. or alkoxyl radicals. b. Both OH and H may react with the substrate but yield different radicals, e.g. OH and H adducts. c. The radical produced by the initial attack may undergo certain processes before reacting with the substrate, e.g. protonation on carbon, acid-base equilibria, water elimination. Under certain conditions more than one form of the radical may react with the substrate. d. The aliphatic radical may react with the substrate by more than one mechanism, e.g. electron transfer and addition, of which only one is spectrophotometrically monitored. e. The aliphatic radical may react with the substrate slowly and undergo radical-radical decay as a competing process.

In many of the above cases, the reaction observed constitutes the main process (>90%) so that the rate constant measured can be taken to represent that reaction. However, when the observation is made on a less dominant process, the experimental rate constant may not represent the real value for that reaction. In these cases, the experimental values are reported in the tables and marked "uncorrected". The corrected value can sometimes be derived after considering the unobserved reactions.

## Arrangement of the Tables

A numbered list of the radicals for which reaction rate data have been compiled precedes the tables. Numbers have been assigned to the radicals for use in numbering the reactions in subsequent tables, to identify the radical and to provide a means of generating an index. A name and line formula are listed for most of the radicals. Line formulae for radicals containing rings are written with dashes at the left and right to indicate the cycle, as  $\text{-O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2\text{-}$  for the radical from dioxane (2,5-dioxacyclohexyl, 104). When more than one radical species may be involved the radicals are identified only by their origin, as "Radicals from glucose + OH".

Tables 1-3 contain the rate data for the reactions of the radicals with various substrates. Inorganic ions and molecules are grouped in table 1, in alphabetic order by the symbol for the main element. Organic substrates are listed alphabetically by name in table 2. Table 3 contains proton transfer reactions of radicals with acids, bases and buffer components. Entries for each substrate are numbered, and each reaction has an entry number including both the number of the substrate and the radical number. For example, the reaction of oxygen with methyl radical is 93.001, while oxygen with carboxymethyl radical is 93.035. Reactions of substrates with radicals in more than one acid-base form are listed in the tables under the same entry number.

Table 4 contains a few representative radical-radical reactions whose rates have been determined in most cases by measuring the second-order decay rate of the optical absorption ( $2k/\epsilon$ ). The molar absorptivity,  $\epsilon$ , used to calculate the rate constant  $k$  is given in the *Comments* column.

Products are listed only for those reactions for which evidence has been obtained on the nature of the products. The error limits given are those reported in the original papers; lacking such a statement the uncertainty in the rate constant should be assumed to be  $\pm 25\%$ . Many of the second order rate constants are derived from pseudo-first order rates for the decay of the transient radical or the substrate or the build-up of a product. Rate constants for certain reactions have been used as reference values to determine values for other reactions, which are labeled "rel."; in those cases the reference values are listed in the *Comments* column. The label "uncor." is used for observed values of  $k$  measured in systems in which more than one process is occurring.

Temperature is assumed to be ambient unless otherwise specified. The *Method* column indicates the method for generation of the transient and in the *Comments* column the method of measurement is indicated along with the source of the transient and other information such as activation energy,  $pK$ , etc. When the symbol R is used in a *Comment*, such as in  $G(R)$  or  $k(R + R)$ , it refers to the radical of the same entry. *References* are indicated by the serial number from the Radiation Chemistry Data Center bibliographic data base, from which the reference list was generated. The first two digits of the serial number represent the year in which the work was published.

*Indexes* have been produced from tables 1-3 for reactions of (1) the radicals in order of their number and for (2) the radicals and substrates by molecular formula. The indexes follow the tables.

## Acknowledgments

The assistance of Dr. W. Phillip Helman in providing programs for indexing and text processing is gratefully acknowledged. We also wish to thank Dr. Norman Lichtin and Dr. Guillermo Ferraudi for their useful comments on tables 3 and 1, respectively.

The Radiation Laboratory is operated under Contract DE-AC02-76ER00038 with the Department of Energy. The Radiation Chemistry Data Center is supported jointly by the Office of Standard Reference Data of the National Bureau of Standards and of the Office of Basic Energy Sciences and the Office of Health and Environmental Research of the Department of Energy. This is Radiation Laboratory Document No. NDRL-2196.



## List of Abbreviations and Symbols

<i>A</i>	frequency factor	<i>G</i>	radiation yield (molecules per 100 eV)
Ac	acetyl	gly	glycine or glycinate
ala	alanine	$\Delta H^\ddagger$	activation enthalpy
abs.	absorption	<i>I</i>	ionic strength
abstr.	abstraction	irradn.	irradiation
addn.	addition	<i>K</i>	equilibrium constant
alk.	alkaline	<i>k</i>	rate constant
bpy	2,2'-bipyridine	<i>k<sub>f</sub></i>	specific rate of the forward reaction
BuOH	butanol	<i>k<sub>r</sub></i>	specific rate of the reverse reaction
calcd.	calculated	<i>L</i>	ligand
chem.	chemical	Me	methyl
c.k.	competition kinetics	meas.	measured
concn.	concentration	MeOH	methanol
condy.	conductivity	NaLS	sodium dodecylsulfate
contg.	containing	obs.	observed
cor.	corrected	o.d.	optical density
CTAB	hexadecyltrimethylammonium bromide	p.b.k.	product buildup kinetics
detd.	determined	phot.	photolysis
dien	diethylenetriamine	$pK_a$	negative logarithm of the acid dissociation constant, e.g., where $AH + H_2O \rightleftharpoons A^- + H_3O^+$
d.k.	decay kinetics (decay of radical absorption and bleaching of substrate absorption)	PNAP	<i>p</i> -nitroacetophenone
<i>e</i> -r.	electron radiolysis	PNBPA	<i>p</i> -nitrobenzoato(pentaammine)cobalt(III) ion
$\epsilon$	extinction coefficient (molar absorptivity)	p.r.	pulse radiolysis
$E_a$	activation energy	PrOH	propanol
en	ethylenediamine	R	radical
esr	electron spin resonance	redn.	reduction
estd.	estimated	rel.	relative
EtOH	ethanol	RNO	<i>N,N</i> -dimethyl- <i>p</i> -nitrosoaniline
Et <sub>2</sub> O	ethyl ether	$\Delta S^\ddagger$	activation entropy
f.p.	flash photolysis	satd.	saturated
formn.	formation	soln.	solution
$\gamma$ -r.	gamma radiolysis		

## List of Aliphatic Radicals

Reactions of the following radicals are included in these tables; comments within the tables identify the compounds from which the radicals were derived. Only one acid-base form is listed below but different forms may be used in the tables, depending on pH. The entry numbers for the reactions in the tables include the three-digit numbers listed below following the decimal point. Thus, the first reaction in the tables is numbered 1.016 and involves the radical from *tert*-

butanol,  $\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$ , which is 016 in the following list. The numbers have been used to create an index to reactions of the various radicals; the index follows the tables. Radicals herein are restricted to carbon radicals and have been ordered from the most *hydrogen* substitution on carbon to the most *oxygen* substitution. That is, radicals  $\cdot\text{CXYZ}$  are ordered as X, Y, and Z progressively represent H, C, halogens, N, and O.

001	Methyl	$\cdot\text{CH}_3$
002	Ethyl	$\cdot\text{CH}_2\text{CH}_3$
002a	1-Methylethyl	$(\text{CH}_3)_2\dot{\text{C}}\text{H}$
002b	1,1-Dimethylethyl	$(\text{CH}_3)_3\dot{\text{C}}$
003	Radicals from pentane	$\cdot\text{C}_5\text{H}_{11}$
004	Cyclopentyl	$c\text{-}\dot{\text{C}}_5\text{H}_9$
005	2-Chloroethyl	$\cdot\text{CH}_2\text{CH}_2\text{Cl}$
006	2-Carboxy-2,2-dimethylethyl	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{CO}_2\text{H}$
007	Radicals from oleate + OH	R
008	Radicals from linoleate + OH	R
009	Radicals from linolenate + OH	R
010	Radicals from arachidonate + OH	R
011	2-Aminoethyl	$\cdot\text{CH}_2\text{CH}_2\text{NH}_2$
012	2-Amino-2,2-dimethylethyl	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{NH}_2$
013	2-Amino-2-carboxy-2-methylethyl	$\cdot\text{CH}_2\text{C}(\text{CH}_3)(\text{NH}_2)\text{CO}_2\text{H}$
014	2-Hydroxyethyl	$\cdot\text{CH}_2\text{CH}_2\text{OH}$
015	2-Hydroxypropyl	$\cdot\text{CH}_2\text{CHOHCH}_3$
016	2-Hydroxy-2,2-dimethylethyl	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$
016a	Radicals from cyclopentene + OH	R
017	2-Carboxy-2-hydroxy-2-methylethyl	$\cdot\text{CH}_2\text{C}(\text{CH}_3)(\text{OH})\text{CO}_2\text{H}$
018	Radicals from allyl alcohol + OH	R
019	Radicals from crotyl alcohol + OH	R
020	Formylmethyl	$\cdot\text{CH}_2\text{CHO}$
021	1-Formylethyl	$\text{CH}_3\dot{\text{C}}\text{HCHO}$
022	2-Hydroxyethenyl	$\cdot\text{CH}=\text{CHOH}$
023	2-Oxopropyl	$\cdot\text{CH}_2\text{COCH}_3$
024	1-Acetyethyl	$\text{CH}_3\text{CO}\dot{\text{C}}\text{HCH}_3$
025	2-Oxocyclohexyl	$-(\text{CH}_2)_4\dot{\text{C}}\text{HCO}-$
026	1-Formyl-2-hydroxyethyl	$\text{HOCH}_2\dot{\text{C}}\text{HCHO}$
027	4-Hydroxy-2-oxobutyl	$\text{HOCH}_2\text{CH}_2\text{CO}\dot{\text{C}}\text{H}_2$
028	1-Formyl-4-hydroxybutyl	$\text{HOCH}_2(\text{CH}_2)_2\dot{\text{C}}\text{HCHO}$
029	1-Formyl-5-hydroxypentyl	$\text{HOCH}_2(\text{CH}_2)_3\dot{\text{C}}\text{HCHO}$
030	Radicals from <i>meso</i> -erythritol	R
031	Radicals from xylitol	R
032	2-Oxo-3,4,5-trihydroxycyclohexyl	$-(\text{CHOH})_3\text{CH}_2\dot{\text{C}}\text{HCO}-$
033	Radicals from sorbitol	R
034	Carbamoylmethyl	$\cdot\text{CH}_2\text{CONH}_2$
035	Carboxymethyl	$\cdot\text{CH}_2\text{CO}_2\text{H}$
036	Methoxycarbonylmethyl	$\cdot\text{CH}_2\text{CO}_2\text{CH}_3$
037	1-Carboxyethyl	$\text{CH}_3\dot{\text{C}}\text{HCO}_2\text{H}$
038	1-Carboxypropyl	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HCO}_2\text{H}$
039	1-Carboxy-2-hydroxyethyl	$\text{HOCH}_2\dot{\text{C}}\text{HCO}_2\text{H}$
040	1-Carboxy-2-hydroxypropyl	$\text{CH}_3\text{CHOH}\dot{\text{C}}\text{HCO}_2\text{H}$
040a	1,2-Dicarboxyethyl	$\text{HO}_2\dot{\text{C}}\text{HCH}_2\text{CO}_2\text{H}$
041	1,2-Dicarboxy-2-hydroxyethyl	$\text{HO}_2\text{CCHOH}\dot{\text{C}}\text{HCO}_2\text{H}$

042	Dicarboxymethyl	$\cdot\text{CH}(\text{CO}_2\text{H})_2$
043	Chloromethyl	$\cdot\text{CH}_2\text{Cl}$
044	Difluoromethyl	$\cdot\text{CHF}_2$
045	Dichloromethyl	$\cdot\text{CHCl}_2$
046	Dichlorofluoromethyl	$\cdot\text{CFCl}_2$
047	Trifluoromethyl	$\cdot\text{CF}_3$
048	Trichloromethyl	$\cdot\text{CCl}_3$
049	1-Aminoethyl	$\text{CH}_3\dot{\text{C}}\text{H}\text{NH}_2$
050	1-Amino-1-methylethyl	$(\text{CH}_3)_2\dot{\text{C}}\text{NH}_2$
051	1-( <i>N,N</i> -Diethylamino)ethyl	$\text{CH}_3\dot{\text{C}}\text{H}\text{N}(\text{C}_2\text{H}_5)_2$
052	Triethylammonioethyl	$\text{CH}_3\dot{\text{C}}\text{H}\text{N}^+(\text{C}_2\text{H}_5)_3$
053	( <i>N</i> -Acetylamino)methyl	$\text{CH}_3\text{CONH}\dot{\text{C}}\text{H}_2$
054	( <i>N</i> -Formyl- <i>N</i> -methylamino)methyl	$\text{HCON}(\text{CH}_3)\dot{\text{C}}\text{H}_2$
055	( <i>N</i> -Acetyl- <i>N</i> -methylamino)methyl	$\text{CH}_3\text{CON}(\text{CH}_3)\dot{\text{C}}\text{H}_2$
056	Amino(carboxy)methyl	$\text{NH}_2\dot{\text{C}}\text{HCO}_2\text{H}$
057	Amino(carbamoyl)methyl	$\text{NH}_2\dot{\text{C}}\text{HCONH}_2$
058	Radicals from glycylglycinamide + OH	R
059	Carboxy( <i>N</i> -methylamino)methyl	$\text{CH}_3\text{NH}\dot{\text{C}}\text{HCO}_2\text{H}$
060	<i>N</i> -Acetylamino(carboxy)methyl	$\text{CH}_3\text{CONH}\dot{\text{C}}\text{HCO}_2\text{H}$
061	<i>N</i> -Acetyl- <i>N</i> -methylamino(carboxy)methyl	$\text{CH}_3\text{CON}(\text{CH}_3)\dot{\text{C}}\text{HCO}_2\text{H}$
062	Radical from glycine anhydride + OH	R
063	Radical from alanine anhydride + OH	R
064	Radical from sarcosine anhydride + OH	R
065	Radicals from diglycine + OH	R
066	Radicals from glycylsarcosine + OH	R
067	Radicals from acetyldiglycine + OH	R
068	Radicals from triglycine + OH	R
069	Radicals from acetyltriglycine + OH	R
070	Radicals from acetyltrialanine + OH	R
071	Radicals from acetyltrisarcosine + OH	R
072	Radicals from acetylserineamide + OH	R
073	Radicals from acetylaspargine + OH	R
074	Radical from 2-pyrrolidone-5-carboxylic acid	R
075	Hydroxymethyl	$\cdot\text{CH}_2\text{OH}$
076	1-Hydroxyethyl	$\text{CH}_3\dot{\text{C}}\text{HOH}$
077	1-Hydroxypropyl	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH}$
078	1-Hydroxy-1-methylethyl	$(\text{CH}_3)_2\dot{\text{C}}\text{OH}$
079	1-Hydroxybutyl	$\text{CH}_3(\text{CH}_2)_2\dot{\text{C}}\text{HOH}$
080	1-Hydroxy-1-methylpropyl	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{OHCH}_3$
081	1-Hydroxy-2-methylpropyl	$(\text{CH}_3)_2\text{CH}\dot{\text{C}}\text{HOH}$
082	1-Hydroxypentyl	$\text{CH}_3(\text{CH}_2)_3\dot{\text{C}}\text{HOH}$
083	(Cyclohexyl)hydroxymethyl	$c\text{-C}_6\text{H}_{11}\dot{\text{C}}\text{HOH}$
084	1-Hydroxycyclobutyl	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_2-$
085	1-Hydroxycyclopentyl	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_3-$
086	1-Hydroxycyclohexyl	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_4-$
087	1-Hydroxycycloheptyl	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_5-$
088	1-Hydroxycyclooctyl	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_6-$
089	2,2,2-Trifluoro-1-hydroxyethyl	$\cdot\text{CHOHCF}_3$
090	Acetoxymethyl	$\text{CH}_3\text{CO}_2\dot{\text{C}}\text{H}_2$
091	1-Sulfatoethyl	$\text{CH}_3\dot{\text{C}}\text{HOSO}_3^-$
092	Radicals from sodium dodecylsulfate	R
092a	Methoxymethyl	$\cdot\text{CH}_2\text{OCH}_3$
093	1-Ethoxyethyl	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5$
093a	1,1-Dimethylethoxymethyl	$\cdot\text{CH}_2\text{OC}(\text{CH}_3)_3$
094	2-Tetrahydrofuryl	$-\text{O}\dot{\text{C}}\text{H}(\text{CH}_2)_3-$
094a	(Methoxymethoxy)methyl	$\cdot\text{CH}_2\text{OCH}_2\text{OCH}_3$
095	1,4-Dihydroxybutyl	$\cdot\text{CHOHCH}_2\text{CH}_2\text{CH}_2\text{OH}$
096	1,3-Dihydroxy-2,2-di(hydroxymethyl)propyl	$\cdot\text{CHOHC}(\text{CH}_2\text{OH})_3$



097	2-Carboxy-1-hydroxy-1-methylethyl	$\text{CH}_3\dot{\text{C}}\text{OHCH}_2\text{CO}_2\text{H}$
098	2-Amino-1-hydroxyethyl	$\text{NH}_2\text{CH}_2\dot{\text{C}}\text{HOH}$
099	2-Amino-1-phosphoryloxyethyl	$\text{NH}_2\text{CH}_2\dot{\text{C}}\text{HOPO}_2\text{OH}$
100	2-Amino-2-carboxy-1-hydroxyethyl	$\cdot\text{CHOHCH}(\text{NH}_2)\text{CO}_2\text{H}$
101	1,2-Dihydroxyethyl	$\cdot\text{CHOHCH}_2\text{OH}$
102	1,2-Dihydroxypropyl	$\cdot\text{CHOHCHOHCH}_3$
103	1,2-Dihydroxy-1-methylpropyl	$\text{CH}_3\dot{\text{C}}\text{OHCHOHCH}_3$
104	2,5-Dioxacyclohexyl	$-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2-$
105	Radicals from polyethyleneglycol	$-\text{O}\dot{\text{C}}\text{HCH}_2(\text{OCH}_2\text{CH}_2)_n-$
106	1,2,3-Trihydroxypropyl	$\cdot\text{CHOHCHOHCH}_2\text{OH}$ (+ $\text{CH}_2\text{OH}\dot{\text{C}}\text{OHCH}_2\text{OH}$ )
107	1-Hydroxy-1-methyl-2-oxopropyl	$\text{CH}_3\text{CO}\dot{\text{C}}(\text{O}^-)\text{CH}_3$
108	Radicals from deoxyribose + OH	R
109	Radicals from ribose + OH	R
110	Radicals from ribose phosphate + OH	R
111	Radicals from glucose + OH	R
112	Radicals from sucrose + OH	R
113	Radical from ascorbate + OH	$\cdot\text{A}^-$
114	(Carbamoyl)hydroxymethyl	$\cdot\text{CHOHCONH}_2$
115	(Carboxy)hydroxymethyl	$\cdot\text{CHOHCO}_2\text{H}$
116	1-Carbamoyl-1-hydroxyethyl	$\text{CH}_3\dot{\text{C}}\text{OHCONH}_2$
117	1-Carboxy-1-hydroxyethyl	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2\text{H}$
118	1-Hydroxy-1-(methoxycarbonyl)ethyl	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2\text{CH}_3$
119	(Dicarboxy)hydroxymethyl	$\text{HO}_2\text{C}\dot{\text{C}}\text{OHCO}_2\text{H}$
120	1,2-Dicarboxy-1-hydroxyethyl	$\text{HO}_2\text{C}\dot{\text{C}}\text{OHCH}_2\text{CO}_2\text{H}$
121	1,2-Dicarboxy-1,2-dihydroxyethyl	$\text{HO}_2\text{C}\dot{\text{C}}\text{OHCHOHCO}_2\text{H}$
122	1,3-Dicarboxy-1-hydroxypropyl	$\text{HO}_2\text{C}\dot{\text{C}}\text{OH}(\text{CH}_2)_2\text{CO}_2\text{H}$
123	Dihydroxymethyl	$\cdot\text{CH}(\text{OH})_2$
124	Dimethoxymethyl	$\cdot\text{CH}(\text{OCH}_3)_2$
125	2,5-Dioxacyclopentyl	$-\text{O}\dot{\text{C}}\text{HO}(\text{CH}_2)_2-$
126	2,2,2-Trichloro-1,1-dihydroxyethyl	$\text{CCl}_3\dot{\text{C}}(\text{OH})_2$
127	Carbamoyl	$\cdot\text{CONH}_2$
128	Electron adduct of acrylamide	$[\text{CH}_2\text{CHCONH}_2]\cdot^-$
129	Electron adduct of crotonamide	$[\text{CH}_3\text{CHCHCONH}_2]\cdot^-$
130	Electron adduct of methacrylamide	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CONH}_2]\cdot^-$
131	Electron adduct of $\beta,\beta$ -dimethylacrylamide	$[(\text{CH}_3)_2\text{CCHCONH}_2]\cdot^-$
132	Electron adduct of <i>N,N</i> -dimethylacrylamide	$[\text{CH}_2\text{CHCON}(\text{CH}_3)_2]\cdot^-$
133	Electron adduct of Diamide	R
134	Electron adduct of acrylate	$[\text{CH}_2\text{CHCO}_2\text{H}]\cdot^-$
135	Electron adduct of crotonate	$[\text{CH}_3\text{CHCHCO}_2\text{H}]\cdot^-$
136	Electron adduct of methacrylate	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CO}_2\text{H}]\cdot^-$
137	Electron adduct of $\beta,\beta$ -dimethylacrylate	$[(\text{CH}_3)_2\text{CCHCO}_2\text{H}]\cdot^-$
138	Electron adduct of methyl methacrylate	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CO}_2\text{CH}_3]\cdot^-$
139	Electron adduct of sorbate	$[\text{CH}_3\text{CHCHCHCHCO}_2\text{H}]\cdot^-$
141	Electron adduct of dimethyl fumarate	R
142	Electron adduct of glycine anhydride	R
143	Electron adduct of alanine anhydride	R
144	Electron adduct of sarcosine anhydride	R
146	Electron adduct of acetylglycylglycinamide	R
147	Carboxyl	$\cdot\text{CO}_2^-/\cdot\text{CO}_2\text{H}$
148	Electron adduct of methyl acetate	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{OCH}_3$

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
<b>Ag</b>						
1.	<b>Silver(I) ion</b>					
1.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Ag}^+ \rightarrow \text{Ag}^0$		nat	p.r.	No electron transfer to give $\text{Ag}^0$ in 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	78A410
1.075	$\cdot\text{CH}_2\text{OH} + \text{Ag}^+ \rightarrow \text{Ag}^0$		nat	p.r.	No electron transfer to give $\text{Ag}^0$ in 0.1 mol L <sup>-1</sup> MeOH.	78A410
1.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Ag}^+ \rightarrow \text{Ag}^0$		nat	p.r.	No electron transfer to give $\text{Ag}^0$ in 0.1 mol L <sup>-1</sup> 2-PrOH.	78A410
1.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{Ag}^+ \rightarrow \text{Ag}^0$ + $\text{H}^+ + \text{CH}_2=\text{CHCONH}_2$	$(1.1 \pm 0.4) \times 10^8$	~6	p.r.	P.b.k. at 313 nm in soln. contg. acrylamide.	70G052
1.147	$\cdot\text{CO}_2^- + \text{Ag}^+ \rightarrow \text{AgCO}_2$ (+ $\text{Ag}^+$ ) $\rightarrow \text{Ag}_2\text{CO}_2^+$		nat	p.r.	$\text{Ag}^+$ was reduced in 10 <sup>-2</sup> mol L <sup>-1</sup> formate soln. contg. $\text{CO}_2$ by a complex mechanism.	78A410
2.	<b>Silver(I) ion, complex with Ag(0)</b>					
2.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Ag}_2^+ \rightarrow$ $2 \text{Ag}^+ + \text{OH}^- + \text{CH}_2=\text{C}(\text{CH}_3)_2$	$2.0 \times 10^9$	nat	p.r.	Calcd. from best fit for increase in condy. as function of time in 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH-5 x 10 <sup>-4</sup> mol L <sup>-1</sup> $\text{AgClO}_4$ soln.	78A410
2.075	$\cdot\text{CH}_2\text{OH} + \text{Ag}_2^+ \rightarrow \text{Ag}_2\text{CH}_2\text{OH}^+$ $\rightarrow \text{CH}_2\text{O} + \text{H}^+ + \text{Ag}_2$	$5.0 \times 10^9$	nat	p.r.	Calcd. from increase in condy. as function of time in 0.1 mol L <sup>-1</sup> MeOH-5 x 10 <sup>-4</sup> mol L <sup>-1</sup> $\text{AgClO}_4$ soln. assuming values for $k(\text{R} + \text{R})$ , $[\text{Ag}_2^+]$ , $[\text{R}]$ , and $[\text{H}_2\text{O}_2]$ .	78A410
2.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Ag}_2^+ \rightarrow$ $\text{CH}_3\text{CHO} + \text{H}^+ + \text{Ag}_2$	$1.0 \times 10^9$		p.r.	Calcd. from increase in condy. as function of time in EtOH- $\text{AgClO}_4$ soln. assuming values for $k(\text{R} + \text{R})$ , $[\text{Ag}_2^+]$ , $[\text{R}]$ , and $[\text{H}_2\text{O}_2]$ .	78A410
2.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Ag}_2^+ \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{Ag}_2$	$2.5 \times 10^9$		p.r.	Calcd. from increase in condy. as function of time in 0.1 mol L <sup>-1</sup> 2-PrOH-5 x 10 <sup>-4</sup> mol L <sup>-1</sup> $\text{AgClO}_4$ soln. assuming values for $k(\text{R} + \text{R})$ , $[\text{Ag}_2^+]$ , $[\text{R}]$ , and $[\text{H}_2\text{O}_2]$ .	78A410
<b>Br</b>						
3.	<b>Bromate ion</b>					
3.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{BrO}_3^-$ $\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{BrO}_3^-$	$< 5 \times 10^6$ $(3.0 \pm 0.5) \times 10^7$	6 11.8	p.r.	D.k.; radical from ethanol.	72A018
<b>Cd</b>						
4.	<b>Cadmium(I) ions</b>					
4.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Cd}^+ \rightarrow$ $\text{Cd}^{2+} + (\text{CH}_3)_2\text{C}=\text{CH}_2 + \text{OH}^-$	$\sim 1 \times 10^9$	nat	p.r.	Calcd. from opt. and condy. studies in <i>tert</i> -BuOH- $\text{Cd}^{2+}$ soln. assuming values for competing reactions.	75A064
	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Cd}_2^{2+} \rightarrow \text{OH}^- +$ $\text{Cd}^+ + (\text{CH}_3)_2\text{C}=\text{CH}_2 + \text{Cd}^{2+}$	$\sim 1 \times 10^9$	nat	p.r.	See above.	75A064
4.075	$\cdot\text{CH}_2\text{OH} + \text{Cd}^+ \rightarrow \text{CdCH}_2\text{OH}^+$	$2 \times 10^8$	nat	p.r.	D.k. at 300 nm ( $\text{Cd}^+$ ) in soln. contg. MeOH and $\text{Cd}^{2+}$ , as well as condy. and p.b.k. at 240 nm ( $\text{Cd}_2^{2+}$ ); assumed $k(\text{Cd}^+ + \text{H}_2\text{O}_2) = 1.5 \times 10^9$ , $k(\text{R} + \text{R}) = 1.2 \times 10^9$ .	75A064

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
4.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Cd}^+ \rightarrow \text{CdCHOHCH}_3^+$	$(1.3 \pm 0.6) \times 10^9$	nat	p.r.	D.k. at 300 nm ( $\text{Cd}^+$ ) in soln. contg. EtOH and $\text{Cd}^{2+}$ , as well as condy. and p.b.k. at 240 nm ( $\text{Cd}_2^{2+}$ ); assumed $k(\text{Cd}^+ + \text{H}_2\text{O}_2) = 1.5 \times 10^9$ , $k(\text{R} + \text{R}) = 1.15 \times 10^9$ and $2k(\text{Cd}^+ + \text{Cd}^+) = 3.0 \times 10^9$ .	75A064
4.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Cd}^+ \rightarrow \text{CdC}(\text{CH}_3)_2\text{OH}^+$	$(2.4 \pm 1.2) \times 10^9$	nat	p.r.	D.k. at 300 nm ( $\text{Cd}^+$ ) in soln. contg. 2-PrOH and $\text{Cd}^{2+}$ , as well as condy. and p.b.k. at 240 nm ( $\text{Cd}_2^{2+}$ ); assumed $k(\text{Cd}^+ + \text{H}_2\text{O}_2) = 1.5 \times 10^9$ , $k(\text{R} + \text{R}) = 7 \times 10^8$ and $2k(\text{Cd}^+ + \text{Cd}^+) = 3.0 \times 10^9$ .	75A064
4.105	$-\text{O}\dot{\text{C}}\text{HCH}_2(\text{OCH}_2\text{CH}_2\text{O})_n^- + \text{Cd}^+$	$< 2 \times 10^8$	nat	p.r.	D.k. of $\text{Cd}^+$ in the presence of polyethylene glycol entirely by $\text{Cd}^+ + \text{Cd}^+$ ( $2k = 5 \times 10^9$ ); $2k(\text{R} + \text{R}) = 2.2 \times 10^7$ .	75A153
<b>5. Cadmium(II) ion</b>						
5.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Cd}^{2+}$	$< 10^6$	nat	p.r.	Est. from lack of increase in $\text{Cd}^+$ in 0.1 mol L <sup>-1</sup> $\text{Cd}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	75A027
5.075	$\cdot\text{CH}_2\text{OH} + \text{Cd}^{2+}$	$< 2.5 \times 10^5$	nat	p.r.	No reaction.	75A153
		$< 10^2$	nat	p.r.	Est. from lack of increase in $\text{Cd}^+$ in 0.1 mol L <sup>-1</sup> $\text{Cd}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> MeOH.	75A027
5.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Cd}^{2+}$	$< 2.5 \times 10^5$	nat	p.r.	No reaction.	75A153
5.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Cd}^{2+}$	$< 10^6$		p.r.	Est. from lack of increase in $\text{Cd}^+$ in 0.1 mol L <sup>-1</sup> $\text{Cd}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> 2-PrOH.	75A027
5.101	$\cdot\text{CHOHCH}_2\text{OH} + \text{Cd}^{2+}$	$< 2.5 \times 10^5$	nat	p.r.	No reaction.	75A153
5.105	$-\text{O}\dot{\text{C}}\text{HCH}_2(\text{OCH}_2\text{CH}_2\text{O})_n^- + \text{Cd}^{2+}$	$< 2.5 \times 10^5$	nat	p.r.	No reaction obs. in polyethylene glycol soln.	75A153
5.106	$\cdot\text{CHOHCHOHCH}_2\text{OH} + \text{Cd}^{2+}$	$< 2.5 \times 10^5$	nat	p.r.	No reaction.	75A153
5.147	$\cdot\text{CO}_2^- + \text{Cd}^{2+} \rightarrow \text{Cd}^+ + \text{CO}_2$	$\sim 10^5$		p.r.	Est. from increase in $\text{Cd}^+$ in 0.1 mol L <sup>-1</sup> $\text{Cd}^{2+}$ soln. contg. 0.1 mol L <sup>-1</sup> $\text{HCO}_2^-$ , $\text{CO}_2$ -satd.	75A027
		$(5.1 \pm 0.3) \times 10^6$	nat	p.r.	No details given.	75A153
<b>Co</b>						
<b>6. Cobalt(II) ions</b>						
6.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Co}^{2+} \rightarrow \text{Co}^+$	$< 10^6$	nat	p.r.	Est. from lack of increase in $\text{Co}^+$ in 0.1 mol L <sup>-1</sup> $\text{Co}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	75A027
6.075	$\cdot\text{CH}_2\text{OH} + \text{Co}^{2+} \rightarrow \text{Co}^+$	$< 10^2$	nat	p.r.	Est. from lack of increase in $\text{Co}^+$ in 0.1 mol L <sup>-1</sup> $\text{Co}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> MeOH.	75A027
6.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}^{2+} \rightarrow \text{Co}^+$	$< 10^6$	nat	p.r.	Est. from lack of increase in $\text{Co}^+$ in 0.1 mol L <sup>-1</sup> $\text{Co}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> 2-PrOH.	75A027
6.147	$\cdot\text{CO}_2^- + \text{Co}^{2+} \rightarrow \text{CO}_2 + \text{Co}^+$	$10^2 < k < 10^5$	nat	p.r.	Est. from lack of increase in $\text{Co}^+$ in 0.1 mol L <sup>-1</sup> $\text{Co}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> formate, as well as $\gamma$ -r. expts. (73G039).	75A027
<b>7. Diaqua(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(II) ion</b>						
7.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CoL}(\text{H}_2\text{O})_2^{2+} \rightarrow \text{Co(I)}$	$5.5 \times 10^9$	1.25, 6.5	p.r.	P.b.k. in 2 mol L <sup>-1</sup> 2-PrOH.	76A001
7.147	$\cdot\text{CO}_2^- + \text{CoL}(\text{H}_2\text{O})_2^{2+} \rightarrow \text{Co(I)}$	$4.7 \times 10^9$	6.5	p.r.	P.b.k. in 0.1 mol L <sup>-1</sup> formate.	76A001



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
8.	<b>Diaqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,14-diene)cobalt(II) ion</b>					
8.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CoL}(\text{H}_2\text{O})_2^{2+} \rightarrow \text{Co(I)}$	$\leq 10^7$	6.5	p.r.	No reaction in 2 mol L <sup>-1</sup> 2-PrOH.	76A001
8.147	$\cdot\text{CO}_2^- + \text{CoL}(\text{H}_2\text{O})_2^{2+} \rightarrow \text{Co(I)}$	$\leq 10^7$	6.5	p.r.	No reaction in 0.1 mol L <sup>-1</sup> formate.	76A001
9.	<b>Diaqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(II) ion</b>					
9.001	$\cdot\text{CH}_3 + \text{CoL}(\text{OH}_2)_2^{2+} \rightarrow$ $\text{CH}_3\text{Co}^{\text{III}}\text{L}(\text{OH}_2)_2^{2+}$	$\sim 7 \times 10^8$  (1 to 2) $\times 10^8$	1 9.5	f.p. p.r.	Radical from $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_3^{2+}$ in $\text{HClO}_4$ soln. D.k. at 330 nm ( $\text{Co}^{\text{II}}$ ) as well as p.b.k. at 280 nm in Ar-satd. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH soln. contg. $5 \times 10^{-3}$ mol L <sup>-1</sup> $\text{CH}_3\text{I}$ .	74F644 76A203
9.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{CoL}(\text{OH}_2)_2^{2+}$	$< 1 \times 10^7$		p.r.	No reaction obs.	78A200
9.020	$\cdot\text{CH}_2\text{CHO} + \text{CoL}(\text{OH}_2)_2^{2+} \rightarrow$ $\text{Co}^{\text{III}}\text{LCH}_2\text{CHO}$	$8 \times 10^7$	1-3, 7-10	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 1 mol L <sup>-1</sup> ethylene glycol.	78A200
9.035	$\cdot\text{CH}_2\text{CO}_2^- + \text{CoL}(\text{OH}_2)_2^{2+}$	$< 1 \times 10^7$		p.r.	No reaction obs.	78A200
9.075	$\cdot\text{CH}_2\text{OH} + \text{CoL}(\text{OH}_2)_2^{2+} \rightarrow$ $\text{Co}^{\text{III}}\text{LCH}_2\text{OH}$	$(7 \pm 1) \times 10^7$	1-6	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	78A200
9.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{CoL}(\text{OH}_2)_2^{2+} \rightarrow$ $\text{Co}^{\text{III}}\text{LCHOHCH}_3$	$(3.0 \pm 0.4) \times 10^7$	1-6	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 1 mol L <sup>-1</sup> EtOH.	78A200
9.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CoL}(\text{OH}_2)_2^{2+}$	$\leq 10^7$	6.5	p.r.	No reaction obs. in soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	76A001
		$< 10^7$		p.r.	No reaction obs.	78A200
9.089	$\cdot\text{CHOHCF}_3 + \text{CoL}(\text{OH}_2)_2^{2+} \rightarrow$ $\text{Co}^{\text{III}}\text{LCHOHCF}_3$	$(1 \pm 0.5) \times 10^7$	6	p.r.	P.b.k.	78A200
9.098	$\text{NH}_3^+\text{CH}_2\dot{\text{C}}\text{HOH} + \text{CoL}(\text{OH}_2)_2^{2+}$	$4.0 \times 10^7$	7.0	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-aminoethanol; rate pH dependent.	78A200
9.101	$\cdot\text{CHOHCH}_2\text{OH} + \text{CoL}(\text{OH}_2)_2^{2+} \rightarrow$ $\text{Co}^{\text{III}}\text{LCHOHCH}_2\text{OH}$	$\sim 1 \times 10^7$	3-7	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 1 mol L <sup>-1</sup> ethylene glycol.	78A200
9.147	$\cdot\text{CO}_2^- + \text{CoL}(\text{OH}_2)_2^{2+}$	$\leq 10^7$	6.5	p.r.	No reaction obs. in soln. contg. 0.1 mol L <sup>-1</sup> formate.	76A001
10.	<b>Cobal(II)amin (Vitamin B12r)</b>					
10.001	$\cdot\text{CH}_3 + \text{B12r} \rightarrow \text{methylcobalamin}$	$\sim 1.5 \times 10^9$		f.p.	D.k. following irradiation-induced $\text{Co}-\text{CH}_3$ homolysis of methylcobalamin; back reaction.	77F005
10.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{B12r} \rightarrow \text{B12a}$ (B12a See 54.)	$2.4 \times 10^8$	5	p.r.	P.b.k. at 350-60 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	75A169
10.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{B12r} \rightarrow$ $\text{B12s} (\text{Co}^{\text{I}}) + (\text{CH}_3)_2\text{CO} + \text{H}^+$	$(4 \pm 2) \times 10^9$		f.p.	D.k. at 474 nm in 90% 2-PrOH soln. of adenosyl cobalamin after homolysis of the adenosyl group.	79A158
10.101	$\cdot\text{CHOHCH}_2\text{OH} + \text{B12r} \rightarrow \text{B12a}$ (B12a See 54.)	$2.4 \times 10^8$		p.r.	P.b.k. at 525 nm as well as d.k. at 400 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.05 mol L <sup>-1</sup> ethylene glycol.	75A169
10.147	$\cdot\text{CO}_2^- + \text{B12r} \rightarrow \text{CO}_2 + \text{B12s} (\text{Co}^{\text{I}})$	$8.2 \times 10^8$	9.2	p.r.	D.k. at 311 and 478 nm as well as p.b.k. at 386 and 280 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	74A105
10a.	<b>3,10,17,24-Tetrasulfophthalocyaninecobalt(II) ion dimer</b>					
10a.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + (\text{Co}^{\text{II}}\text{pts})_2^{8-} \rightarrow$ addn.	$2.2 \times 10^8$	9	p.r.	P.b.k. at 440 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1-0.2 mol L <sup>-1</sup> <i>tert</i> -BuOH; mechanism suggested to involve addn. followed by splitting to give $(\text{RCo}^{\text{III}}\text{pts})_2^{5-}$ .	80A146
10a.075	$\cdot\text{CH}_2\text{OH} + (\text{Co}^{\text{II}}\text{pts})_2^{8-} \rightarrow$ electron transfer	$2.2 \times 10^8$	9	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1-0.2 mol L <sup>-1</sup> MeOH; mechanism suggested to involve ligand-radical formation, followed by metal oxid. and dimer splitting to give $(\text{Co}^{\text{I}}\text{pts})_2^{7-}$ .	80A146
10a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + (\text{Co}^{\text{II}}\text{pts})_2^{8-} \rightarrow$ electron transfer	$1.5 \times 10^9$	9	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 2-PrOH; see above.	80A146

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
11.	<b>Nitrilotriacetatocobalt(II) ion</b>					
11.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Co}^{\text{II}}\text{NTA} \rightarrow$ $[\text{NTACoCH}_2\text{C}(\text{CH}_3)_2\text{OH}]^-$	$(1.1 \pm 0.2) \times 10^7$	7	p.r.	P.b.k. at 330 nm in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> <i>tert</i> -BuOH.	79A255
11.147	$\cdot\text{CO}_2^- + \text{Co}^{\text{II}}\text{NTA} \rightarrow [\text{NTACo}(\text{CO}_2)]^{2-}$	$(7.3 \pm 0.7) \times 10^7$	7	p.r.	P.b.k. at 270 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	79A255
	<b>Cobalt(III) ions</b>					
12.	<b>Hexaamminecobalt(III) ion</b>					
12.035	$\cdot\text{CH}_2\text{CO}_2^- + \text{Co}(\text{NH}_3)_6^{3+}$	$< 3.2 \times 10^6$	7.3	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> acetate.	72A018
	$\cdot\text{CH}_2\text{CO}_2\text{H} + \text{Co}(\text{NH}_3)_6^{3+}$	$< 1.0 \times 10^7$	2.5	p.r.	D.k. in Ar-satd. soln. contg. 0.01 mol L <sup>-1</sup> acetic acid.	72A018
12.042	$\cdot\text{CH}(\text{CO}_2\text{H})_2 + \text{Co}(\text{NH}_3)_6^{3+}$	$< 8.0 \times 10^6$	2.5	p.r.	D.k. in Ar-satd. soln. contg. 0.01 mol L <sup>-1</sup> malonic acid.	72A018
12.054	$\text{HCON}(\text{CH}_3)\dot{\text{C}}\text{H}_2 + \text{Co}(\text{NH}_3)_6^{3+}$	$< 1.0 \times 10^7$	6.0	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> dimethylformamide.	72A018
12.056	$\text{NH}_3^+\dot{\text{C}}\text{HCO}_2^- + \text{Co}(\text{NH}_3)_6^{3+}$	$< 1.2 \times 10^7$	5.7	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> glycine.	72A018
12.075	$\cdot\text{CH}_2\text{O}^- + \text{Co}(\text{NH}_3)_6^{3+} \rightarrow$ $\text{CH}_2\text{O} + \text{Co}(\text{NH}_3)_6^{2+}$	$(9.0 \pm 1.4) \times 10^9$	12	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> MeOH.	72A018
	$\cdot\text{CH}_2\text{OH} + \text{Co}(\text{NH}_3)_6^{3+} \rightarrow$ $\text{CH}_2\text{O} + \text{H}^+ + \text{Co}(\text{NH}_3)_6^{2+}$	$(1.4 \pm 0.2) \times 10^8$	5-6	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> MeOH.	72A018
		$(1.4 \pm 0.2) \times 10^8$	6.1	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	77A100
		$(6 \pm 0.9) \times 10^7$	4.9			
		$(4.5 \pm 0.7) \times 10^7$	4.5			
		$< 1 \times 10^7$	3.5			
12.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{Co}(\text{NH}_3)_6^{3+}$	$(8.5 \pm 1.3) \times 10^9$	12	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> EtOH; <i>e</i> -transfer.	72A018
	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{NH}_3)_6^{3+}$	$(5.2 \pm 0.8) \times 10^7$	5-6			
12.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Co}(\text{NH}_3)_6^{3+}$	$(5.0 \pm 0.8) \times 10^9$	12	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> 2-PrOH; <i>e</i> -transfer.	72A018
	$(\text{CH}_3)_2\dot{\text{C}}\text{HOH} + \text{Co}(\text{NH}_3)_6^{3+}$	$(1.3 \pm 0.2) \times 10^7$	5-6			
12.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{Co}(\text{NH}_3)_6^{3+}$	$< 5 \times 10^6$	5.5-6	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. ethyl ether.	77A100
12.104	$-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2- + \text{Co}(\text{NH}_3)_6^{3+}$	$< 2 \times 10^6$	3.5-4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. dioxane.	77A100
12.107	$\text{CH}_3\text{CO}\dot{\text{C}}(\text{O}^-)\text{CH}_3 + \text{Co}(\text{NH}_3)_6^{3+}$	$< 1.8 \times 10^5$	5.5	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> biacetyl.	72A018
	$\text{CH}_3\text{CO}\dot{\text{C}}(\text{OH})\text{CH}_3 + \text{Co}(\text{NH}_3)_6^{3+}$	$< 2.0 \times 10^5$	1.2			
12.117	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{CO}_2^- + \text{Co}(\text{NH}_3)_6^{3+}$	$2.4 \times 10^{10}$	12.0	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> lactate; <i>e</i> -transfer.	72A018
	$\text{CH}_3\dot{\text{C}}(\text{OH})\text{CO}_2^- + \text{Co}(\text{NH}_3)_6^{3+}$	$7.0 \times 10^6$	6.0			
12.127	$\cdot\text{CONH}_2 + \text{Co}(\text{NH}_3)_6^{3+}$	$< 3.6 \times 10^7$	6.2	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> formamide.	72A018
12.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_6^{3+} \rightarrow$ $\text{CO}_2 + \text{Co}(\text{NH}_3)_6^{2+}$	$4.0 \times 10^7$	4.8	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.01 mol L <sup>-1</sup> formate.	72A018
		$(1.1 \pm 0.3) \times 10^8$ (rel.)	6.9	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate rel. to $k(\cdot\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ . [26.147]	73A075
13.	<b>Pentaammineaquacobalt(III) ion</b>					
13.075	$\cdot\text{CH}_2\text{OH} + \text{Co}(\text{NH}_3)_5\text{OH}_2^{3+} \rightarrow$ Co(II)	$(1.5 \pm 0.5) \times 10^6$ (rel.)	3.5-4	$\gamma$ -r.	Calcd. from $G(\text{Co}^{2+})$ , dose rate and $k(\text{R} + \text{R}) = 2.4 \times 10^9$ .	77A100
13.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{OH}_2^{3+} \rightarrow$ Co(II)	$(1.7 \pm 0.3) \times 10^8$ (rel.)	5.2	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate; rel. to $k(\cdot\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ . [26.147]	73A075
14.	<b>Pentaamminehydroxycobalt(III) ion</b>					
14.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{OH}^{2+} \rightarrow$ Co(II)	$< (3 \pm 1.5) \times 10^7$ (rel.)	7.8	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate; rel. to $k(\cdot\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ . [26.147]	73A075

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
15.	<b>Pentaamminebromocobalt(III) ion</b>					
15.075	$\cdot\text{CH}_2\text{OH} + \text{Co}(\text{NH}_3)_5\text{Br}^{2+}$	$(9.0 \pm 1.4) \times 10^7$ $(2.5 \pm 0.4) \times 10^7$ $(2.0 \pm 0.3) \times 10^7$ $(1.8 \pm 0.3) \times 10^7$ $(1.8 \pm 0.3) \times 10^7$ $(1.8 \pm 0.3) \times 10^7$	6.1 4.9 4.5 3.5 2.0 1.0	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	77A100
15.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{NH}_3)_5\text{Br}^{2+}$	$(1.5 \pm 0.2) \times 10^8$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH.	77A100
15.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{NH}_3)_5\text{Br}^{2+}$	$(3.0 \pm 0.45) \times 10^8$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100
15.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{Co}(\text{NH}_3)_5\text{Br}^{2+}$	$(1.6 \pm 0.2) \times 10^8$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. ethyl ether.	77A100
15.104	$-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2- + \text{Co}(\text{NH}_3)_5\text{Br}^{2+}$	$\leq 2 \times 10^6$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. dioxane.	77A100
16.	<b>Pentaamminechlorocobalt(III) ion</b>					
16.075	$\cdot\text{CH}_2\text{OH} + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	$(3 \pm 1) \times 10^6$ (rel.)	3.5–4	$\gamma$ -r	Est. from effect of complex concn. on $G(\text{Co}^{2+})$ assuming $2k(\text{R} + \text{R}) = 2.4 \times 10^9$ .	77A100
16.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	$(3.0 \pm 0.4) \times 10^6$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH.	77A100
16.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	$(4.0 \pm 0.6) \times 10^7$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100
16.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	$(1.4 \pm 0.2) \times 10^7$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. ethyl ether.	77A100
16.104	$-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2- + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	$\leq 2 \times 10^6$	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. dioxane.	77A100
16.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	$(1.5 \pm 0.3) \times 10^8$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
17.	<b>Pentaamminefluorocobalt(III) ion</b>					
17.075	$\cdot\text{CH}_2\text{OH} + \text{Co}(\text{NH}_3)_5\text{F}^{2+}$	$(5.5 \pm 2) \times 10^5$ (rel.)	3.5–4	$\gamma$ -r.	Est. from effect of complex concn. on $G(\text{Co}^{2+})$ assuming $2k(\text{R} + \text{R}) = 2.4 \times 10^9$ .	77A100
18.	<b>Pentaammine(nitrito-N)cobalt(III) ion</b>					
18.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{NO}_2^{2+}$	$< (2 \pm 1) \times 10^7$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
19.	<b>Pentaammine(nitrato-O)cobalt(III) ion</b>					
19.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{NO}_3^{2+}$	$(2.1 \pm 0.3) \times 10^8$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
20.	<b>(Acetato)pentaamminecobalt(III) ion</b>					
20.001	$\cdot\text{CH}_3 + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_3^{2+} \rightarrow$ $\text{CH}_4 + \text{Co}^{2+} + 4\text{NH}_4^+ + \text{NH}_2^+ +$ $\text{CH}_3\text{CO}_2\text{H}$	$(4 \pm 1) \times 10^3$ (rel.)		phot.	Est. from intensity dependence of $\text{CH}_4$ and $\text{C}_2\text{H}_6$ formn. assuming $k(\text{R} + \text{R}) = 2 \times 10^{10}$ .	71F579
20.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_3^{2+}$	$(1.1 \pm 0.3) \times 10^8$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
21.	<b>Pentaammine(phenylacetato)cobalt(III) ion</b>					
21.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_2\text{C}_6\text{H}_5^{2+}$	$(7.0 \pm 2) \times 10^7$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
22. 22.147	<b>Pentaammine(4-nitrophenylacetato)cobalt(III) ion</b> $\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_2\text{C}_6\text{H}_4\text{NO}_2^{2+}$	$(1.2 \pm 0.1) \times 10^9$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
23. 23.147	<b>Pentaammine(benzoato)cobalt(III) ion</b> $\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_5^{2+}$	$(4.5 \pm 2) \times 10^7$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
24. 24.078	<b>Pentaammine(2-nitrobenzoato)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{NO}_2^{2+}$	$1.7 \times 10^9$	7.0	p.r.	P.b.k.	77A027
24.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{NO}_2^{2+}$	$2.0 \times 10^9$	7.0	p.r.	P.b.k.	77A027
25. 25.078	<b>Pentaammine(3-nitrobenzoato)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{NO}_2^{2+}$	$1.5 \times 10^9$	7.0	p.r.	P.b.k.	77A027
25.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{NO}_2^{2+}$	$1.5 \times 10^9$	7.0	p.r.	P.b.k.	77A027
26. 26.078	<b>Pentaammine(4-nitrobenzoato)cobalt(III) ion (PNBPA)</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{NO}_2^{2+} \rightarrow$ $(\text{CH}_3)_2\dot{\text{C}}\text{O} + \text{H}^+ +$ $\text{Co}^{\text{III}}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\dot{\text{N}}\text{O}_2^-$	$2.6 \times 10^9$ $2.6 \times 10^9$	4.5 7.0	p.r. p.r.	Condy. in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> 2-PrOH. P.b.k.	74A002 77A027
26.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{NO}_2^{2+} \rightarrow$ $\text{CO}_2 + \text{Co}^{\text{III}}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\dot{\text{N}}\text{O}_2^-$	$1.9 \times 10^9$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	72G340 77A027
27. 27.078	<b>Pentaammine(2,4-dinitrobenzoato)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2^{2+}$	$4.3 \times 10^9$	7.0	p.r.	P.b.k.	77A027
27.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2^{2+}$	$7.5 \times 10^9$	7.0	p.r.	P.b.k.	77A027
28. 28.078	<b>Pentaammine(3,5-dinitrobenzoato)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2^{2+}$	$2.9 \times 10^9$	7.0	p.r.	P.b.k.	77A027
28.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2^{2+}$	$8.1 \times 10^9$	7.0	p.r.	P.b.k.	77A027
29. 29.078	<b>Hexaamminebis(<math>\mu</math>-hydroxy)[<math>\mu</math>-(4-nitrobenzoato)]dicobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \dots\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\dot{\text{C}}\text{O} + \text{H}^+ + \dots\dot{\text{N}}\text{O}_2^-$	$(2 \pm 0.4) \times 10^9$ $(1.4 \pm 0.4) \times 10^9$	0.7– 6.0	p.r. p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH. P.b.k. at 370 nm in soln. contg. 1% 2-PrOH; $\Delta H^\ddagger = 3.1 \pm 2.0$ kcal mol <sup>-1</sup> (13 kJ mol <sup>-1</sup> ); $k$ decreases with concn. of 2-PrOH; $k = 1.25 \times 10^9$ in 99% D <sub>2</sub> O– 1% 2-PrOH.	78A108 80A066
29.101	$\cdot\text{CHOHCH}_2\text{OH} + \dots\text{NO}_2 \rightarrow \dots\dot{\text{N}}\text{O}_2$	$(8 \pm 2) \times 10^7$		p.r.	P.b.k. at 370 nm in soln. of 50% H <sub>2</sub> O–50% ethylene glycol; $\Delta H^\ddagger =$ $5.4 \pm 0.3$ kcal mol <sup>-1</sup> (21 kJ mol <sup>-1</sup> )	80A066
30. 30.147	<b>Pentaammine(4-cyanobenzoato)cobalt(III) ion</b> $\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_4\text{CN}^{2+}$	$(4.6 \pm 2) \times 10^7$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
31. 31.078	<b>Pentaammine(pyridine)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{NH}_3)_5\text{C}_5\text{H}_5\text{N}^{3+} \rightarrow$ $\text{Co}^{2+} + 5\text{NH}_3 + \text{C}_5\text{H}_5\text{N}$ $+ (\text{CH}_3)_2\dot{\text{C}}\text{O} + \text{H}^+$	$\sim 10^9$	1–7	$\gamma$ -r.	Est. from $G(\text{Co}^{2+})$ in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	79A213

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
31.147	$\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{C}_5\text{H}_5\text{N}^{3+}$	$(3.3 \pm 0.4) \times 10^8$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
32. 32.147	<b>Pentaammine(pyridinecarboxylato)cobalt(III) ion</b> $\cdot\text{CO}_2^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_5\text{H}_4\text{N}^{2+}$	$(5.1 \pm 2) \times 10^7$ (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
33. 33.078	<b>Pentaammine(pyrazinecarboxylato)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_4\text{H}_3\text{N}_2^{2+} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \dots\dot{\text{N}}\text{H}$	$(4.0 \pm 0.6) \times 10^8$	0-5	p.r.	P.b.k. in N <sub>2</sub> O or Ar-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	78A222
34. 34.078	<b>Tetraammine(pyrazinecarboxylato-O,N)cobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} +$ $\text{Co}(\text{NH}_3)_4\text{O}_2\text{CC}_4\text{H}_3\text{N}_2^{2+} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{Co}^{\text{II}}$	$(9 \pm 2) \times 10^8$	5	p.r.	P.b.k. in N <sub>2</sub> O or Ar-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	78A222
35. 35.078	<b>Hexaamminebis(μ-hydroxy)[μ-(pyrazinecarboxylato)](dicobalt(III) ion</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \dots\text{N} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \dots\dot{\text{N}}\text{H}$	$(4.2 \pm 0.4) \times 10^8$	0-6	p.r.	P.b.k. in N <sub>2</sub> O or Ar-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	78A222
36. 36.075 36.076 36.078	<b>Tris(ethylenediamine)cobalt(III) ion</b> $\cdot\text{CH}_2\text{OH} + \text{Co}(\text{en})_3^{3+}$ $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{en})_3^{3+}$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{en})_3^{3+}$	no reaction no reaction no reaction	3.5-4 3.5-4 3.5-4	p.r. p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH. D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH. D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100 77A100 77A100
37. 37.075 37.076 37.078	<b>cis-Amminechlorobis(ethylenediamine)cobalt(III) ion</b> $\cdot\text{CH}_2\text{OH} + \text{Co}(\text{NH}_3\text{Cl}(\text{en})_2)^{2+}$ $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{NH}_3\text{Cl}(\text{en})_2)^{2+}$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{NH}_3\text{Cl}(\text{en})_2)^{2+}$	$<2 \times 10^6$ $4.2 \times 10^6$ $2.2 \times 10^7$	3.5-4 3.5-4 3.5-4	p.r. p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH. D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH. D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100 77A100 77A100
37.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{Co}(\text{NH}_3\text{Cl}(\text{en})_2)^{2+}$	$4.6 \times 10^6$	3.5-4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. Et <sub>2</sub> O.	77A100
38. 38.075 38.076 38.078	<b>cis-Aquachlorobis(ethylenediamine)cobalt(III) ion</b> $\cdot\text{CH}_2\text{OH} + \text{Co}(\text{OH}_2\text{Cl}(\text{en})_2)^{2+}$ $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{OH}_2\text{Cl}(\text{en})_2)^{2+}$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{OH}_2\text{Cl}(\text{en})_2)^{2+}$	$(1.8 \pm 0.5) \times 10^6$ $2.0 \times 10^7$ $8.2 \times 10^7$	3.5-4 3.5-4 3.5-4	p.r. p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH. D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH. D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100 77A100 77A100
38.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{Co}(\text{OH}_2\text{Cl}(\text{en})_2)^{2+}$	$3.5 \times 10^7$	3.5-4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. Et <sub>2</sub> O.	77A100
39. 39.075 39.076 39.078	<b>trans-Dibromobis(ethylenediamine)cobalt(III) ion</b> $\cdot\text{CH}_2\text{OH} + \text{Co}(\text{en})_2\text{Br}_2^+$ $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Co}(\text{en})_2\text{Br}_2^+$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Co}(\text{en})_2\text{Br}_2^+$	$2.6 \times 10^8$ $5.7 \times 10^8$ $6.8 \times 10^8$	3.5-4 3.5-4 3.5-4	p.r. p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH. D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH. D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100 77A100 77A100
39.093 39.104	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{Co}(\text{en})_2\text{Br}_2^+$ $-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2- + \text{Co}(\text{en})_2\text{Br}_2^+$	$6.5 \times 10^8$ $4.4 \times 10^8$	3.5-4 3.5-4	p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. Et <sub>2</sub> O. D.k. in N <sub>2</sub> O-satd. soln. contg. dioxane.	77A100 77A100
40. 40.075 40.076 40.078	<b>cis-Bromobis(ethylenediamine)fluorocobalt(III) ion</b> $\cdot\text{CH}_2\text{OH} + \text{CoBr}(\text{en})_2\text{F}^+$ $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{CoBr}(\text{en})_2\text{F}^+$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CoBr}(\text{en})_2\text{F}^+$	$<2 \times 10^6$ $2.8 \times 10^7$ $1.1 \times 10^8$	3.5-4 3.5-4 3.5-4	p.r. p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH. D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH. D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	77A100 77A100 77A100
40.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{CoBr}(\text{en})_2\text{F}^+$	$4.9 \times 10^7$	3.5-4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. Et <sub>2</sub> O.	77A100
41. 41.075	<b>cis-Dichlorobis(ethylenediamine)cobalt(III) ion</b> $\cdot\text{CH}_2\text{OH} + \text{CoCl}_2(\text{en})_2^+$	$<5 \times 10^6$	3.5-4	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH.	77A100

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
41.076	CH <sub>3</sub> $\dot{\text{C}}$ HOH + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	3.8 x 10 <sup>7</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. EtOH.	77A100
41.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	1.0 x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 2-PrOH.	77A100
42.	<b>trans-Dichlorobis(ethylenediamine)cobalt(III) ion</b>					
42.075	·CH <sub>2</sub> OH + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	(8 ± 2) x 10 <sup>6</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. MeOH.	77A100
42.076	CH <sub>3</sub> $\dot{\text{C}}$ HOH + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	1.5 x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. EtOH.	77A100
42.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	3.8 x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 2-PrOH.	77A100
42.093	CH <sub>3</sub> $\dot{\text{C}}$ HOCH <sub>2</sub> H <sub>5</sub> + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	1.5 x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. Et <sub>2</sub> O.	77A100
42.104	–OCHCH <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> – + CoCl <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	3.5 x 10 <sup>7</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. dioxane.	77A100
43.	<b>trans-Bis(ethylenediamine)difluorocobalt(III) ion</b>					
43.075	·CH <sub>2</sub> OH + CoF <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	no reaction	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. MeOH.	77A100
43.076	CH <sub>3</sub> $\dot{\text{C}}$ HOH + CoF <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	no reaction	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. EtOH.	77A100
43.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + CoF <sub>2</sub> (en) <sub>2</sub> <sup>+</sup>	no reaction	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 2-PrOH.	77A100
44.	<b>Tris(2,2'-bipyridine)cobalt(III) ion</b>					
44.075	·CH <sub>2</sub> OH + Co(bpy) <sub>3</sub> <sup>3+</sup> → Co(bpy) <sub>3</sub> <sup>2+</sup>	2 x 10 <sup>8</sup>	1,7	p.r.	P.b.k. at 326 nm in N <sub>2</sub> O–satd. soln. contg. 0.25 mol L <sup>-1</sup> MeOH.	79A034
44.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + Co(bpy) <sub>3</sub> <sup>3+</sup> → Co(bpy) <sub>3</sub> <sup>2+</sup>	(2.5 ± 0.3) x 10 <sup>9</sup>	0.5, 7.8	p.r.	P.b.k. at 330 nm in N <sub>2</sub> O–satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	79A034
44.147	·CO <sub>2</sub> <sup>-</sup> + Co(bpy) <sub>3</sub> <sup>3+</sup> → Co(bpy) <sub>3</sub> <sup>2+</sup>	(7.6 ± 0.2) x 10 <sup>9</sup> (rel.)	6.9	p.r.	C.k. with PNBPA assuming $k(\text{CO}_2^- + \text{PNBPA}) = 1.9 \times 10^9$ in N <sub>2</sub> O–satd. soln. contg. 0.1 mol L <sup>-1</sup> formate. [26.147]	73A075
		(7.8 ± 0.5) x 10 <sup>9</sup>	6.9	p.r.	P.b.k. at 330 nm in N <sub>2</sub> O–satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	79A034
45.	<b>Tris(1,10-phenanthroline)cobalt(III) ion</b>					
45.075	·CH <sub>2</sub> OH + Co(phen) <sub>3</sub> <sup>3+</sup> → addn. to ligand	8 x 10 <sup>8</sup>	1–7	p.r.	P.b.k. at 475 nm in N <sub>2</sub> O–satd. soln. contg. 2.5 mol L <sup>-1</sup> MeOH; c.k. with O <sub>2</sub> gave (7 ± 1) x 10 <sup>8</sup> assuming $k(\text{R} + \text{O}_2) = 4.2 \times 10^9$ .	80A227
45.076	CH <sub>3</sub> $\dot{\text{C}}$ HOH + Co(phen) <sub>3</sub> <sup>3+</sup> → addn. to ligand	3.8 x 10 <sup>9</sup>	1–7	p.r.	P.b.k. at 475 nm in N <sub>2</sub> O–satd. soln. contg. 0.25 mol L <sup>-1</sup> EtOH.	80A227
45.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + Co(phen) <sub>3</sub> <sup>3+</sup> → Co(phen) <sub>3</sub> <sup>2+</sup>	4.6 x 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 360 nm in N <sub>2</sub> O–satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	79A034
45a.	<b>Tris(5,6-dimethyl-1,10-phenanthroline)cobalt(III) ion</b>					
45a.075	·CH <sub>2</sub> OH + Co(5,6-Me <sub>2</sub> phen) <sub>3</sub> <sup>3+</sup> → Co(5,6-Me <sub>2</sub> phen) <sub>3</sub> <sup>2+</sup>	4.9 x 10 <sup>8</sup>	~7	p.r.	P.b.k., as well as d.k. at 380 nm in N <sub>2</sub> O–satd. soln. contg. 2.5 mol L <sup>-1</sup> MeOH.	80A227
45a.076	CH <sub>3</sub> $\dot{\text{C}}$ HOH + Co(5,6-Me <sub>2</sub> phen) <sub>3</sub> <sup>3+</sup> → Co(5,6-Me <sub>2</sub> phen) <sub>3</sub> <sup>2+</sup>	3.1 x 10 <sup>9</sup>	~7	p.r.	P.b.k., as well as d.k. at 380 nm in N <sub>2</sub> O–satd. soln. contg. 0.25 mol L <sup>-1</sup> EtOH.	80A227
45a.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + Co(5,6-Me <sub>2</sub> phen) <sub>3</sub> <sup>3+</sup> → Co(5,6-Me <sub>2</sub> phen) <sub>3</sub> <sup>2+</sup>	3.2 x 10 <sup>9</sup>	~7	p.r.	P.b.k., as well as d.k. at 380 nm in N <sub>2</sub> O–satd. soln. contg. 0.25 mol L <sup>-1</sup> 2-PrOH; same product from e <sub>aq</sub> <sup>-</sup> reaction.	80A227
46.	<b>Diaqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(III) ion</b>					
46.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + Co <sup>III</sup> → Co <sup>II</sup>	2.0 x 10 <sup>8</sup>	2.0	p.r.	P.b.k.	76A203
46.147	·CO <sub>2</sub> <sup>-</sup> + Co <sup>III</sup> → Co <sup>II</sup>	8.1 x 10 <sup>8</sup>	2.5	p.r.	P.b.k.	76A203
47.	<b>Aqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)hydroxycobalt(III) ion</b>					
47.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + Co <sup>III</sup> → Co <sup>II</sup>	1.1 x 10 <sup>8</sup>	6.0	p.r.	P.b.k.	76A203
48.	<b>(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)dihydroxycobalt(III) ion</b>					
48.078	(CH <sub>3</sub> ) <sub>2</sub> $\dot{\text{C}}$ OH + Co <sup>III</sup> → Co <sup>II</sup>	1.1 x 10 <sup>8</sup>	10.0	p.r.	P.b.k.	76A203



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
49.	<b>Dichloro(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(III) ion</b>					
49.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Co <sup>III</sup> → Co <sup>II</sup>	7.0 × 10 <sup>8</sup>	1.0	p.r.	P.b.k.	76A203
49.147	·CO <sub>2</sub> <sup>-</sup> + Co <sup>III</sup> → Co <sup>II</sup>	1.1 × 10 <sup>9</sup>	2.5	p.r.	P.b.k.	76A203
50.	<b>Diaqua(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(III) ion</b>					
50.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Co <sup>III</sup> → Co <sup>II</sup>	1.9 × 10 <sup>9</sup>	1.0	p.r.	P.b.k.	76A203
50.147	·CO <sub>2</sub> <sup>-</sup> + Co <sup>III</sup> → Co <sup>II</sup>	6.4 × 10 <sup>9</sup>	2.5	p.r.	P.b.k.	76A203
51.	<b>Aquahydroxy(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(III) ion</b>					
51.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Co <sup>III</sup> → Co <sup>II</sup>	5.5 × 10 <sup>8</sup>	5.0	p.r.	P.b.k.	76A203
52.	<b>Dihydroxy(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(III) ion</b>					
52.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Co <sup>III</sup> → Co <sup>II</sup>	3.3 × 10 <sup>8</sup>	9.0	p.r.	P.b.k.	76A203
53.	<b>Cyanocob(III)alamin(Vitamin B12)</b>					
53.147	·CO <sub>2</sub> <sup>-</sup> + B12 → B12r (B12r See 10.)	1.2 × 10 <sup>9</sup>	6.0	p.r.	P.b.k. at 530 nm as well as d.k. at 390 nm in soln. contg. 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH, satd. with CO <sub>2</sub> .	73G116
		no reaction		p.r.	No change in o.d. in N <sub>2</sub> O or CO <sub>2</sub> -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate, or CO <sub>2</sub> -satd. soln. contg. 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	74A105
54.	<b>Hydroxocob(III)alamin(Vitamin B12a)</b>					
54.147	·CO <sub>2</sub> <sup>-</sup> + B12a → B12r (B12r See 10.)	1.45 × 10 <sup>9</sup>	9.2	p.r.	D.k. at 350 nm as well as p.b.k. at 310 nm in CO <sub>2</sub> -satd. soln. contg. <i>tert</i> -BuOH.	74A105
<b>Cr</b>						
55.	<b>Chromium(II) ion</b>					
55.006	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CO <sub>2</sub> H + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CO <sub>2</sub> H	1.1 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. pivalic acid-HClO <sub>4</sub> .	74A146
55.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH	1.0 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. <i>tert</i> -BuOH-HClO <sub>4</sub> .	74A146
55.020	·CH <sub>2</sub> CHO + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> CH(OH) <sub>2</sub>	(3.5 ± 0.5) × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. ethylene glycol-HClO <sub>4</sub> ; ·CHOHCH <sub>2</sub> OH loses water in acid soln.	74A146
55.035	·CH <sub>2</sub> CO <sub>2</sub> H + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> CO <sub>2</sub> H	2.5 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. acetic acid-HClO <sub>4</sub> .	74A146
55.037	CH <sub>3</sub> ĊHCO <sub>2</sub> H + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH(CH <sub>3</sub> )CO <sub>2</sub> H	1.1 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. propionic acid-HClO <sub>4</sub> .	74A146
55.042	·CH(CO <sub>2</sub> H) <sub>2</sub> + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH(CO <sub>2</sub> H) <sub>2</sub>	6.0 × 10 <sup>7</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. malonic acid-HClO <sub>4</sub> .	74A146
55.054	HCON(CH <sub>3</sub> )CH <sub>2</sub> + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> N(CH <sub>3</sub> )CHO	(1.1 ± 0.1) × 10 <sup>8</sup>	~5	p.r.	P.b.k. in Ar or N <sub>2</sub> O-satd. soln. contg. dimethylformamide.	74A146
55.075	·CH <sub>2</sub> OH + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> OH	1.6 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. MeOH-HClO <sub>4</sub> .	74A146
55.076	CH <sub>3</sub> ĊHOH + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CHOHCH <sub>3</sub>	7.9 × 10 <sup>7</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. EtOH-HClO <sub>4</sub> .	74A146
55.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> C(CH <sub>3</sub> ) <sub>2</sub> OH	5.1 × 10 <sup>7</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. 2-PrOH-HClO <sub>4</sub> .	74A146
55.093	CH <sub>3</sub> ĊHOC <sub>2</sub> H <sub>5</sub> + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH(CH <sub>3</sub> )OC <sub>2</sub> H <sub>5</sub>	3.4 × 10 <sup>7</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. Et <sub>2</sub> O-HClO <sub>4</sub> .	74A146
55.101	·CHOHCH <sub>2</sub> OH + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CH <sub>2</sub> CH(OH) <sub>2</sub>	(1.5 ± 0.2) × 10 <sup>8</sup>	3.0–4.5	p.r.	P.b.k. in Ar-satd. soln. contg. ethylene glycol-HClO <sub>4</sub> ; product same as in 55.020.	74A146
55.104	·OĊHCH <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> · + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> R	1.0 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. dioxane-HClO <sub>4</sub> .	74A146
55.115	·CHOHCO <sub>2</sub> H + Cr(H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> → (H <sub>2</sub> O) <sub>5</sub> Cr <sup>III</sup> CHOHCO <sub>2</sub> H	1.4 × 10 <sup>8</sup>	~1	p.r.	P.b.k. in Ar-satd. soln. contg. glycolic acid-HClO <sub>4</sub> .	74A146

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
55.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2\text{H} + \text{Cr}(\text{H}_2\text{O})_6^{2+} \rightarrow$ $(\text{H}_2\text{O})_5\text{Cr}^{\text{III}}\text{COH}(\text{CH}_3)\text{CO}_2\text{H}$	$9.2 \times 10^7$	$\sim 1$	p.r.	P.b.k. in Ar-satd. soln. contg. lactic acid-HClO <sub>4</sub> .	74A146
55.127	$\cdot\text{CONH}_2 + \text{Cr}(\text{H}_2\text{O})_6^{2+} \rightarrow$ $(\text{H}_2\text{O})_5\text{Cr}^{\text{III}}\text{CONH}_2$	$(6.5 \pm 0.6) \times 10^8$	$\sim 5$	p.r.	P.b.k. in Ar- or N <sub>2</sub> O-satd. soln. contg. formamide.	74A146
55.147	$\cdot\text{CO}_2^-(\text{CO}_2\text{H}) + \text{Cr}^{\text{II}} \rightarrow \text{Cr}^{\text{III}}\text{CO}_2^-$	$(1.1 \pm 0.1) \times 10^9$	1.4	p.r.	D.k. in soln. contg. 1 mol L <sup>-1</sup> formic acid; product spectrum similar to products containing C-Cr bonds [74A146].	73A057
56.	<b>Chromium(III) ion</b>					
56.147	$\cdot\text{CO}_2^-(\text{CO}_2\text{H}) + \text{Cr}^{\text{III}}$	no reaction	1.4	p.r.	D.k. in soln. contg. 1 mol L <sup>-1</sup> formic acid.	73A057
	<b>Cu</b>					
57.	<b>Copper(I) ions</b>					
57.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{Cu}^+ \rightarrow \text{Cu}^{\text{II}}\text{CH}_2\text{CH}_2\text{OH}^+ \rightarrow \text{Cu}^{2+} + \text{C}_2\text{H}_4 + \text{OH}^-$	$(1.9 \pm 0.6) \times 10^{10}$	4.5	p.r.	Est. from p.b.k. in N <sub>2</sub> O-satd. soln. contg. ethylene and Cu <sup>2+</sup> .	78A322
57.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Cu}^+ \rightarrow \text{Cu}^{\text{II}}\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}^+ \rightarrow \text{Cu}^{2+} + \text{OH}^- + \text{CH}_2=\text{C}(\text{CH}_3)_2$	$(2.6 \pm 0.9) \times 10^{10}$	4.5	p.r.	Est. from p.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH and Cu <sup>2+</sup> .	78A322
57.075	$\cdot\text{CH}_2\text{OH} + \text{Cu}^+ \rightleftharpoons \text{Cu}^{\text{II}}\text{CH}_2\text{OH}^+$	$\sim 10^{10}$	4.5	p.r.	Est. from growth and decay of absorption in soln. contg. MeOH and Cu <sup>2+</sup> ; $K \approx 10^4 \text{ dm}^3 \text{ mol}^{-1}$ .	78A322
		$\sim 6 \times 10^9$	4.0	p.r.	Est. from growth and decay of absorption in soln. contg. MeOH and Cu <sup>2+</sup> ; $K \approx 5 \times 10^3 \text{ dm}^3 \text{ mol}^{-1}$ .	80A278
57.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Cu}^+$	$\sim 5 \times 10^9$	4.5	p.r.	Est. from growth and decay of absorption in soln. contg. 2-ProOH and Cu <sup>2+</sup> .	78A322
58.	<b>Copper(I) ion, complex with ethylene</b>					
58.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{CuC}_2\text{H}_4^+ \rightarrow 2\text{C}_2\text{H}_4 + \text{Cu}^{2+} + \text{OH}^-$	$(7.8 \pm 2.5) \times 10^7$	4.5	p.r.	Est. from effect of [C <sub>2</sub> H <sub>4</sub> ] on rate of formn. and decay of CuC <sub>2</sub> H <sub>4</sub> <sup>+</sup> at 227 nm; radical from OH + ethylene.	78A322
58.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{CuC}_2\text{H}_4^+ \rightarrow \text{C}_2\text{H}_4 + \text{CH}_2=\text{C}(\text{CH}_3)_2 + \text{Cu}^{2+} + \text{OH}^-$	$(5.3 \pm 1.6) \times 10^7$	4.5	p.r.	Est. from effect of [C <sub>2</sub> H <sub>4</sub> ] on rate of formn. and decay of CuC <sub>2</sub> H <sub>4</sub> <sup>+</sup> at 227 nm in soln. contg. <i>tert</i> -BuOH.	78A322
58a.	<b>Ethylenediaminetetraacetatocuprate(I) ion</b>					
58a.079	$\text{CH}_3\text{CH}_2\text{CH}_2\dot{\text{C}}\text{HOH} + \text{CuEDTA}^{3-} \rightarrow \text{RCuEDTA}^{3-}$	$(4 \text{ to } 5) \times 10^9$	7.5	p.r.	P.b.k. at 440 nm in soln. contg. 0.01 mol L <sup>-1</sup> CuEDTA <sup>2-</sup> and 0.06 mol L <sup>-1</sup> 1-BuOH assuming $2k(\text{R} + \text{R}) = 10^9$ .	80A153
59.	<b>Copper(II) ions</b>					
59.001	$\cdot\text{CH}_3 + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{CH}_3\text{Cu}^{2+}$	$(7.4 \pm 0.6) \times 10^5$	$\sim 2$	f.p.	P.b.k. in 10 <sup>-2</sup> mol L <sup>-1</sup> HClO <sub>4</sub> soln. contg. 0.01–0.2 mol L <sup>-1</sup> Cu <sup>2+</sup> and Co(NH <sub>3</sub> ) <sub>5</sub> O <sub>2</sub> CCH <sub>3</sub> <sup>2+</sup> .	78F301
59.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^{\text{III}}-\text{CH}_2\text{CH}_2\text{OH}^+ \rightarrow \text{Cu}^+ + \text{prod.}$	$(1.9 \pm 0.4) \times 10^7$ $(2.2 \pm 0.4) \times 10^7$	4.5 2	p.r.	P.b.k. at 270 nm (Cu <sup>I</sup> C <sub>2</sub> H <sub>4</sub> ) in N <sub>2</sub> O-contg. ethylene (10 <sup>-3</sup> mol L <sup>-1</sup> ) soln.; $\cdot\text{C}_2\text{H}_5$ is present at pH 4.5 ( $\sim 10\%$ ) and pH 2 ( $\sim 50\%$ ).	78A322
		$(3 \pm 1) \times 10^7$	6	p.r.	P.b.k. in soln. contg. (1–5) $\times 10^{-4}$ mol L <sup>-1</sup> CuSO <sub>4</sub> satd. with ethylene-N <sub>2</sub> O (1:1).	80A277
59.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^{\text{III}}-\text{R} \rightarrow \text{Cu}^+ + \text{prod.}$	$(2.7 \pm 0.5) \times 10^6$ $(3.2 \pm 0.6) \times 10^6$	4.5 3	p.r.	P.b.k. at 320 nm (Cu <sup>I</sup> CH <sub>2</sub> CHCONH <sub>2</sub> ) in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH and acrylamide.	78A322
		$(5 \pm 2) \times 10^6$	6	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. (1–5) $\times 10^{-4}$ mol L <sup>-1</sup> CuSO <sub>4</sub> and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	80A277

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
59.035	$\cdot\text{CH}_2\text{CO}_2^- + \text{Cu}^{2+} \rightarrow \text{Cu}^{\text{III}}\text{CH}_2\text{CO}_2^+ \rightarrow \text{Cu}^+ + \text{prod.}$	$(6.4 \pm 1.5) \times 10^8$	6.0	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> acetate soln.	77A025
59.037	$\text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^{\text{III}}\text{-R} \rightarrow \text{Cu}^+ + \text{prod.}$	$(5 \pm 0.5) \times 10^8$	6	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. (2-10) $\times 10^{-5}$ mol L <sup>-1</sup> CuSO <sub>4</sub> and 10 <sup>-2</sup> mol L <sup>-1</sup> propionate.	80A277
59.041	$^-\text{O}_2\text{CCHOH}\dot{\text{C}}\text{HCO}_2^- + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^{\text{III}}\text{-R} \rightarrow \text{Cu}^+ + \text{prod.}$	$(8 \pm 0.3) \times 10^8$	6	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. (5-10) $\times 10^{-5}$ mol L <sup>-1</sup> CuSO <sub>4</sub> and (2.5-10) $\times 10^{-4}$ mol L <sup>-1</sup> fumarate; pH dependent, $k = 8.0 \times 10^8$ , $1.8 \times 10^8$ , $1.2 \times 10^8$ , $6.4 \times 10^7$ , $3.1 \times 10^7$ , and $< 2 \times 10^7$ at pH 5.0, 3.5, 3.0, 2.6, 2.1, and 1.5, resp.	80A277
59.048	$\cdot\text{CCl}_3 (+ \cdot\text{CHCl}_2) + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^{\text{III}}\text{-CCl}_3 \rightarrow \text{Cu}^+ + \text{prod.}$	$(3 \pm 1.5) \times 10^7$	6	p.r.	P.b.k. in Ar-satd. soln. contg. (2-10) $\times 10^{-5}$ mol L <sup>-1</sup> CuSO <sub>4</sub> and (1-10) $\times 10^{-2}$ mol L <sup>-1</sup> CHCl <sub>3</sub> ; $G(\cdot\text{CHCl}_2) = G(e_{\text{aq}}^-)$ and $G(\cdot\text{CCl}_3) = G(\text{OH})$ .	80A277
59.075	$\cdot\text{CH}_2\text{OH} + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^+ + \text{CH}_2\text{O} + \text{H}^+$	$(1.1 \pm 0.2) \times 10^8$	5-6	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> MeOH soln.	72A018
		$(1.6 \pm 0.3) \times 10^8$	2-5	p.r.	P.b.k. at 320 nm (Cu <sup>I</sup> CH <sub>2</sub> CHCONH <sub>2</sub> ) in soln. contg. MeOH and acrylamide (10 <sup>-2</sup> mol L <sup>-1</sup> ).	78A322
		$(1.9 \pm 0.4) \times 10^8$	$\leq 3$	p.r.	D.k.	78A322
		$(1.6 \pm 0.2) \times 10^8$	6	p.r.	D.k. at $\sim 300$ nm (radical) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	80A277
59.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^+ + \text{CH}_3\text{CHO} + \text{H}^+$	$(7.4 \pm 1.1) \times 10^7$	5-6	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> EtOH soln.	72A018
		$(9.4 \pm 1.9) \times 10^7$	2-5	p.r.	P.b.k. at 320 nm (Cu <sup>I</sup> CH <sub>2</sub> CHCONH <sub>2</sub> ) in soln. contg. EtOH and acrylamide (10 <sup>-2</sup> mol L <sup>-1</sup> ).	78A322
		$(9 \pm 2) \times 10^7$	6	p.r.	D.k. at $\sim 300$ nm (radical) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	80A277
59.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{Cu}^+ +$	$(4.5 \pm 0.7) \times 10^7$	5-6	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> 2-PrOH soln.	72A018
		$(5.2 \pm 1.0) \times 10^7$	2-5	p.r.	P.b.k. at 320 nm (Cu <sup>I</sup> CH <sub>2</sub> CHCONH <sub>2</sub> ) in soln. contg. 2-PrOH and acrylamide (10 <sup>-2</sup> mol L <sup>-1</sup> ).	78A322
		$(5.0 \pm 1.0) \times 10^7$	$\leq 3$	p.r.	D.k.	78A322
		$(5.0 \pm 1.5) \times 10^7$	6	p.r.	D.k. at $\sim 300$ nm (radical) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	80A277
59.113	$\cdot\text{A}^- + \text{Cu}_{\text{aq}}^{2+}$ (A <sup>-</sup> = Radical from ascorbate)	$\leq 10^6$		phot.	Est.	78A394
59.147	$\cdot\text{CO}_2^- + \text{Cu}_{\text{aq}}^{2+} \rightarrow \text{CO}_2 + \text{Cu}^+$	$(1.5 \pm 0.3) \times 10^8$	6.8	p.r.	D.k. at 260 nm in N <sub>2</sub> O-satd. formate (0.1 mol L <sup>-1</sup> ) soln.	78A176
60.	<b>(5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)copper(II) ion</b>					
60.075	$\cdot\text{CH}_2\text{OH} + \text{Cu}^{\text{II}}\text{L} \rightarrow \text{Cu}^{\text{I}}\text{L}$	$\leq 2 \times 10^6$	3.5-10	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH; product ident. with that from $e_{\text{aq}}^-$ .	80A189
	$\cdot\text{CH}_2\text{O}^- + \text{Cu}^{\text{II}}\text{L} \rightarrow \text{Cu}^{\text{I}}\text{L}$	$9.0 \times 10^8$	12.0	p.r.	P.b.k. at 410 nm; no reduction obs. in neutral soln.; in methanolic soln. $k = 2.2 \times 10^4$ was detd. by f.phot. (p.b.k. at 415 nm)[77F639].	76A039
60.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Cu}^{\text{II}}\text{L} \rightarrow \text{Cu}^{\text{I}}\text{L}$	$\leq 2 \times 10^6$	3.5-10	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; product ident. with that from $e_{\text{aq}}^-$ .	80A189
60.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Cu}^{\text{II}}\text{L} \rightarrow \text{Cu}^{\text{I}}\text{L}$	$\leq 2 \times 10^6$	3.5-10	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; product ident. with that from $e_{\text{aq}}^-$ .	80A189
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Cu}^{\text{II}}\text{L} \rightarrow \text{Cu}^{\text{I}}\text{L}$	$9.0 \times 10^8$	12.5	p.r.	P.b.k. at 410 nm; no redn. obs. in neutral soln.	76A039



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
60.147	$\cdot\text{CO}_2^- + \text{Cu}^{\text{II}}\text{L} \rightarrow \text{Cu}^{\text{I}}\text{L}$	$2.3 \times 10^9$	7.0	p.r.	P.b.k. at 410 nm.	76A039
61.	<b>Glycylglycylglycinatocopper(II) complex</b>					
61.147	$\cdot\text{CO}_2^- + \text{Cu}^{\text{II}}(\text{gly}_3) \rightarrow \text{Cu}(\text{I}) + \text{CO}_2$	$(2.8 \pm 0.3) \times 10^8$	9.1	p.r.	D.k. at 550 nm ( $\text{Cu}^{\text{II}}$ ) in $\text{N}_2\text{O}$ -satd. formate ( $10^{-2}$ mol L <sup>-1</sup> ) soln. contg. $\text{Cu}(\text{II})$ and $\text{gly}_3$ in 1:5, 1:3 and 1:2 ratio.	76A016
62.	<b>(Oxidized glutathione)copper(II) complex</b>					
62.147	$\cdot\text{CO}_2^- + \text{Cu}^{\text{II}}(\text{GSSG})_n \rightarrow \text{Cu}(\text{I}) + \text{CO}_2$	$(1.0 \pm 0.2) \times 10^8$	11	p.r.	D.k. at 595 nm ( $\text{Cu}^{\text{II}}$ ) in $\text{N}_2\text{O}$ -satd. soln. contg. formate ion; 10% of the $\cdot\text{CO}_2^-$ reacted with the disulfide $\rightarrow \text{GSSG}^-$ (p.b.k. at 410 nm).	76A016
63.	<b>Histidinecopper(II) complex</b>					
63.147	$\cdot\text{CO}_2^- + \text{Cu}(\text{II}) \rightarrow \text{Cu}(\text{I})$	$(4.1 \pm 0.4) \times 10^8$	11.0	p.r.	D.k. at 600 nm ( $\text{Cu}^{\text{II}}$ ) in $\text{N}_2\text{O}$ -satd. soln. contg. formate.	77A138
64.	<b>Glycylhistidinecopper(II) complex</b>					
64.147	$\cdot\text{CO}_2^- + \text{Cu}(\text{II})$	$(4.5 \pm 0.3) \times 10^8$ $(1.6 \pm 0.2) \times 10^7$	6.6 11.0	p.r.	D.k. at 565 nm ( $\text{Cu}^{\text{II}}$ ) in $\text{N}_2\text{O}$ -satd. soln. contg. formate.	77A138
65.	<b><math>\beta</math>-Alanylhistidinecopper(II) complex</b>					
65.147	$\cdot\text{CO}_2^- + \text{Cu}(\text{II})$	$(3.5 \pm 0.4) \times 10^8$	7.5–11	p.r.	D.k. at 600 nm ( $\text{Cu}^{\text{II}}$ ) in $\text{N}_2\text{O}$ -satd. soln. contg. formate.	77A138
65a.	<b>Ethylenediaminetetraacetatocuprate(II) ion</b>					
65a.079	$\text{CH}_3\text{CH}_2\text{CH}_2\dot{\text{C}}\text{HOH} + \text{CuEDTA}^{2-}$	$(0.5 \text{ to } 1) \times 10^6$	7.5	p.r.	Calcd. from concn. effect on p.b.k. for $\text{RCuEDTA}^{3-}$ ; see 58a.079.	80A153
<b>Eu</b>						
66.	<b>Europium(III) ion</b>					
66.147	$\cdot\text{CO}_2\text{H}(\cdot\text{CO}_2^-) + \text{Eu}(\text{III}) \rightarrow \text{Eu}(\text{II})$	$\geq 7 \times 10^6$	1.4	p.r.	Est. from p.b.k. at 250 nm ( $\text{Eu}^{\text{II}}$ ).	73A057
<b>Fe</b>						
67.	<b>Iron(II) ion</b>					
67.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Fe}(\text{H}_2\text{O})_6^{2+}$	$2.9 \times 10^6$	$\sim 2$	p.r.	Est. from change in abs. at 250 nm; product suggested to be $\text{Fe}(\text{H}_2\text{O})_5\text{R}^{2+}$ .	75A166
<i>Deuteroheme See 223.</i>						
68.	<b>Iron(III) ions</b>					
68.075	$\cdot\text{CH}_2\text{OH} + \text{Fe}^{3+} \rightarrow \text{Fe}^{2+} + \text{CH}_2\text{O} + \text{H}^+$	$1 \times 10^8$ (rel.) $(8 \pm 2) \times 10^7$	$\sim 1$	$\gamma$ -r.	C.k. rel. to $k(\text{R} + \text{C}(\text{NO}_2)_4) = 5 \times 10^9$ ; obs. $\text{C}(\text{NO}_2)_3^-$ and $\text{Fe}^{2+}$ yields. Unpubl. data, C.N.Barnes and G.V.Buxton.	77G411 78A322
68.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Fe}^{3+} \rightarrow \text{Fe}^{2+} + \text{H}^+ + \text{CH}_3\text{CHO}$	$2.7 \times 10^8$ (rel.) $(3.8 \pm 0.2) \times 10^8$	$\sim 1$	$\gamma$ -r.	C.k. rel. to $k(\text{R} + \text{C}(\text{NO}_2)_4) = 5 \times 10^9$ ; obs. $\text{C}(\text{NO}_2)_3^-$ and $\text{Fe}^{2+}$ yields. Unpubl. data, C.N.Barnes and G.V.Buxton.	77G411 78A322
68.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Fe}^{3+} \rightarrow \text{Fe}^{2+} + \text{H}^+ + (\text{CH}_3)_2\text{CO}$	$(4.5 \pm 0.4) \times 10^8$ $1.8 \times 10^8$ (rel.) $(5.8 \pm 0.3) \times 10^8$	$\sim 1$ $\sim 1$	p.r. $\gamma$ -r.	D.k. at 270 nm in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH, $2 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{ClO}_4)_3$ and 0.5 mol L <sup>-1</sup> $\text{HClO}_4$ ; $\mu = 1$ mol L <sup>-1</sup> . C.k. rel. to $k(\text{R} + \text{C}(\text{NO}_2)_4) = 5 \times 10^9$ ; obs. $\text{C}(\text{NO}_2)_3^-$ and $\text{Fe}^{2+}$ yields. Unpubl. data, C.N.Barnes and G.V.Buxton.	74A074 77G411 78A322

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
69.	<b>Ferricyanide ion</b>					
69.001	$\cdot\text{CH}_3 + \text{Fe}(\text{CN})_6^{3-}$	$(5 \pm 1) \times 10^6$	6–7	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. dimethyl sulfoxide.	81A003
69.002	$\cdot\text{CH}_2\text{CH}_3 + \text{Fe}(\text{CN})_6^{3-}$	$(5.0 \pm 0.5) \times 10^7$	6–7	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. diethyl sulfoxide.	81A003
69.002a	$(\text{CH}_3)_2\dot{\text{C}}\text{H} + \text{Fe}(\text{CN})_6^{3-}$	$(1.25 \pm 0.5) \times 10^9$	6–7	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. di-(1-methylethyl) sulfoxide.	81A003
69.002b	$(\text{CH}_3)_3\dot{\text{C}} + \text{Fe}(\text{CN})_6^{3-}$	$(3.6 \pm 1.5) \times 10^9$	6–7	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. di-( <i>tert</i> -butyl) sulfoxide.	81A003
69.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{Fe}(\text{CN})_6^{3-}$	$\sim 10^7$	7	p.r.	D.k. in soln. of 10 <sup>-2</sup> mol L <sup>-1</sup> ethylene contg. 10 <sup>-2</sup> mol L <sup>-1</sup> N <sub>2</sub> O.	69G522
69.035	$\cdot\text{CH}_2\text{CO}_2\text{H} + \text{Fe}(\text{CN})_6^{3-}$	$2 \times 10^6$	3.3	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. acetic acid.	81A003
69.043	$\cdot\text{CH}_2\text{Cl} + \text{Fe}(\text{CN})_6^{3-}$	$< 5 \times 10^5$	6–7	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. methylene chloride.	81A003
69.045	$\cdot\text{CHCl}_2 + \text{Fe}(\text{CN})_6^{3-}$	$< 5 \times 10^5$	6–7	p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. chloroform.	81A003
69.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{Fe}(\text{CN})_6^{3-}$	$1 \times 10^9$		p.r.	D.k. at 420 nm in soln. contg. glycine.	76A082
69.075	$\cdot\text{CH}_2\text{OH} + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{CH}_2\text{O} + \text{H}^+ + \text{Fe}(\text{CN})_6^{4-}$	$4.0 \times 10^9$	7	p.r.	D.k. at 410 nm (ferricyanide) in soln. contg. MeOH.	68G308 69G522
		$(4.2 \pm 0.4) \times 10^9$	$\sim 6$	p.r.	Condy. change in N <sub>2</sub> O–satd. soln. contg. MeOH.	70G254
	$\cdot\text{CH}_2\text{O}^- + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{CH}_2\text{O} + \text{Fe}(\text{CN})_6^{4-}$	$3.1 \times 10^9$	13	p.r.	D.k. at 410 nm (ferricyanide) in soln. contg. MeOH.	68G308
69.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Fe}(\text{CN})_6^{3-}$	$5.3 \times 10^9$	7	p.r.	D.k. at 410 nm in 0.1 mol L <sup>-1</sup> EtOH soln.	69G522
		$4 \times 10^9$		p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. EtOH.	79N061
69.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{Fe}(\text{CN})_6^{3-}$	$3.7 \times 10^9$	7	p.r.	D.k. at 410 nm in 0.1 mol L <sup>-1</sup> 1-PrOH soln.	69G522
69.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Fe}(\text{CN})_6^{3-}$	$4.7 \times 10^9$	7	p.r.	D.k. at 410 nm in 0.1 mol L <sup>-1</sup> 2-PrOH soln.	69G522
		$(5.6 \pm 0.6) \times 10^9$	7.0	p.r.	D.k. at 420 nm.	73A104
69.080	$\text{CH}_3\dot{\text{C}}\text{OHCH}_2\text{CH}_3 + \text{Fe}(\text{CN})_6^{3-}$	$4.8 \times 10^9$	7	p.r.	D.k. at 410 nm in 0.1 mol L <sup>-1</sup> 2-BuOH soln.	69G522
69.081	$(\text{CH}_3)_2\text{CH}\dot{\text{C}}\text{HOH} + \text{Fe}(\text{CN})_6^{3-}$	$3.0 \times 10^9$	7	p.r.	D.k. at 410 nm in 0.3 mol L <sup>-1</sup> 2-methyl-1-propanol soln.	69G522
69.082	$\text{CH}_3(\text{CH}_2)_3\dot{\text{C}}\text{HOH} + \text{Fe}(\text{CN})_6^{3-}$	$(9 \pm 2) \times 10^8$		p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. 1-pentanol.	79N061
69.084	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_2- + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{R}=\text{O} + \text{Fe}(\text{CN})_6^{4-} + \text{H}^+$	$2.6 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclobutanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; $k = 5.3 \times 10^9$ at 64°C; $E_a = 2.1$ kcal mol <sup>-1</sup> (8.8 kJ mol <sup>-1</sup> ).	76A103
69.085	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_3- + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{R}=\text{O} + \text{Fe}(\text{CN})_6^{4-} + \text{H}^+$	$2.3 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclopentanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; $k = 5.4 \times 10^9$ at 64°C; $E_a = 2.6$ kcal mol <sup>-1</sup> (10.9 kJ mol <sup>-1</sup> ).	76A103
69.086	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_4- + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{R}=\text{O} + \text{Fe}(\text{CN})_6^{4-} + \text{H}^+$	$1.8 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclohexanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; $k = 5.0 \times 10^9$ at 64°C; $E_a = 2.9$ kcal mol <sup>-1</sup> (12.1 kJ mol <sup>-1</sup> ).	76A103
		$(1.0 \pm 0.2) \times 10^9$		p.r.	D.k. at 420 nm in N <sub>2</sub> O–satd. soln. contg. cyclohexanol.	79N061
69.087	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_5- + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{R}=\text{O} + \text{Fe}(\text{CN})_6^{4-} + \text{H}^+$	$2.1 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cycloheptanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; $k = 5.4 \times 10^9$ at 64°C; $E_a = 2.8$ kcal mol <sup>-1</sup> (11.7 kJ mol <sup>-1</sup> ).	76A103
69.088	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_6- + \text{Fe}(\text{CN})_6^{3-} \rightarrow \text{R}=\text{O} + \text{Fe}(\text{CN})_6^{4-} + \text{H}^+$	$2.2 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclooctanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; $k = 5.4 \times 10^9$ at 64°C; $E_a = 2.8$ kcal mol <sup>-1</sup> (11.7 kJ mol <sup>-1</sup> ).	76A103

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
69.091	$\text{CH}_3\dot{\text{C}}\text{HOSO}_3^- + \text{Fe}(\text{CN})_6^{3-}$	$\approx 2 \times 10^8$		p.r.	D.k. at 420 nm in N <sub>2</sub> O-satd. soln. contg. ethylsulfate ion; values are given also for radicals produced from reaction of OH with higher alkyl sulfates.	79N061
69.092a	$\cdot\text{CH}_2\text{OCH}_3 + \text{Fe}(\text{CN})_6^{3-}$	$(4.3 \pm 0.23) \times 10^9$	6-7	p.r.	D.k. at 420 nm in N <sub>2</sub> O-satd. soln. contg. dimethyl ether.	81A003
69.097	$\text{CH}_3(\dot{\text{C}}\text{O})\text{CH}_2\text{CO}_2^- + \text{Fe}(\text{CN})_6^{3-}$	$(7.3 \pm 0.7) \times 10^8$	7.0	p.r.	D.k. at 420 nm in soln. contg. acetoacetate ion (radical by reaction with $e_{\text{aq}}^-$ ).	73A104
69.100	$\cdot\text{CHOHCH}(\text{NH}_2)\text{CO}_2^- + \text{Fe}(\text{CN})_6^{3-}$	$3.2 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> serine.	69G522
69.101	$\cdot\text{CHOHCH}_2\text{OH} + \text{Fe}(\text{CN})_6^{3-}$	$3.6 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> ethylene glycol.	69G522
69.105	$\text{R} + \text{Fe}(\text{CN})_6^{3-}$ (R = Radicals from polyethylene oxide)	$2.1 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. ~2% polyethylene oxide.	69G522
69.106	$\cdot\text{CHOHCHOHCH}_2\text{OH} + \text{Fe}(\text{CN})_6^{3-}$ (+ $\text{CH}_2\dot{\text{C}}\text{OHCH}_2\text{OH}$ )	$3.3 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> glycerol.	69G522
69.108	$\text{R} + \text{Fe}(\text{CN})_6^{3-}$ (R = Radicals from deoxyribose)	$2.8 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> deoxyribose	71G618
69.111	$\text{R} + \text{Fe}(\text{CN})_6^{3-}$ (R = Radicals from glucose)	$1.4 \times 10^9$ $1.9 \times 10^9$	7	p.r. p.r.	Soln. contains 2-deoxy-D-ribose. D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> glucose.	78A175 69G522
69.115	$\cdot\text{CHOHCO}_2^- + \text{Fe}(\text{CN})_6^{3-}$	$5 \times 10^8$	7	p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> glycolate ion.	69G522
	$\cdot\text{CHOHCO}_2\text{H} + \text{Fe}(\text{CN})_6^{3-}$	$1.0 \times 10^8$	3.4	p.r.	D.k. at 420 nm in N <sub>2</sub> O-satd. soln. contg. glycolate.	81A003
69.117	$\cdot\text{CH}(\text{O})\text{CO}_2^- + \text{Fe}(\text{CN})_6^{3-}$ $\text{CH}_3\dot{\text{C}}\text{OHCO}_2^- + \text{Fe}(\text{CN})_6^{3-}$	$7.5 \times 10^8$ $1.5 \times 10^9$	11.5 7	p.r. p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> lactate ion.	69G522
69.147	$\cdot\text{CO}_2^- + \text{Fe}(\text{CN})_6^{3-} \rightarrow$ $\text{CO}_2 + \text{Fe}(\text{CN})_6^{4-}$	$1.06 \times 10^9$	7	p.r.	D.k. at 410 nm in soln. contg. 0.3 mol L <sup>-1</sup> formate ion.	69G522
70.	<b>Pentacyanonitrosylferrate(III) ion</b>					
70.075	$\cdot\text{CH}_2\text{OH} + \text{Fe}(\text{CN})_5\text{NO}^{2-} \rightarrow$ $\text{CH}_2\text{O} + \text{H}^+ + \text{Fe}(\text{CN})_5\dot{\text{N}}\text{O}^{3-}$	$6.7 \times 10^8$	8.5	p.r.	P.b.k. in N <sub>2</sub> O-satd. 0.5 mol L <sup>-1</sup> MeOH soln.	77A120
70.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Fe}(\text{CN})_5\text{NO}^{2-}$	$2.9 \times 10^9$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. 0.5 mol L <sup>-1</sup> 2-PrOH soln.	77A120
70.147	$\cdot\text{CO}_2^- + \text{Fe}(\text{CN})_5\dot{\text{N}}\text{O}^{2-} \rightarrow$ $\text{CO}_2 + \text{Fe}(\text{CN})_5\dot{\text{N}}\text{O}^{3-}$	$3.7 \times 10^8$	7	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. $2 \times 10^{-2}$ mol L <sup>-1</sup> formate soln.; $I = 0.02$ .	69G052
		$4.0 \times 10^8$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> formate soln.	77A120
71.	<b>Pentacyanonitrosylferrate(III) radical ion (electron adduct)</b>					
71.006	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{CO}_2^- + \text{Fe}(\text{CN})_5\dot{\text{N}}\text{O}^{3-} \rightarrow$ $\text{Fe}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_2\text{CO}_2^{4-}$	$(2.0 \pm 0.3) \times 10^9$	6.0- 7.5	p.r.	D.k. at 380 nm (nitrosyl radical, $\epsilon = 3.5 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> pivalate ion and $5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{CN})_5\text{NO}^{2-}$ .	79A134
71.012	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{NH}_3^+ + \text{Fe}(\text{CN})_5\dot{\text{N}}\text{O}^{3-} \rightarrow$ $\text{Fe}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_2\text{NH}_3^{2-}$	$(1.2 \pm 0.2) \times 10^{10}$	6.0- 7.5	p.r.	D.k. at 380 nm (nitrosyl radical, $\epsilon = 3.5 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> <i>tert</i> -butylamine and $5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{CN})_5\text{NO}^{2-}$ .	79A134
71.013	$\cdot\text{CH}_2\text{C}(\text{CH}_3)(\text{NH}_3^+)\text{CO}_2^- +$ $\text{Fe}(\text{CN})_5\dot{\text{N}}\text{O}^{3-} \rightarrow$ $\text{Fe}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)(\text{NH}_3)\text{CO}_2^{3-}$	$(1.6 \pm 0.2) \times 10^9$	6.0- 7.5	p.r.	D.k. at 380 nm (nitrosyl radical, $\epsilon = 3.5 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> 2-aminoisobutyrate ion and $5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{CN})_5\text{NO}^{2-}$ .	79A134



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
71.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Fe}(\text{CN})_5\text{NO}^{3-} \rightarrow \text{Fe}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}^{3-}$	$(2.5 \pm 0.4) \times 10^9$	6.0–7.5	p.r.	D.k. at 380 nm (nitrosyl radical, $\epsilon = 3.5 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> <i>tert</i> -BuOH and $5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{CN})_5\text{NO}^{2-}$ .	79A134
71.017	$\cdot\text{CH}_2\text{COH}(\text{CH}_3)\text{CO}_2^- + \text{Fe}(\text{CN})_5\text{NO}^{3-} \rightarrow \text{Fe}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{COH}(\text{CH}_3)\text{CO}_2^{4-}$	$(6.3 \pm 0.9) \times 10^8$	6.0–7.5	p.r.	D.k. at 380 nm (nitrosyl radical, $\epsilon = 3.5 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> 2-hydroxyisobutyrate ion and $5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{CN})_5\text{NO}^{2-}$ .	79A134
71.055	$\text{CH}_3\text{CON}(\text{CH}_3)\dot{\text{C}}\text{H}_2 + \text{Fe}(\text{CN})_5\text{NO}^{3-} \rightarrow \text{Fe}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{N}(\text{CH}_3)\text{C}(\text{O})\text{CH}_3^{3-}$	$(3.5 \pm 0.5) \times 10^{10}$	6.0–7.5	p.r.	D.k. at 380 nm (nitrosyl radical, $\epsilon = 3.5 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> <i>N,N</i> -dimethylacetamide and $5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Fe}(\text{CN})_5\text{NO}^{2-}$ .	79A134
72.	<b>Tris(1,10-phenanthroline)iron(III) ion</b>					
72.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Fe}(\text{phen})_3^{3+} \rightarrow \text{Fe}(\text{phen})_3^{2+}$	$\sim 10^7$	acid	p.r.	P.b.k. in soln. contg. <i>tert</i> -BuOH, H <sub>2</sub> SO <sub>4</sub> and HClO <sub>4</sub> ; slower process follows fast H reaction.	79A174
	<i>Deuterohemin See 222.</i>					
	<i>Hemin See 224.</i>					
	<i>Hemin c See 225.</i>					
	<i>Methemerythrin See 256.</i>					
	<b>Ga</b>					
73.	<b>Gallium(II) ions</b>					
73.075	$\cdot\text{CH}_2\text{OH} + \text{Ga}^{2+} \rightarrow \text{CH}_2\text{O} + \text{H}^+ + \text{Ga}^+$	$1.1 \times 10^9$	2.3	p.r.	D.k. in soln. contg. MeOH; Ga(II) formed from $e_{\text{aq}}^- + \text{Ga(III)}$ ; cor. to $I = 0$ .	79A190
	$\cdot\text{CH}_2\text{OH} + \text{Ga}(\text{OH})^+ \rightarrow \text{CH}_2\text{O} + \text{H}_2\text{O} + \text{Ga}^+$	$1.0 \times 10^9$	2.9			
	$\cdot\text{CH}_2\text{O}^- + \text{Ga}(\text{OH})_6^{4+} \rightarrow \text{prod.}$	$7.8 \times 10^8$	12.0			
73.076	$\cdot\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{Ga}(\text{OH})_6^{4+} \rightarrow \text{prod.}$	$1.2 \times 10^9$	12.0	p.r.	D.k. in soln. contg. EtOH; Ga(II) from $e_{\text{aq}}^- + \text{Ga(III)}$ ; cor. to $I = 0$ .	79A190
73.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Ga}(\text{OH})_6^{4+} \rightarrow \text{prod.}$	$1.7 \times 10^9$	12.0	p.r.	D.k. in soln. contg. 2-PrOH; Ga(II) from $e_{\text{aq}}^- + \text{Ga(III)}$ ; cor. to $I = 0$ .	79A190
	<b>Hg</b>					
74.	<b>Mercuric bromide</b>					
74.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HgBr}_2 \rightarrow \text{HgBr} + \text{Br}^- + \text{H}^+ + (\text{CH}_3)_2\text{CO}$	$(2.4 \pm 0.6) \times 10^9$		p.r.	P.b.k. in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> 2-PrOH soln.	76A042
74.147	$\cdot\text{CO}_2^- + \text{HgBr}_2 \rightarrow \text{CO}_2 + \text{Hg(I)}$	$< 9 \times 10^9$		p.r.	P.b.k. at 360 nm in soln. contg. formate ion; based on max. optical density in various solns.; rel. rates are $\cdot\text{CO}_2^-$ 1.0, $(\text{CH}_3)_2\dot{\text{C}}\text{OH}$ 0.89, $\text{CH}_3\dot{\text{C}}\text{HOH}$ 0.87, and $\cdot\text{CH}_2\text{OH}$ 0.63.	76A087
75.	<b>Mercuric chloride</b>					
75.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HgCl}_2 \rightarrow \text{HgCl} + \text{Cl}^- + \text{H}^+ + (\text{CH}_3)_2\text{CO}$	$(2.0 \pm 0.2) \times 10^9$		p.r.	P.b.k. in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> 2-PrOH soln.	73G043 76A042
76.	<b>Mercuric iodide</b>					
76.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{HgI}_2$	$(7 \pm 2.5) \times 10^8$		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. EtOH.	78A165

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
76.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + HgI <sub>2</sub> → HgI + I <sup>-</sup> + H <sup>+</sup> + (CH <sub>3</sub> ) <sub>2</sub> CO	(2.0 ± 0.5) × 10 <sup>9</sup> (1.0 ± 0.5) × 10 <sup>9</sup>		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A042
76.147	·CO <sub>2</sub> <sup>-</sup> + HgI <sub>2</sub>	(3.0 ± 1.0) × 10 <sup>9</sup>		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate ion.	78A165
77.	<b>Mercurous cyanide</b>					
77.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + HgCN	1.6 × 10 <sup>9</sup>		p.r.	Calcd. from increase in decay rate at 285 nm in soln. contg. <i>tert</i> -BuOH and Hg(CN) <sub>2</sub> ; 2 <i>k</i> (HgCN + HgCN) = 3.4 × 10 <sup>9</sup> .	75A203
77.075	·CH <sub>2</sub> OH + HgCN	4.0 × 10 <sup>9</sup>		p.r.	Calcd. from increase in decay rate at 285 nm in soln. contg. MeOH and Hg(CN) <sub>2</sub> ; 2 <i>k</i> (HgCN + HgCN) = 3.4 × 10 <sup>9</sup> .	75A203
77.076	CH <sub>3</sub> ĊHOH + HgCN	3.9 × 10 <sup>9</sup>		p.r.	Calcd. from increase in decay rate at 285 nm in soln. contg. EtOH and Hg(CN) <sub>2</sub> ; 2 <i>k</i> (HgCN + HgCN) = 3.4 × 10 <sup>9</sup> .	75A203
77.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + HgCN	2.4 × 10 <sup>9</sup>		p.r.	Calcd. from increase in decay rate at 285 nm in soln. contg. 2-PrOH and Hg(CN) <sub>2</sub> ; 2 <i>k</i> (HgCN + HgCN) = 3.4 × 10 <sup>9</sup> .	75A203
78.	<b>Mercuric cyanide</b>					
78.147	·CO <sub>2</sub> <sup>-</sup> + Hg(CN) <sub>2</sub> → CO <sub>2</sub> + Hg(I)	(3.4 ± 0.2) × 10 <sup>9</sup>		p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. formate.	75A203
79.	<b>Mercuric thiocyanate</b>					
79.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Hg(SCN) <sub>2</sub> → HgSCN + SCN <sup>-</sup> + H <sup>+</sup> + (CH <sub>3</sub> ) <sub>2</sub> CO	(2.2 ± 0.5) × 10 <sup>9</sup>		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A042
<b>I</b>						
80.	<b>Iodine</b>					
80.001	·CH <sub>3</sub> + I <sub>2</sub> → ·[CH <sub>3</sub> I <sub>2</sub> ]	6.0 × 10 <sup>9</sup> (rel.)		p.r.	C.k. with O <sub>2</sub> ; radical from MeI; rel. to <i>k</i> (CH <sub>3</sub> + O <sub>2</sub> ) = 4.7 × 10 <sup>9</sup> .	67G041
81.	<b>Iodate ion</b>					
81.076	CH <sub>3</sub> ĊHOH + IO <sub>3</sub> <sup>-</sup> CH <sub>3</sub> ĊHO <sup>-</sup> + IO <sub>3</sub> <sup>-</sup>	< 5 × 10 <sup>6</sup> (7.5 ± 0.2) × 10 <sup>8</sup>	6 11.8	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. EtOH.	72A018
<b>Ir</b>						
<b>Iridium(IV) ions</b>						
81a.	<b>Hexachloroiridate(IV) ion</b>					
81a.001	·CH <sub>3</sub> + IrCl <sub>6</sub> <sup>2-</sup>	(1.15 ± 0.5) × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.; radical from dimethyl sulfoxide.	81A003
81a.002	·CH <sub>2</sub> CH <sub>3</sub> + IrCl <sub>6</sub> <sup>2-</sup>	(3.1 ± 0.8) × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.; radical from diethyl sulfoxide.	81A003
81a.002a	(CH <sub>3</sub> ) <sub>2</sub> ĊH + IrCl <sub>6</sub> <sup>2-</sup>	(3.6 ± 0.4) × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.; radical from di-(1-methylethyl) sulfoxide.	81A003
81a.002b	(CH <sub>3</sub> ) <sub>3</sub> Ċ + IrCl <sub>6</sub> <sup>2-</sup>	(3.8 ± 1.1) × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.; radical from di-( <i>tert</i> -butyl) sulfoxide.	81A003
81a.005	·CH <sub>2</sub> CH <sub>2</sub> Cl + IrCl <sub>6</sub> <sup>2-</sup>	~ 1 × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln. contg. 1,2-dichloroethane.	81A003
81a.014	·CH <sub>2</sub> CH <sub>2</sub> OH + IrCl <sub>6</sub> <sup>2-</sup>	~ 2 × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln. contg. ethylene.	81A003
81a.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + IrCl <sub>6</sub> <sup>2-</sup>	(1.2 ± 0.16) × 10 <sup>9</sup>	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH.	81A003
81a.020	·CH <sub>2</sub> CHO + IrCl <sub>6</sub> <sup>2-</sup>	1.7 × 10 <sup>9</sup>	7	p.r.	D.k. at 490 nm in soln. contg. 2-chloroethanol.	81A003

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
81a.035	$\cdot\text{CH}_2\text{CO}_2^- + \text{IrCl}_6^{2-}$	$4.2 \times 10^8$	7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
	$\cdot\text{CH}_2\text{CO}_2\text{H} + \text{IrCl}_6^{2-}$	$1.4 \times 10^9$	3.3		contg. acetate.	
81a.040a	$\cdot\text{O}_2\dot{\text{C}}\text{CHCH}_2\text{CO}_2^- + \text{IrCl}_6^{2-}$	$1.1 \times 10^8$	7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
	$\text{HO}_2\dot{\text{C}}\text{CHCH}_2\text{CO}_2\text{H} + \text{IrCl}_6^{2-}$	$4.6 \times 10^8$	3.3		contg. succinate.	
81a.043	$\cdot\text{CH}_2\text{Cl} + \text{IrCl}_6^{2-}$	$\sim 1 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. dichloromethane.	
81a.048	$\cdot\text{CCl}_3 + \text{IrCl}_6^{2-}$	$2.8 \times 10^7$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. carbon tetrachloride.	
81a.075	$\cdot\text{CH}_2\text{OH} + \text{IrCl}_6^{2-}$	$6.0 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. MeOH.	
81a.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{IrCl}_6^{2-}$	$4.5 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. EtOH.	
81a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{IrCl}_6^{2-}$	$4.7 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. 2-PrOH.	
81a.092	$\cdot\text{CH}_2\text{OCH}_3 + \text{IrCl}_6^{2-}$	$6.5 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. dimethyl ether.	
81a.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{IrCl}_6^{2-}$	$5.7 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. diethyl ether.	
81a.104	$-\text{CH}_2\dot{\text{C}}\text{HOCH}_2\text{CH}_2\text{O}- + \text{IrCl}_6^{2-}$	$5.4 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. dioxane.	
81a.115	$\cdot\text{CHOHCO}_2\text{H} + \text{IrCl}_6^{2-}$	$2.3 \times 10^9$	3.3	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
	$\cdot\text{CHOHCO}_2^- + \text{IrCl}_6^{2-}$	$2.0 \times 10^9$	6.9		contg. glycolate.	
	$\cdot\text{CH}(\text{O}^-)\text{CO}_2^- + \text{IrCl}_6^{2-}$	$1.8 \times 10^9$	10.7			
81a.147	$\cdot\text{CO}_2^- + \text{IrCl}_6^{2-}$	$1.7 \times 10^9$	6-7	p.r.	D.k. at 490 nm in N <sub>2</sub> O-satd. soln.	81A003
					contg. formate.	
<b>Mn</b>						
82.	<b>Manganese(II) ions</b>					
82.147	$\cdot\text{CO}_2^- + \text{Mn}^{2+} \rightarrow \text{CO}_2 + \text{Mn}^+$	$< 2 \times 10^5$		p.r.	No effect of Mn <sup>2+</sup> on d.k. of $\cdot\text{CO}_2^-$ at 280 or 256 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> formate.	76A109
83.	<b>Permanganate ion</b>					
83.001	$\cdot\text{CH}_3 + \text{MnO}_4^-$	$1.05 \times 10^9$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln.; radical from dimethyl sulfoxide.	81A003
83.002	$\cdot\text{CH}_2\text{CH}_3 + \text{MnO}_4^-$	$\sim 2 \times 10^9$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. radical from diethyl sulfoxide.	81A003
83.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{MnO}_4^-$	$1.3 \times 10^9$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH.	81A003
83.043	$\cdot\text{CH}_2\text{Cl} + \text{MnO}_4^-$	$\sim 1 \times 10^9$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. contg. dichloromethane.	81A033
83.045	$\cdot\text{CHCl}_2 + \text{MnO}_4^-$	$\sim 1 \times 10^9$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. contg. chloroform.	81A003
83.048	$\cdot\text{CCl}_3 + \text{MnO}_4^-$	$\sim 4 \times 10^8$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. contg. carbon tetrachloride.	81A003
83.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{MnO}_4^-$	$(4.2 \pm 0.4) \times 10^9$	7.0	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73A104
83.097	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{CH}_2\text{CO}_2^- + \text{MnO}_4^-$	$(4.8 \pm 0.5) \times 10^9$	9.2	p.r.	D.k. at 545 nm in Ar-satd. soln. contg. acetoacetate and <i>tert</i> -BuOH.	73A104
83.104	$-\text{CH}_2\dot{\text{C}}\text{HOCH}_2\text{CH}_2\text{O}- + \text{MnO}_4^-$	$(6.5 \pm 1.3) \times 10^9$	6-7	p.r.	D.k. at 545 nm in N <sub>2</sub> O-satd. soln. contg. dioxane.	81A003
<b>Nitrogen compounds</b>						
84.	<b>Hydroxylamine</b>					
84.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NH}_2\text{OH} \rightarrow \cdot\text{NH}_2$	$\leq 3 \times 10^3$	$< 2$	chem.	Est. from esr in Ti(III)-NH <sub>2</sub> OH soln. contg. 2-PrOH.	76D419



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
85.	<b>Nitrous oxide</b>					
85.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{N}_2\text{O}$	$(3.8 \pm 0.4) \times 10^4$ (rel.)	13.5	$\gamma$ -r.	Est. from $G(\text{acetone})$ based on an assumed mechanism and values for competing reactions.	72G167
86.	<b>Nitrite ion</b>					
86.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{NO}_2^-$	$< 5 \times 10^6$	6	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. EtOH soln.	72A018
87.	<b>Nitrate ion</b>					
87.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{NO}_3^-$	$< 5 \times 10^6$	6	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. EtOH soln.	72A018
87.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NO}_3^-$	28		$\gamma$ -r.	Calcd. from $G$ for destruction of $\text{NO}_3^-$ in soln. of 0.1–0.2 mol L <sup>-1</sup> acetone, 0.2–0.4 mol L <sup>-1</sup> 2-PrOH and $2.5 \times 10^{-4}$ mol L <sup>-1</sup> $\text{Ag}^+$ and $10^{-3}$ mol L <sup>-1</sup> Na dodecylsulfate assuming $G(\text{R}) = 5.8$ and $2k(\text{R} + \text{R}) = 1.4 \times 10^9$ .	80N062
	<b>Ni</b>					
88.	<b>Nickel(I) ion</b>					
88.004	$c\text{-}\dot{\text{C}}_5\text{H}_9 + \text{Ni}^+ \rightarrow \text{C}_5\text{H}_9\text{Ni}^+$	$2.8 \times 10^9$		p.r.	D.k. at 300 nm (as well as p.b.k.) in soln. contg. $\text{NiSO}_4$ and cyclopentane.	74A037
88.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Ni}^+ \rightarrow \text{Ni}^{2+} + \text{CH}_2=\text{C}(\text{CH}_3)_2 + \text{OH}^-$	$3 \times 10^9$		p.r.	D.k. at 300 nm ( $\text{Ni}^+$ ) in soln. contg. $\text{NiSO}_4$ and <i>tert</i> -BuOH; assume $G(\text{Ni}^+) = G(\text{R}) = 2.7$ .	74A037
88.075	$\cdot\text{CH}_2\text{OH} + \text{Ni}^+ \rightarrow \text{NiCH}_2\text{OH}^+$	$4.2 \times 10^9$		p.r.	D.k. at 300 nm (as well as p.b.k. at 250 nm); assume $G(\text{Ni}^+) = G(\text{R}) = 3.2$ ; soln. contains $\text{NiSO}_4$ and MeOH.	74A037
88.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Ni}^+ \rightarrow \text{NiCHOHCH}_3^+$	$2.3 \times 10^9$		p.r.	D.k. at 300 nm (as well as p.b.k.) in soln. cont. $\text{NiSO}_4$ and EtOH.	74A037
88.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Ni}^+ \rightarrow \text{NiC}(\text{OH})(\text{CH}_3)_2^+$	$1.4 \times 10^9$		p.r.	D.k. at 300 nm (as well as p.b.k.) in soln. contg. $\text{NiSO}_4$ and 2-PrOH.	74A037
88.147	$\cdot\text{CO}_2^- + \text{Ni}^+ \rightarrow \text{NiCO}_2$	$6.6 \times 10^9$	5.0	p.r.	D.k. at 300 nm (as well as p.b.k.) in soln. contg. $\text{NiSO}_4$ and formate ion.	74A037
89.	<b>Nickel(II) ions</b>					
89.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Ni}^{2+}$	$< 10^6$		p.r.	Est. from lack of increase in $\text{Ni}^+$ in 0.1 mol L <sup>-1</sup> $\text{Ni}^{2+}$ on addn. of 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	75A027
89.075	$\cdot\text{CH}_2\text{OH} + \text{Ni}^{2+}$	$< 10^2$		p.r.	Est. from lack of increase in $\text{Ni}^+$ in 0.1 mol L <sup>-1</sup> $\text{Ni}^{2+}$ on addn. of 0.1 mol L <sup>-1</sup> MeOH.	75A027
89.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Ni}^{2+}$	$< 10^6$		p.r.	Est. from lack of increase in $\text{Ni}^+$ in 0.1 mol L <sup>-1</sup> $\text{Ni}^{2+}$ on addn. of 0.1 mol L <sup>-1</sup> 2-PrOH.	75A027
89.147	$\cdot\text{CO}_2^- + \text{Ni}^{2+} \rightarrow \text{CO}_2 + \text{Ni}^+$	$10^2 < k < 10^5$		p.r.	Est. from lack of increase in $\text{Ni}^+$ in 0.1 mol L <sup>-1</sup> $\text{Ni}^{2+}$ on addn. of 0.1 mol L <sup>-1</sup> formate ion.	75A027
90.	<b>Tetracyanonickelate(II) ion</b>					
90.147	$\cdot\text{CO}_2^- + \text{Ni}(\text{CN})_4^{2-} \rightarrow \text{CO}_2 + \text{Ni}(\text{CN})_4^{3-}$	$(1.2 \pm 0.1) \times 10^9$		p.r.	P.b.k. at 240 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate ion.	74A072
91.	<b>(5,7,7,12,12,14-Hexamethyl-1,4,8,11-tetraazacyclotetradecane)nickel(II) ion</b>					
91.147	$\cdot\text{CO}_2^- + \text{Ni}^{\text{II}}\text{L} \rightarrow \text{Ni}^{\text{I}}\text{L} + \text{CO}_2$	$5.7 \times 10^9$	7.0	p.r.	P.b.k. in Ar-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate; $\text{Ni}(\text{I})$ is also formed by reaction of $e_{\text{aq}}^-$ .	76A039
92.	<b>(5,7,7,12,12,14)Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)nickel(II) ion</b>					
92.147	$\cdot\text{CO}_2^- + \text{Ni}^{\text{II}}\text{L} \rightarrow \text{Ni}^{\text{I}}\text{L} + \text{CO}_2$	$6.7 \times 10^9$	7.0	p.r.	P.b.k. in Ar-satd. 0.1 mol L <sup>-1</sup> formate.	76A039

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
93.	<b>Oxygen</b>					
93.001	$\cdot\text{CH}_3 + \text{O}_2 \rightarrow \text{CH}_3\text{O}_2\cdot$	$(4.7 \pm 0.7) \times 10^9$	nat	p.r.	P.b.k. at 260 nm in soln. contg. $\text{CH}_3\text{Br}$ , $\text{O}_2$ and $\text{N}_2\text{O}$ or $\text{SCN}^-$ ; $E_a = 3.5 \pm 0.5$ kcal mol <sup>-1</sup> (14.6 kJ mol <sup>-1</sup> ).	67G041
		$(3.2 \pm 0.4) \times 10^8$	5.5	p.r.	P.b.k. at 250 nm in soln. contg. $\text{CH}_4$ and $\text{O}_2$ .	72G445
93.002	$\cdot\text{CH}_2\text{CH}_3 + \text{O}_2 \rightarrow \text{C}_2\text{H}_5\text{O}_2\cdot$	$(2.9 \pm 0.8) \times 10^9$	nat	p.r.	P.b.k. at 270 nm in soln. contg. $6 \times 10^{-5}$ mol L <sup>-1</sup> $\text{O}_2$ and ethane.	75A055
93.004	$c\text{-}\dot{\text{C}}_5\text{H}_9 + \text{O}_2 \rightarrow \text{RO}_2\cdot$	$(4.9 \pm 0.6) \times 10^9$	7	p.r.	P.b.k. at 270 nm in $\text{N}_2\text{O}$ -satd. soln. contg. cyclopentane.	74A051
93.007	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radicals from oleate)	$1.0 \times 10^9$	13	p.r.	P.b.k. at 280 nm in $\text{O}_2$ - $\text{N}_2\text{O}$ -contg. $\sim 0.4$ - $0.8 \times 10^{-3}$ mol L <sup>-1</sup> soap soln.	78A365
93.008	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radicals from linoleate)	$3 \times 10^8$	10.5, 13	p.r.	D.k. at 280 nm in $\text{O}_2$ - $\text{N}_2\text{O}$ -contg. soln. of $10^{-5}$ mol L <sup>-1</sup> soap and 0.02 mol L <sup>-1</sup> phosphate.	78A365
93.009	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radicals from linolenate)	$3 \times 10^8$	10.5, 13	p.r.	D.k. at 280 nm in $\text{O}_2$ - $\text{N}_2\text{O}$ -contg. soln. of $0.4$ - $0.8 \times 10^{-3}$ mol L <sup>-1</sup> soap.	78A365
93.010	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radicals from arachidonate)	$2 \times 10^8$	13	p.r.	D.k. at 280 nm in $\text{O}_2$ - $\text{N}_2\text{O}$ -contg. soln. of $\sim 0.4$ - $0.8 \times 10^{-3}$ mol L <sup>-1</sup> soap.	78A365
93.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{O}_2 \rightarrow \cdot\text{O}_2\text{CH}_2\text{CH}_2\text{OH}$	$(6.6 \pm 1.3) \times 10^9$	1	p.r.	P.b.k. at 240 nm; radical from OH addn. in soln. contg. ethylene- $\text{O}_2$ (99:1); includes $\text{O}_2$ reaction with $\cdot\text{C}_2\text{H}_5$ from H addn.	67G269
93.035	$\cdot\text{CH}_2\text{CO}_2^- + \text{O}_2 \rightarrow \cdot\text{O}_2\text{CH}_2\text{CO}_2^-$	$(5.0 \pm 0.8) \times 10^8$	4-9	p.r.	D.k. at 350 nm in $\text{N}_2\text{O}$ -satd. soln. contg. acetate ion.	73A052
		$(2.1 \pm 0.3) \times 10^9$	8.2	p.r.	D.k. at 366 nm in $\text{N}_2\text{O}$ -satd. 0.01 mol L <sup>-1</sup> acetate soln.; cor. for $k(\text{R} + \text{R}) = 5.5 \times 10^8$ .	76A082
93.036	$\cdot\text{CH}_2\text{CO}_2\text{CH}_3 + \text{O}_2 \rightarrow \cdot\text{O}_2\text{CH}_2\text{CO}_2\text{CH}_3$	$(1.8 \pm 0.2) \times 10^9$	4	p.r.	D.k. at 340 nm in soln. contg. methyl chloroacetate and formate ion.	78A402
93.052	$\text{CH}_3\dot{\text{C}}\text{HN}^+(\text{C}_2\text{H}_5)_3 + \text{O}_2 \rightarrow \text{addn.}$		13	p.r.	Est. that $k(\text{R} + \text{R})/k(\text{R} + \text{O}_2) \approx 10$ .	78A095
93.057	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{O}_2$	$\sim 1 \times 10^9$	7.9	p.r.	D.k. at 307 nm in $\text{O}_2$ - $\text{N}_2\text{O}$ -satd. 0.2 mol L <sup>-1</sup> glycine soln.; c.k. with ferricyanide also gave $\sim 10^9$ assuming $2k(\text{R} + \text{ferricyanide}) = 1 \times 10^9$ .	76A082
93.062	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radical from glycine anhydride)	$(1.2 \pm 0.1) \times 10^9$ $(2.8 \pm 0.3) \times 10^8$	5.0 12.0	p.r.	D.k. at 360 nm in $\text{N}_2\text{O}$ -satd. soln. contg. glycine anhydride.	71G554
93.063	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radical from alanine anhydride)	$(1.0 \pm 0.2) \times 10^9$ $(1.1 \pm 0.2) \times 10^9$	5.4 12.0	p.r.	D.k. at 360 nm in $\text{N}_2\text{O}$ -satd. soln. alanine anhydride.	71G554
93.064	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radical from sarcosine anhydride)	$(9 \pm 2) \times 10^8$	5.2	p.r.	D.k. at 360 nm in $\text{N}_2\text{O}$ -satd. soln. contg. sarcosine anhydride.	71G554
93.067	$\text{R} + \text{O}_2 \rightarrow \text{RO}_2\cdot$ (R = Radicals from acetyldiglycine)	$(5.5 \pm 0.8) \times 10^8$	5.5	p.r.	D.k. at 330 nm in $\text{N}_2\text{O}$ -satd. soln. contg. acetyldiglycine.	73A052
93.075	$\cdot\text{CH}_2\text{OH} + \text{O}_2 \rightarrow \cdot\text{O}_2\text{CH}_2\text{OH}$	$4.9 \times 10^9$ (rel.)	7	p.r.	C.k. in 0.1 mol L <sup>-1</sup> MeOH soln. contg. ferricyanide; $k(\text{R} + \text{ferricyanide}) = 4.0 \times 10^9$ .	69G522
		$(4.2 \pm 0.5) \times 10^9$	10.7	p.r.	Calcd. from abs. at 248 nm assuming $k[\text{O}_2] = \text{rate of } \text{O}_2^- \text{ formn. from peroxy radical.}$	74A099
93.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{O}_2 \rightarrow \cdot\text{O}_2\text{CHOHCH}_3$	$3.6 \times 10^8$ (rel.)	7	p.r.	C.k. with RNO [ $k(\text{R} + \text{RNO}) = 2.4 \times 10^9$ ].	69G156
		$4.6 \times 10^9$ (rel.)	7	p.r.	C.k. in 0.1 mol L <sup>-1</sup> EtOH soln. contg. ferricyanide; $k(\text{R} + \text{ferricyanide}) = 5.3 \times 10^9$ .	69G522
93.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{O}_2$	$4.7 \times 10^9$ (rel.)	7	p.r.	C.k. in 0.1 mol L <sup>-1</sup> 1-PrOH soln. contg. ferricyanide; $k(\text{R} + \text{ferricyanide}) = 5.3 \times 10^9$ .	69G522

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
93.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + O <sub>2</sub> → (CH <sub>3</sub> ) <sub>2</sub> C(OH)O <sub>2</sub> ·	6.2 x 10 <sup>8</sup> (rel.) 4.2 x 10 <sup>9</sup> (rel.) 3.5 x 10 <sup>9</sup> (rel.) (4.5 ± 0.5) x 10 <sup>9</sup>	7 7 5-6 ~1	p.r. p.r. p.r. p.r.	C.k. with RNO [ $k(R + RNO) = 3.2 \times 10^9$ ]. C.k. in 0.1 mol L <sup>-1</sup> 2-PrOH soln. contg. ferricyanide; $k(R + \text{ferricyanide}) = 4.7 \times 10^9$ . C.k. in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> 2-PrOH soln. with <i>p</i> -nitroacetophenone; $k(R + \text{PNAP}) = 3.8 \times 10^9$ . D.k. at 290-300 nm in air-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 0.5 mol L <sup>-1</sup> HClO <sub>4</sub> .	69G156 69G522 71G618 74A074
93.080	CH <sub>3</sub> CH <sub>2</sub> ĊOHCH <sub>3</sub> + O <sub>2</sub>	4.0 x 10 <sup>9</sup> (rel.)	7	p.r.	C.k. in soln. contg. 0.3 mol L <sup>-1</sup> 2-BuOH with ferricyanide; $k(R + \text{ferricyanide}) = 4.8 \times 10^9$ .	69G522
93.081	(CH <sub>3</sub> ) <sub>2</sub> CHĊHOH + O <sub>2</sub>	3.4 x 10 <sup>9</sup> (rel.)	7	p.r.	C.k. in soln. contg. 0.3 mol L <sup>-1</sup> 2-methyl-1-propanol with ferricyanide; $k(R + \text{ferricyanide}) = 3.0 \times 10^9$ .	69G522
93.090	CH <sub>3</sub> CO <sub>2</sub> ĊH <sub>2</sub> + O <sub>2</sub> → CH <sub>3</sub> CO <sub>2</sub> CH <sub>2</sub> O <sub>2</sub> ·	(1.35 ± 0.15) x 10 <sup>10</sup>	6.4	p.r.	P.b.k. at 260 nm as well as d.k. at 260 nm in soln. contg. methyl acetate and N <sub>2</sub> O.	78A402
93.100	·CHOHCH(NH <sub>2</sub> )CO <sub>2</sub> <sup>-</sup> + O <sub>2</sub>	2.4 x 10 <sup>9</sup> (rel.)	7	p.r.	C.k. in 0.3 mol L <sup>-1</sup> serine soln. contg. ferricyanide; $k(R + \text{ferricyanide}) = 3.2 \times 10^9$ .	69G522
93.101	·CHOHCH <sub>2</sub> OH + O <sub>2</sub>	3.2 x 10 <sup>9</sup> (rel.)	7	p.r.	C.k. in soln. contg. 0.3 mol L <sup>-1</sup> ethylene glycol with ferricyanide; $k(R + \text{ferricyanide}) = 3.6 \times 10^9$ .	69G522
93.105	R + O <sub>2</sub> (R = Radicals from polyethylene oxide)	(1.9 - 2.9) x 10 <sup>9</sup>	7	p.r.	C.k. in soln. contg. ~2% polyethylene oxide with ferricyanide; $k(R + \text{ferricyanide}) = 2.1 \times 10^9$ .	69G522
93.106	·CHOHCHOHCH <sub>2</sub> OH + O <sub>2</sub> (+ CH <sub>2</sub> OHĊOHCH <sub>2</sub> OH)	3.3 x 10 <sup>9</sup> (rel.)	7	p.r.	C.k. in soln. contg. 0.4 mol L <sup>-1</sup> glycerol with ferricyanide; $k(R + \text{ferricyanide}) = 3.3 \times 10^9$ .	69G522
93.108	R + O <sub>2</sub> (R = Radicals from deoxyribose)	2.0 x 10 <sup>9</sup> (rel.)	5-6	p.r.	C.k. with ferricyanide in N <sub>2</sub> O-satd. soln. contg. deoxyribose; rel. to $k(R + \text{ferricyanide}) = 2.8 \times 10^9$ .	71G618
93.111	R + O <sub>2</sub> (R = Radicals from glucose)	1.6 x 10 <sup>9</sup> (rel.)	7	p.r.	C.k. in soln. contg. 0.3 mol L <sup>-1</sup> glucose and ferricyanide; $k(R + \text{ferricyanide}) = 1.9 \times 10^9$ .	69G522
93.113	·A <sup>-</sup> + O <sub>2</sub> (A = Radical from ascorbate)	< 5 x 10 <sup>2</sup> ~ 6 x 10 <sup>2</sup>	8.6	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate ion.	75A240
93.115	·CHOHCO <sub>2</sub> <sup>-</sup> + O <sub>2</sub>	1.8 x 10 <sup>9</sup> (rel.)	7	γ-r. p.r.	Obs. rate of consumption of O <sub>2</sub> . C.k. in 0.3 mol L <sup>-1</sup> glycolate soln. contg. ferricyanide; $k(R + \text{ferricyanide}) = 5 \times 10^8$ .	78A417 69G522
93.117	CH <sub>3</sub> ĊOHCO <sub>2</sub> <sup>-</sup> + O <sub>2</sub> → CH <sub>3</sub> C(ĊO <sub>2</sub> )(OH)CO <sub>2</sub> <sup>-</sup>	2.6 x 10 <sup>9</sup> (rel.) (3.5 ± 0.5) x 10 <sup>8</sup>	7 7.3	p.r. p.r.	C.k. in 0.3 mol L <sup>-1</sup> lactate soln. contg. ferricyanide; $k(R + \text{ferricyanide}) = 1.5 \times 10^9$ . D.k. at 270 nm in N <sub>2</sub> O-satd. soln. contg. lactate ion.	69G522 73A052
93.123	·CH(OH) <sub>2</sub> + O <sub>2</sub> → ·O <sub>2</sub> CH(OH) <sub>2</sub> ⇌ ·O <sub>2</sub> CH(OH)O <sup>-</sup> + H <sup>+</sup>	7.7 x 10 <sup>8</sup> (4.5 ± 0.7) x 10 <sup>9</sup>	5.7 3.5-6.5	p.r. p.r.	D.k. at 250 nm in N <sub>2</sub> O-satd. soln. contg. formaldehyde, as well as condy. increase. Condy. increase in oxygenated soln. contg. 5 x 10 <sup>-3</sup> mol L <sup>-1</sup> formaldehyde; products are formate and H <sub>2</sub> O <sub>2</sub> .	71G424 80A282
93.126	CCl <sub>3</sub> Ċ(OH) <sub>2</sub> + O <sub>2</sub> → ·O <sub>2</sub> C(OH) <sub>2</sub> CCl <sub>3</sub>	(1.0 ± 0.2) x 10 <sup>8</sup>	5.9	p.r.	D.k. at 250 nm in N <sub>2</sub> O-satd. soln. contg. chloral.	73G062
93.141	R + O <sub>2</sub> (R = Electron adduct of dimethyl fumarate)	5.3 x 10 <sup>9</sup> 2.2 x 10 <sup>9</sup>	11	p.r. p.r.	D.k. D.k.	67G729 73G097



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
93.147	$\cdot\text{CO}_2^- + \text{O}_2 \rightarrow \text{CO}_2 + \text{O}_2^-$	$7.7 \times 10^8$ (rel.)	7	p.r.	C.k. with RNO in N <sub>2</sub> O-std. soln. contg. formate; $k(\text{R} + \text{RNO}) = 1.84 \times 10^9$ .	69G156
		$2.4 \times 10^9$ (rel.)	7	p.r.	C.k. with ferricyanide in 0.3 mol L <sup>-1</sup> formate soln.; $k(\text{R} + \text{ferricyanide}) = 1.06 \times 10^9$ .	69G522
		$(2.0 \pm 0.4) \times 10^9$	8.0	p.r.	P.b.k. at 260 nm in 0.1 mol L <sup>-1</sup> formate soln. satd. with oxygen.	76A072
		$(4.2 \pm 0.4) \times 10^9$	6.8	p.r.	D.k. at 270 nm and 300 nm in 0.18 mol L <sup>-1</sup> formate ion.	76A132
		$1 \times 10^9$ (rel.)	1-4	p.r.	Est. from buildup of C(NO <sub>2</sub> ) <sub>3</sub> <sup>-</sup> in O <sub>2</sub> -satd. 0.01 mol L <sup>-1</sup> HCO <sub>2</sub> H contg. C(NO <sub>2</sub> ) <sub>4</sub> ; rel. to $k(\text{R} + \text{TNM}) = 4 \times 10^9$ .(336.147)	78A177
94.	<b>Hydrogen peroxide</b>					
94.001	$\cdot\text{CH}_3 + \text{H}_2\text{O}_2$	$<10^6$	nat	p.r.	Addn. of 10 <sup>-3</sup> and 10 <sup>-2</sup> mol L <sup>-1</sup> H <sub>2</sub> O <sub>2</sub> had no effect on second order d.k. of $\cdot\text{CH}_3$ . Suggested that reaction is slow compared to $\text{R} + \text{R}$ and may form HO <sub>2</sub> .	75A055
94.003	$\cdot\text{C}_5\text{H}_{11} + \text{H}_2\text{O}_2$	$3.4 \times 10^4$ (rel.)		$\gamma$ -r.	Est. from dose rate effect on G(H <sub>2</sub> O <sub>2</sub> ), assuming $2k(\text{R} + \text{R}) = 2 \times 10^9$ in soln. contg. $5 \times 10^{-3}$ mol L <sup>-1</sup> pentane and 10 <sup>-2</sup> mol L <sup>-1</sup> N <sub>2</sub> O.	76A254
94.004	$c\text{-}\dot{\text{C}}_5\text{H}_9 + \text{H}_2\text{O}_2$	$4.6 \times 10^4$ (rel.)		$\gamma$ -r.	Est. from dose rate effect on G(H <sub>2</sub> O <sub>2</sub> ), assuming $2k(\text{R} + \text{R}) = 1.5 \times 10^9$ in soln. contg. $2.5 \times 10^{-3}$ mol L <sup>-1</sup> cyclopentane and 10 <sup>-2</sup> mol L <sup>-1</sup> N <sub>2</sub> O.	76A254
94.016a	$\text{R} + \text{H}_2\text{O}_2$ (R = Radicals from cyclopentene + OH, 73% addn., 23% abstraction)	$3.7 \times 10^4$ (rel.)		$\gamma$ -r.	Est. from dose rate effect on G(H <sub>2</sub> O <sub>2</sub> ), assuming $2k(\text{R} + \text{R}) = 1.5 \times 10^9$ in soln. contg. $8.2 \times 10^{-3}$ mol L <sup>-1</sup> cyclopentene and 10 <sup>-2</sup> mol L <sup>-1</sup> N <sub>2</sub> O.	76A254
94.075	$\cdot\text{CH}_2\text{OH} + \text{H}_2\text{O}_2 \rightarrow \text{CH}_2\text{O} + \text{H}_2\text{O} + \text{OH}$	$(4.0 \pm 0.4) \times 10^4$ (rel.)		$\gamma$ -r.	Calcd. from dose rate effect on H <sub>2</sub> O <sub>2</sub> redn. in 0.99 mol L <sup>-1</sup> MeOH soln. assuming $2k(\text{R} + \text{R}) = 2.4 \times 10^9$ .	70G338
		$(2.3 \pm 0.8) \times 10^4$ (rel.)		chem.	Calcd. from esr of Ti(III)-H <sub>2</sub> O <sub>2</sub> soln. contg. MeOH and <i>tert</i> -BuOH; $2k(\text{R} + \text{R}) = 2 \times 10^9$ .	74D144
94.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{H}_2\text{O}_2 \rightarrow \text{CH}_3\text{CHO} + \text{H}_2\text{O} + \text{OH}$	$1.5 \times 10^5$ (rel.)		$\gamma$ -r.	Calcd. from dose rate effect on G(H <sub>2</sub> O <sub>2</sub> ) in soln. contg. EtOH assuming $2k(\text{R} + \text{R}) = 2.0 \times 10^9$ .	67G094
94.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}_2\text{O}_2 \rightarrow \text{CH}_3\text{COCH}_3 + \text{H}_2\text{O} + \text{OH}$	$5 \times 10^5$ (rel.)	1,7	phot.	Esr; effect of [H <sub>2</sub> O <sub>2</sub> ] on [R] in soln. contg. 2-PrOH and acetone; $2k(\text{R} + \text{R}) = 1.4 \times 10^9$ .	71D227
94.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{H}_2\text{O}_2$	$(5.5 \pm 1.1) \times 10^4$ (rel.)		chem.	Calcd. from esr in Ti(III)-H <sub>2</sub> O <sub>2</sub> soln. contg. Et <sub>2</sub> O; assumed $2k(\text{R} + \text{R}) = 3 \times 10^9$ .	74D144
94.094	$-\text{O}\dot{\text{C}}\text{H}(\text{CH}_2)_3- + \text{H}_2\text{O}_2$	$3.0 \times 10^4$ (rel.)		chem.	Calcd. from esr in Ti(III)-H <sub>2</sub> O <sub>2</sub> soln. contg. tetrahydrofuran; assumed $2k(\text{R} + \text{R}) = 3 \times 10^9$ .	74D144
94.104	$-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2- + \text{H}_2\text{O}_2$	$(3.0 \pm 0.6) \times 10^4$ (rel.)		chem.	Calcd. from esr in Ti(III)-H <sub>2</sub> O <sub>2</sub> soln. contg. dioxane and <i>tert</i> -BuOH; assumed $2k(\text{R} + \text{R}) = 2 \times 10^9$ .	74D144
94.124	$\cdot\text{CH}(\text{OCH}_3)_2 + \text{H}_2\text{O}_2$	$(1.1 \pm 0.3) \times 10^6$ (rel.)		chem.	Calcd. from esr in Ti(III)-H <sub>2</sub> O <sub>2</sub> soln. contg. dimethoxymethane; assumed $2k(\text{R} + \text{R}) = 2 \times 10^9$ .	74D144
94.125	$-\text{O}\dot{\text{C}}\text{HO}(\text{CH}_2)_2- + \text{H}_2\text{O}_2$	$>6 \times 10^4$ (rel.)		chem.	Calcd. from esr in Ti(III)-H <sub>2</sub> O <sub>2</sub> soln. contg. dioxolane; assumed $2k(\text{R} + \text{R}) = 3 \times 10^9$ .	74D144
94.147	$\cdot\text{CO}_2^- + \text{H}_2\text{O}_2$	$<7 \times 10^5$ (rel.)	7	phot.	Calcd. from assumed chain mechanism in CO-H <sub>2</sub> O <sub>2</sub> soln.; $k < 2.2 \times 10^6$ assuming $2k(\text{R} + \text{R}) < 10^{10}$ , recalcd. in 74D144 assuming $2k(\text{R} + \text{R}) = 3 \times 10^9$ .	63F005

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
<b>Pb</b>						
95.	<b>Lead(I) ions</b>					
95.075	$\cdot\text{CH}_2\text{OH} + \text{Pb}^+ \rightarrow \text{PbCH}_2\text{OH}^+$	$2.9 \times 10^9$		p.r.	D.k. at 300 nm (Pb <sup>+</sup> ) in soln. contg. Pb <sup>2+</sup> and MeOH, knowing initial [R] and [Pb <sup>+</sup> ]; cor. for R + R.	76A170
95.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Pb}^+ \rightarrow \text{PbCHOHCH}_3^+$	$1.7 \times 10^9$		p.r.	D.k. at 300 nm (Pb <sup>+</sup> ) in soln. contg. Pb <sup>2+</sup> and EtOH, knowing initial [R] and [Pb <sup>+</sup> ]; cor. for R + R.	76A170
95.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pb}^+ \rightarrow \text{PbC}(\text{CH}_3)_2\text{OH}^+$	$1.1 \times 10^9$		p.r.	D.k. at 300 nm (Pb <sup>+</sup> ) in soln. contg. $5 \times 10^{-4}$ mol L <sup>-1</sup> Pb <sup>2+</sup> and $5 \times 10^{-2}$ mol L <sup>-1</sup> 2-PrOH knowing initial [R] and [Pb <sup>+</sup> ]; cor. for R + R.	76A170
95.096	$\cdot\text{CHOHC}(\text{CH}_2\text{OH})_3 + \text{Pb}^+ \rightarrow \text{PbCHOHC}(\text{CH}_2\text{OH})_3^+$	$7.0 \times 10^8$		p.r.	D.k. at 300 nm (Pb <sup>+</sup> ) in soln. contg. Pb <sup>2+</sup> and pentaerythritol knowing initial [R] and [Pb <sup>+</sup> ]; cor. for R + R.	76A170
96.	<b>Lead(II) ions</b>					
96.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pb}^{2+} \rightarrow (\text{CH}_3)_2\text{CO} + \text{Pb}^+ + \text{H}^+$	$3.0 \times 10^4$	5		P.b.k. at 300 nm in 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> Pb <sup>2+</sup> during 100 $\mu$ s after pulse; Pb <sup>+</sup> formed initially by $e_{\text{aq}}^- + \text{Pb}^{2+}$ .	76A170
<b>Pt</b>						
97.	<b>trans-Dichlorobis(ethylenediamine)platinum(IV) ion</b>					
97.075	$\cdot\text{CH}_2\text{OH} + \text{Pt}(\text{en})_2\text{Cl}_2^{2+} \rightarrow \text{Pt(III)}$	$(6.9 \pm 1.0) \times 10^8$	7	p.r.	P.b.k. at 240–340 nm in N <sub>2</sub> O-satd. soln. contg. 1–4 mol L <sup>-1</sup> MeOH.	75A188
97.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pt}(\text{en})_2\text{Cl}_2^{2+} \rightarrow \text{Pt(III)}$	$(8.1 \pm 1.1) \times 10^8$	7	p.r.	P.b.k. at 260–340 nm in 2 mol L <sup>-1</sup> 2-PrOH soln.; same product as from $e_{\text{aq}}^-$ reaction.	75A188
<b>Rh</b>						
98.	<b>Tris(2,2'-bipyridine)rhodium(III) ion</b>					
98.075	$\cdot\text{CH}_2\text{OH} + \text{Rh}(\text{bpy})_3^{3+}$	$(2.2 \pm 0.2) \times 10^8$	7	p.r.	P.b.k.	74A167
98.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Rh}(\text{bpy})_3^{3+}$	$(2.9 \pm 0.3) \times 10^9$	7	p.r.	P.b.k.; same product as from $e_{\text{aq}}^-$ .	74A167
98.147	$\cdot\text{CO}_2^- + \text{Rh}(\text{bpy})_3^{3+}$	$(6.2 \pm 0.6) \times 10^9$	7	p.r.	P.b.k.; same product as from $e_{\text{aq}}^-$ .	74A167
98a.	<b>Tris(1,10-phenanthroline)rhodium(III) ion</b>					
98a.075	$\cdot\text{CH}_2\text{OH} + \text{Rh}(\text{phen})_3^{3+} \rightarrow$ addn. to ligand	$4 \times 10^8$		p.r.	P.b.k. at 475 nm in N <sub>2</sub> O-satd. soln. contg. 2.5 mol L <sup>-1</sup> MeOH.	80A227
98a.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Rh}(\text{phen})_3^{3+} \rightarrow$ addn. to ligand	$2.5 \times 10^9$		p.r.	P.b.k. at 475 nm in N <sub>2</sub> O-satd. soln. contg. 0.25 mol L <sup>-1</sup> EtOH.	80A227
98a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Rh}(\text{phen})_3^{3+} \rightarrow \text{Rh}(\text{phen})_3^{2+}$	$3.2 \times 10^9$	~7	p.r.	P.b.k. at 365 nm in N <sub>2</sub> O-satd. soln. contg. 0.25 mol L <sup>-1</sup> 2-PrOH; same product as with $e_{\text{aq}}^-$ .	80A227
<b>Ru</b>						
<b>Ruthenium(II) ions</b>						
99.	<b>Tris(2,2'-bipyridine)ruthenium(II) ion</b>					
99.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Ru}(\text{bpy})_3^{2+}$	$< 10^6$	7	p.r.	P.b.k.; no reduction.	78A068
99.075	$\cdot\text{CH}_2\text{O}^- + \text{Ru}(\text{bpy})_3^{2+} \rightarrow \text{Ru}(\text{bpy})_3^+$	$2.9 \times 10^9$	11–13	p.r.	Same product as from $e_{\text{aq}}^-$ reaction; suggested to be $[\text{Ru}(\text{bpy})_2(\text{bpy}^-)]^+$ [72G381].	78A068
99.076	$\cdot\text{CH}_2\text{OH} + \text{Ru}(\text{bpy})_3^{2+}$	$< 10^6$	7	p.r.	No reduction.	78A068
	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{Ru}(\text{bpy})_3^{2+}$	$7.0 \times 10^9$	11–13	p.r.	Same product as from $e_{\text{aq}}^-$ reaction.	78A068
	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Ru}(\text{bpy})_3^{2+}$	$< 10^6$	7	p.r.	No reduction	78A068

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
99.078	(CH <sub>3</sub> ) <sub>2</sub> ĊO <sup>-</sup> + Ru(bpy) <sub>3</sub> <sup>2+</sup> → Ru(bpy) <sub>3</sub> <sup>+</sup> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + Ru(bpy) <sub>3</sub> <sup>2+</sup>	4.9 x 10 <sup>9</sup> <10 <sup>6</sup> ~1.5 x 10 <sup>8</sup>	11–13 7 1	p.r. p.r. p.r.	Same product as from e <sub>aq</sub> <sup>-</sup> reaction. No reduction Radical addn. to ligand.	78A068 78A068 80A227
99.147	•CO <sub>2</sub> <sup>-</sup> + Ru(bpy) <sub>3</sub> <sup>2+</sup>	<10 <sup>6</sup>	7	p.r.	No reduction.	78A068
99a.	<b>Tris(1,10-phenanthroline)ruthenium(II) ion</b>					
99a.075	•CH <sub>2</sub> O <sup>-</sup> + Ru(phen) <sub>3</sub> <sup>2+</sup> → Ru(phen) <sub>3</sub> <sup>+</sup>	5.2 x 10 <sup>9</sup>	11	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2.5 mol L <sup>-1</sup> MeOH.	80A227
99a.076	CH <sub>3</sub> ĊHO <sup>-</sup> + Ru(phen) <sub>3</sub> <sup>2+</sup> → Ru(phen) <sub>3</sub> <sup>+</sup>	5.9 x 10 <sup>9</sup>	12	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.25 mol L <sup>-1</sup> EtOH; $k \approx 10^8$ at pH 1.	80A227
99a.078	(CH <sub>3</sub> ) <sub>2</sub> ĊO <sup>-</sup> + Ru(phen) <sub>3</sub> <sup>2+</sup> → Ru(phen) <sub>3</sub> <sup>+</sup> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + Ru(phen) <sub>3</sub> <sup>2+</sup>	3.7 x 10 <sup>9</sup> 8 x 10 <sup>8</sup>	13 1	p.r. p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.25 mol L <sup>-1</sup> 2-PrOH. P.b.k. in Ar-satd. soln.	80A227 80A227
	<b>Ruthenium(III) ions</b>					
100.	<b>Hexaammineruthenium(III) ion</b>					
100.035	•CH <sub>2</sub> CO <sub>2</sub> H + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	<3.0 x 10 <sup>7</sup> <5 x 10 <sup>6</sup>	2.5 3.9	p.r. p.r.	D.k. in Ar-satd. 0.01 mol L <sup>-1</sup> acetic acid soln. D.k. in N <sub>2</sub> O-satd. 0.5–1 mol L <sup>-1</sup> acetic acid soln.	72A018 77A100
	•CH <sub>2</sub> CO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	<1 x 10 <sup>7</sup>	7.3	p.r.	D.k. in N <sub>2</sub> O-satd. 0.01 mol L <sup>-1</sup> acetate soln.	72A018
100.042	•CH(CO <sub>2</sub> H) <sub>2</sub> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	<1.1 x 10 <sup>8</sup>	2.5	p.r.	D.k. in Ar-satd. 0.01 mol L <sup>-1</sup> malonic acid soln.	72A018
100.054	HCON(CH <sub>3</sub> )ĊH <sub>2</sub> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	<3.0 x 10 <sup>7</sup>	6.0	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> dimethylformamide soln.; e <sup>-</sup> -transfer.	72A018
100.056	NH <sub>2</sub> ĊHCO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	(4.0 ± 0.6) x 10 <sup>8</sup>	5.7	p.r.	D.k. in N <sub>2</sub> O-satd. 0.01 mol L <sup>-1</sup> glycine soln.; e <sup>-</sup> -transfer.	72A018
100.075	•CH <sub>2</sub> OH + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup> → CH <sub>2</sub> O + H <sup>+</sup> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>2+</sup>	(4.1 ± 0.6) x 10 <sup>7</sup>	5–6	p.r.	D.k. in N <sub>2</sub> O-satd. 0.01 mol L <sup>-1</sup> MeOH soln.	72A018
100.076	CH <sub>3</sub> ĊCHO + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	(5.5 ± 1.6) x 10 <sup>8</sup>	5–6	p.r.	D.k. in N <sub>2</sub> O-satd. 0.01 mol L <sup>-1</sup> EtOH soln.; e <sup>-</sup> -transfer.	72A018
100.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	(9.2 ± 1.4) x 10 <sup>8</sup>	5–6	p.r.	D.k. in N <sub>2</sub> O-satd. 0.01 mol L <sup>-1</sup> 2-PrOH soln.; e <sup>-</sup> -transfer.	72A018
100.093	CH <sub>3</sub> ĊHOC <sub>2</sub> H <sub>5</sub> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	(1.0 ± 0.15) x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. 0.5–1 mol L <sup>-1</sup> ethyl ether soln.	77A100
100.104	•OĊHCH <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> • + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	(5.0 ± 0.75) x 10 <sup>6</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. 0.5–1 mol L <sup>-1</sup> dioxane soln.	77A100
100.107	CH <sub>3</sub> COĊ(OH)CH <sub>3</sub> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	<3.8 x 10 <sup>6</sup> 2.0 x 10 <sup>9</sup>	1.2 5.5	p.r. p.r.	D.k. in Ar-satd. 10 <sup>-3</sup> mol L <sup>-1</sup> biacetyl soln. D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> biacetyl soln.	72A018 72A018
100.117	CH <sub>3</sub> ĊOHCO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	(2.5 ± 0.4) x 10 <sup>9</sup>	6.0	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> lactate ion; e <sup>-</sup> -transfer.	72A018
100.127	•CONH <sub>2</sub> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	<7.0 x 10 <sup>7</sup>	6.2	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> formamide soln.; e <sup>-</sup> -transfer.	72A018
100.147	•CO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup> → CO <sub>2</sub> + Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>2+</sup>	(2.0 ± 0.6) x 10 <sup>9</sup>	4.8	p.r.	D.k. in N <sub>2</sub> O-satd. 10 <sup>-2</sup> mol L <sup>-1</sup> formate soln.; e <sup>-</sup> -transfer.	72A018
101.	<b>Bromopentaammineruthenium(III) ion</b>					
101.035	•CH <sub>2</sub> CO <sub>2</sub> H + Ru(NH <sub>3</sub> ) <sub>5</sub> Br <sup>2+</sup>	(4.6 ± 0.7) x 10 <sup>8</sup>	3.9	p.r.	D.k. in N <sub>2</sub> O-satd. 0.5–1 mol L <sup>-1</sup> acetic acid.	77A100
101.093	CH <sub>3</sub> ĊHOC <sub>2</sub> H <sub>5</sub> + Ru(NH <sub>3</sub> ) <sub>5</sub> Br <sup>2+</sup>	(5.8 ± 0.9) x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. 0.5–1 mol L <sup>-1</sup> ethyl ether soln.	77A100
101.104	•OĊHCH <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> • + Ru(NH <sub>3</sub> ) <sub>5</sub> Br <sup>2+</sup>	(2.7 ± 0.4) x 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O-satd. 0.5–1 mol L <sup>-1</sup> dioxane soln.	77A100
102.	<b>Chloropentaammineruthenium(III) ion</b>					
102.035	•CH <sub>2</sub> CO <sub>2</sub> H + Ru(NH <sub>3</sub> ) <sub>5</sub> Cl <sup>2+</sup>	(4.0 ± 0.6) x 10 <sup>7</sup>	3.9	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.5–1 mol L <sup>-1</sup> acetic acid.	77A100
102.075	•CH <sub>2</sub> OH + Ru(NH <sub>3</sub> ) <sub>5</sub> Cl <sup>2+</sup>	(1.2 ± 0.2) x 10 <sup>8</sup>	3.4–5	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.5–1 mol L <sup>-1</sup> methanol.	77A100



TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
102.076	CH <sub>3</sub> ĊHOH + Ru(NH <sub>3</sub> ) <sub>5</sub> Cl <sup>2+</sup>	(8.0 ± 1.2) × 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 0.5–1 mol L <sup>-1</sup> ethanol.	77A100
102.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Ru(NH <sub>3</sub> ) <sub>5</sub> Cl <sup>2+</sup>	(1.3 ± 0.2) × 10 <sup>9</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 0.5–1 mol L <sup>-1</sup> 2-propanol.	77A100
102.093	CH <sub>3</sub> ĊHOC <sub>2</sub> H <sub>5</sub> + Ru(NH <sub>3</sub> ) <sub>5</sub> Cl <sup>2+</sup>	(2.6 ± 0.4) × 10 <sup>8</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 0.5–1 mol L <sup>-1</sup> ethyl ether.	77A100
102.104	–OĊHCH <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> – + Ru(NH <sub>3</sub> ) <sub>5</sub> Cl <sup>2+</sup>	(8.3 ± 1.2) × 10 <sup>7</sup>	3.5–4	p.r.	D.k. in N <sub>2</sub> O–satd. soln. contg. 0.5–1 mol L <sup>-1</sup> dioxane.	77A100
103.	<b>Nitrosopentaammineruthenium(III) ion</b>					
103.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> → Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> + (CH <sub>3</sub> ) <sub>2</sub> CO + H <sup>+</sup>	5.5 × 10 <sup>8</sup>	5.0	p.r.	P.b.k. at 280 nm in N <sub>2</sub> O–satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	75A049
103.147	·CO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> → Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> + CO <sub>2</sub>	3.1 × 10 <sup>9</sup>	6.6	p.r.	P.b.k. at 280 nm in N <sub>2</sub> O–satd. soln. contg. 0.5 mol L <sup>-1</sup> formate ion.	75A049
104.	<b>Nitrosopentaammineruthenium(III) radical ion (electron adduct)</b>					
104.006	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> → [Ru(NH <sub>3</sub> ) <sub>5</sub> N(O)CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CO <sub>2</sub> ] <sup>+</sup>	(2.9 ± 0.4) × 10 <sup>9</sup>	6.0– 7.5	p.r.	D.k. at 280 nm (nitrosyl radical, ε = 3.8 × 10 <sup>3</sup> L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> pivalate ion and 5 × 10 <sup>-4</sup> mol L <sup>-1</sup> Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> .	79A134
104.012	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> NH <sub>3</sub> <sup>+</sup> + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> → [Ru(NH <sub>3</sub> ) <sub>5</sub> N(O)CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> NH <sub>3</sub> ] <sup>3+</sup>	(2.0 ± 0.3) × 10 <sup>9</sup>	6.0– 7.5	p.r.	D.k. at 280 nm (nitrosyl radical, ε = 3.8 × 10 <sup>3</sup> L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> <i>tert</i> -BuNH <sub>2</sub> and 5 × 10 <sup>-4</sup> mol L <sup>-1</sup> Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> .	79A134
104.013	·CH <sub>2</sub> C(CH <sub>3</sub> )(NH <sub>3</sub> <sup>+</sup> )CO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> → [Ru(NH <sub>3</sub> ) <sub>5</sub> N(O)CH <sub>2</sub> C(CH <sub>3</sub> )(NH <sub>3</sub> )CO <sub>2</sub> ] <sup>2+</sup>	(3.1 ± 0.5) × 10 <sup>9</sup>	6.0– 7.5	p.r.	D.k. at 280 nm (nitrosyl radical, ε = 3.8 × 10 <sup>3</sup> L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> 2-aminoisobutyrate ion and 5 × 10 <sup>-4</sup> mol L <sup>-1</sup> Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> .	79A134
104.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> → [Ru(NH <sub>3</sub> ) <sub>5</sub> N(O)CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH] <sup>2+</sup>	(3.7 ± 0.6) × 10 <sup>9</sup>	6.0– 7.5	p.r.	D.k. at 280 nm (nitrosyl radical, ε = 3.8 × 10 <sup>3</sup> L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> <i>tert</i> -BuOH and 5 × 10 <sup>-4</sup> mol L <sup>-1</sup> Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> .	79A134
104.017	·CH <sub>2</sub> C(CH <sub>3</sub> )(OH)CO <sub>2</sub> <sup>-</sup> + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> → [Ru(NH <sub>3</sub> ) <sub>5</sub> N(O)CH <sub>2</sub> C(CH <sub>3</sub> )(OH)CO <sub>2</sub> ] <sup>+</sup>	(3.0 ± 0.4) × 10 <sup>9</sup>	6.0– 7.5	p.r.	D.k. at 280 nm (nitrosyl radical, ε = 3.8 × 10 <sup>3</sup> L mol <sup>-1</sup> cm <sup>-1</sup> ) as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> 2-hydroxyisobutyrate ion and 4 × 10 <sup>-4</sup> mol L <sup>-1</sup> Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> .	79A134
104.055	CH <sub>3</sub> CON(CH <sub>3</sub> )ĊH <sub>2</sub> + Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>2+</sup> → [Ru(NH <sub>3</sub> ) <sub>5</sub> N(O)CH <sub>2</sub> N(CH <sub>3</sub> )C(O)CH <sub>3</sub> ] <sup>2+</sup>	(2.9 ± 0.4) × 10 <sup>9</sup>	6.0– 7.5	p.r.	D.k. at 280 nm (nitrosyl radical, ε = 3.8 × 10 <sup>3</sup> L mol <sup>-1</sup> cm <sup>-1</sup> ), as well as p.b.k. in soln. contg. 0.3 mol L <sup>-1</sup> <i>N,N</i> -dimethylacetamide and 5 × 10 <sup>-4</sup> mol L <sup>-1</sup> Ru(NH <sub>3</sub> ) <sub>5</sub> NO <sup>3+</sup> .	79A134
105.	<b>Tris(2,2'-bipyridine)ruthenium(III) ion</b>					
105.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + Ru(bpy) <sub>3</sub> <sup>3+</sup> → Ru(bpy) <sub>3</sub> <sup>2+</sup>	1.3 × 10 <sup>8</sup> (1.9 ± 0.2) × 10 <sup>8</sup>	acid 4.6	p.r. p.r.	P.b.k. in N <sub>2</sub> O–satd. soln. contg. <i>tert</i> -BuOH.	72G462 78A070
<b>Sulfur compounds</b>						
106.	<b>Hydrogen sulfide</b>					
106.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + H <sub>2</sub> S ⇌ H <sub>2</sub> ŜCH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH → (CH <sub>3</sub> ) <sub>3</sub> COH + ·SH		6	p.r.	Effect of [H <sub>2</sub> S] on p.b.k. at 380 nm (complex) gave $K = 1 \times 10^{-3}$ mol L <sup>-1</sup> ; $k(\text{complex} \rightarrow \text{tert-BuOH} + \cdot\text{SH}) = 1.3 \times 10^5 \text{ s}^{-1}$ .	67G262
106.075	·CH <sub>2</sub> OH + H <sub>2</sub> S ⇌ H <sub>2</sub> ŜCH <sub>2</sub> OH → CH <sub>3</sub> OH + ·SH		6	p.r.	Effect of [H <sub>2</sub> S] on p.b.k. at 360 nm (complex) gave $K = 5.3 \times 10^{-4}$ mol L <sup>-1</sup> ; $k(\text{complex} \rightarrow \text{MeOH} + \cdot\text{SH}) = 2.3 \times 10^5 \text{ s}^{-1}$ .	67G262

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
106.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{H}_2\text{S} \rightleftharpoons \text{H}_2\dot{\text{S}}\text{CHOHCH}_3$ $\rightarrow \text{CH}_3\text{CH}_2\text{OH} + \cdot\text{SH}$		6	p.r.	Effect of $[\text{H}_2\text{S}]$ on p.b.k. at 380 nm (complex) gave $K = 6.4 \times 10^{-4}$ mol L <sup>-1</sup> ; $k(\text{complex} \rightarrow \text{EtOH} + \cdot\text{SH}) = 4.6 \times 10^5 \text{ s}^{-1}$ .	67G262
106.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{H}_2\text{S} \rightleftharpoons$ $\text{H}_2\dot{\text{S}}\text{CHOHCH}_2\text{CH}_3$ $\rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + \cdot\text{SH}$		6	p.r.	Effect of $[\text{H}_2\text{S}]$ on p.b.k. at 380 nm (complex) gave $K = 5.7 \times 10^{-4}$ mol L <sup>-1</sup> ; $k(\text{complex} \rightarrow 1\text{-PrOH} + \cdot\text{SH}) = 3.8 \times 10^5 \text{ s}^{-1}$ .	67G262
106.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}_2\text{S} \rightleftharpoons$ $\text{H}_2\dot{\text{S}}\text{COH}(\text{CH}_3)_2 \rightarrow$ $(\text{CH}_3)_2\text{CHOH} + \cdot\text{SH}$		6	p.r.	Effect of $[\text{H}_2\text{S}]$ on p.b.k. at 380 nm (complex) gave $K = 1.1 \times 10^{-3}$ mol L <sup>-1</sup> ; $k(\text{complex} \rightarrow 2\text{-PrOH} + \cdot\text{SH}) = 8.6 \times 10^5 \text{ s}^{-1}$ .	67G262
107.	<b>Sulfur dioxide</b>					
107.147	$\cdot\text{CO}_2^- + \text{SO}_2 \rightarrow \text{SO}_2^- + \text{CO}_2$	$(7.6 \pm 1) \times 10^8$	3.1	p.r.	P.b.k. at 320 nm in soln. contg. 1 mol L <sup>-1</sup> formate; $2k(\text{R} + \text{R}) = 7.6 \times 10^8$ .	75A118
108.	<b>Tetrathionate ion</b>					
108.147	$\cdot\text{CO}_2^- + \text{S}_4\text{O}_6^{2-} \rightarrow \text{CO}_2 + \text{S}_4\text{O}_6^{3-}$	$5.8 \times 10^7$		p.r.	P.b.k. at 370 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate, as well as d.k. at 280 nm.	73A027
	<b>Sc</b>					
109.	<b>Scandium(III) ions</b>					
109.147	$\cdot\text{CO}_2^-(\cdot\text{CO}_2\text{H}) + \text{Sc(III)}$		1.4	p.r.	No reaction in 1 mol L <sup>-1</sup> formic acid and $1 \times 10^{-2}$ mol L <sup>-1</sup> Sc(III).	73A057
	<b>Ti</b>					
110.	<b>Titanium(III) ions</b>					
110.035	$\cdot\text{CH}_2\text{CO}_2\text{H} + \text{Ti}^{3+} \rightarrow \text{Ti}^{3+}\text{-CH}_2\text{CO}_2\text{H}$ $\rightarrow \text{Ti(IV)}$	$(4 \pm 1) \times 10^6$	0.5	p.r.	Complex formn. deduced from transient spectra.	79A341
110.042	$\cdot\text{CH}(\text{CO}_2\text{H})_2 + \text{Ti(III)} \rightarrow \text{Ti(IV)}$	$8 \times 10^6$	~0	e-r.	Calcd. from effect of $[\text{Ti}^{\text{III}}]$ on on esr signal intensity; $2k(\text{R} + \text{R}) \approx 10^9$ .	73G249
110.075	$\cdot\text{CH}_2\text{OH} + \text{Ti(III)}$			p.r.	No reaction in soln. contg. 1 mol L <sup>-1</sup> MeOH and $10^{-2}$ mol L <sup>-1</sup> Ti(III).	73A057
110.147	$\cdot\text{CO}_2^- + \text{Ti(III)} \rightarrow \text{Ti(II)}$	$\sim 5 \times 10^6$	1.4	p.r.	D.k. in 1 mol L <sup>-1</sup> formic acid soln.; $\text{p}K_a(\cdot\text{CO}_2\text{H}) = 1.4$ .	73A057
	$\cdot\text{CO}_2\text{H}(\cdot\text{CO}_2^-) + \text{Ti}^{3+} \rightarrow \text{Ti}^{3+}\text{-CO}_2\text{H}$ $\rightarrow \text{Ti(II)}$	$(4 \pm 1) \times 10^6$	0.5	p.r.	Complex formn. deduced from transient spectra.	79A341
	<b>Tl</b>					
111.	<b>Thallium(I) ion</b>					
111.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{Tl}^+ \rightarrow \text{Tl}_2^+$	$1.5 \times 10^9$	13	p.r.	P.b.k. at 420 nm in soln. contg. 0.1 mol L <sup>-1</sup> EtOH and $2 \times 10^{-4}$ mol L <sup>-1</sup> Tl <sup>+</sup> ; no reaction in neutral or acid soln.	80A123
111.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Tl}^+ \rightarrow \text{Tl}_2^+$	$3.0 \times 10^9$	13	p.r.	P.b.k. at 420 nm in soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH and $1 \times 10^{-4}$ mol L <sup>-1</sup> Tl <sup>+</sup> ; no reaction in neutral or acid soln. See also 79G191.	80A123
111.147	$\cdot\text{CO}_2^- + \text{Tl}^+ \rightarrow \text{Tl}_2^+$	$2.3 \times 10^6$	13	p.r.	P.b.k. at 420 nm in soln. contg. 1 mol L <sup>-1</sup> formate and $1.5 \times 10^{-2}$ mol L <sup>-1</sup> Tl <sup>+</sup> ; reaction also obs. for neutral and acid soln.	80A123

TABLE 1. Rate constants for reactions of aliphatic radicals with inorganic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
111a.	<b>Thallium(I) ion, complex with Tl(O)</b>					
111a.075	$\cdot\text{CH}_2\text{OH} + \text{Tl}_2^+ + \text{H}^+ \rightarrow \text{CH}_3\text{OH} + 2 \text{Tl}^+$	$4.0 \times 10^9$	6	p.r.	Calcd. from d.k. at 420 nm. and condy. change in soln. contg. alcohol and $\text{Tl}^+$ assuming values for $2k(\text{Tl}_2^+ + \text{Tl}_2^+)$ , $k(\text{Tl}_2^+ + \text{H}_2\text{O}_2)$ and $2k(\text{R} + \text{R})$ .	80A123
111a.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Tl}_2^+ + \text{H}^+ \rightarrow \text{CH}_3\text{CH}_2\text{OH} + 2 \text{Tl}^+$	$3.0 \times 10^9$	6	p.r.	See above.	80A123
111a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Tl}_2^+ + \text{H}^+ \rightarrow (\text{CH}_3)_2\text{CHOH} + 2 \text{Tl}^+$	$3.0 \times 10^9$	6	p.r.	See above.	80A123
	<b>Yb</b>					
112.	<b>Ytterbium(III) ions</b>					
112.147	$\cdot\text{CO}_2^- + (\text{CO}_2\text{H}) + \text{Yb(III)}$		1.4	p.r.	No reaction in soln. contg. 1 mol L <sup>-1</sup> formic acid and 10 <sup>-2</sup> mol L <sup>-1</sup> Yb(III).	73A057
	<b>Zn</b>					
113.	<b>Zinc(I) ion</b>					
113.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Zn}^+ (+ \text{H}^+) \rightarrow \text{Zn}^{2+} + (\text{CH}_3)_3\text{COH}$	$(1.0 \pm 0.3) \times 10^9$	7	p.r.	Calcd. from d.k. at 310 nm ( $\text{Zn}^+$ ) in <i>tert</i> -BuOH-ZnSO <sub>4</sub> soln. assuming $k(\text{H} + \text{H}) = 1.3 \times 10^{10}$ , $k(\text{Zn}^+ + \text{Zn}^+) = 4 \times 10^8$ , $k(\text{R} + \text{R}) = 6.5 \times 10^8$ and $k(\text{Zn}^+ + \text{H}) = 2.8 \times 10^9$ .	77A011
113.075	$\cdot\text{CH}_2\text{OH} + \text{Zn}^+ (+ \text{H}^+) \rightarrow \text{Zn}^{2+} + \text{CH}_3\text{OH}$	$(2.5 \pm 0.3) \times 10^9$	7	p.r.	Calcd. from d.k. at 310 nm ( $\text{Zn}^+$ ) in N <sub>2</sub> O-MeOH-ZnSO <sub>4</sub> soln. taking $k(\text{Zn}^+ + \text{Zn}^+) = 4.5 \times 10^8$ , $k(\text{Zn}^+ + \text{H}_2\text{O}_2) = 2.4 \times 10^9$ , $k(\text{Zn}^+ + \text{N}_2\text{O}) = 1.6 \times 10^7$ and $k(\text{R} + \text{R}) = 1.5 \times 10^9$ .	77A011
113.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Zn}^+ (+ \text{H}^+) \rightarrow \text{Zn}^{2+} + (\text{CH}_3)_2\text{CHOH}$	$(1.3 \pm 0.25) \times 10^9$	~3,7	p.r.	Calcd. from d.k. at 310 nm ( $\text{Zn}^+$ ) in 2-PrOH-ZnSO <sub>4</sub> soln. taking $k(\text{Zn}^+ + \text{Zn}^+) = 4.5 \times 10^8$ .	77A011
113.147	$\cdot\text{CO}_2^- + \text{Zn}^+ (+ \text{H}^+) \rightarrow \text{Zn}^{2+} + \text{HCO}_2^-$	$\sim 4 \times 10^9$		p.r.	Est. from first-order decay at 310 nm ( $\text{Zn}^+$ ) in formate-ZnSO <sub>4</sub> soln.	77A011
114.	<b>Zinc(II) ion</b>					
114.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Zn}^{2+}$	$< 10^6$		p.r.	Est. from lack of increase in $\text{Zn}^+$ in 0.1 mol L <sup>-1</sup> $\text{Zn}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	75A027
114.075	$\cdot\text{CH}_2\text{OH} + \text{Zn}^{2+}$	$< 10^2$		p.r.	Est. from lack of increase in $\text{Zn}^+$ in 0.1 mol L <sup>-1</sup> $\text{Zn}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> MeOH.	75A027
114.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Zn}^{2+}$	$< 10^6$		p.r.	Est. from lack of increase in $\text{Zn}^+$ in 2-PrOH.	75A027
114.147	$\cdot\text{CO}_2^- + \text{Zn}^{2+}$	$< 10^2$		p.r.	Est. from lack of increase in $\text{Zn}^+$ in 0.1 mol L <sup>-1</sup> $\text{Zn}^{2+}$ soln. upon addn. of 0.1 mol L <sup>-1</sup> formate ion.	75A027
		$< 2 \times 10^4$		p.r.	No reaction in 0.5 mol L <sup>-1</sup> ZnSO <sub>4</sub> .	77A011
<i>Zinc hematoporphyrin See 221.</i>						
<i>Zinc(II)-Insulin See 233a.</i>						



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
115.	<b>Acetic acid</b>					
115.001	$\cdot\text{CH}_3 + \text{CH}_3\text{CO}_2\text{H} \rightarrow$ $\text{CH}_4 + \cdot\text{CH}_2\text{CO}_2\text{H}$	$2 \times 10^2$	$\sim 1$	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
116.	<b>Acetone</b>					
116.001	$\cdot\text{CH}_3 + \text{CH}_3\text{COCH}_3 \rightarrow$ $\text{CH}_4 + \cdot\text{CH}_2\text{COCH}_3$	$0.99[k(\text{R} + \text{R})]^{0.5}$ $1.00[k(\text{R} + \text{R})]^{0.5}$	nat nat	phot. phot.	Est. from quantum yields of CH <sub>4</sub> , CO, and C <sub>2</sub> H <sub>6</sub> . Obs. yields of CH <sub>4</sub> and C <sub>2</sub> H <sub>6</sub> ; $\log k$ (in cm <sup>3</sup> mol <sup>-1</sup> s <sup>-1</sup> ) = [(4.47 $\pm$ 0.05) - (1335 $\pm$ 18)/T] - 0.5 $\log k(\text{R} + \text{R})$ .	60F004 69F176
117.	<b>Acetonitrile</b>					
117.001	$\cdot\text{CH}_3 + \text{CH}_3\text{CN} \rightarrow$ $\text{CH}_4 + \cdot\text{CH}_2\text{CN}$	$< 3 \times 10^2$	$\sim 1$	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr.	75D188
118.	<b>Acetophenone</b>					
118.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow$ $\text{CH}_3\text{CHO} + (\text{C}_6\text{H}_5\text{COCH}_3)^{\cdot-}$	$1.1 \times 10^9$	13	p.r.	P.b.k. at 440 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> EtOH; pH dependent, $k$ extrapolated to pH 14.	73G122
118.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow$ $\text{CH}_3\text{COCH}_3 + (\text{C}_6\text{H}_5\text{COCH}_3)^{\cdot-}$	$7.8 \times 10^8$ $9 \times 10^8$	13 13	p.r. p.r.	P.b.k. at 440 nm in acetone soln. P.b.k. at 440 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH; pH dependent, $k$ extrapolated to pH 14.	67G729 73G122
118.142	$\text{R} + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow (\text{C}_6\text{H}_5\text{COCH}_3)^{\cdot-}$ (R = Electron adduct of glycine anhydride)	$(2.3 \pm 0.3) \times 10^9$ $(2.0 \pm 0.3) \times 10^9$	5.2 12.3	p.r.	P.b.k. at 280 nm (320 at pH 12.3) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
118.143	$\text{R} + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow (\text{C}_6\text{H}_5\text{COCH}_3)^{\cdot-}$ (R = Electron adduct of alanine anhydride)	$(2.0 \pm 0.3) \times 10^9$ $(1.5 \pm 0.4) \times 10^9$	5.2 12.2	p.r.	P.b.k. at 280 nm (320 at pH 12.2) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
118.144	$\text{R} + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow (\text{C}_6\text{H}_5\text{COCH}_3)^{\cdot-}$ (R = Electron adduct of sarcosine anhydride)	$(2.0 \pm 0.2) \times 10^9$ $(2.1 \pm 0.2) \times 10^9$	5.2 12.4	p.r.	P.b.k. at 280 nm (320 at pH 12.4) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
118.147	$\cdot\text{CO}_2^- + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow$ $\text{CO}_2 + (\text{C}_6\text{H}_5\text{COCH}_3)^{\cdot-}$	$1 \times 10^7$	12	p.r.	P.b.k. at 440 nm.	68G308
119.	<b>Acetylene</b>					
119.022	$\cdot\text{CH}=\text{CHOH} + \text{CH}=\text{CH}$	$(8.8 \pm 0.9) \times 10^6$	4.0, 6.0	p.r.	Radical from OH + C <sub>2</sub> H <sub>2</sub> ; p.b.k. at 290 nm; $E_s = 25.1$ kJ mol <sup>-1</sup> ; product thought to be hydroxybutadienyl which reacts to give hydroxyhexatrienyl (p.b.k. at 260 nm, $k = 5 \times 10^6$ L mol <sup>-1</sup> s <sup>-1</sup> ).	78A007
120.	<b>3-Acetylpyridine</b>					
120.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3\text{COC}_5\text{H}_4\text{NH}^+ \rightarrow$ $e^-$ -transfer	$(8.6 \pm 1.7) \times 10^9$	0.6	p.r.	P.b.k.; no $e^-$ -transfer in neutral soln.; $pK_a = 4.9$ .	74A089
121.	<b>Acridine</b>					
121.075	$\cdot\text{CH}_2\text{OH} + \text{C}_{13}\text{H}_9\text{NH}^+$	$5.0 \times 10^8$	2.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. MeOH; $\sim 30\%$ $e^-$ -transfer; no reaction at pH 7.6; $pK_a = 5.6$ .	74A127
121.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_{13}\text{H}_9\text{N} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \cdot\text{A}^-$	$3 \times 10^9$	13	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	79A305
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_{13}\text{H}_9\text{N} \rightarrow$ $\cdot\text{AR} + \cdot\text{A}$ (50% redn., 50% addn.)	$3.0 \times 10^8$ $\sim 3 \times 10^8$	7.6 7-9	p.r. p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $\sim 40\%$ $e^-$ -transfer.	74A127 79A305
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_{13}\text{H}_9\text{NH}^+ \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{AH}$	$3.7 \times 10^9$ $3.3 \times 10^9$	2.0 4	p.r. p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $\sim 90\%$ $e^-$ -transfer.	74A127 79A305

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
121.147	$\cdot\text{CO}_2^- + \text{AH}^+ \rightarrow \cdot\text{AH}$	$\sim 3 \times 10^8$	7	p.r.	P.b.k.; at pH 13 adduct is formed ( $\cdot\text{CO}_2^- + \text{A} \rightarrow \cdot\text{ACO}_2^-$ ).	79A305
122. 122.147	<b>Acriflavin (3,10-Diamino-10-methylacridinium chloride)</b> $\cdot\text{CO}_2^- + \text{A}$	$(3.7 \pm 0.4) \times 10^8$		p.r.	D.k. (dye) in Ar-satd. 0.1 mol L <sup>-1</sup> formate; same product as concurrent fast reaction with $e_{\text{aq}}^-$ .	70G241
123. 123.016	<b>Acrylamide</b> $\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{CH}_2=\text{CHCONH}_2 \rightarrow$ $\text{HOC}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\dot{\text{C}}\text{HCONH}_2$	$(4.5 \pm 1.4) \times 10^7$ $(6.8 \pm 2) \times 10^7$ (rel.)	4.5 3	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH and Cu <sup>2+</sup> ; rel. to $k(\text{R} + \text{Cu}^{2+}) = 2.7 \times 10^6$ at pH 4.5 and $3.2 \times 10^6$ at pH 3.	78A322
123.075	$\cdot\text{CH}_2\text{OH} + \text{CH}_2=\text{CHCONH}_2 \rightarrow$ $\text{HOCH}_2\text{CH}_2\dot{\text{C}}\text{HCONH}_2$	$(2.6 \pm 0.8) \times 10^7$ (rel.)	2-5	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. MeOH and Cu <sup>2+</sup> ; rel. to $k(\text{R} + \text{Cu}^{2+}) = 1.6 \times 10^8$ .	78A322
123.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{CH}_2=\text{CHCONH}_2 \rightarrow$ $\text{CH}_3\text{CHOHCH}_2\dot{\text{C}}\text{HCONH}_2$	$(2.6 \pm 0.8) \times 10^7$ (rel.)	2-5	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. EtOH and Cu <sup>2+</sup> ; rel. to $k(\text{R} + \text{Cu}^{2+}) = 9.4 \times 10^7$ .	78A322
123.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_2=\text{CHCONH}_2 \rightarrow$ $(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\dot{\text{C}}\text{HCONH}_2$	$(4.1 \pm 1.2) \times 10^7$ (rel.)	2-5	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH and Cu <sup>2+</sup> ; rel. to $k(\text{R} + \text{Cu}^{2+}) = 5.2 \times 10^7$ .	78A322
123.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{CH}_2=\text{CHCONH}_2 \rightarrow$ $\text{HOC}(\text{NH}_2)=\text{CH}(\text{CH}_2)_2\dot{\text{C}}\text{HCONH}_2$	$(2.5 \pm 0.4) \times 10^5$	$\sim 6$	p.r.	D.k. at 275 nm.	70G052
123.147	$\cdot\text{CO}_2^- + \text{CH}_2=\text{CHCONH}_2 \rightarrow \text{addn.}$	$\sim (4 \pm 1) \times 10^7$	$\sim 5$	p.r.	Electron transfer not obs.	70G052
124. 124.078	<b>Adenosine</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{A} \rightarrow$ $e^-$ -transfer $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{A}$	$4.7 \times 10^7$ $< 10^6$ $< 10^6$	2.2 7.0 13.6	p.r.	P.b.k.	75A060
125. 125.147	<b>Adenosine monophosphate</b> $\cdot\text{CO}_2^- + \text{A}$	$< 10^6$	8.3	p.r.	P.b.k. at 550 nm (A <sup>-</sup> ) in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> formate.	68G441
126. 126.147	<b>Alloxan</b> $\cdot\text{CO}_2^- + \text{A} (+ \text{H}^+) \rightarrow$ $\text{CO}_2 + \cdot\text{AH}$	$(3.7 \pm 1.1) \times 10^7$	4.8	p.r.	P.b.k. at 310 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate; cor. for decay of $\cdot\text{CO}_2^-$ .	79A327 80A197
127. 127.078	<b>2-Amino-5-nitrothiazole</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{NX}^- + \text{H}^+$	$2.0 \times 10^9$		p.r.	P.b.k.	76A075
128. 128.020	<b>4-Aminophenoxide ion</b> $\cdot\text{CH}_2\text{CHO} + \text{NH}_2\text{C}_6\text{H}_4\text{O}^- \rightarrow$ $\text{CH}_3\text{CHO} + \text{NH}_2\text{C}_6\text{H}_4\text{O}\cdot$	$2.1 \times 10^9$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
129. 129.078	<b>4-Aminopyrimidine</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{A}$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{A}$		0.8- 13.0	p.r.	No $e^-$ -transfer.	77A034
130. 130.036	<b>Aniline</b> $\cdot\text{CF}_3 + \text{C}_6\text{H}_5\text{NH}_2 \rightarrow \text{addn.}$	$(4.7 \pm 0.9) \times 10^6$	9-10	p.r.	P.b.k. at 330 nm in N <sub>2</sub> O-satd. soln. contg. CF <sub>3</sub> Cl; c.k. rel. to $k(\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ gave $k = 5.6 \times 10^6$ .	70G407
131. 131.078	<b>9,10-Anthraquinone</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{A} \rightarrow$ semiquinone	$(1.6 \pm 0.1) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73A104

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
131.097	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{CH}_2\text{CO}_2^- + \text{A} \rightarrow$ semiquinone	$(6.7 \pm 0.7) \times 10^8$	9.2	p.r.	P.b.k. at 400 nm in Ar-satd. soln. contg. acetoacetate and 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	73A104
132.	<b>9,10-Anthraquinone-1-sulfonate ion</b>					
132.147	$\cdot\text{CO}_2^- + \text{A} \rightarrow$ semiquinone	$1.0 \times 10^9$ $3.3 \times 10^9$	3 7	p.r.	P.b.k.	72G391
133.	<b>9,10-Anthraquinone-2-sulfonate ion</b>					
133.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{A} \rightarrow$ semiquinone	$(2.2 \pm 0.2) \times 10^9$	8.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine.	73A104
133.068	$\text{R} + \text{A} \rightarrow$ semiquinone (R = Radicals from triglycine)	$(1.4 \pm 0.1) \times 10^9$	10	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. triglycine.	73A104
133.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{A} \rightarrow$ semiquinone	$(3.0 \pm 0.3) \times 10^9$ $(5.6 \pm 0.6) \times 10^9$	7.0	p.r. p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH. P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	73A104 76A070
133.097	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{CH}_2\text{CO}_2^- + \text{A} \rightarrow$ semiquinone	$(2.1 \pm 0.2) \times 10^9$	9.2	p.r.	P.b.k. at 400 nm. in Ar-satd. soln. contg. acetoacetate and <i>tert</i> -BuOH.	73A104
133.115	$\cdot\text{CHOHCO}_2^- + \text{A} \rightarrow$ semiquinone	$(7.1 \pm 0.7) \times 10^8$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycolate.	73A104
133.147	$\cdot\text{CO}_2^- + \text{A} \rightarrow$ semiquinone	$2.8 \times 10^9$ $3.1 \times 10^9$ $(1.6 \pm 0.2) \times 10^9$	3 7 7	p.r. p.r.	P.b.k. in soln. contg. formate. P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. formate.	72G391 73A104
134.	<b>9,10-Anthraquinone-2,6-disulfonate ion</b>					
134.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{A} \rightarrow$ semiquinone	$(2.6 \pm 0.3) \times 10^9$	8.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine.	73A104
134.068	$\text{R} + \text{A} \rightarrow$ semiquinone (R = Radicals from triglycine)	$(1.8 \pm 0.2) \times 10^9$	10	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. triglycine.	73A104
134.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{A} \rightarrow$ semiquinone	$(4.6 \pm 0.5) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73A104
134.097	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{CH}_2\text{CO}_2^- + \text{A} \rightarrow$ semiquinone	$(7.2 \pm 0.7) \times 10^8$	9.2	p.r.	P.b.k. at 400 nm. in Ar-satd. soln. contg. acetoacetate and <i>tert</i> -BuOH.	73A104
134.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2\text{H} + \text{A}$	$3.0 \times 10^9$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. $5 \times 10^{-3}$ mol L <sup>-1</sup> lactate; 58% <i>e</i> - transfer.	75A051
134.147	$\cdot\text{CO}_2^- + \text{A} \rightarrow$ semiquinone	$(2.4 \pm 0.2) \times 10^9$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate.	73A104
135.	<b>Ascorbate ion</b>					
135.020	$\cdot\text{CH}_2\text{CHO} + \text{A}^- (+ \text{H}_2\text{O}) \rightarrow$ $\text{CH}_3\text{CHO} + \cdot\text{A}^- + \text{OH}^-$	$8.8 \times 10^7$	7	p.r.	Radical from ClCH <sub>2</sub> CH <sub>2</sub> OH by reaction with OH and HCl elimination.	79A051
135.042	$\cdot\text{CH}(\text{CO}_2^-)_2 + \text{AH}^- \rightarrow$ $\text{CH}_2(\text{CO}_2^-)_2 + \cdot\text{A}^-$	$(1.3 \pm 0.1) \times 10^7$	7.5	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.8 mol L <sup>-1</sup> malonate.	73R006
135.075	$\cdot\text{CH}_2\text{OH} + \text{AH}^-$	$< 10^6$		p.r.		77A036
135.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{AH}^- \rightarrow$ $(\text{CH}_3)_2\text{CHOH} + \cdot\text{A}^-$	$(1.2 \pm 0.1) \times 10^6$	6.1	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> acetone and 2 mol L <sup>-1</sup> 2-PrOH.	73R006
136.	<b>Azobenzene</b>					
136.075	$\cdot\text{CH}_2\text{O}^- + \text{C}_6\text{H}_5\text{N}=\text{NC}_6\text{H}_5 \rightarrow$ $\text{CH}_2\text{O} + [\text{C}_6\text{H}_5\text{N}=\text{NC}_6\text{H}_5]^-$	$1 \times 10^9$	14	p.r.	P.b.k. at 380 nm in soln. contg. MeOH; $k$ same for <i>syn</i> - and <i>anti</i> - isomers.	77A169
136.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{N}=\text{NC}_6\text{H}_5 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + [\text{C}_6\text{H}_5\text{N}=\text{NC}_6\text{H}_5]^-$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_6\text{H}_5\text{N}=\text{NC}_6\text{H}_5 \rightarrow$ $(\text{CH}_3)_2\text{CO} + [\text{C}_6\text{H}_5\text{N}=\text{NC}_6\text{H}_5]^-$	$4 \times 10^8$ $2 \times 10^9$	$\sim 7$ 14	p.r. p.r.	P.b.k. at 380 nm in soln. contg. 2-PrOH; <i>anti</i> - isomer. P.b.k. at 380 nm in soln. contg. 2-PrOH; <i>anti</i> -isomer.	77A169 77A169
137.	<b>Benzophenone</b>					
137.075	$\cdot\text{CH}_2\text{O}^- + (\text{C}_6\text{H}_5)_2\text{CO} \rightarrow$ $\text{CH}_2\text{O} + (\text{C}_6\text{H}_5)_2\dot{\text{C}}\text{O}^-$	$1.2 \times 10^8$	13	p.r.	P.b.k. at 610 nm in N <sub>2</sub> O-satd. soln. 1 mol L <sup>-1</sup> MeOH.	75A125



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
137.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + (\text{C}_6\text{H}_5)_2\text{CO} \rightarrow \text{CH}_3\text{CHO} + (\text{C}_6\text{H}_5)_2\dot{\text{C}}\text{O}^-$	$1 \times 10^9$	13	p.r.	P.b.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. EtOH; $E_a \approx 3.1$ kcal mol <sup>-1</sup> (13 kJ mol <sup>-1</sup> ).	74A010
137.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + (\text{C}_6\text{H}_5)_2\text{CO} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5)_2\dot{\text{C}}\text{O}^- \rightarrow$	$2.6 \times 10^8$	13	p.r.	P.b.k. at 610 nm in N <sub>2</sub> O-satd. soln.	75A125
		$1.2 \times 10^9$	12	p.r.	P.b.k.	68G308
		$(1.6 \pm 0.2) \times 10^9$	13.1	p.r.	P.b.k. at 605 nm in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
		$7.0 \times 10^8$	13	p.r.	P.b.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	75A125
137.142	$\text{R} + (\text{C}_6\text{H}_5)_2\text{CO} \rightarrow (\text{C}_6\text{H}_5)_2\dot{\text{C}}\text{O}^-$ (R = Electron adduct of glycine anhydride)	$(2.2 \pm 0.3) \times 10^9$	5.5	p.r.	P.b.k. at 330 nm (320 at pH 12.3) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
		$(2.5 \pm 0.4) \times 10^9$	12.3			
137.143	$\text{R} + (\text{C}_6\text{H}_5)_2\text{CO} \rightarrow (\text{C}_6\text{H}_5)_2\dot{\text{C}}\text{O}^-$ (R = Electron adduct of alanine anhydride)	$(1.6 \pm 0.2) \times 10^9$	5.2	p.r.	P.b.k. at 330 nm (320 at pH 12.2) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
		$(1.9 \pm 0.3) \times 10^9$	12.2			
137.144	$\text{R} + (\text{C}_6\text{H}_5)_2\text{CO} \rightarrow (\text{C}_6\text{H}_5)_2\dot{\text{C}}\text{O}^-$ (R = Electron adduct of sarcosine anhydride)	$(2.3 \pm 0.2) \times 10^9$	5.2	p.r.	P.b.k. at 330 nm (320 at pH 12.2) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
		$(2.4 \pm 0.2) \times 10^9$	12.2			
138.	<b>Benzoquinone</b>					
138.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{Q} \rightarrow \text{semiquinone}$	$(3.9 \pm 0.4) \times 10^9$	8.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine.	73A104
138.062	$\text{R} + \text{Q} \rightarrow \text{semiquinone}$ (R = Radical from glycine anhydride)	$2.2 \times 10^9$	10.5	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. glycine anhydride; 87% <i>e</i> -transfer.	73A052
138.068	$\text{R} + \text{Q} \rightarrow \text{semiquinone}$ (R = Radicals from triglycine)	$(2.5 \pm 0.2) \times 10^9$	10	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. triglycine	73A104
138.075	$\cdot\text{CH}_2\text{OH} + \text{Q} \rightarrow \text{CH}_2\text{O} + \text{H}^+ + \cdot\text{Q}^-$	$6.1 \times 10^9$	nat	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	71G619
		$4.8 \times 10^9$	6.2	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	73G049
138.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Q} \rightarrow \text{CH}_3\text{CHO} + \text{H}^+ + \cdot\text{Q}^-$	$4.5 \times 10^9$	nat	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> EtOH.	71G619
138.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{Q}^-$	$5.0 \times 10^9$	nat	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	71G619
		$(5.4 \pm 0.5) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73A104
		$5 \times 10^9$	6.9	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> 2-PrOH.	73G049
		$5.0 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> acetone and 1 mol L <sup>-1</sup> 2-PrOH.	73G125
				p.r.	P.b.k. at 430 nm in soln. contg. 0.1 mol L <sup>-1</sup> sodium dodecylsulfate $k(\text{R} + \text{micelle}) = 9.9 \times 10^8$ .	78A261
138.092	$\text{R} + \text{Q} \rightarrow \cdot\text{Q}^-$ (R = Radicals from sodium dodecylsulfate)	$1.2 \times 10^9$		p.r.		78A261
138.108	$\text{R} + \text{Q} \rightarrow \text{semiquinone}$ (R = Radicals from deoxyribose)	$2.7 \times 10^9$	5.6	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. deoxyribose.	73G049
138.115	$\cdot\text{CHOHCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$2.2 \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycolate 72% <i>e</i> -transfer.	73A052 73A104
138.116	$\text{CH}_3\dot{\text{C}}\text{OHCONH}_2 + \text{Q} \rightarrow \cdot\text{Q}^-$	$2.0 \times 10^9$ $3.6 \times 10^9$	5.0 7.3	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. lactamide.	73A052
138.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$6.5 \times 10^9$ $\sim 7.0 \times 10^9$	7.3 10.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. lactate; 97% <i>e</i> -transfer.	73A052
138.147	$\cdot\text{CO}_2^- + \text{Q} \rightarrow \text{CO}_2 + \cdot\text{Q}^-$	$6.6 \times 10^9$	$\sim 7$	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	71G619
		$7 \times 10^9$	6.9	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	73G049
		$(6.6 \pm 0.7) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. formate.	73A104
139.	<b>3-Benzoyl-1-methylpyridinium ion</b>					
139.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N}^+\text{CH}_3 \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COHC}_5\text{H}_4\text{NCH}_3)^+$	$(2.3 \pm 0.3) \times 10^9$	5.0, 1.0	p.r.	P.b.k. at 530 nm in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
140.	<b>2-Benzoylpyridine</b>					
140.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{NH}^+ \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COHC}_5\text{H}_4\text{NH})^+$	$(3.0 \pm 0.3) \times 10^9$	1.0	p.r.	P.b.k. at 500 nm, as well as at 330 nm, in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{NH})\cdot$	$(1.5 \pm 0.2) \times 10^8$	5.9	p.r.	P.b.k. at 500 nm, as well as at 330 nm, in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N})\cdot^-$	$(2.3 \pm 0.2) \times 10^9$	13.2	p.r.	P.b.k. at 550 nm, as well as at 330 nm, in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
141.	<b>3-Benzoylpyridine</b>					
141.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{NH}^+ \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COHC}_5\text{H}_4\text{NH})^+$	$(1.7 \pm 0.2) \times 10^9$	0.9	p.r.	P.b.k. at 500 nm, as well as at 330 nm (at pH 0.5 giving $k = 2.3 \times 10^9$ ) in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{NH})\cdot$	$(1.0 \pm 0.2) \times 10^8$	5.1	p.r.	P.b.k. at 530 nm, as well as at 330 nm in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N})\cdot^-$	$(2.0 \pm 0.2) \times 10^9$	13.0	p.r.	P.b.k. at 605 nm, as well as at 341 nm, in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
142.	<b>4-Benzoylpyridine</b>					
142.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{NH}^+ \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COHC}_5\text{H}_4\text{NH})^+$	$(2.5 \pm 0.2) \times 10^9$	0.7	p.r.	P.b.k. at 510 nm in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{NH})\cdot$	$(2.4 \pm 0.4) \times 10^8$	7.7	p.r.	P.b.k. at 500 nm in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N} \rightarrow (\text{CH}_3)_2\text{CO} + (\text{C}_6\text{H}_5\text{COC}_5\text{H}_4\text{N})\cdot^-$	$(2.5 \pm 0.2) \times 10^9$	13.2	p.r.	P.b.k. at 575 nm in N <sub>2</sub> O-satd. soln. contg. 2 mol L <sup>-1</sup> 2-PrOH.	72G359
<i>Benzyl viologen See 1,1'-Dibenzyl-4,4'-bipyridinium ion 171.</i>						
<i>Biacetyl See 2,3-Butanedione 148.</i>						
143.	<b>2,2'-Bipyridine</b>					
143.075	$\cdot\text{CH}_2\text{OH} + \text{bpy}$	$<10^6$	7	p.r.	P.b.k.	79A148
	$\cdot\text{CH}_2\text{O}^- + \text{bpy}$	$<10^6$	alk.	p.r.	P.b.k.	79A148
143.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{bpy}$	$<10^6$	7	p.r.	P.b.k.	79A148
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{bpyH}^+ \rightarrow (\text{CH}_3)_2\text{CO} + \text{bpyH}_2^{+ \cdot}$	$3.5 \times 10^8$	1.4, 3.7	p.r.	P.b.k. at 375 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> 2-PrOH.	79A148
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{bpy} \rightarrow (\text{CH}_3)_2\text{CO} + \text{bpy}\cdot^-$	$1.3 \times 10^8$	13.0	p.r.	P.b.k. at 365 nm in soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH and 0.1 mol L <sup>-1</sup> acetone.	79A148
143.147	$\cdot\text{CO}_2^- + \text{bpy}$	$<10^6$	7	p.r.	P.b.k.	79A148
	$\cdot\text{CO}_2^- + \text{bpyH}^+$	$5.0 \times 10^8$	4.4	p.r.	P.b.k. at 375 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	79A148
144.	<b>1,1'-Bis(2-hydroxyethyl)-4,4'-bipyridinium ion</b>					
144.147	$\cdot\text{CO}_2^- + \text{BP}^{2+} \rightarrow \text{CO}_2 + \cdot\text{BP}^+$	$1.9 \times 10^{10}$	6.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	78A321
145.	<b>2-Bromo-5-nitrothiazole</b>					
145.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{NX}^-$	$3.0 \times 10^9$		p.r.	P.b.k.	76A075
146.	<b>5-Bromouracil</b>					
146.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{BrU}$	$2 \times 10^7$ (rel.)		p.r.	C.k. rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
146.147	$\cdot\text{CO}_2^- + \text{BrU} \rightarrow \text{CO}_2 + \text{Br}^- + \text{U}\cdot$	$>1.0 \times 10^8$		p.r.	P.b.k. at 440 nm (BrUCO <sub>2</sub> <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. formate.	69G826

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
147.	<b>Butadiene</b>					
147.001	$\cdot\text{CH}_3 + \text{C}_4\text{H}_6 \rightarrow \text{addn.}$	$1.25 \times 10^6$ (rel.)		$\gamma$ -r.	C.k. with MeOH in soln. contg. MeI; obs. $G(\text{CH}_4)$ ; rel. to $k(\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
147.047	$\cdot\text{CF}_3 + \text{C}_4\text{H}_6 \rightarrow \text{addn.}$	$(5.8 \pm 1.0) \times 10^8$ (rel.)	9-10	$\gamma$ -r.	C.k. with 2-PrOH in soln. contg. $\text{CF}_3\text{Cl}$ ; rel. to $k(\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ .	70G407
148.	<b>2,3-Butanedione</b>					
148.075	$\cdot\text{CH}_2\text{OH} + \text{CH}_3\text{COCOCH}_3 (\text{B}) \rightarrow \text{CH}_2\text{O} + \text{H}^+ + \cdot\text{B}^-$	$1.1 \times 10^8$ $(6.5 \pm 1) \times 10^7$		p.r.	P.b.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> MeOH.	68G249
			5-6	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> MeOH.	72A018
148.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{B} \rightarrow \text{CH}_3\text{CHO} + \text{H}^+ + \cdot\text{B}^-$	$5.6 \times 10^8$		p.r.	P.b.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> EtOH.	68G249
148.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{B} \rightarrow \text{CH}_3\text{CH}_2\text{CHO} + \text{H}^+ + \cdot\text{B}^-$	$6.8 \times 10^8$		p.r.	P.b.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 1-PrOH.	68G249
148.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{B} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{B}^-$	$8.6 \times 10^8$ $6.0 \times 10^8$		p.r.	P.b.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> 2-PrOH.	68G249
			5-6	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> 2-PrOH.	72A018
148.080	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{OHCH}_3 + \text{B} \rightarrow \text{CH}_3\text{COCH}_2\text{CH}_3 + \text{H}^+ + \cdot\text{B}^-$	$7.2 \times 10^8$		p.r.	P.b.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 2-BuOH.	68G249
148.083	$c\text{-C}_6\text{H}_{11}\dot{\text{C}}\text{HOH} + \text{B} \rightarrow \text{C}_6\text{H}_{11}\text{CHO} + \text{H}^+ + \cdot\text{B}^-$	$2.6 \times 10^8$		p.r.	P.b.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. cyclohexanemethanol.	68G249
148.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2^- + \text{B}$	$2.8 \times 10^7$	6.0	p.r.	D.k. in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> lactate.	72A018
149.	<b>1-Butene</b>					
149.001	$\cdot\text{CH}_3 + \text{C}_4\text{H}_8 \rightarrow \text{addn.}$	$3.0 \times 10^4$ (rel.)		$\gamma$ -r.	C.k. with MeOH in soln. contg. MeI; obs. $G(\text{CH}_4)$ ; rel. to $k(\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
149.047	$\cdot\text{CF}_3 + \text{C}_4\text{H}_8 \rightarrow \text{addn.}$	$(5.3 \pm 0.7) \times 10^7$ (rel.)	9-10	$\gamma$ -r.	C.k. with 2-PrOH; rel. to $k(\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ .	70G407
<i>1,1'-Butylene-2,2'-bipyridinium ion See Tetramethylene-2,2'-bipyridinium ion 341.</i>						
150.	<b>Camphor</b>					
150.147	$\cdot\text{CO}_2^- + \text{RC}=\text{O}$	$\leq 10^6$	13	p.r.	D.k. of $\cdot\text{CO}_2^-$ at 260 nm. in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate and 0.1 mol L <sup>-1</sup> NaOH same in the presence and absence of $10^{-3}$ mol L <sup>-1</sup> camphor.	79A191
151.	<b>3-Carbamoyl-1-methylpyridinium ion</b>					
151.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CONH}_2$	$(3.6 \pm 0.7) \times 10^8$	9.5	p.r.	P.b.k. (pyridinyl radical) in $\text{N}_2\text{O}$ -satd. soln. contg. 1-2 mol L <sup>-1</sup> 2-PrOH.	74A089
151.147	$\cdot\text{CO}_2^- + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CONH}_2$	$4.6 \times 10^9$	8.5	p.r.	P.b.k. at 420 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	68G441
152.	<b>Carbon tetrachloride</b>					
152.075	$\cdot\text{CH}_2\text{OH} + \text{CCl}_4 \rightarrow \cdot\text{CCl}_3 + \text{Cl}^- + \text{CH}_2\text{O} + \text{H}^+$	$\leq 10^6$		p.r.	Condy.; $G(\text{Cl}^-) \approx 0.3G(\cdot\text{CH}_2\text{OH})$ .	71G778
152.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CCl}_4 \rightarrow \cdot\text{CCl}_3 + \text{Cl}^- + (\text{CH}_3)_2\text{CO} + \text{H}^+$	$1.0 \times 10^8$ $7 \times 10^8$ (rel.)		p.r.	Condy.; build-up of HCl in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	71G778
				p.r.	C.k. with <i>p</i> -nitroacetophenone in soln. contg. 2-PrOH and acetone; rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140 73A150
152.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{CCl}_4 \rightarrow \text{Cl}^-$	$2.5 \times 10^7$		p.r.	Condy.; buildup of HCl in $\text{N}_2\text{O}$ -satd. soln. contg. $\text{Et}_2\text{O}$ .	71G778



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
152.094	$\text{-O}\dot{\text{C}}\text{H}(\text{CH}_2)_3\text{-} + \text{CCl}_4 \rightarrow \text{Cl}^-$	$2.0 \times 10^7$		p.r.	Condy.; buildup of HCl in N <sub>2</sub> O-satd. soln. contg. tetrahydrofuran.	71G778
152.147	$\cdot\text{CO}_2^- + \text{CCl}_4$	no reaction		p.r.	Condy.	71G778
153.	<b>3-Carboxy-1-methylpyridinium ion</b>					
153.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CO}_2^-$	$1.0 \times 10^8$	9.2	p.r.	P.b.k. (pyridinyl radical) at 400 nm in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	74A106
154.	<b>4-Carboxy-1-methylpyridinium ion</b>					
154.075	$\cdot\text{CH}_2\text{O}^- + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CO}_2^-$	$3.8 \times 10^9$	12.7	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	79A051
154.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CO}_2^-$	$3.8 \times 10^9$	12.7	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	79A051
154.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CO}_2^-$	$1.5 \times 10^9$	8.6	p.r.	P.b.k. at 400 nm in soln contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	79A106
<i>Catechol See 1,2-Dihydroxybenzene 176.</i>						
155.	<b>Chloral hydrate</b>					
155.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Cl}_3\text{CH}(\text{OH})_2$	$1 \times 10^6$ (rel.)		p.r.	C.k. with <i>p</i> -nitroacetophenone in soln. contg. 2-PrOH and acetone; rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140 73A150
<i>Chloranil See Tetrachlorobenzoquinone 338.</i>						
156.	<b>Chloroacetic acid</b>					
156.001	$\cdot\text{CH}_3 + \text{ClCH}_2\text{CO}_2\text{H} \rightarrow \text{CH}_4 + \text{Cl}\dot{\text{C}}\text{HCO}_2\text{H}$	$3.0 \times 10^3$	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
156a.	<b><i>m</i>-Chlorobenzonitrile</b>					
156a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{ClC}_6\text{H}_4\text{CN} \rightarrow (\text{CH}_3)_2\dot{\text{C}}\text{O} + (\text{ClC}_6\text{H}_4\text{CN})^-$	$(2 \pm 0.5) \times 10^8$	13	p.r.	P.b.k. at 310 nm (radical anion); $k < 10^7$ at pH 7.	81A001
157.	<b>Chloroform</b>					
157.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CHCl}_3$	$< 1 \times 10^7$ (rel.)		p.r.	C.k. with <i>p</i> -nitroacetophenone in soln. contg. 2-PrOH and acetone; rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
158.	<b><i>p</i>-Chloronitrobenzene</b>					
158.111	$\text{R} + \text{ClC}_6\text{H}_4\text{NO}_2$ (R = Radicals from glucose)	$4.0 \times 10^8$		p.r.		77R167
158.147	$\cdot\text{CO}_2^- + \text{ClC}_6\text{H}_4\text{NO}_2$	$3 \times 10^8$		p.r.		77R167
159.	<b>Crystal Violet</b>					
159.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{dye}$	$1.2 \times 10^9$	7	p.r.	D.k. at 525 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 56% <i>e</i> -transfer.	73A078
159.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{dye}$	$2.3 \times 10^9$	7	p.r.	D.k. at 525 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 84% <i>e</i> -transfer.	73A078
159.147	$\cdot\text{CO}_2^- + \text{dye} \rightarrow \text{CO}_2 + \text{dye}^-$	$1.5 \times 10^9$	7	p.r.	D.k. at 520 nm as well as p.b.k. at ~400 nm.	73A078
160.	<b>Cyanoacetic acid</b>					
160.001	$\cdot\text{CH}_3 + \text{CNCH}_2\text{CO}_2\text{H} \rightarrow \text{CH}_4 + \text{CN}\dot{\text{C}}\text{HCO}_2\text{H}$	$> 6.6 \times 10^3$	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr and values for competing reactions.	75D188

*2-Cyanoethanol See 3-Hydroxypropionitrile 230.*

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
160a.	<b>Cystamine</b>					
160a.142	R + Cys → <i>e</i> -transfer (R = Electron adduct of glycine anhydride)	(1.2 ± 0.3) × 10 <sup>8</sup>	5.7, 11.0	p.r.	D.k. at 265 nm (R) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
160a.143	R + Cys → <i>e</i> -transfer (R = Electron adduct of alanine anhydride)	(1.1 ± 0.2) × 10 <sup>8</sup>	5.1, 11.4	p.r.	D.k. at 265 nm (R) in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
161.	<b>Cysteamine</b>					
161.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + RSH → (CH <sub>3</sub> ) <sub>3</sub> COH + RS·	1.8 × 10 <sup>7</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH.	68G132
161.018	R' + RSH (R' = Radicals from allyl alcohol)	< 1 × 10 <sup>7</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. allyl alcohol.	68G132
161.033	·CH <sub>2</sub> COCH <sub>3</sub> + RSH → CH <sub>3</sub> COCH <sub>3</sub> + RS·	~4 × 10 <sup>8</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. acetone.	68G132
161.075	·CH <sub>2</sub> OH + RSH → CH <sub>3</sub> OH + RS·	6.8 × 10 <sup>7</sup>	7.6	p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. MeOH; $k \approx 1 \times 10^7$ at pH 12.	68G132
161.076	CH <sub>3</sub> ·CHOH + RSH → CH <sub>3</sub> CH <sub>2</sub> OH + RS·	2.9 × 10 <sup>7</sup> 1.4 × 10 <sup>8</sup>	7	p.r. p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH. P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. EtOH.	72R003 68G132
161.078	(CH <sub>3</sub> ) <sub>2</sub> ·CHOH + RSH → (CH <sub>3</sub> ) <sub>2</sub> CHOH + RS·	4.2 × 10 <sup>8</sup> 2.0 × 10 <sup>8</sup>	7	p.r. p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 2-PrOH. D.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	68G132 72R003
161.079	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> ·CHOH + RSH → CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH + RS·	8.2 × 10 <sup>7</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 1-BuOH.	68G132
161.081	(CH <sub>3</sub> ) <sub>2</sub> CH·CHOH + RSH → (CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> OH + RS·	1.4 × 10 <sup>8</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 2-methyl-1-propanol.	68G132
161.095	·CHOHCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH + RSH → (CH <sub>2</sub> CH <sub>2</sub> OH) <sub>2</sub> + RS·	1.1 × 10 <sup>8</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 1,4-butanediol.	68G132
161.111	R' + RSH → R'H + RS· (R' = Radicals from glucose)	3.2 × 10 <sup>7</sup>		p.r.	P.b.k. at 410 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. glucose.	68A132
162.	<b>Cysteine</b>					
162.075	·CH <sub>2</sub> OH + RSH → CH <sub>3</sub> OH + RS·	0.5[k(CH <sub>2</sub> OH + O <sub>2</sub> )] 4.2 × 10 <sup>7</sup>		γ-r. 7	C.k. with O <sub>2</sub> ; obs. CH <sub>2</sub> O and glycol yields. D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH.	69G524 72R003
162.062	R + Cys → no <i>e</i> -transfer (R = Radicals from glycine anhydride + OH)	< 10 <sup>7</sup>	6,10, 12	p.r. p.r.	No change in d.k. at 265 nm on addn. of cysteine in N <sub>2</sub> O-satd. soln.	71G554
162.142	R + Cys → <i>e</i> -transfer (R = Electron adduct of glycine anhydride)	(2.1 ± 0.4) × 10 <sup>8</sup>	6.4	p.r.	D.k. at 265 nm in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
162.143	R + Cys → <i>e</i> -transfer (R = Electron adduct of alanine anhydride)	(1.4 ± 0.3) × 10 <sup>8</sup>	7.4	p.r.	D.k. at 265 nm in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
162.144	R + Cys → <i>e</i> -transfer (R = Electron adduct of sarcosine anhydride)	(1.5 ± 0.3) × 10 <sup>8</sup>	7.0	p.r.	D.k. at 265 nm in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
163.	<b>Cytochrome c</b>					
163.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + Fe <sup>3+</sup> cyt c			p.r.	No reaction; p.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> <i>tert</i> -BuOH.	78A288
163.062	R + Fe <sup>3+</sup> cyt c (R = Radical from glycine anhydride)	< 10 <sup>7</sup>	6.8	p.r.	P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. glycine anhydride.	78A288
163.075	·CH <sub>2</sub> OH + Fe <sup>3+</sup> cyt c → Fe <sup>2+</sup> cyt c	(3.0 ± 0.5) × 10 <sup>7</sup>	7	p.r.	Abs. change at 550 and 435 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 100% redn.	79A153

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L. mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
163.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{Fe}^{2+}\text{cyt c}$	$2.4 \times 10^8$ $(1.4 \pm 0.2) \times 10^8$  $1.8 \times 10^8$	2.0 7  $\sim 7$	p.r. p.r.  p.r.	 Abs. change at 550 and 435 nm in N <sub>2</sub> O-satd. soln. contg. EtOH; 100% redn. Abs. change at 550 and 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; see also 74A007 and 77A128.	77A128 79A153  79A312
163.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{Fe}^{2+}\text{cyt c}$	$3.8 \times 10^8$ $1.6 \times 10^8$	7 9.3	p.r.	P.b.k. at 550 nm; radical from 2-PrOH.	75A012
163.096	$\cdot\text{CHOHC}(\text{CH}_2\text{OH})_3 + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{Fe}^{2+}\text{cyt c}$	$<10^6$ $1.4 \times 10^8$ $1.6 \times 10^8$	5.6 9.1 9.8	p.r.	P.b.k. at 550 nm; radical from pentaerythritol.	75A012
163.106	$\cdot\text{CHOHCHOHCH}_2\text{OH} + \text{Fe}^{3+}\text{cyt c} \rightarrow (+\text{CH}_2\text{OHCHOHCH}_2\text{OH})$	$2.5 \times 10^6$	7	p.r.	P.b.k. at 550 nm; radicals from glycerol.	75A012
163.113	$\cdot\text{A}^- + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{A} + \text{Fe}^{2+}\text{cyt c}$ ( $\cdot\text{A}^-$ = radical from ascorbate ion)	$6.6 \times 10^3$	7.4	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate	75A240
163.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2^- + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{Fe}^{2+}\text{cyt c}$	$(2.4 \pm 0.2) \times 10^8$  $2.3 \times 10^8$ $2.5 \times 10^8$	  7 11	p.r. p.r.	P.b.k. at 550 nm in soln. contg. lactate. P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> lactate.	74A007 75A012
163.120	$\cdot\text{O}_2\dot{\text{C}}\text{OHCH}_2\text{CO}_2^- + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{Fe}^{2+}\text{cyt c}$	$(8.5 \pm 0.8) \times 10^7$	6.4	p.r.	P.b.k. at 550 nm in malate soln.; $E_a = 3$ kcal mol <sup>-1</sup> (12 kJ mol <sup>-1</sup> ).	74A007
163.121	$\cdot\text{O}_2\dot{\text{C}}\text{OHCHOHCO}_2^- + \text{Fe}^{3+}\text{cyt c}$	$1.7 \times 10^8$	7	p.r.	P.b.k. at 550 nm in tartrate soln.	75A012
163.142	$\text{R} + \text{Fe}^{3+}\text{cyt c}$ (R = Electron adduct of glycine anhydride)	$8 \times 10^8$	6.8	p.r.	P.b.k. at 500 nm in soln. contg. glycine anhydride.	78A288
163.147	$\cdot\text{CO}_2^- + \text{Fe}^{3+}\text{cyt c} \rightarrow \text{CO}_2 + \text{Fe}^{2+}\text{cyt c}$	$\sim 5 \times 10^8$ $\sim 3 \times 10^9$ $6.9 \times 10^8$ $2.5 \times 10^8$ $1.0 \times 10^9$ $6.3 \times 10^8$	$>4$ $<4$ 7 10.8 6.2 8.7	p.r. p.r. p.r.	P.b.k. at 550 nm in soln. contg. 0.03–1 mol L <sup>-1</sup> formate; concn. effect. P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. formate. Abs. change at 450 and 550 nm in 0.1 mol L <sup>-1</sup> formate soln.; at pH 6.2 $E_a = 2.7$ kcal mol <sup>-1</sup> and $A = 1.0 \times 10^{11}$ ; at pH 8.7 $E_a = 3.0$ and $A = 1.0 \times 10^{11}$ .	71G327 75A012 76A127
		$(7 \pm 0.5) \times 10^8$  $7.4 \times 10^8$ $9.4 \times 10^8$ $1.3 \times 10^9$  $1.0 \times 10^8$	7.4  6.7 2.0 7.0  $\sim 7$	p.r. p.r. p.r.	P.b.k. in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> formate; $k = (5.0 \pm 0.6) \times 10^8$ in basic soln.  P.b.k. at 550 nm in 1 atm N <sub>2</sub> O, and 0.01 mol L <sup>-1</sup> formate. D.k. in formate soln.; $I = 0.1$ .	77A096 77A128 78A288 79A312
164.	<b>Cytochrome c, acetylated</b>					
164.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Ac-cyt c}$	$2.5 \times 10^8$	$\sim 7$	p.r.	Abs. change at 550 and 450 nm in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> EtOH soln.; $I = 0.005$ .	79A312
164.147	$\cdot\text{CO}_2^- + \text{Ac-cyt c}$	$1.5 \times 10^9$	$\sim 7$	p.r.	D.k. at 550 nm in formate soln.; $I = 0.1$ .	79A312
165.	<b>Cytochrome c, carboxymethylated</b>					
165.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Cxm-cyt c}$	$3.5 \times 10^8$	$\sim 7$	p.r.	Abs. change at 550 nm and 450 nm in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> EtOH soln.	79A312
165.121	$\cdot\text{O}_2\dot{\text{C}}\text{OHCHOHCO}_2^- + \text{Cxm-cyt c}$	$2.8 \times 10^7$	7	p.r.	Radical from tartrate ion	78A288
165.147	$\cdot\text{CO}_2^- + \text{Cxm-cyt c}$	$1.4 \times 10^8$ $1.3 \times 10^8$	7 $\sim 7$	p.r. p.r.	Radical from formate ion. D.k. at 550 nm in formate soln.; $I = 0.1$ .	78A288 79A312
167.	<b>Cytochrome c, succinylated</b>					
167.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Suc-cyt c}$	$1.8 \times 10^9$	$\sim 7$	p.r.	Abs. change at 550 and 450 nm in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> EtOH soln.	79A312



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
167.147	$\cdot\text{CO}_2^- + \text{Suc-cyt c}$	$4.0 \times 10^9$	$\sim 7$	p.r.	D.k. at 550 nm in formate soln.; $I = 0.1$ .	79A312
168.	<b>Cytochrome c<sub>3</sub></b>					
168.147	$\cdot\text{CO}_2^- + \text{cyt c}_3$	$2.1 \times 10^8$	8.1	p.r.	D.k. in N <sub>2</sub> O-satd. formate soln.	78A232
169.	<b>Cytochrome P450</b>					
169.147	$\cdot\text{CO}_2^- + \text{cyt P450}$				No redn. obs. in N <sub>2</sub> O-satd. soln. contg. formate.	79A036
<i>Deuteroheme See 223.</i>						
<i>Deuterohemin See 222.</i>						
<i>Deuteroporphyrin dimethyl ester See 221a.</i>						
170.	<b>Diamide (Tetramethyldiazenedicarboxamide)</b>					
170.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{D} \rightarrow \text{D}^\cdot + \text{H}^+ + (\text{CH}_3)_2\text{CO}$	$\sim 2.5 \times 10^9$		p.r.	P.b.k. at 400 nm.	75A194
170.147	$\cdot\text{CO}_2^- + \text{D} \rightarrow \text{D}^\cdot + \text{CO}_2$ $\text{D} = (\text{CH}_3)_2\text{NCON}=\text{NCON}(\text{CH}_3)_2$	$\sim 2.5 \times 10^9$		p.r.	P.b.k. at 400 nm.	75A194
171.	<b>1,1'-Dibenzyl-4,4'-bipyridinium ion (Benzyl viologen)</b>					
171.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{BV}^{2+} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{BV}^\cdot+$	$(3.0 \pm 0.1) 10^9$		p.r.	P.b.k. at 400 or 600 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> 2-PrOH.	76A070
171.147	$\cdot\text{CO}_2^- + \text{BV}^{2+} \rightarrow \text{CO}_2 + \text{BV}^\cdot+$	$6.7 \times 10^9$	7.0	p.r.	P.b.k. at 680 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	76A169
		$1.7 \times 10^{10}$	6.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	78A321
172.	<b>1,1'-Di(carboxyethyl)-4,4'-bipyridinium ion</b>					
172.147	$\cdot\text{CO}_2^- + \text{BP}^{2+} \rightarrow \text{CO}_2 + \text{BP}^\cdot+$	$2.0 \times 10^9$	7.0	p.r.	P.b.k. at 680 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	76A169
173.	<b>Dichloroindophenol</b>					
173.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{dye}$	$3.6 \times 10^9$	7	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 87% <i>e</i> -transfer.	73A078
173.075	$\cdot\text{CH}_2\text{OH} + \text{dye}$	$3.2 \times 10^9$	7	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 99% <i>e</i> -transfer.	73A078
173.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{dye}$	$4.4 \times 10^9$	7	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 88% <i>e</i> -transfer.	73A078
173.147	$\cdot\text{CO}_2^- + \text{dye}$	$3.5 \times 10^9$	7	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. formate, as well as p.b.k. at $\sim 400$ nm; 100% <i>e</i> -transfer.	73A078
174.	<b>Dichloromethane</b>					
174.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_2\text{Cl}_2 \rightarrow \cdot\text{CH}_2\text{Cl} + (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{Cl}^-$			$\gamma$ -r.	No redn. in soln. contg. $1.5 \times 10^{-3}$ mol L <sup>-1</sup> Na dodecyl sulfate, 0.2 mol L <sup>-1</sup> 2-PrOH and 0.1 mol L <sup>-1</sup> acetone.	79G191
		$\approx 10^6$		p.r.		80A011
175.	<b>1,1'-Di(4-cyanophenyl)-4,4'-bipyridinium ion</b>					
175.147	$\cdot\text{CO}_2^- + \text{BP}^{2+} \rightarrow \text{CO}_2 + \text{BP}^\cdot+$	$1.4 \times 10^{10}$	6.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	78A321
<i>Diethyl ketone See 3-Pentanone 305.</i>						
176.	<b>1,2-Dihydroxybenzene (Catechol)</b>					
176.020	$\cdot\text{CH}_2\text{CHO} + \text{HOC}_6\text{H}_4\text{O}^- \rightarrow \text{CH}_3\text{CHO} + \cdot\text{OC}_6\text{H}_4\text{O}^\cdot$	$7.4 \times 10^8$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
177.	<b>1,3-Dihydroxybenzene (Resorcinol)</b>					

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
177.020	$\cdot\text{CH}_2\text{CHO} + \text{HOC}_6\text{H}_4\text{O}^- \rightarrow \text{CH}_3\text{CHO} + \cdot\text{C}_6\text{H}_4\text{O}\cdot$	$1.6 \times 10^9$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
<i>1,4-Dihydroxybenzene See Hydroquinone 227.</i>						
178.	<b>1,3-Dihydroxy-2-nitrobenzene</b>					
178.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + (\text{OH})_2\text{C}_6\text{H}_3\text{NO}_2 \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + (\text{OH})_2\text{C}_6\text{H}_3\dot{\text{N}}\text{O}_2^-$	$6.8 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
179.	<b>2-(3,4-Dihydroxyphenyl)ethylamine (Dopamine)</b>					
179.113	$\cdot\text{A}^- + (\text{OH})_2\text{C}_6\text{H}_3\text{CH}_2\text{CH}_2\text{NH}_2 \rightarrow (\cdot\text{A}^- = \text{Radical from ascorbate})$	$(3.6 \pm 0.4) \times 10^2$	8.4	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate.	75A240
180.	<b>2,3-Dimethylbenzoquinone</b>					
180.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{Q}\cdot^-$	$3.5 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
181.	<b>2,5-Dimethylbenzoquinone</b>					
181.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{Q}\cdot^-$	$3.9 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
181.097	$\text{CH}_3\dot{\text{C}}(\text{O}^-)\text{CH}_2\text{CO}_2^- + \text{Q} \rightarrow \text{Q}\cdot^-$	$3.3 \times 10^9$	9.2	p.r.	P.b.k. at 400 nm in Ar-satd. soln. contg. <i>tert</i> -BuOH + acetoacetate ion.	73A104
182.	<b>2,6-Dimethylbenzoquinone</b>					
182.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{Q}\cdot^-$	$4.2 \times 10^9$		p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
183.	<b>1,1'-Dimethyl-4,4'-bipyridinium ion (Methyl viologen)</b>					
183.075	$\cdot\text{CH}_2\text{OH} + \text{MV}^{2+} \rightarrow \text{CH}_2\text{O} + \text{H}^+ + \text{MV}\cdot^+$	$3 \times 10^8$		p.r.	N <sub>2</sub> O-satd. 3% MeOH soln.	77A177
183.147	$\cdot\text{CO}_2^- + \text{MV}^{2+} \rightarrow \text{CO}_2 + \text{MV}\cdot^+$	$1.5 \times 10^{10}$		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	73A074
184.	<b>Dimethyl fumarate</b>					
184.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3\text{O}_2\text{CCH}=\text{CHCO}_2\text{CH}_3$	$4.0 \times 10^9$ (uncor.)	7.0	p.r.	12% <i>e</i> -transfer.	73G097
184.147	$\cdot\text{CO}_2^- + \text{CH}_3\text{O}_2\text{CCH}=\text{CHCO}_2\text{CH}_3$	$9 \times 10^8$ (uncor.)	7.0	p.r.	>80% <i>e</i> -transfer.	73G097
185.	<b>2,3-Dimethylnaphthoquinone</b>					
185.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{Q}\cdot^-$	$3.9 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
185a.	<b>N,N-Dimethyl-4-nitrosoaniline (RNO)</b>					
185a.075	$\cdot\text{CH}_2\text{OH} + \text{RNO}$	$(7.9 \pm 0.3) \times 10^8$	7	p.r.	D.k. at 440 nm in N <sub>2</sub> O-satd. soln., contg. MeOH.	69G156
185a.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{RNO}$	$(2.4 \pm 0.04) \times 10^9$	7	p.r.	D.k. at 440 nm in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> EtOH.	69G156
185a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{RNO}$	$(3.2 \pm 0.05) \times 10^9$	7	p.r.	D.k. at 440 nm in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	69G156
185a.147	$\cdot\text{CO}_2^- + \text{RNO}$	$(1.8 \pm 0.01) \times 10^9$	7	p.r.	D.k. at 440 nm in N <sub>2</sub> O-satd. soln. contg. 0.25 mol L <sup>-1</sup> formate.	69G156
186.	<b>3,5-Dinitroanisole</b>					
186.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{dNA} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{dNA}\cdot^-$	$2.5 \times 10^9$	7	p.r.	P.b.k. at 310 nm in soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; reaction observed as a slower process following fast $e_{\text{aq}}^-$ reaction.	79A176
187.	<b>1,2-Dinitrobenzene</b>					
187.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_4(\text{NO}_2)_2 \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + [\text{C}_6\text{H}_4(\text{NO}_2)_2]\cdot^-$	$2.9 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
188.	<b>1,3-Dinitrobenzene</b>					
188.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_4(\text{NO}_2)_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + [\text{C}_6\text{H}_4(\text{NO}_2)_2]^-$	$3.6 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
189.	<b>1,4-Dinitrobenzene</b>					
189.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_4(\text{NO}_2)_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + [\text{C}_6\text{H}_4(\text{NO}_2)_2]^-$	$3.2 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
190.	<b>2,4-Dinitrobenzoate ion</b>					
190.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ +$ $[{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$2.9 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
190.147	$\cdot\text{CO}_2^- + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $\text{CO}_2 + [{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$1.8 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate.	76A111
191.	<b>2,5-Dinitrobenzoate ion</b>					
191.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ +$ $[{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$3.3 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
191.147	$\cdot\text{CO}_2^- + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $\text{CO}_2 + [{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$1.9 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate.	76A111
192.	<b>3,4-Dinitrobenzoate ion</b>					
192.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ +$ $[{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$3.2 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
192.147	$\cdot\text{CO}_2^- + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $\text{CO}_2 + [{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$1.8 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate.	76A111
193.	<b>3,5-Dinitrobenzoate ion</b>					
193.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ +$ $[{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$3.1 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
193.147	$\cdot\text{CO}_2^- + {}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2 \rightarrow$ $\text{CO}_2 + [{}^-\text{O}_2\text{CC}_6\text{H}_3(\text{NO}_2)_2]^-$	$2.5 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate.	76A111
194.	<b>3,5-Dinitrobenzonitrile</b>					
194.111	$\text{R} + (\text{NO}_2)_2\text{C}_6\text{H}_3\text{CN} \rightarrow$ $[(\text{NO}_2)_2\text{C}_6\text{H}_3\text{CN}]^-$ (R = Radicals from glucose)	$1.0 \times 10^9$		p.r.		77R167
195.	<b>1-(2,4-Dinitrophenyl)pyridinium ion</b>					
195.100	$\text{R} + (\text{NO}_2)_2\text{C}_6\text{H}_4\text{N}^+\text{C}_5\text{H}_5 \rightarrow$ $[(\text{NO}_2)_2\text{C}_6\text{H}_4\text{N}^+\text{C}_5\text{H}_5]^-$ (R = Radicals from glucose)	$1.0 \times 10^9$		p.r.		77R167
195.147	$\cdot\text{CO}_2^- + (\text{NO}_2)_2\text{C}_6\text{H}_4\text{N}^+\text{C}_5\text{H}_5 \rightarrow$ $[(\text{NO}_2)_2\text{C}_6\text{H}_4\text{N}^+\text{C}_5\text{H}_5]^-$	$4 \times 10^8$		p.r.		77R167
196.	<b>1,1'-Diphenyl-4,4'-bipyridinium ion</b>					
196.147	$\cdot\text{CO}_2^- + \text{BP}^{2+} \rightarrow \text{CO}_2 + \text{BP}^{\cdot+}$	$1.3 \times 10^{10}$	6.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate ion.	78A321
197.	<b>Dithiothreitol</b>					
197.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} +$ $\text{HSCH}_2\text{CHOHCHOHCH}_2\text{SH} \rightarrow$ $(\text{CH}_3)_3\text{COH} + \text{RS}\cdot (\rightarrow \text{RSSR}^-)$	$(6.8 \pm 0.6) \times 10^7$	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	73A020
197.075	$\cdot\text{CH}_2\text{OH} + \text{RSH} \rightarrow$ $\text{CH}_3\text{OH} + \text{RS}\cdot$	$(6.8 \pm 0.6) \times 10^7$	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	73A020
197.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{RSH} \rightarrow$ $(\text{CH}_3)_2\text{CHOH} + \text{RS}\cdot$	$(2.1 \pm 0.2) \times 10^8$	7	p.r.	P.b.k. at 390 nm in soln. of 2 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73A020

Dopamine See 2-(3,4-Dihydroxyphenyl)ethylamine 179.



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
198. 198.078	<b>Duroquinone</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{Q}^-$	$4.0 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
199. 199.076	<b>Eosin</b> $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{S} \rightarrow$ $\text{CH}_3\text{CHO} + \cdot\text{S}^- + \text{H}^+$	$(1.1 \pm 0.2) \times 10^9$	8.5– 9.0	p.r.	P.b.k. at 405 nm in soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> EtOH and 10 <sup>-3</sup> mol L <sup>-1</sup> H <sub>2</sub> O <sub>2</sub> .	67G038
199.147	$\cdot\text{CO}_2^- + \text{S} \rightarrow \text{CO}_2 + \cdot\text{S}^- + \text{H}^+$	$(2.5 \pm 0.5) \times 10^8$	8.5– 9.0	p.r.	P.b.k. at 405 nm in soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> HCO <sub>2</sub> <sup>-</sup> and 10 <sup>-3</sup> mol L <sup>-1</sup> H <sub>2</sub> O <sub>2</sub> .	67G038
200. 200.001	<b>Ethanol</b> $\cdot\text{CH}_3 + \text{CH}_3\text{CH}_2\text{OH} \rightarrow$ $\text{CH}_4 + \text{CH}_3\dot{\text{C}}\text{HOH}$	$5.9 \times 10^2$ (rel.)		$\gamma$ -r.	C.k. with O <sub>2</sub> ; radical from MeI; rel. to $k(\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
200.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{C}_2\text{H}_5\text{OH} \rightarrow$ $\text{C}_2\text{H}_5\text{OH} + \text{CH}_3\dot{\text{C}}\text{HOH}$	$16 \pm 3$		$\gamma$ -r.	Calcd. from obs. yields and assumed mechanism in oxid. of EtOH by H <sub>2</sub> O <sub>2</sub> .	70G338
200.047	$\cdot\text{CF}_3 + \text{C}_2\text{H}_5\text{OH} \rightarrow$ $\text{CF}_3\text{H} + \text{C}_2\text{H}_4\text{OH}$	$(4.6 \pm 0.5) \times 10^4$ (rel.)	9–10	$\gamma$ -r.	C.k. with $\cdot\text{CF}_3$ addn. to propylene; obs. CF <sub>3</sub> H yield; rel. to $k(\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ .	70G407
201. 201.001	<b>Ethyl acetate</b> $\cdot\text{CH}_3 + \text{CH}_3\text{CO}_2\text{Et} \rightarrow$ $\text{CH}_4 + \text{CH}_3\text{CO}_2\dot{\text{C}}\text{HCH}_3$	$<1.7 \times 10^3$	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
202. 202.001	<b>Ethylene</b> $\cdot\text{CH}_3 + \text{CH}_2=\text{CH}_2 \rightarrow \cdot\text{CH}_2\text{CH}_2\text{CH}_3$	$4.9 \times 10^3$ (rel.)		$\gamma$ -r.	C.k. with 2-PrOH in soln. contg. MeI; obs. $G(\text{CH}_4)$ ; rel. to $k(\cdot\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
202.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{CH}_2=\text{CH}_2 \rightarrow$ $\cdot\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	$(1.6 \pm 0.5) \times 10^6$ (rel.) $(4.1 \pm 1.2) \times 10^6$ (rel.)  $(3 \pm 2) \times 10^4$ (rel.)	4.5  2	p.r.  $\gamma$ -r.	Radical from OH + ethylene; contains ~10% (pH 4.5) and ~50% (pH 2) $\cdot\text{C}_2\text{H}_5$ ; obs. rate of addn. of Cu <sup>+</sup> to ethylene (Cu <sup>+</sup> from OH + Cu <sup>2+</sup> in N <sub>2</sub> O-satd. soln.); rel. to $k(\text{R} + \text{Cu}^{2+}) = 1.9 \times 10^7$ (pH 4.5) and $2.2 \times 10^7$ (pH 2). Calcd. from dose rate effect on yields of $\alpha,\omega$ -diols in N <sub>2</sub> O-satd. soln. contg. ethylene assuming termination rate constant of chain reaction is $5 \times 10^8$ .	78A322  80A054
202.046	$\cdot\text{CFCl}_2 + \text{CH}_2=\text{CH}_2 \rightarrow \text{CFCl}_2\text{CH}_2\dot{\text{C}}\text{H}_2$	$\sim 4 \times 10^7$	5.9	p.r.	Est. from condy meas. in soln. contg. CFCl <sub>3</sub> .	71G026
202.047	$\cdot\text{CF}_3 + \text{CH}_2=\text{CH}_2 \rightarrow \cdot\text{CH}_2\text{CH}_2\text{CF}_3$	$(4.0 \pm 0.6) \times 10^7$ (rel.)  $\sim 7 \times 10^8$	9–10  5.9	$\gamma$ -r.  p.r.	C.k. with 2-PrOH (H abstr. by CF <sub>3</sub> ); obs. CF <sub>3</sub> H yield; rel. to $k(\cdot\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ . Est. from condy. meas. in soln. contg. CF <sub>3</sub> Cl.	70G407  71G026
203. 203.147	<b>1,1'-Ethylene-2,2'-bipyridinium ion</b> $\cdot\text{CO}_2^- + \text{BP}^{2+} \rightarrow \text{CO}_2 + \cdot\text{BP}^+$	$4.0 \times 10^9$  $1.2 \times 10^{10}$	7.0  6.8	p.r.  p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate ion. P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate ion.	76A169  78A321
204. 204.075	<b>Ethylene glycol</b> $\cdot\text{CH}_2\text{OH} + \text{HOCH}_2\text{CH}_2\text{OH} \rightarrow$ $\text{CH}_3\text{OH} + \text{HO}\dot{\text{C}}\text{HCH}_2\text{OH}$	$\sim 10^3$ – $10^4$		$\gamma$ -r.	Est. from dose effect on yields in MeOH soln.	71G929

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
205.	<b>Ethyl ether</b>					
205.048	$\cdot\text{CCl}_3 + \text{CH}_3\text{CH}_2\text{OEt} \rightarrow \text{CHCl}_3 + \text{CH}_3\dot{\text{C}}\text{HOEt}$	35		$\gamma$ -r.	Calcd. from dependence of $G(\text{Cl}^-)$ on dose rate in soln. contg. $2 \times 10^{-3}$ mol L <sup>-1</sup> $\text{CCl}_4$ and 0.5 mol L <sup>-1</sup> ethyl ether; assume $2k(\text{R} + \text{R}) = 10^9$ .	71G778
206.	<b>N-Ethylmaleimide (NEM)</b>					
206.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{NEM} \rightarrow \cdot\text{NEM-R}$			p.r.	No $e$ -transfer; very low addn. rate; radical from <i>tert</i> -BuOH.	72G144
206.053	$\text{CH}_3\text{CONH}\dot{\text{C}}\text{H}_2 + \text{NEM} \rightarrow \cdot\text{NEM-R}$	$\sim 10^8$	6-7	p.r.	100% addn. based on abs. spectra; radical from <i>N</i> -methylacetamide.	72G144
206.075	$\cdot\text{CH}_2\text{OH} + \text{NEM} \rightarrow \cdot\text{NEM}^- + \cdot\text{NEM-CH}_2\text{OH}$	$2.4 \times 10^9$ (uncor.)	6-7	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. MeOH; 15% $e$ -transfer, 85% addn. based on abs. spectra.	72G144
206.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NEM} \rightarrow \cdot\text{NEM}^- + \cdot\text{NEM-R}$	$5.0 \times 10^9$ (uncor.)	6-7	p.r.	47% $e$ -transfer, 50% addn. based on abs. spectra; radical from 2-PrOH.	72G144
206.109	$\text{R} + \text{NEM}$ (R = Radicals from ribose)	$2.1 \times 10^9$ (uncor.)	6-7	p.r.	30% $e$ -transfer, 20% addn. based on abs. spectra.	72G144
206.141	$\text{R} + \text{NEM} \rightarrow \cdot\text{NEM}^-$ (R = Electron adduct of dimethyl fumarate)	$2.2 \times 10^9$		p.r.	D.k.	73G097
206.147	$\cdot\text{CO}_2^- + \text{NEM} \rightarrow \text{CO}_2 + \cdot\text{NEM}^-$	$5.4 \times 10^9$	6-7	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. formate; 100% $e$ -transfer based on abs. spectra.	72G144
207.	<b>4-Ethylphenoxide ion</b>					
207.020	$\cdot\text{CH}_2\text{CHO} + \text{C}_2\text{H}_5\text{C}_6\text{H}_4\text{O}^- (+ \text{H}_2\text{O}) \rightarrow \text{CH}_3\text{CHO} + \text{OH}^- + \text{C}_2\text{H}_5\text{C}_6\text{H}_4\text{O}\cdot$	$7.0 \times 10^7$	$\sim 11.5$	p.r.	Soln. contg. $\text{N}_2\text{O}$ and ethylene glycol.	79A051
208.	<b>Ferredoxin</b>					
208.147	$\cdot\text{CO}_2^- + \text{ferredoxin}$	$(8.0 \pm 0.7) \times 10^7$	7.5	p.r.	D.k. at 420 nm in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	73A064
<i>Flagyl See Metronidazole 271.</i>						
<i>Flavin mononucleotide (FMN) See Riboflavin 5'-phosphate 334.</i>						
209.	<b>Fluorescein</b>					
209.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Dye} \rightarrow \text{semiquinone}$	$4.5 \times 10^8$	10.8	p.r.	D.k. at 500 nm in $\text{N}_2\text{O}$ -satd. soln. contg. EtOH.	73A078
209.147	$\cdot\text{CO}_2^- + \text{Dye} \rightarrow \text{semiquinone}$	$(2.6 \pm 0.9) \times 10^7$	10.4	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> formate.	68G172
210.	<b>Folic acid</b>					
210.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{FH}_2^+$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{FH} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{FH}^-$	$1.1 \times 10^9$ $4.0 \times 10^8$	$\sim 0.5$ 6.0	p.r.	P.b.k.; structure and abs. spectra of reduced species dependent on pH.	76A060
211.	<b>Formaldehyde</b>					
211.075	$\cdot\text{CH}_2\text{OH} + \text{CH}_2(\text{OH})_2 \rightarrow \text{CH}_3\text{OH} + \cdot\text{CH}(\text{OH})_2$	$\sim 10^4$ - $10^5$		$\gamma$ -r.	Est. from dose effects on yields in soln. contg. MeOH.	71G929
212.	<b>Formate ion</b>					
212.047	$\cdot\text{CF}_3 + \text{HCO}_2^- \rightarrow \text{CF}_3\text{H} + \cdot\text{CO}_2^-$	$(3.4 \pm 0.7) \times 10^5$	9-10	p.r.	P.b.k. at 270 nm ( $\text{CO}_2^-$ ); radical from $\text{CF}_3\text{Cl} + e_{\text{aq}}^-$ .	70G407
213.	<b>N-Formylkynurenine</b>					
213.147	$\cdot\text{CO}_2^- + \text{FK} (+ \text{H}^+) \rightarrow \text{CO}_2 + \cdot\text{FKH}$	$> 3 \times 10^7$	7.6	p.r.	P.b.k. (semiquinone) in $\text{N}_2\text{O}$ -satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	75F361

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
214.	<b>Fumarate ion</b>					
214.047	$\cdot\text{CF}_3 + \cdot\text{O}_2\text{CCH}=\text{CHCO}_2^- \rightarrow$ $\cdot\text{O}_2\dot{\text{C}}\text{CHCH}(\text{CF}_3)\text{CO}_2^-$	$\sim 10^8$		<i>e-r.</i>	Est. from esr.	71G284
214.075	$\cdot\text{CH}_2\text{OH} + \cdot\text{O}_2\text{CCH}=\text{CHCO}_2^- \rightarrow$ $\cdot\text{O}_2\dot{\text{C}}\text{CHCH}(\text{CH}_2\text{OH})\text{CO}_2^-$	$\sim 10^7$		<i>e-r.</i>	Est. by esr from dose rate effect on $\cdot\text{CH}_2\text{OH}$ and adduct radical (in 1 mol L <sup>-1</sup> MeOH and 0.01 mol L <sup>-1</sup> fumarate) assuming $2k(\text{R} + \text{R}) = 2.4 \times 10^9$ ; other radicals adding to fumarate ion were $\cdot\text{CH}_3$ , $\cdot\text{CHOHCH}_3$ , $(\text{CH}_3)_2\dot{\text{C}}\text{OH}$ , $\cdot\text{CHOHCH}_2\text{OH}$ , $\cdot\text{CH}_2\text{NH}_2$ , $\cdot\text{CH}_2\text{Cl}$ , $\cdot\text{CHClCO}_2^-$ , $\cdot\text{CO}_2^-$ , $\cdot\text{CF}_3$ , and $\cdot\text{CH}(\text{CO}_2^-)\text{CH}_2\text{CO}_2^-$ .	71G284
214.113	$\cdot\text{A}^- + \cdot\text{O}_2\text{CCH}=\text{CHCO}_2^-$ ( $\cdot\text{A}^-$ = ascorbate radical)	$< 10$	8.7	<i>p.r.</i>	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate ion.	75A240
214.147	$\cdot\text{CO}_2^- + \cdot\text{O}_2\text{CCH}=\text{CHCO}_2\text{H}$	$2.0 \times 10^7$ (uncor.)	4.0	<i>p.r.</i>	$> 30\%$ <i>e</i> -transfer; no <i>e</i> -transfer at pH 10.0.	73G097
215.	<b>Fumaric acid</b>					
215.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HO}_2\text{CCH}=\text{CHCO}_2\text{H}$	$9.0 \times 10^8$ (uncor.)	0.5	<i>p.r.</i>	14% <i>e</i> -transfer.	73G097
215a.	<b>Glutathione</b>					
215a.142	$\text{R} + \text{GSH} \rightarrow e\text{-transfer}$ ( $\text{R}$ = Electron adduct of glycine anhydride)	$(1.8 \pm 0.3) \times 10^8$	6.4, 7.4	<i>p.r.</i>	D.k. at 265 nm in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
216.	<b>Glutathione disulfide</b>					
216.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{GSSG}$	$\leq 10^7$	7	<i>p.r.</i>	No 420 nm abs. ( $\text{RSSR}^-$ ) obs. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	72G388
216.142	$\text{R} + \text{GSSG} \rightarrow (\text{GSSG})^-$ ( $\text{R}$ = Electron adduct of glycine anhydride)	$(4.0 \pm 0.4) \times 10^7$	6.2	<i>p.r.</i>	D.k. at 265 nm in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
216.147	$\cdot\text{CO}_2^- + \text{GSSG}$	$\leq 10^7$		<i>p.r.</i>	No 420 nm abs. ( $\text{RSSR}^-$ ) obs. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	72G388
217.	<b>Glycine</b>					
217.001	$\cdot\text{CH}_3 + \text{H}_2\text{NCH}_2\text{CO}_2^- \rightarrow$ $\text{CH}_4 + \text{H}_2\text{N}\dot{\text{C}}\text{HCO}_2^-$ $\cdot\text{CH}_3 + \text{H}_3\text{N}^+\text{CH}_2\text{CO}_2^- \rightarrow$ $\sim 4$ (rel.)	$\sim 1.2 \times 10^2$ (rel.) $\sim 4$ (rel.)	$\sim 10$	phot.	$\cdot\text{CH}_3$ from cumene hydroperoxide; obs. formn. of CH <sub>4</sub> and C <sub>2</sub> H <sub>6</sub> ; assume $2k(\text{R} + \text{R}) = 2 \times 10^9$ ; $k_{\text{H}}/k_{\text{D}} = 10.5$ at pH 11.5 [74F528].	70F280
218.	<b>Glycine anhydride</b>					
218.147	$\cdot\text{CO}_2^- + \cdot\text{NHCH}_2\text{CONHCH}_2\text{C}(\text{O})-$	$< 10^7$	6.2	<i>p.r.</i>	No <i>e</i> -transfer obs. in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	71G554
219.	<b>Glycolic acid</b>					
219.001	$\cdot\text{CH}_3 + \text{HOCH}_2\text{CO}_2\text{H} \rightarrow$ $\text{CH}_4 + \text{HO}\dot{\text{C}}\text{HCO}_2\text{H}$	$3.6 \times 10^3$	$\sim 1$	chem.	Radical from dimethylsulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
220.	<b>Hematoporphyrin</b>					
220.075	$\cdot\text{CH}_2\text{OH} + \text{PH}_2 \rightarrow \cdot\text{PH}_2^-$ $\cdot\text{CH}_2\text{O}^- + \text{PH}_2 \rightarrow \cdot\text{PH}_2^-$	$< 1 \times 10^7$ $(3.3 \pm 0.5) \times 10^8$	7.0 13.0	<i>p.r.</i>	P.b.k. at 600–650 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	74A040
220.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{PH}_2 \rightarrow \cdot\text{PH}_2^-$	$(7.0 \pm 1.0) \times 10^8$	13	<i>p.r.</i>	P.b.k. at 600–650 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	74A040
220.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PH}_2 \rightarrow \cdot\text{PH}_2^-$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{PH}_2 \rightarrow \cdot\text{PH}_2^-$	$(2.4 \pm 0.4) \times 10^8$ $(1.1 \pm 0.2) \times 10^9$	7.0 13.0	<i>p.r.</i>	P.b.k. at 600–650 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	74A040
220.147	$\cdot\text{CO}_2^- + \text{PH}_2 \rightarrow \cdot\text{PH}_2^-$	$(4 \pm 1) \times 10^7$	13.0	<i>p.r.</i>	P.b.k. at 600–650 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	74A040



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
221.	<b>Zinc hematoporphyrin</b>					
221.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{ZnP} \rightarrow (\text{CH}_3)_2\text{CO} + \text{ZnP}^-$	$(1.0 \pm 0.1) \times 10^9$	13.0	p.r.	P.b.k. at 600–650 nm in N <sub>2</sub> O–satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	74A040
221a.	<b>Deuteroporphyrin dimethyl ester</b>					
221a.001	$\cdot\text{CH}_3 + \text{DP}$	$(1 \text{ to } 2) \times 10^7$	~13	p.r.	Est. from abs. change in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH, 0.7 mol L <sup>-1</sup> acetone and $2.3 \times 10^{-2}$ mol L <sup>-1</sup> CH <sub>3</sub> I; cor. for $\cdot\text{CH}_3 + \cdot\text{CH}_3$ .	81A002
221a.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{DP} \rightarrow \text{DP}^-$	$(6 \pm 1) \times 10^8$	~13	p.r.	Abs. change in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH and 0.7 mol L <sup>-1</sup> acetone.	81A002
<i>Deuteroporphyrin (ferri) See Deuterohemin 222.</i>						
<i>Deuteroporphyrin (ferro) See Deuteroheme 223.</i>						
222.	<b>Deuterohemin</b>					
222.001	$\cdot\text{CH}_3 + \text{DH}(\text{Fe}^{\text{III}}) \rightarrow \text{addn.}$	$(2.3 \pm 0.5) \times 10^9$ $\sim 1.3 \times 10^9$	~13 7	p.r.	Abs. change in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH and $4.1 \times 10^{-3}$ mol L <sup>-1</sup> methyl iodide.	81A002
222.050	$\cdot\text{CCl}_3 + \text{DH}(\text{Fe}^{\text{III}})$	$\leq 10^6$	7.2	p.r.	No spectral change in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH, $5 \times 10^{-3}$ mol L <sup>-1</sup> CCl <sub>4</sub> and $10^{-4}$ mol L <sup>-1</sup> deuterohemin.	80A011
222.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{DH}(\text{Fe}^{\text{III}}) \rightarrow \text{DH}(\text{Fe}^{\text{II}})$	$3.7 \times 10^8$	7.2	p.r.	D.k. in Ar–satd. soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH and $3.4 \times 10^{-2}$ mol L <sup>-1</sup> acetone.	80A011
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{DH}(\text{Fe}^{\text{III}}) \rightarrow \text{DH}(\text{Fe}^{\text{II}})$	$(9 \pm 1) \times 10^8$	~13	p.r.	D.k. in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH.	81A002
223.	<b>Deuteroheme</b>					
223.001	$\cdot\text{CH}_3 + \text{DH}(\text{Fe}^{\text{II}}) \rightarrow \text{addn.}$	$(3.9 \pm 0.5) \times 10^9$	7	p.r.	Abs. change in 6.5 mol L <sup>-1</sup> 2-PrOH soln. of deuterohemin chemically reduced by dithionite to which methyl chloride was added.	81A002
223.043	$\cdot\text{CH}_2\text{Cl} + \text{DH}(\text{Fe}^{\text{II}})$	$2 \times 10^9$	7.2	p.r.	Abs. change in soln. of deuterohemin chemically reduced by dithionite to which $4 \times 10^{-2}$ mol L <sup>-1</sup> CH <sub>2</sub> Cl <sub>2</sub> was added.	80A011
223.048	$\cdot\text{CCl}_3 + \text{DH}(\text{Fe}^{\text{II}})$	$(2 \pm 1) \times 10^9$	7.2	p.r.	Abs. change in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH, $3.2 \times 10^{-2}$ mol L <sup>-1</sup> acetone, CCl <sub>4</sub> , and deuterohemin, the latter reduced by (CH <sub>3</sub> ) <sub>2</sub> COH.	80A011
223.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{DH}(\text{Fe}^{\text{II}})$	$(6.3 \pm 0.5) \times 10^8$	7	p.r.	Abs. change in 6.5 mol L <sup>-1</sup> 2-PrOH soln. of deuterohemin chemically reduced by dithionite to which 0.7 mol L <sup>-1</sup> acetone was added.	81A002
224.	<b>Hemin (Protoferriheme IX)</b>					
224.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{H}(\text{Fe}^{\text{III}}) \rightarrow \text{H}(\text{Fe}^{\text{II}})$	$(9.0 \pm 1.0) \times 10^8$	13	p.r.	Soln. 30% EtOH, N <sub>2</sub> O–satd.	74A040
	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{H}(\text{Fe}^{\text{III}}) \rightarrow \text{H}(\text{Fe}^{\text{II}})$	$7.7 \times 10^8$	7	p.r.	Redn. obs. at 560 nm in N <sub>2</sub> O–satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	77A128
		$(5.6 \pm 0.6) \times 10^8$ $(1.6 \pm 0.6) \times 10^9$	9.2 4.0	p.r.	D.k. at 490 nm in N <sub>2</sub> O–satd. soln. contg. 1 mol L <sup>-1</sup> EtOH and 0.2 mol L <sup>-1</sup> Na dodecylsulfate (micelles).	78A033
225.	<b>Hemin c</b>					
225.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}(\text{Fe}^{\text{III}}) \rightarrow \text{H}(\text{Fe}^{\text{II}})$	$(2.8 \pm 0.6) \times 10^9$	7.0	p.r.	P.b.k. at 413 nm in N <sub>2</sub> O–satd. soln. contg. 0.04 mol L <sup>-1</sup> 2-PrOH.	75A241
225.096	$\cdot\text{CHOHC}(\text{CH}_2\text{OH})_3 + \text{H}(\text{Fe}^{\text{III}}) \rightarrow \text{H}(\text{Fe}^{\text{II}})$	$(3.0 \pm 0.6) \times 10^8$ $(2.8 \pm 0.6) \times 10^8$	7.0 11.8	p.r.	P.b.k. at 413 nm in N <sub>2</sub> O–satd. soln. contg. 0.25 mol L <sup>-1</sup> pentaerythritol.	75A241

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
225.106	CH <sub>2</sub> OH $\dot{\text{C}}$ OHCH <sub>2</sub> OH + H(Fe <sup>III</sup> ) $\rightarrow$ H(Fe <sup>II</sup> ) (+ CH <sub>2</sub> OHCHOH $\dot{\text{C}}$ OH)	(1.3 $\pm$ 0.3) $\times 10^9$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> glycerol.	75A241
225.117	CH <sub>3</sub> $\dot{\text{C}}$ OHCO <sub>2</sub> <sup>-</sup> + H(Fe <sup>III</sup> ) $\rightarrow$ H(Fe <sup>II</sup> )	(5.6 $\pm$ 1.1) $\times 10^8$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.025 mol L <sup>-1</sup> lactate.	75A241
225.121	O <sub>2</sub> CCHOH $\dot{\text{C}}$ OHCO <sub>2</sub> <sup>-</sup> + H(Fe <sup>III</sup> ) $\rightarrow$ H(Fe <sup>II</sup> )	(8.1 $\pm$ 1.6) $\times 10^7$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.02 mol L <sup>-1</sup> tartrate.	75A241
225.147	$\cdot\text{CO}_2^-$ + H(Fe <sup>III</sup> ) $\rightarrow$ H(Fe <sup>II</sup> )	(1.3 $\pm$ 0.3) $\times 10^9$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.02 mol L <sup>-1</sup> formate.	75A241
226.	<b>Hemoglobin</b>					
226.075	$\cdot\text{CH}_2\text{OH}$ + H(Fe <sup>III</sup> ) $\rightarrow$ H(Fe <sup>II</sup> )	(9.5 $\pm$ 1.5) $\times 10^6$	7	p.r.	Abs. change at 550 and 435 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 100% redn.	79A153
226.076	CH <sub>3</sub> $\dot{\text{C}}$ HOH + H(Fe <sup>III</sup> ) $\rightarrow$ H(Fe <sup>II</sup> )	(4.0 $\pm$ 0.4) $\times 10^7$	7	p.r.	Abs. change at 550 and 435 nm in N <sub>2</sub> O-satd. soln. contg. EtOH; 100% redn.	79A153
227.	<b>Hydroquinone</b>					
227.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$ + HOC <sub>6</sub> H <sub>4</sub> OH $\rightarrow$ (CH <sub>3</sub> ) <sub>3</sub> COH + $\cdot\text{OC}_6\text{H}_4\cdot$	$\leq 2 \times 10^6$	11.5	p.r.	Semiquinone not formed in <i>tert</i> -BuOH soln.	79A051
227.020	$\cdot\text{CH}_2\text{CHO}$ + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> $\rightarrow$ CH <sub>3</sub> CHO + $\cdot\text{OC}_6\text{H}_4\cdot$	(2.2 $\pm$ 0.1) $\times 10^9$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol or 2-methoxy- or 2-ethoxy-ethanol.	79A051
227.021	$\cdot\text{CH}_2\text{CHO}$ + HOC <sub>6</sub> H <sub>4</sub> OH	$\leq 2 \times 10^6$	7.2	p.r.	Radical from 2-chloroethanol.	79A051
227.023	CH <sub>3</sub> $\dot{\text{C}}$ HCHO + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(1.2 $\pm$ 0.1) $\times 10^9$	$\sim 11.5$	p.r.	P.b.k. at 430 nm (semiquinone) in N <sub>2</sub> O-satd. soln. contg. 1,2-propanediol; radical mixture.	79A051
227.024	CH <sub>3</sub> CO $\dot{\text{C}}$ HCH <sub>3</sub> + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(5.6 $\pm$ 0.5) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 2,3-butanediol.	79A051
227.025	-(CH <sub>2</sub> ) <sub>4</sub> $\dot{\text{C}}$ HCO- + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(5.5 $\pm$ 1.9) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 1,2-cyclohexanediol.	79A051
227.026	HOCH <sub>2</sub> $\dot{\text{C}}$ HCHO + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> (and HOCH <sub>2</sub> CO $\dot{\text{C}}$ H <sub>2</sub> )	(1.3 $\pm$ 0.1) $\times 10^9$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 2,3-epoxypropanol; radical also obtained from glycerol, $k = (1.5 \pm 0.1) \times 10^9$ and glycerol 2-phosphate, $k = (1.7 \pm 0.2) \times 10^9$ .	79A051
227.027	HOCH <sub>2</sub> CH <sub>2</sub> CO $\dot{\text{C}}$ H <sub>2</sub> + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(6.2 $\pm$ 0.8) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 2-hydroxyfuran.	79A051
227.028	HO(CH <sub>2</sub> ) <sub>3</sub> $\dot{\text{C}}$ HCHO + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(8.6 $\pm$ 1.2) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 2-(hydroxymethyl)furan.	79A051
227.029	HO(CH <sub>2</sub> ) <sub>4</sub> $\dot{\text{C}}$ HCHO + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(5.2 $\pm$ 0.9) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. 2-(hydroxymethyl)pyran.	79A051
227.030	R + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> (R = Radicals from erythritol)	(1.3 $\pm$ 0.3) $\times 10^9$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. <i>meso</i> -erythritol.	79A051
227.031	R + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> (R = Radicals from xylitol)	(1.2 $\pm$ 0.1) $\times 10^9$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. xylitol.	79A051
227.032	-(CHOH) <sub>3</sub> CH <sub>2</sub> $\dot{\text{C}}$ HCO- + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup>	(6.4 $\pm$ 1.1) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. <i>meso</i> -inositol.	79A051
227.033	R + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> (R = Radicals from sorbitol)	(9.8 $\pm$ 1.3) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. sorbitol.	79A051
227.109	R + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> (R = Radicals from ribose)	(9.6 $\pm$ 1) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. ribose.	79A051
227.111	R + HOC <sub>6</sub> H <sub>4</sub> O <sup>-</sup> (R = Radicals from glucose)	(7.1 $\pm$ 0.9) $\times 10^8$	$\sim 11.5$	p.r.	P.b.k. at 430 nm in N <sub>2</sub> O-satd. soln. contg. glucose.	79A051

1-(2-Hydroxyethyl)-2-methyl-5-nitroimidazole See Metronidazole 271.

1-(2-Hydroxy-3-methoxypropyl)-2-nitroimidazole See Misonidazole 272.

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
228.	<b>2-Hydroxy-1,4-naphthoquinone</b>					
228.042	$R + Q \rightarrow \cdot Q^-$ (R = Radicals from triglycine)	$1.9 \times 10^9$	10	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. triglycine.	73A104
228.056	$NH_2\dot{C}HCO_2^- + Q \rightarrow \cdot Q^-$	$(3.1 \pm 0.3) \times 10^9$	8.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine.	73A104
228.078	$(CH_3)_2\dot{C}OH + Q \rightarrow \cdot Q^-$	$(3.4 \pm 0.3) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73A104
228.097	$CH_3\dot{C}(O^-)CH_2CO_2^- + Q \rightarrow \cdot Q^-$	$(1.5 \pm 0.2) \times 10^9$	9.2	p.r.	P.b.k. at 400 nm in Ar-satd. soln. contg. acetate ion and <i>tert</i> -BuOH.	73A104
228.115	$\cdot CHOHCO_2^- + Q \rightarrow \cdot Q^-$	$(9.1 \pm 0.9) \times 10^8$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycolate.	73A104
228.147	$\cdot CO_2^- + Q \rightarrow \cdot Q^-$	$(1.95 \pm 0.2) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. formate.	73A104
229.	<b>3-Hydroxy-2-nitrobenzoate ion</b>					
229.078	$(CH_3)_2\dot{C}OH + ^-O_2CC_6H_3(OH)NO_2 \rightarrow$ $(CH_3)_2\dot{C}O + H^+ + ^-O_2CC_6H_3(OH)\dot{N}O_2^-$	$3.3 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
230.	<b>3-Hydroxypropionitrile</b>					
230.001	$\cdot CH_3 + CNCH_2CH_2OH \rightarrow$ $CH_4 + CNCH_2\dot{C}HOH$	$< 1.6 \times 10^3$	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est from esr meas. and values for competing reactions.	75D188
<i>4-Hydroxy-2,2,6,6-tetramethyl-1-piperidinyloxy See Tetramethylhydroxypiperidinyloxy (TMPN) 343.</i>						
231.	<b>Indigo disulfonate</b>					
231.056	$NH_2\dot{C}HCO_2^- + Dye$	$2.8 \times 10^9$	9	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 71% <i>e</i> -transfer.	73A078
231.075	$\cdot CH_2OH + Dye$	$2.0 \times 10^9$ $1.9 \times 10^9$	7 9	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; <i>e</i> -transfer 75% at pH 7 and 62% at pH 9.	73A078
231.078	$(CH_3)_2\dot{C}OH + Dye$	$4.0 \times 10^9$	7.9	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 88% <i>e</i> -transfer.	73A078
231.147	$\cdot CO_2^- + Dye \rightarrow$ $CO_2 + Dye^-$	$2.0 \times 10^9$ $2.1 \times 10^9$	7.0 7.0	p.r. p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. formate. P.b.k. at 400 nm.	73A078 73A078
232.	<b>Indigo tetrasulfonate</b>					
232.056	$NH_2\dot{C}HCO_2^- + Dye$	$2.6 \times 10^9$	7.0	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 78% <i>e</i> -transfer.	73A078
232.075	$\cdot CH_2OH + Dye$	$3.0 \times 10^9$	7	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. ssoln. contg. MeOH; 80% <i>e</i> -transfer.	73A078
232.078	$(CH_3)_2\dot{C}OH + Dye$	$4.2 \times 10^9$	7	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 89% <i>e</i> -transfer.	73A078
233.	<b>Indophenol</b>					
233.075	$\cdot CH_2OH + Dye$	$3.1 \times 10^9$	9	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 92% <i>e</i> -transfer.	73A078
233.078	$(CH_3)_2\dot{C}OH + Dye$	$4.0 \times 10^9$	9	p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 86% <i>e</i> -transfer.	73A078
233.147	$\cdot CO_2^- + Dye \rightarrow$ $CO_2 + Dye^-$	$2.8 \times 10^9$ $2.7 \times 10^9$	9.0 9.0	p.r. p.r.	D.k. at 610 nm in N <sub>2</sub> O-satd. soln. contg. formate. P.b.k. at ~400 nm.	73A078 73A078
233a.	<b>Zinc(II) Insulin</b>					
233a.147	$\cdot CO_2^- + Zn(II)Insulin$	$6 \times 10^8$	9.0	p.r.	P.b.k. at 420 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> formate; based on monomer concn.	80A204



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
234. 234.078	<b>Iodoacetamide</b> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + ICH <sub>2</sub> CONH <sub>2</sub>	4 × 10 <sup>8</sup> (rel.)		p.r.	C.k. with <i>p</i> -nitroacetophenone in soln. contg. 2-PrOH and acetone; rel. to $k(R + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
235. 235.078	<b>Iodoacetate ion</b> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + ICH <sub>2</sub> CO <sub>2</sub> <sup>-</sup>	7 × 10 <sup>7</sup> (rel.)		p.r.	C.k. with <i>p</i> -nitroacetophenone in soln. contg. 2-PrOH and acetone; rel. to $k(R + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
236. 236.016	<b>Iodoacetic acid</b> ·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ICH <sub>2</sub> CO <sub>2</sub> H → ICH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ·CH <sub>2</sub> CO <sub>2</sub> H	(1.3 ± 0.3) × 10 <sup>7</sup> (rel.)	1	chem.	Est. from esr meas. in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln. contg. <i>tert</i> -BuOH; rel. to $2k(\cdot\text{CH}_2\text{CO}_2\text{H} + \cdot\text{CH}_2\text{CO}_2\text{H}) = 1.8 \times 10^9$ .	74D286
236.075	·CH <sub>2</sub> OH + ICH <sub>2</sub> CO <sub>2</sub> H → ·CH <sub>2</sub> CO <sub>2</sub> H + prod.	(2.1 ± 0.7) × 10 <sup>8</sup> (rel.)  < 3.5 × 10 <sup>8</sup>	1	chem.  p.r.	Est. from esr meas. in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln. contg. MeOH; rel. to $2k(\cdot\text{CH}_2\text{CO}_2\text{H} + \cdot\text{CH}_2\text{CO}_2\text{H}) = 1.8 \times 10^9$ .  P.b.k. at 330 nm.	74D286  74D286
237. 237.016	<b>Iodoacetonitrile</b> ·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ICH <sub>2</sub> CN → ICH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ·CH <sub>2</sub> CN	2.0 × 10 <sup>7</sup> (rel.)	1	chem.	Est. from esr meas. in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln. contg. <i>tert</i> -BuOH; rel. to $2k(\cdot\text{CH}_2\text{CN} + \cdot\text{CH}_2\text{CN}) = 2 \times 10^9$ .	74D286
238. 238.016	<b>Iodoethanol</b> ·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ICH <sub>2</sub> CH <sub>2</sub> OH → ICH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ·CH <sub>2</sub> CH <sub>2</sub> OH	1.5 × 10 <sup>5</sup> (rel.)	1	chem.	Est. from esr meas in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln. contg. <i>tert</i> -BuOH; rel. to $2k(\cdot\text{CH}_2\text{CH}_2\text{OH} + \cdot\text{CH}_2\text{CH}_2\text{OH}) = 1.9 \times 10^9$ .	74D286
238a. 238a.078	<b>Iodomethane</b> (CH <sub>3</sub> ) <sub>2</sub> ĊO <sup>-</sup> + CH <sub>3</sub> I → ·CH <sub>3</sub> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + CH <sub>3</sub> I	(1.1 ± 0.1) × 10 <sup>8</sup> < 10 <sup>5</sup>	~13 7	p.r.	C.k. with PNAP in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH and 0.68 mol L <sup>-1</sup> acetone assuming $k = 2.2 \times 10^9$ (279.078).	81A002
239. 239.016	<b>3-Iodopropionic acid</b> ·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ICH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H → ICH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + ·CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	1.8 × 10 <sup>5</sup> (rel.)	1	chem.	Est. from esr meas. in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln. contg. <i>tert</i> -BuOH; rel. to $2k(\cdot\text{CH}_2\text{CH}_2\text{CO}_2\text{H} + \cdot\text{CH}_2\text{CH}_2\text{CO}_2\text{H}) = 2.4 \times 10^9$ .	74D286
239.075	·CH <sub>2</sub> OH + ICH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H → ·CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H + prod.	1.0 × 10 <sup>5</sup> (rel.)	1	chem.	Est. from esr meas. in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln. contg. MeOH; rel. to $2k(\cdot\text{CH}_2\text{CH}_2\text{CO}_2\text{H} + \cdot\text{CH}_2\text{CH}_2\text{CO}_2\text{H}) = 2.4 \times 10^9$ .	74D286
240. 240.147	<b>3-Iodotyrosine</b> ·CO <sub>2</sub> <sup>-</sup> + IC <sub>6</sub> H <sub>3</sub> (OH)CH <sub>2</sub> CH(·NH <sub>3</sub> )CO <sub>2</sub> H → CO <sub>2</sub> + I <sup>-</sup> + ·Ar	(1.3 ± 0.1) × 10 <sup>5</sup>	3.0	X-r.	Est. from dependence of tyrosine yields on irradiation time in soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> formate assuming $2k(R + R) = 5.0 \times 10^8$ .	72G610
241. 241.001	<b>Isobutylene</b> ·CH <sub>3</sub> + (CH <sub>3</sub> ) <sub>2</sub> C=CH <sub>2</sub> → addn.	3.9 × 10 <sup>4</sup> (rel.)		γ-r.	C.k. with MeOH in soln. contg. MeI obs. $G(\text{CH}_3)$ ; rel. to $k(\cdot\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
242. 242.001	<b>Isobutyric acid</b> ·CH <sub>3</sub> + (CH <sub>3</sub> ) <sub>2</sub> CHCO <sub>2</sub> H → CH <sub>4</sub> + (CH <sub>3</sub> ) <sub>2</sub> ĊCO <sub>2</sub> H	9.0 × 10 <sup>3</sup> (rel.)	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> -soln.; est. from esr meas. and values for competing reactions.	75D188

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
243. 243.001	<b>Isobutyronitrile</b> $\cdot\text{CH}_3 + (\text{CH}_3)_2\text{CHCN} \rightarrow$ $\text{CH}_4 + (\text{CH}_3)_2\dot{\text{C}}\text{CN}$	$4.5 \times 10^3$ (rel.)	$\sim 1$	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> -soln.; est. from esr meas. and values for competing reactions.	75D188
244. 244.078	<b>Isonicotinamide</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NH}^+ \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{NH}$	$(3.1 \pm 0.6) \times 10^9$	0.7	p.r.	P.b.k. (pyridinyl radical) in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	74A089
245. 245.078	<b>Isonicotinic acid</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}^+\text{NC}_5\text{H}_4\text{CO}_2\text{H}$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}^+\text{NC}_5\text{H}_4\text{CO}_2^-$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NC}_5\text{H}_4\text{CO}_2^-$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{NC}_5\text{H}_4\text{CO}_2^-$	$2.0 \times 10^9$ $8.5 \times 10^8$ $< 1 \times 10^6$ $2 \times 10^8$	0.4 3.2 9.0 13.3	p.r.	P.b.k. at $\sim 400$ nm (pyridinyl radical) in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	74A106
<i>Isopropanol See 2-Propanol 314.</i>						
246. 246.113	<b><math>\alpha</math>-Ketoglutarate ion</b> $\cdot\text{A}^- + \cdot\text{O}_2\text{CCH}_2\text{CH}_2\text{COCO}_2^-$ ( $\cdot\text{A}^-$ = ascorbate radical)	$< 10$	9.7	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate ion.	75A240
247. 247.113	<b>Lactate ion</b> $\cdot\text{A}^- + \text{CH}_3\text{CHOHCO}_2^-$ ( $\cdot\text{A}^-$ = ascorbate radical)	$< 10$	8.6	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate ion	75A240
248. 248.001	<b>Lactic acid</b> $\cdot\text{CH}_3 + \text{CH}_3\text{CHOHCO}_2\text{H} \rightarrow$ $\text{CH}_4 + \text{CH}_3\dot{\text{C}}\text{OHCO}_2\text{H}$	$1.2 \times 10^4$	$\sim 1$	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
249. 249.076	<b>Lipoate ion</b> $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{RSSR} \rightarrow$ $(\text{RSSR})^- + \text{CH}_3\text{CHO} + \text{H}^+$	$1.0 \times 10^8$	7	p.r.	P.b.k. at 410 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	70G560
249.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{RSSR} \rightarrow$ $(\text{RSSR})^- + (\text{CH}_3)_2\text{CO} + \text{H}^+$	$1.8 \times 10^8$	7	p.r.	P.b.k. at 410 nm in soln. contg. 1 mol L <sup>-1</sup> acetone and 2-PrOH.	70G560
249.147	$\cdot\text{CO}_2^- + \text{RSSR} \rightarrow$ $(\text{RSSR})^- + \text{CO}_2$	$5.5 \times 10^8$ $5.6 \times 10^8$	7 6.1–9.2	p.r. p.r.	P.b.k. at 410 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate ion. P.b.k. at 410 nm; $k = 9 \times 10^8$ at pH 3.	70G560 75A195
$\text{RSSR} = -\text{SCH}_2\text{CH}_2\text{CH}[(\text{CH}_2)_4\text{CO}_2^-]\text{S}-$						
250. 250.078	<b>Lumazine</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{LH}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{LH}_3\cdot$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{LH}^- \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{LH}_2\cdot^-$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{L}^{2-} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{LH}\cdot^{2-}$	$(1.3 \pm 0.1) \times 10^9$ $(1.7 \pm 0.2) \times 10^8$ (uncor.) $(1.0 \pm 0.2) \times 10^9$ (uncor.)	0.8, 5.1 9.5 14.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. solns. of 1–2 mol L <sup>-1</sup> 2-PrOH; 100%, 70% and 85% <i>e</i> -transfer at acid pH and pH 9.5 and 14.0, resp.	75A056
251. 251.147	<b>Lysozyme</b> $\cdot\text{CO}_2^- + \text{lys} \rightarrow$ $\text{CO}_2 + (-\text{SS})^-$	$\sim 5 \times 10^8$	$\sim 6$	p.r.	P.b.k. at 420 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate and $2 \times 10^{-4}$ mol L <sup>-1</sup> lysozyme; value from graph; $k$ pH-dependent.	75A080
252. 252.078	<b>Maleic acid</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HO}_2\text{CCH}=\text{CHCO}_2\text{H}$	$2.2 \times 10^8$ (uncor.)	0.5	p.r.	18% <i>e</i> -transfer; no <i>e</i> -transfer at pH 10.0 (dianion).	73G097
252.147	$\cdot\text{CO}_2^- + \text{HO}_2\text{CCH}=\text{CHCO}_2^-$	$1.1 \times 10^8$ (uncor.)	5.2	p.r.	$\geq 65\%$ <i>e</i> -transfer; no <i>e</i> -transfer at pH 10.5 (dianion).	73G097

*Menaquinone See 2-Methylnaphthoquinone 262.*

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
253.	<b>2-Mercaptoethanol</b>					
253.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{HO}(\text{CH}_2)_2\text{SH} \rightarrow$ $\text{CH}_3\text{CH}_2\text{OH} + \text{HO}(\text{CH}_2)_2\text{S}\cdot$	$(4.7 \pm 0.7) \times 10^7$	10	p.r.	P.b.k. (RSSR <sup>-</sup> ) in Ar-satd. soln.; radical from $e_{\text{aq}}^- + \text{HO}(\text{CH}_2)_2\text{SH}$ .	69G553
253.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{HO}(\text{CH}_2)_2\text{SH} \rightarrow$ $(\text{CH}_3)_3\text{COH} + \text{HO}(\text{CH}_2)_2\text{S}\cdot$	$(8.2 \pm 1.2) \times 10^7$	10	p.r.	P.b.k. at 420 nm (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	69G553
253.075	$\cdot\text{CH}_2\text{OH} + \text{HO}(\text{CH}_2)_2\text{SH} \rightarrow$ $\text{CH}_3\text{OH} + \text{HO}(\text{CH}_2)_2\text{S}\cdot$	$(1.3 \pm 0.2) \times 10^8$	10	p.r.	P.b.k. (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	69G553
253.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{HO}(\text{CH}_2)_2\text{SH} \rightarrow$ $\text{CH}_3\text{CH}_2\text{OH} + \text{HO}(\text{CH}_2)_2\text{S}\cdot$	$(2.3 \pm 0.3) \times 10^8$	10	p.r.	P.b.k. (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	69G553
253.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HO}(\text{CH}_2)_2\text{SH} \rightarrow$ $(\text{CH}_3)_2\text{CHOH} + \text{HO}(\text{CH}_2)_2\text{S}\cdot$	$(5.1 \pm 0.8) \times 10^8$	10	p.r.	P.b.k. (RSSR <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	69G553
253a.	<b>3-Mercaptopropionate ion</b>					
253a.142	$\text{R} + \text{HSCH}_2\text{CH}_2\text{CO}_2^- \rightarrow e^-$ -transfer (R = Electron adduct of glycine anhydride)	$(3.0 \pm 0.3) \times 10^8$	5.3,	p.r.	D.k. at 265 nm in soln. contg. 1 mol L <sup>-1</sup> <i>tert</i> -BuOH.	71G554
254.	<b>Methanethiol</b>					
254.001	$\cdot\text{CH}_3 + \text{CH}_3\text{SH} \rightarrow \text{CH}_4 + \text{CH}_3\text{S}\cdot$	$(7.4 \pm 1) \times 10^7$	11	p.r.	P.b.k. (RSSR <sup>-</sup> ) in Ar-satd. soln.; radical from $e_{\text{aq}}^-$ reaction with CH <sub>3</sub> SH.	69G553
255.	<b>Methanol</b>					
255.001	$\cdot\text{CH}_3 + \text{CH}_3\text{OH} \rightarrow \text{CH}_4 + \cdot\text{CH}_2\text{OH}$	$2.2 \times 10^2$ (rel.) $\geq 1 \times 10^2$		$\gamma$ -r. 1 phot.	C.k. with O <sub>2</sub> ; radical from MeI; rel. to $k(\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ . Radical from $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_3^{2+}$ ; est. from effect of addn. of MeOH on CH <sub>4</sub> and C <sub>2</sub> H <sub>6</sub> yields.	67G041 71F579
255.047	$\cdot\text{CF}_3 + \text{CH}_3\text{OH} \rightarrow \text{CF}_3\text{H} + \cdot\text{CH}_2\text{OH}$	$(8.1 \pm 1.2) \times 10^3$ (rel.)	9-10	$\gamma$ -r.	C.k. with addn. of $\cdot\text{CF}_3$ to propylene; obs. $G(\text{CF}_3\text{H})$ ; rel. to $k(\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ .	70G407
255.113	$\cdot\text{A}^- + \text{CH}_3\text{OH}$ ( $\cdot\text{A}^-$ = ascorbate radical)	$< 0.1$	8.8	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate ion.	75A240
256.	<b>Methemerythrin</b>					
256.147	$\cdot\text{CO}_2^- + \text{MFe}^{3+}$	$6.8 \times 10^7$	8.2	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> formate; $I = 0.03$ .	79A204
257.	<b>3-Methoxy-2-nitrotoluene</b>					
257.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{ArNO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{Ar}\dot{\text{N}}\text{O}_2^-$	$2.3 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
258.	<b>4-Methoxyphenoxide ion</b>					
258.020	$\cdot\text{CH}_2\text{CHO} + \text{CH}_3\text{OC}_6\text{H}_4\text{O}^- (+ \text{H}_2\text{O}) \rightarrow$ $\text{CH}_3\text{CHO} + \text{CH}_3\text{C}_6\text{H}_4\text{O}\cdot + \text{OH}^-$	$9.8 \times 10^8$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
259.	<b>2-Methylbenzoquinone</b>					
259.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{Q}^-$	$3.5 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
260.	<b>Methylene blue</b>					
260.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{MB}^+$	$3.7 \times 10^9$	7	p.r.	D.k. at 580 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 86% <i>e</i> -transfer.	73A078
260.075	$\cdot\text{CH}_2\text{OH} + \text{MB}^+$	$3.4 \times 10^9$	7	p.r.	D.k. at 580 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 91% <i>e</i> -transfer.	73A078
260.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{MB}^+$	$(4 \pm 2) \times 10^9$ $4.4 \times 10^9$		p.r. 7 p.r.	D.k. at 580 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 91% <i>e</i> -transfer.	73A150 73A078



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
260.147	$\cdot\text{CO}_2^- + \text{MB}^+ \rightarrow$ $\text{CO}_2 + \text{MB}\cdot$	$5.6 \times 10^9$	$\sim 9$	p.r.	D.k. at 580 nm in soln. contg. 0.1 mol L <sup>-1</sup> formate.	65G396
	$\cdot\text{CO}_2\text{H} + \text{MB}^+ \rightarrow$ $\text{CO}_2 + \text{H}^+ + \text{MB}\cdot$	$\sim 2 \times 10^9$	1.75	p.r.	D.k. at 580 nm.	65G396
	$\cdot\text{CO}_2\text{H} + \text{MBH}_2^{2+} \rightarrow$ $\text{CO}_2 + \cdot\text{MBH}_2^+ + \text{H}^+$	$\sim 10^9$	$\text{H}_0 = -0.8$	p.r.	D.k. at 660 nm.	65G396
	$\cdot\text{CO}_2\text{H} + \text{MBH}_2^{3+} \rightarrow$ $\text{CO}_2 + \cdot\text{MBH}_2^{2+} + \text{H}^+$	$\sim 7 \times 10^8$	$\text{H}_0 = -5.7$	p.r.	D.k. at 420 nm.	65G396
261.	<b>1-Methylguanosine</b>					
261.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{MG}^+ \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{MG}\cdot + \text{H}^+$	$8.0 \times 10^7$	0.5	p.r.	P.b.k.	75A060
<i>Methyl iodide See Iodomethane 238a.</i>						
<i>1-Methylisonicotinic acid See 4-Carboxy-1-methylpyridinium ion 154.</i>						
<i>Methyl mercaptan See Methanethiol 254.</i>						
262.	<b>2-Methyl-1,4-naphthoquinone</b>					
262.011	$\cdot\text{CH}_2\text{CH}_2\text{NH}_2 + \text{Q}$		7.5	p.r.	P.b.k. in soln. contg. ethylamine; 10% <i>e</i> -transfer; see 262.049.	73A047
262.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{Q}$		7.0, 12.4	p.r.	P.b.k. in soln. contg. 0.05 mol L <sup>-1</sup> <i>tert</i> -BuOH; $\leq 5\%$ <i>e</i> -transfer.	73A047
262.034	$\cdot\text{CH}_2\text{CONH}_2$ (or $\text{CH}_3\text{CONH}\cdot$ ) $+ \text{Q} \rightarrow \cdot\text{Q}^-$	$(1.1 \pm 0.1) \times 10^9$ (uncor.)	6.0, 10.9	p.r.	P.b.k. at 395 nm in soln. contg. acetamide; $\sim 17\%$ <i>e</i> -transfer.	73A047
262.049	$\text{CH}_3\dot{\text{C}}\text{HNH}_2 + \text{Q} \rightarrow \cdot\text{Q}^-$	$(3.3 \pm 0.3) \times 10^9$ (uncor.)	11.6	p.r.	P.b.k. at 395 nm in soln. contg. ethylamine; 34% <i>e</i> -transfer.	73A047
262.050	$(\text{CH}_3)_2\dot{\text{C}}\text{NH}_2 + \text{Q} \rightarrow \cdot\text{Q}^-$	$(3.6 \pm 0.4) \times 10^9$ (uncor.)	11.4	p.r.	P.b.k. at 395 nm in soln. contg. isopropylamine; 41% <i>e</i> -transfer; 14% <i>e</i> -transfer at pH 9.0.	73A047
262.051	$\text{CH}_3\dot{\text{C}}\text{HN}(\text{C}_2\text{H}_5)_2 + \text{Q} \rightarrow \cdot\text{Q}^-$	$(4.6 \pm 0.5) \times 10^9$ (uncor.)	11.6	p.r.	P.b.k. at 395 nm in soln. contg. triethylamine; 37% <i>e</i> -transfer; $\sim 10\%$ <i>e</i> -transfer at pH 6.	73A047
262.053	$\text{CH}_3\text{CONH}\dot{\text{C}}\text{H}_2 + \text{Q} \rightarrow \cdot\text{Q}^-$	$(2.0 \pm 0.2) \times 10^9$ (uncor.)	6.0, 10.9	p.r.	P.b.k. at 395 nm in soln. contg. <i>N</i> -methylacetamide; 19% <i>e</i> -transfer.	73A047
262.055	$\text{CH}_3\text{CON}(\text{CH}_3)\dot{\text{C}}\text{H}_2 + \text{Q}$		6.0, 10.9	p.r.	11% <i>e</i> -transfer in soln. contg. <i>N,N</i> -dimethylacetamide.	73A047
262.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$3.8 \times 10^9$ (uncor.)	6.2	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 78% <i>e</i> -transfer.	72R057
		$(5.5 \pm 0.6) \times 10^9$ (uncor.)	5.4	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 71% <i>e</i> -transfer.	73A047
		$(4.0 \pm 0.4) \times 10^9$ (uncor.)	8.0	p.r.	P.b.k. at 395 nm in soln. contg. glycine; 79% <i>e</i> -transfer.	73A047 73A104
262.057	$\text{NH}_3^+\dot{\text{C}}\text{HCONH}_2 + \text{Q} \rightarrow \cdot\text{Q}^-$	$(3.6 \pm 0.4) \times 10^9$ (uncor.)	3.2	p.r.	P.b.k. at 395 nm in soln. contg. glycinamide; 24% <i>e</i> -transfer.	73A047
	$\text{NH}_2\dot{\text{C}}\text{HCONH}_2 + \text{Q}$	$5 \times 10^9$ (uncor.)	8.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycinamide; 38% <i>e</i> -transfer.	72R057
		$(5.4 \pm 0.5) \times 10^9$ (uncor.)	10.4	p.r.	P.b.k. at 395 nm in soln. contg. glycinamide; 41% <i>e</i> -transfer.	73A047
262.058	$\text{R} + \text{Q} \rightarrow \cdot\text{Q}^-$ (R = Radicals from glycyl glycinamide)	$(8.5 \pm 0.9) \times 10^8$ $(8.5 \pm 0.9) \times 10^8$ (uncor.)	6.8 11.0	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycylglycinamide; 25% <i>e</i> -transfer at pH 6.8 (16% <i>e</i> -transfer at pH 5.5 and 45% at pH 11.0).	73A047
262.059	$\text{CH}_3\text{NH}\dot{\text{C}}\text{HCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$(1.1 \pm 0.1) \times 10^9$ (uncor.)	6.0	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. sarcosine; 33% <i>e</i> -transfer.	73A047
		$(1.7 \pm 0.2) \times 10^9$ (uncor.)	12.5	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. sarcosine; 64% <i>e</i> -transfer.	73A047
262.061	$\text{CH}_3\text{CON}(\text{CH}_3)\dot{\text{C}}\text{HCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$(1.3 \pm 0.1) \times 10^9$ $(1.0 \pm 0.1) \times 10^9$ (uncor.)	7.0 12.5	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetylsarcosine; 39% <i>e</i> -transfer at pH 7, 38% at pH 12.5.	73A047

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
262.062	R + Q → ·Q <sup>-</sup> (R = Radical from glycine anhydride)	(4.0 ± 0.4) × 10 <sup>9</sup> (uncor.)	10.9	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycine anhydride; 88% <i>e</i> -transfer; no <i>e</i> -transfer at pH 6.7.	73A047
		5.0 × 10 <sup>9</sup> (uncor.)	10.9	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine anhydride; 85% <i>e</i> -transfer; no <i>e</i> -transfer at pH 7.0.	72R057
262.063	R + Q → ·Q <sup>-</sup> (R = Radical from alanine anhydride)	(3.1 ± 0.3) × 10 <sup>9</sup> (uncor.)	10.9	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. alanine anhydride, 75% <i>e</i> -transfer.	73A047
262.065	R + Q → ·Q <sup>-</sup> (R = Radicals from diglycine)	(1.2 ± 0.1) × 10 <sup>9</sup> (uncor.)	6.6 11.0	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. diglycine; 19% <i>e</i> -transfer at pH 6.6 and 47% at pH 11.	73A047
262.066	R + Q → ·Q <sup>-</sup> (R = Radicals from glycylsarcosine)	(1.0 ± 0.1) × 10 <sup>9</sup> (uncor.)	10.9	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycylsarcosine; 46% <i>e</i> -transfer (20% at pH 6.8).	73A047
262.067	R + Q → ·Q <sup>-</sup> (R = Radicals from acetyldiglycine)	(3.8 ± 0.4) × 10 <sup>9</sup> (uncor.)	12.3	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetyldiglycine; 55% <i>e</i> -transfer.	73A047
262.068	R + Q → ·Q <sup>-</sup> (R = Radicals from triglycine)	(1.8 ± 0.2) × 10 <sup>9</sup> (uncor.)	7.0	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. triglycine; 11% <i>e</i> -transfer.	73A047
		(1.9 ± 0.2) × 10 <sup>9</sup> (uncor.)	10	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. triglycine.	73A104
		(1.8 ± 0.2) × 10 <sup>9</sup> (uncor.)	12.0	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. triglycine; 77% <i>e</i> -transfer.	73A047
262.069	R + Q → ·Q <sup>-</sup> (R = Radicals from acetyltriglycine)	(3.7 ± 0.4) × 10 <sup>9</sup> (uncor.)	12.5	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetyltriglycine; 50% <i>e</i> -transfer.	73A047
262.070	R + Q → ·Q <sup>-</sup> (R = Radicals from acetyltrialanine)	(2.1 ± 0.2) × 10 <sup>9</sup> (uncor.)	6.9	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetyltrialanine; 18% <i>e</i> -transfer.	73A047
		(2.6 ± 0.3) × 10 <sup>9</sup> (uncor.)	12.3	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetyltrialanine; 47% <i>e</i> -transfer.	73A047
262.071	R + Q → ·Q <sup>-</sup> (R = Radicals from acetyltrisarcosine)	(1.3 ± 0.1) × 10 <sup>9</sup> (uncor.)	12.5	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetyltrisarcosine; 39% <i>e</i> -transfer at pH 12.5 and 7.0.	73A047
262.072	R + Q → ·Q <sup>-</sup> (R = Radicals from acetylserineamide)	(1.5 ± 0.2) × 10 <sup>9</sup> (1.9 ± 0.2) × 10 <sup>9</sup> (uncor.)	6, 9, 11.0	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetylserineamide; 52%, 57%, and 68% <i>e</i> -transfer, resp., at pH 6, 9, and 11.	73A047
262.073	R + Q → ·Q <sup>-</sup> (R = Radicals from acetylaspargine)	(1.5 ± 0.2) × 10 <sup>9</sup> (uncor.)	12.5	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. acetylaspargine; 39% <i>e</i> -transfer; ~12% <i>e</i> -transfer at pH 3.2 and 6.0.	73A047
262.075	·CH <sub>2</sub> OH + Q → ·Q <sup>-</sup>	(3.7 ± 0.4) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 395 nm in 0.5 mol L <sup>-1</sup> MeOH; 88% <i>e</i> -transfer.	73A047
	·CH <sub>2</sub> O <sup>-</sup> + Q → ·Q <sup>-</sup>	(4.4 ± 0.4) × 10 <sup>9</sup>	12.4	p.r.	P.b.k. at 395 nm in soln. contg. 0.5 mol L <sup>-1</sup> MeOH; 92% <i>e</i> -transfer.	73A047
262.076	CH <sub>3</sub> ĊHOH + Q → ·Q <sup>-</sup>	(3.8 ± 0.4) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 395 nm in soln. contg. 0.5 mol L <sup>-1</sup> EtOH; 90% <i>e</i> -transfer.	73A047
	CH <sub>3</sub> ĊHO <sup>-</sup> + Q → ·Q <sup>-</sup>	(4.2 ± 0.4) × 10 <sup>9</sup>	12.5	p.r.	P.b.k. at 395 nm in soln. contg. 0.5 mol L <sup>-1</sup> EtOH; 92% <i>e</i> -transfer.	73A047
262.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Q → ·Q <sup>-</sup>	6.2 × 10 <sup>9</sup>	6.9	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 80% <i>e</i> -transfer.	72R057
		(4.2 ± 0.4) × 10 <sup>9</sup>	7	p.r.	P.b.k. in soln. cont. 1 mol L <sup>-1</sup> acetone and 1 mol L <sup>-1</sup> 2-PrOH.	73G125
		(4.1 ± 0.4) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. in soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH; 91% <i>e</i> -transfer.	73A047
		(4.8 ± 0.5) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73A104
		(4.2 ± 0.4) × 10 <sup>9</sup>	12.4	p.r.	P.b.k. in soln. contg. 0.5 mol L <sup>-1</sup> EtOH; 91% <i>e</i> -transfer.	73A047



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
262.079	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CHOH + Q → ·Q <sup>-</sup>	(4.1 ± 0.4) × 10 <sup>9</sup> (4.2 ± 0.4) × 10 <sup>9</sup> (uncor.)	7.0 12.7	p.r.	P.b.k. in soln. contg. 0.05 mol L <sup>-1</sup> 1-BuOH; 32.3% (pH 7) and 40% (pH 12.7) <i>e</i> -transfer.	73A047
262.097	CH <sub>3</sub> Ċ(O <sup>-</sup> )CH <sub>2</sub> CO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup>	(3.7 ± 0.4) × 10 <sup>9</sup>	9.2	p.r.	P.b.k. at 395 nm in soln. contg. acetoacetate and 1 mol L <sup>-1</sup> <i>tert</i> -BuOH; 95% <i>e</i> -transfer.	73A047 73A104
262.108	R + Q → ·Q <sup>-</sup> (R = Radicals from deoxyribose)	4 × 10 <sup>9</sup> (uncor.) (2.1 ± 0.2) × 10 <sup>9</sup> (uncor.)	6.8 6.8	p.r. p.r.	P.b.k. at 400 nm in soln. contg. deoxyribose; 74% <i>e</i> -transfer. P.b.k. in soln. contg. 0.05 mol L <sup>-1</sup> deoxyribose; 81% <i>e</i> -transfer.	72R057 73A047
262.109	R + Q → ·Q <sup>-</sup> (R = Radicals from ribose)	2.6 × 10 <sup>9</sup> (uncor.) (1.4 ± 0.1) × 10 <sup>9</sup> (uncor.)	6.9 6.9	p.r. p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. ribose; 44% <i>e</i> -transfer. P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> ribose; 60% <i>e</i> -transfer.	72R057 73A047
262.110	R + Q → Q <sup>-</sup> (R = Radicals from ribose phosphate)	4.0 × 10 <sup>9</sup> (uncor.)	6.8	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. ribose-5-phosphate; 25% <i>e</i> -transfer.	72R057
262.113	·A <sup>-</sup> + Q → ·Q <sup>-</sup> (·A <sup>-</sup> = Radical from ascorbate)	~1.3 × 10 <sup>9</sup> (3.8 ± 0.4) × 10 <sup>9</sup> (uncor.)	3.3 9.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> ascorbic acid; 43 and 38% <i>e</i> -transfer, resp., at pH 3.3 and 9.0.	73A047
262.114	ĊHO <sup>-</sup> CONH <sub>2</sub> + Q → ·Q <sup>-</sup>	(2.3 ± 0.2) × 10 <sup>9</sup> (uncor.)	7.1	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycolamide; 48% <i>e</i> -transfer.	73A047
262.115	ĊHOHCO <sub>2</sub> H + Q → ·Q <sup>-</sup> ĊHOHCO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup> ĊHO <sup>-</sup> CO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup>	(9.2 ± 0.9) × 10 <sup>8</sup> (uncor.) (1.5 ± 0.2) × 10 <sup>9</sup> (uncor.) (1.6 ± 0.2) × 10 <sup>9</sup> (uncor.)	3.2 6.5 10.6	p.r. p.r. p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycolic acid; 13% <i>e</i> -transfer. P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycolate ion; 69% <i>e</i> -transfer P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. glycolate ion; 77% <i>e</i> -transfer.	73A047 73A047 73A104
262.117	CH <sub>3</sub> ĊOHCO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup> CH <sub>3</sub> Ċ(O <sup>-</sup> )CO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup>	(1.4 ± 0.1) × 10 <sup>9</sup> (uncor.) (1.9 ± 0.2) × 10 <sup>9</sup> (uncor.)	6.5 10.6	p.r. p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. lactate ion; 55% <i>e</i> -transfer. P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. lactate ion; 72% <i>e</i> -transfer (15% at pH 3.2).	73A047 73A047
262.118	CH <sub>3</sub> Ċ(O <sup>-</sup> )CO <sub>2</sub> CH <sub>3</sub> + Q → ·Q <sup>-</sup>	(2.3 ± 0.2) × 10 <sup>9</sup> (uncor.)	10.4	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. methyl lactate; 74% <i>e</i> -transfer (20% at pH 3.2).	73A047
262.119	ĊO <sup>-</sup> (CO <sub>2</sub> ) <sub>2</sub> + Q → ·Q <sup>-</sup>	(2.5 ± 0.3) × 10 <sup>9</sup>	9.2	p.r.	P.b.k. at 395 nm in soln. contg. ketomalonate and 1 mol L <sup>-1</sup> <i>tert</i> -BuOH; 94% <i>e</i> -transfer.	73A047
262.120	·O <sub>2</sub> CĊ(OH)CH <sub>2</sub> CO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup>	(1.1 ± 0.1) × 10 <sup>9</sup> (uncor.) (3.1 ± 0.3) × 10 <sup>9</sup>	7.0 6.2	p.r. p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. malate; 47% <i>e</i> -transfer. P.b.k. at 395 nm in soln. contg. oxaloacetate and 1 mol L <sup>-1</sup> <i>tert</i> -BuOH; 92% <i>e</i> -transfer.	73A047 73A047
262.121	HO <sub>2</sub> CCHOHĊ(OH)CO <sub>2</sub> H + Q → ·Q <sup>-</sup> ·O <sub>2</sub> CCHOHĊ(OH)CO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup>	(7.0 ± 0.7) × 10 <sup>8</sup> (uncor.) (7.0 ± 0.7) × 10 <sup>8</sup> (uncor.)	3.2 11.0	p.r. p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. tartaric acid; 14% <i>e</i> -transfer. P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. tartrate ion; 69% <i>e</i> -transfer.	73A047 73A047
262.122	·O <sub>2</sub> C(CH <sub>2</sub> ) <sub>2</sub> Ċ(O <sup>-</sup> )CO <sub>2</sub> <sup>-</sup> + Q → ·Q <sup>-</sup>	(3.8 ± 0.4) × 10 <sup>9</sup>	10.2	p.r.	P.b.k. at 395 nm in soln. contg. α-ketoglutarate and 1 mol L <sup>-1</sup> <i>tert</i> -BuOH; 94% <i>e</i> -transfer.	73A047
262.142	R + Q → ·Q <sup>-</sup> (R = Electron adduct of glycine anhydride)	(4.9 ± 0.5) × 10 <sup>9</sup>	6.9	p.r.	P.b.k. at 395 nm in soln. contg. glycine anhydride; 98% <i>e</i> -transfer.	73A047
262.143	R + Q → ·Q <sup>-</sup> (R = Electron adduct of alanine anhydride)	(4.8 ± 0.5) × 10 <sup>9</sup>	6.0	p.r.	P.b.k. at 395 nm in soln. contg. alanine anhydride; 92% <i>e</i> -transfer.	73A047
262.144	R + Q → ·Q <sup>-</sup> (R = Electron adduct of sarcosine anhydride)	(4.6 ± 0.5) × 10 <sup>9</sup>	6.0	p.r.	P.b.k. at 395 nm in soln. contg. sarcosine anhydride; 91% <i>e</i> -transfer.	73A047
262.146	R + Q → ·Q <sup>-</sup> (R = Electron adduct of acetylglcylglycinamide)	(2.7 ± 0.3) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 395 nm; 99% <i>e</i> -transfer.	73A047



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
262.147	$\cdot\text{CO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$5.4 \times 10^9$	6.9	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. formate; 100% $e$ -transfer.	72R057
		$(4.8 \pm 0.5) \times 10^9$	7	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. formate.	73A047 73A104
<i>1-Methylnicotinamide See 3-Carbamoyl-1-methylpyridinium ion 151.</i>						
<i>1-Methylnicotinate ion See 3-Carboxy-1-methylpyridinium ion 153.</i>						
<i>3-Methyl-2-nitroanisole See 3-Methoxy-2-nitrotoluene 257.</i>						
263.	<b>3-Methyl-2-nitrobenzoate ion</b>					
263.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NO}_2\text{B}^- \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{NO}_2\text{B}^{2-}$	$1.9 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
264.	<b>2-Methyl-5-nitroimidazole</b>					
264.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NX}^-$	$2.5 \times 10^9$		p.r.	P.b.k.	76A075
265.	<b>3-Methyl-2-nitrophenol</b>					
265.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3(\text{OH})\text{C}_6\text{H}_3\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{CH}_3(\text{OH})\text{C}_6\text{H}_3\dot{\text{N}}\text{O}_2^-$	$2.9 \times 10^8$		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
266.	<b>4-Methylphenoxide ion</b>					
266.020	$\cdot\text{CH}_2\text{CHO} + \text{CH}_3\text{C}_6\text{H}_4\text{O}^- (+ \text{H}_2\text{O}) \rightarrow$ $\text{CH}_3\text{CHO} + \text{CH}_3\text{C}_6\text{H}_4\text{O} \cdot + \text{OH}^-$	$9.0 \times 10^7$	~11.5	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
267.	<b>Methyl propionate</b>					
267.001	$\cdot\text{CH}_3 + \text{C}_2\text{H}_5\text{CO}_2\text{CH}_3 \rightarrow$ $\text{CH}_4 + \text{CH}_3\dot{\text{C}}\text{HCO}_2\text{CH}_3$	$2.9 \times 10^3$ (rel.)	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> -soln.; est. from esr meas. and values of competing reactions.	75D188
268.	<b>3-Methylpterin</b>					
268.075	$\cdot\text{CH}_2\text{OH} + \text{PtH}^+$	$6 \times 10^7$ (uncor.)	0.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. MeOH; ~45% $e$ -transfer.	76A060
268.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Pt}$	$3.2 \times 10^7$ (uncor.)	6.3	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. EtOH; 55% $e$ -transfer.	76A060
268.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PtH}^+$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pt}$	$1.9 \times 10^9$ $2.9 \times 10^8$	0.8 7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 100% $e$ -transfer.	76A060
269.	<b>9-Methylpurine</b>					
269.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{MP} \rightarrow \text{MPH} \cdot$ $\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{MP} \rightarrow \text{MP}^-$	$<2 \times 10^7$ $5.1 \times 10^8$	8.2 13.6	p.r.	P.b.k.; N <sub>2</sub> O-satd. soln. contg. EtOH.	76A060
269.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{MPH}^+ \rightarrow \text{MPH}_3^{2+}$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{MP} \rightarrow \text{MPH} \cdot$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{MP} \rightarrow \text{MP}^-$	$1.9 \times 10^9$ $1.7 \times 10^8$ $8.7 \times 10^8$	~0 8.6 13.6	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	76A060
<i>1-Methyl-3-pyridinecarboxylate ion See 3-Carboxy-1-methylpyridinium ion 153.</i>						
<i>1-Methyl-4-pyridinecarboxylate ion See 4-Carboxy-1-methylpyridinium ion 154.</i>						
<i>Methylviologen See 1,1'-Dimethyl-4,4'-bipyridinium ion 183.</i>						
270.	<b>Metmyoglobin</b>					
270.075	$\cdot\text{CH}_2\text{OH} + \text{M}(\text{Fe}^{\text{III}}) \rightarrow$ $\text{CH}_2\text{O} + \text{H}^+ + \text{M}(\text{Fe}^{\text{II}})$	$(2.4 \pm 0.5) \times 10^7$	7	p.r.	95% redn. of hemoprotein in N <sub>2</sub> O-satd. soln. contg. MeOH; obs. abs. change at 550 and 435 nm.	79A153
270.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{M}(\text{Fe}^{\text{III}})$	$(5.5 \pm 0.5) \times 10^7$	7	p.r.	95% redn. of hemoprotein in N <sub>2</sub> O-satd. soln. contg. EtOH.	79A153
270.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{M}(\text{Fe}^{\text{III}}) \rightarrow \text{M}(\text{Fe}^{\text{II}})$	$\sim 1 \times 10^8$	7.5	p.r.	$G = 2.2$ .	79A371
270.121	$\cdot\text{O}_2\text{CCHOH}\dot{\text{C}}\text{OHCO}_2^- + \text{M}(\text{Fe}^{\text{III}})$	$(3.5 \pm 0.7) \times 10^7$	7	p.r.	Redn. in soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> tartrate.	78A288

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
270.147	$\cdot\text{CO}_2^- + \text{M}(\text{Fe}^{\text{III}})$	$(2.0 \pm 0.4) \times 10^9$	7	p.r.	Redn. in soln. contg. $10^{-2}$ mol L <sup>-1</sup> formate.	78A288
		$2.9 \times 10^9$	8.2	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> formate; $I = 0.03$ .	79A204
271.	<b>Metronidazole [1-(2-Hydroxyethyl)-2-methyl-5-nitroimidazole]</b>					
271.075	$\cdot\text{CH}_2\text{OH} + \text{RNO}_2$	$10^8$		p.r.	D.k. at 320 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	74A135
271.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{RNO}_2$	$2 \times 10^8$		p.r.	D.k. at 320 nm in N <sub>2</sub> O-satd. soln. contg. EtOH.	74A135
271.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{RNO}_2$	$7 \times 10^8$		p.r.	D.k. at 320 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	74A135
271.147	$\cdot\text{CO}_2^- + \text{RNO}_2$	$8 \times 10^8$		p.r.	D.k. at 320 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate ion; also c.k. gave $8 \times 10^8$ rel. to $k(\cdot\text{CO}_2^- + \text{TAN}) = 6 \times 10^8$ [342.147].	74A135
272.	<b>Misonidazole [1-(2-Hydroxy-3-methoxypropyl)-2-nitroimidazole]</b>					
272.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{RNO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{RNO}_2^-$	$(2.8 \pm 0.4) \times 10^9$		p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A070
273.	<b>1,4-Naphthoquinone</b>					
273.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow \cdot\text{Q}^-$	$3.6 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
274.	<b>1,4-Naphthoquinone-2-sulfonate ion</b>					
274.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$(3.3 \pm 0.3) \times 10^9$	8.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycine.	73A104
274.068	$\text{R} + \text{Q} \rightarrow \cdot\text{Q}^-$ (R = Radicals from triglycine)	$(2.0 \pm 0.2) \times 10^9$	10	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. triglycine.	73A104
274.115	$\cdot\text{CHOHCO}_2^- + \text{Q} \rightarrow \cdot\text{Q}^-$	$(1.7 \pm 0.2) \times 10^9$	7.0	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. glycolate ion.	73A104
275.	<b>Nicotinamide</b>					
275.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NH}^+ \rightarrow$ $(\text{CH}_3)_2\text{CO} + \cdot\text{NH} + \text{H}^+$	$(2.1 \pm 0.4) \times 10^8$	0.9	p.r.	P.b.k.; no e-transfer in neutral soln.	74A089
		$4.0 \times 10^8$	1.9	p.r.	P.b.k. at ~400 nm in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	74A106
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{N}$	$< 1 \times 10^6$	9.2	p.r.	P.b.k. at ~400 nm in N <sub>2</sub> O-satd. soln. contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	74A106
276.	<b>Nicotinamide-adenine dinucleotide</b>					
276.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{NAD}^+$	$1.5 \times 10^9$	6.0	p.r.		73A104
276.075	$\cdot\text{CH}_2\text{OH} + \text{NAD}^+$	$1.0 \times 10^9$	6.0	p.r.		73A104
276.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NAD}^+ \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{NAD}$	$1.0 \times 10^9$		p.r.	Est. from buildup and decay of $\cdot\text{NAD}$ at 400 nm in soln. contg. 1 mol L <sup>-1</sup> acetone, 1 mol L <sup>-1</sup> 2-PrOH, $2 \times 10^{-4}$ mol L <sup>-1</sup> O <sub>2</sub> and $2 \times 10^{-5}$ mol L <sup>-1</sup> benzoquinone.	70R013
		$1.0 \times 10^9$	6.0	p.r.		73A104
276.147	$\cdot\text{CO}_2^- + \text{NAD}^+ \rightarrow \text{CO}_2 + \cdot\text{NAD}$	$1.6 \times 10^9$	6.4	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	68G441
277.	<b>Nicotinic acid</b>					
277.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}^+\text{NC}_5\text{H}_4\text{CO}_2\text{H}$	$3.5 \times 10^8$	0.0	p.r.	P.b.k. at ~400 nm (pyridinyl radical) in N <sub>2</sub> O-satd. soln.	74A106
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}^+\text{NC}_5\text{H}_4\text{CO}_2^-$	$1.8 \times 10^8$	3.4			
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NC}_5\text{H}_4\text{CO}_2^-$	$< 1 \times 10^6$	8.2		contg. 0.5 mol L <sup>-1</sup> 2-PrOH.	
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{NC}_5\text{H}_4\text{CO}_2^-$	$> 1 \times 10^8$	13.1			

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
278.	<b>Nifuroxime [<i>anti</i>-5-Nitro-2-furaldoxime]</b>					
278.018	R + NF (R = Radicals from allyl alcohol)	5.1 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 395 nm (NF <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. allyl alcohol; 3% <i>e</i> -transfer.	73A062 73A099
278.019	R + NF (R = Radicals from crotyl alcohol)	5.6 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm (NF <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. crotyl alcohol; 24% <i>e</i> -transfer.	73A099
		2.4 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. crotyl alcohol; 10% <i>e</i> -transfer.	73A052
278.037	CH <sub>3</sub> ĊHCO <sub>2</sub> <sup>-</sup> + NF	4.0 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm (NF <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> propionate ion; 2% <i>e</i> -transfer based on abs. spectra.	73A099
278.038	CH <sub>3</sub> CH <sub>2</sub> ĊHCO <sub>2</sub> <sup>-</sup> + NF	4.9 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm (NF <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> butyrate ion; 2% <i>e</i> -transfer.	73A099
278.039	HOCH <sub>2</sub> ĊHCO <sub>2</sub> <sup>-</sup> + NF	6 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in soln. contg. 4 x 10 <sup>-4</sup> mol L <sup>-1</sup> acrylate ion; 65% <i>e</i> -transfer.	73A099
278.040	CH <sub>3</sub> CHOHĊHCO <sub>2</sub> <sup>-</sup> + NF	5.5 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in soln. contg. crotonate ion; 25% <i>e</i> -transfer.	73A099
278.041	·O <sub>2</sub> CĊHCHOHCO <sub>2</sub> <sup>-</sup> + NF	2.4 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in soln. contg. maleate ion; 3% <i>e</i> -transfer.	73A099
278.042	·CH(CO <sub>2</sub> <sup>-</sup> ) <sub>2</sub> + NF	1.3 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> malonate ion; 3% <i>e</i> -transfer.	73A099
278.075	·CH <sub>2</sub> OH + NF → NF <sup>-</sup>	8 x 10 <sup>8</sup>	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> MeOH; 90% <i>e</i> -transfer.	73A099
278.076	CH <sub>3</sub> ĊHOH + NF → NF <sup>-</sup>	> 1.5 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> EtOH; > 75% <i>e</i> -transfer.	73A099
278.077	CH <sub>3</sub> CH <sub>2</sub> ĊHOH + NF → NF <sup>-</sup>	3.2 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 1-PrOH; 52% <i>e</i> -transfer.	73A062
		3.2 x 10 <sup>9</sup>	7	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. 1-PrOH; 95% <i>e</i> -transfer.	73A099
278.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + NF → NF <sup>-</sup>	3.3 x 10 <sup>9</sup>	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> 2-PrOH; 100% <i>e</i> -transfer.	73A099
278.079	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> ĊHOH + NF → NF <sup>-</sup>	3.5 x 10 <sup>9</sup> 5.1 x 10 <sup>9</sup> (uncor.)	7	p.r. p.r.	P.b.k. P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 1-BuOH; 75% <i>e</i> -transfer.	76A075 73A099
		3.4 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 395 nm in N <sub>2</sub> O-satd. soln. contg. 1-BuOH; 32% <i>e</i> -transfer.	73A062
278.099	NH <sub>3</sub> <sup>+</sup> CH <sub>2</sub> ĊHOPO <sub>3</sub> <sup>2-</sup> + NF	3.0 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> <i>o</i> -phosphoryl-ethanolamine; 1% <i>e</i> -transfer.	73A099
278.106	CH <sub>2</sub> OHCHOHĊHOH + NF (+ CH <sub>2</sub> OHĊOHCH <sub>2</sub> OH)	1.3 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> glycerol; 20% <i>e</i> -transfer.	73A099
278.108	R + NF → NF <sup>-</sup> (R = Radicals from deoxyribose)	1.4 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> deoxyribose; 70% <i>e</i> -transfer.	73A099
278.109	R + NF → NF <sup>-</sup> (R = Radicals from ribose)	1.5 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> ribose; 50% <i>e</i> -transfer.	73A099
278.110	R + NF (R = Radicals from ribose phosphate)	1.7 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> ribose phosphate; 10% <i>e</i> -transfer.	73A099



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
278.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2^- + \text{NF} \rightarrow \text{NF}^-$	$1.5 \times 10^9$	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> lactate ion; 100% $e$ -transfer.	73A099
278.120	$^-\text{O}_2\text{CCH}_2\dot{\text{C}}\text{OHCO}_2^- + \text{NF}$	$1.2 \times 10^8$ (uncor.)	7	p.r.	P.b.k. at 390 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> malate ion; 30% $e$ -transfer.	73A099
278.147	$^-\text{CO}_2 + \text{NF} \rightarrow \text{NF}^-$	$2.7 \times 10^9$		p.r.	P.b.k. at 390 nm in CO <sub>2</sub> -satd. soln. contg. <i>tert</i> -BuOH or 0.2 mol L <sup>-1</sup> formate; 100% $e$ -transfer.	73A099
279.	<b><i>p</i>-Nitroacetophenone</b>					
279.018	R + PNAP (R = Radicals from allyl alcohol)	$2.7 \times 10^9$ (uncor.)	7	p.r.	P.b.k. at 360 and 550 nm (PNAP <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. allyl alcohol; 3% $e$ -transfer.	73A062
279.019	R + PNAP (R = Radicals from crotyl alcohol)	$2.0 \times 10^9$ (uncor.)	7	p.r.	P.b.k. at 360 and 550 nm (PNAP <sup>-</sup> ) in N <sub>2</sub> O-satd. soln. contg. crotyl alcohol; 9% $e$ -transfer.	73A062
279.075	$^-\text{CH}_2\text{OH} + \text{PNAP} \rightarrow \text{addn.}$	$1 \times 10^8$	7	p.r.	P.b.k. at 330 nm in N <sub>2</sub> O-satd. soln. contg. MeOH.	73G122
	$^-\text{CH}_2\text{OH} + \text{PNAP} \rightarrow \text{PNAP}^-$	$1 \times 10^7$	7	p.r.	P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. MeOH.	73G122
	$^-\text{CH}_2\text{O} + \text{PNAP} \rightarrow \text{PNAP}^-$	$(4.7 \pm 0.5) \times 10^9$	13	p.r.	P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. MeOH.	73G122
279.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{PNAP} \rightarrow \text{PNAP}^-$	$(8 \pm 2) \times 10^8$	11	p.r.	P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. EtOH.	73G122
279.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{PNAP}$	$2.4 \times 10^9$ (uncor.)	7	p.r.	P.b.k. at 360 and 550 nm in N <sub>2</sub> O-satd. soln. contg. 1-PrOH; 45% $e$ -transfer.	73A062
279.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PNAP} \rightarrow \text{PNAP}^-$	$(3.8 \pm 0.4) \times 10^9$	11	p.r.	P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	73G122
		$2.2 \times 10^9$	~13	p.r.	P.b.k. at 355 nm in soln. contg. 6.5 mol L <sup>-1</sup> 2-PrOH and 0.68 mol L <sup>-1</sup> acetone.	81A002
279.079	$\text{CH}_3\text{CH}_2\text{CH}_2\dot{\text{C}}\text{HOH} + \text{PNAP}$	$2.2 \times 10^9$ (uncor.)	7	p.r.	P.b.k. at 360 and 550 nm in N <sub>2</sub> O-satd. soln. contg. 1-BuOH; 24% $e$ -transfer.	73A062
279.108	R + PNAP $\rightarrow$ PNAP <sup>-</sup> (R = Radicals from deoxyribose)	$(1.0 \pm 0.3) \times 10^9$		p.r.	P.b.k.; 47% $e$ -transfer.	75A207
279.111	R + PNAP $\rightarrow$ PNAP <sup>-</sup> (R = Radicals from glucose)	$9 \times 10^8$		p.r.		77R167
279.147	$^-\text{CO}_2 + \text{PNAP} \rightarrow \text{PNAP}^-$	$(1.0 \pm 0.1) \times 10^9$	10	p.r.	P.b.k. at 550 nm in N <sub>2</sub> O-satd. soln. contg. formate ion.	73G122
		$7 \times 10^8$		p.r.	Radical from formate.	77R167
280.	<b><i>o</i>-Nitroaniline</b>					
280.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NH}_2\text{C}_6\text{H}_4\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NH}_2\text{C}_6\text{H}_4\dot{\text{N}}\text{O}_2^-$	$9.2 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
281.	<b><i>m</i>-Nitroaniline</b>					
281.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NH}_2\text{C}_6\text{H}_4\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NH}_2\text{C}_6\text{H}_4\dot{\text{N}}\text{O}_2^-$	$1.5 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
282.	<b><i>p</i>-Nitroaniline</b>					
282.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NH}_2\text{C}_6\text{H}_4\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NH}_2\text{C}_6\text{H}_4\dot{\text{N}}\text{O}_2^-$	$7.2 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
		$(1.9 \pm 0.2) \times 10^9$	10.7	p.r.	P.b.k. at 380 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	77A118
283.	<b>Nitrobenzene</b>					
283.018	R + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> (R = Radicals from allyl alcohol)	~ $1.5 \times 10^9$ (uncor.)	7	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. allyl alcohol; 5% $e$ -transfer.	73A062

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
283.019	R + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> (R = Radicals from crotyl alcohol)	~1 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. crotyl alcohol; 8% <i>e</i> -transfer.	73A062
283.075	·CH <sub>2</sub> OH + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>2</sub> O + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	<10 <sup>7</sup>	7	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	66G432
		(6.0 ± 0.9) x 10 <sup>7</sup>	5-6	p.r.	N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	72A018
	·CH <sub>2</sub> O <sup>-</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>2</sub> O + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	2.7 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	66G432
283.076	CH <sub>3</sub> ·CHOH + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>3</sub> CHO + H <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.3 x 10 <sup>8</sup>	7	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	66G432
		(2.4 ± 0.7) x 10 <sup>8</sup>	5-6	p.r.	N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	72A018
	CH <sub>3</sub> ·CHO <sup>-</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>3</sub> CHO + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.1 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH.	66G432
		3 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. EtOH; <i>E</i> <sub>a</sub> ≈ 3.3 kcal mol <sup>-1</sup> (14 kJ mol <sup>-1</sup> ).	74A010
283.077	CH <sub>3</sub> CH <sub>2</sub> ·CHOH + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>3</sub> CH <sub>2</sub> CHO + H <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.5 x 10 <sup>8</sup>	7	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 1-PrOH; 68% <i>e</i> - transfer.	66G432
		1.0 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 1-PrOH; 60% <i>e</i> -transfer.	73A062
	CH <sub>3</sub> CH <sub>2</sub> ·CHO <sup>-</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>3</sub> CH <sub>2</sub> CHO + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.1 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 1-PrOH.	66G432
283.078	(CH <sub>3</sub> ) <sub>2</sub> ·CHOH + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → (CH <sub>3</sub> ) <sub>2</sub> CO + H <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	1.6 x 10 <sup>9</sup>	7	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	66G432
	(CH <sub>3</sub> ) <sub>2</sub> ·CO <sup>-</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → (CH <sub>3</sub> ) <sub>2</sub> CO + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.0 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	66G432
283.079	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> ·CHOH + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CHO + H <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	4.0 x 10 <sup>8</sup>	7	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 1-BuOH; 35% <i>e</i> - transfer.	66G432
		1.1 x 10 <sup>9</sup> (uncor.)	7	p.r.	P.b.k. at 400 nm in N <sub>2</sub> O-satd. soln. contg. 1-BuOH; 31% <i>e</i> -transfer;	73A062
	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> ·CHO <sup>-</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CHO + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.1 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 1-BuOH.	66G432
283.081	(CH <sub>3</sub> ) <sub>2</sub> CH·CHOH + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → (CH <sub>3</sub> ) <sub>2</sub> CHCHO + H <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	3.9 x 10 <sup>8</sup>	7	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-methyl-1-propanol; 39% <i>e</i> -transfer.	66G432
	(CH <sub>3</sub> ) <sub>2</sub> CH·CHO <sup>-</sup> + C <sub>6</sub> H <sub>5</sub> NO → (CH <sub>3</sub> ) <sub>2</sub> CHCHO + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	2.9 x 10 <sup>9</sup>	13	p.r.	P.b.k. at 285 nm in N <sub>2</sub> O-satd. soln. contg. 2-methyl-1-propanol.	66G432
283.084	-CH <sub>2</sub> ·COH(CH <sub>2</sub> ) <sub>2</sub> - + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → R=O + H <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	1.5 x 10 <sup>9</sup>		p.r.	P.b.k. at 280 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclobutanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; <i>k</i> = 3.2 x 10 <sup>9</sup> at 64°C; <i>E</i> <sub>a</sub> = 2.3 kcal mol <sup>-1</sup> (9.6 kJ mol <sup>-1</sup> ).	76A103
283.085	-CH <sub>2</sub> ·COH(CH <sub>2</sub> ) <sub>3</sub> - + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → R=O + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	1.4 x 10 <sup>9</sup>		p.r.	P.b.k. at 280 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclopentanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; <i>k</i> = 5.2 x 10 <sup>9</sup> at 64°C; <i>E</i> <sub>a</sub> = 4.0 kcal mol <sup>-1</sup> (17 kJ mol <sup>-1</sup> ).	76A103
283.086	-CH <sub>2</sub> ·COH(CH <sub>2</sub> ) <sub>4</sub> - + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → R=O + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	7 x 10 <sup>8</sup>		p.r.	P.b.k. at 280 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cyclohexanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; <i>k</i> = 2.0 x 10 <sup>9</sup> at 64°C <i>E</i> <sub>a</sub> = 3.1 kcal mol <sup>-1</sup> (13 kJ mol <sup>-1</sup> ).	76A103
283.087	-CH <sub>2</sub> ·COH(CH <sub>2</sub> ) <sub>5</sub> - + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> → R=O + C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> <sup>-</sup>	1.4 x 10 <sup>9</sup>		p.r.	P.b.k. at 280 nm in soln. contg. 10 <sup>-3</sup> mol L <sup>-1</sup> cycloheptanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; <i>k</i> = 4.9 x 10 <sup>9</sup> at 64°C; <i>E</i> <sub>a</sub> = 3.8 kcal mol <sup>-1</sup> (16 kJ mol <sup>-1</sup> ).	76A103

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
283.088	$-\text{CH}_2\dot{\text{C}}\text{OH}(\text{CH}_2)_6^- + \text{C}_6\text{H}_5\text{NO}_2 \rightarrow$ $\text{R}=\text{O} + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}_2^-$	$1.6 \times 10^9$		p.r.	P.b.k. at 280 nm in soln. contg. $10^{-3}$ mol L <sup>-1</sup> cyclooctanone and 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH at 2°C; $k = 5.2 \times 10^9$ at 64°C; $E_a = 3.7$ kcal mol <sup>-1</sup> (15.5 kJ mol <sup>-1</sup> ).	76A103
283.093	$\text{CH}_3\dot{\text{C}}\text{HOC}_2\text{H}_5 + \text{C}_6\text{H}_5\text{NO}_2$	$2.5 \times 10^8$		p.r.	D.k.; radical from ethyl ether.	77A100
283.104	$-\text{O}\dot{\text{C}}\text{HCH}_2\text{O}(\text{CH}_2)_2^- + \text{C}_6\text{H}_5\text{NO}_2$	$2.5 \times 10^7$		p.r.	D.k.; radical from dioxane.	77A100
283.117	$\text{CH}_3\dot{\text{C}}\text{OHCO}_2^- + \text{C}_6\text{H}_5\text{NO}_2$	$(6.5 \pm 2) \times 10^7$	6.0	p.r.	N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> lactate ion.	72A018
283.123	$\cdot\text{CH}(\text{OH})_2 + \text{C}_6\text{H}_5\text{NO}_2 \rightarrow$ $\text{HCO}_2^- + 2 \text{H}^+ + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}_2^-$	$1.9 \times 10^9$	5.8	p.r.	P.b.k. at 290 nm in N <sub>2</sub> O-satd. soln. contg. $10^{-2}$ mol L <sup>-1</sup> formaldehyde; also condy.	71G424
	$\cdot\text{CH}(\text{OH})\text{O}^- + \text{C}_6\text{H}_5\text{NO}_2 \rightarrow$ $\text{HCO}_2^- + \text{H}^+ + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}_2^-$	$4.5 \times 10^9$	12			
283.147	$\cdot\text{CO}_2^- + \text{C}_6\text{H}_5\text{NO}_2 \rightarrow$ $\text{CO}_2 + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}_2^-$	$(1.0 \pm 0.2) \times 10^9$ $(5.6 \pm 1) \times 10^8$ $(5.8 \pm 0.6) \times 10^8$ $(7.5 \pm 0.7) \times 10^8$ $(4.6 \pm 0.5) \times 10^8$	6-7 ~3 9.4 2.5 0	p.r. p.r.	P.b.k. at 295 nm in N <sub>2</sub> O-satd. soln. contg. $10^{-3}$ mol L <sup>-1</sup> formate. P.b.k.	70G303 73G085
284.	<b><i>o</i>-Nitrobenzoate ion</b>					
284.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2^- \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \dot{\text{N}}\text{O}_2^-\text{C}_6\text{H}_4\text{CO}_2^-$	$5.4 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; at pH 0.8 $k$ was the same $\pm$ 20-30%.	76A111
284.147	$\cdot\text{CO}_2^- + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2^- \rightarrow$ $\text{CO}_2 + \dot{\text{N}}\text{O}_2^-\text{C}_6\text{H}_4\text{CO}_2^-$	$2.4 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate; at pH 0.8 $k$ was the same $\pm$ 20-30%.	76A111
285.	<b><i>m</i>-Nitrobenzoate ion</b>					
285.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2^- \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \dot{\text{N}}\text{O}_2^-\text{C}_6\text{H}_4\text{CO}_2^-$	$9.0 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; at pH 0.8 $k$ was the same $\pm$ 20-30%.	76A111
285.147	$\cdot\text{CO}_2^- + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2^- \rightarrow$ $\text{CO}_2 + \dot{\text{N}}\text{O}_2^-\text{C}_6\text{H}_4\text{CO}_2^-$	$6.3 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate; at pH 0.8 $k$ was the same $\pm$ 20-30%.	76A111
286.	<b><i>p</i>-Nitrobenzoate ion</b>					
286.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2^- \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \dot{\text{N}}\text{O}_2^-\text{C}_6\text{H}_4\text{CO}_2^-$	$2.1 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; at pH 0.8 $k$ was the same $\pm$ 20-30%.	76A111
286.147	$\cdot\text{CO}_2^- + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2^- \rightarrow$ $\text{CO}_2 + \dot{\text{N}}\text{O}_2^-\text{C}_6\text{H}_4\text{CO}_2^-$	$8.0 \times 10^8$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate; at pH 0.8 $k$ was the same $\pm$ 20-30%.	76A111
287.	<b><i>p</i>-Nitrobenzonitrile</b>					
287.111	$\text{R} + \text{NO}_2\text{C}_6\text{H}_4\text{CN}$ (R = Radicals from glucose)	$1.0 \times 10^9$		p.r.		77R167
287a.	<b>Nitro blue tetrazolium</b>					
287a.147	$\cdot\text{CO}_2^- + \text{NBT}^{2+} \rightarrow \text{CO}_2 + \text{NBT}^{\cdot+}$	$(6.4 \pm 0.2) \times 10^9$	10	p.r.	P.b.k. at 405 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	80A085
<i>anti</i> -5-Nitro-2-furaldoxime see Nifuroxime 278.						
288.	<b>5-Nitrofuroate ion</b>					
288.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \dots\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \dot{\text{N}}\text{O}_2^-$	$1.5 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	73G114
289.	<b>2-Nitroimidazole</b>					
289.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NX}^-$	$3.5 \times 10^9$		p.r.	P.b.k.	76A075
290.	<b>4-Nitroimidazole</b>					
290.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NX}^-$	$3.5 \times 10^9$		p.r.	P.b.k.	76A075



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
291.	<b>2-Nitroisophthalate ion</b>					
291.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + (\text{O}_2\text{C})_2\text{C}_6\text{H}_3\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + (\text{O}_2\text{C})_2\text{C}_6\text{H}_3\dot{\text{N}}\text{O}_2^-$	$1.5 \times 10^8$	7	p.r.	P.b.k. at 310 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
292.	<b>Nitromethane</b>					
292.047	$\cdot\text{CF}_3 + \text{CH}_2\text{NO}_2^- \rightarrow$ $\text{CF}_3\text{CH}_2\dot{\text{N}}\text{O}_2^-$	$5 \times 10^8$ (rel.)		$\gamma$ -r.	C.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and CF <sub>3</sub> Cl; rel. to $k(\text{CF}_3 + 2\text{-PrOH}) = 10^5$ ; product radical identified by esr.	72G244
293.	<b>p-Nitroperoxybenzoic acid</b>					
293.075	$\cdot\text{CH}_2\text{OH} + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2\text{OH} \rightarrow$ $\text{CH}_2\text{O} + \text{H}^+ + \dot{\text{N}}\text{O}_2\text{C}_6\text{H}_4\text{CO}_2\text{OH}$ (+ adduct formn.)	$2 \times 10^8$		p.r.	P.b.k. at 330 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 20% e-transfer based on condy. change.	74A078
293.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NO}_2\text{C}_6\text{H}_4\text{CO}_2\text{OH} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \dot{\text{N}}\text{O}_2\text{C}_6\text{H}_4\text{CO}_2\text{OH}$	$3.3 \times 10^9$	5.0	p.r.	P.b.k. at 330 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	74A078
294.	<b>o-Nitrophenol</b>					
294.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HOC}_6\text{H}_4\text{NO}_2 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{HOC}_6\text{H}_4\text{NO}_2^-$	$2.6 \times 10^9$	1	p.r.	P.b.k. in soln. contg. 2-PrOH.	69G270
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{OC}_6\text{H}_4\text{NO}_2 \rightarrow$	$1.4 \times 10^9$	13	p.r.	P.b.k. in soln. contg. 2-PrOH.	69G270
295.	<b>2-Nitropyrrole</b>					
295.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NX}^-$	$2.0 \times 10^9$		p.r.	P.b.k.	76A075
296.	<b>3-Nitropyrrole</b>					
296.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NX}^-$	$2.0 \times 10^9$		p.r.	P.b.k.	76A075
<i>2-Nitroresorcinol See 1,3-Dihydroxy-2-nitrobenzene 178.</i>						
297.	<b>Nitrosobenzene</b>					
297.075	$\cdot\text{CH}_2\text{OH} + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CH}_2\text{O} + \text{C}_6\text{H}_5\dot{\text{N}}\text{OH}$	$3.2 \times 10^9$	7.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> MeOH.	66G433
	$\cdot\text{CH}_2\text{O}^- + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CH}_2\text{O} + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}^-$	$6.8 \times 10^9$	13.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> MeOH.	66G433
297.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CH}_3\text{CHO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{OH}$	$3.9 \times 10^9$	7.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> EtOH.	66G433
	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CH}_3\text{CHO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}^-$	$6.4 \times 10^9$	13.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> EtOH.	66G433
297.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CH}_3\text{CH}_2\text{CHO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{OH}$	$4.0 \times 10^9$	7.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> 1-PrOH; includes $\beta$ -alcohol radical reaction.	66G433
297.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{OH}$	$5.0 \times 10^9$	7.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> 2-PrOH.	66G433
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}^-$	$7.0 \times 10^9$	13.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> 2-PrOH.	66G433
297.079	$\text{CH}_3\text{CH}_2\text{CH}_2\dot{\text{C}}\text{HOH} + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{OH}$	$4.0 \times 10^9$	7.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> 1-BuOH; includes $\beta$ - and $\gamma$ -alcohol radical reaction.	66G433
297.081	$(\text{CH}_3)_2\text{CH}\dot{\text{C}}\text{HOH} + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $(\text{CH}_3)_2\text{CHCHO} + \text{C}_6\text{H}_5\dot{\text{N}}\text{OH}$	$4.0 \times 10^9$	7.0	p.r.	P.b.k. at 450 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> 2-methyl-1-propanol; includes $\beta$ -alcohol radical reaction.	66G433
297.147	$\cdot\text{CO}_2^- + \text{C}_6\text{H}_5\text{NO} \rightarrow$ $\text{CO}_2 + \text{C}_6\text{H}_5\dot{\text{N}}\text{O}^-$	$4.0 \times 10^9$		p.r.	P.b.k. at 450 nm in soln. contg. formate ion.	66G433
<i>p-Nitroso-N,N-dimethylaniline See N,N-Dimethyl-4-nitrosoaniline, 186a.</i>						
298.	<b>2-Nitrothiophene</b>					
298.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NX}^-$	$3.0 \times 10^9$		p.r.	P.b.k.	76A075

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
299. 299.078	<b>3-Nitrothiophene</b> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + NX → (CH <sub>3</sub> ) <sub>2</sub> CO + H <sup>+</sup> + NX <sup>-</sup>	2.0 × 10 <sup>9</sup>		p.r.	P.b.k.	76A075
300. 300.078	<b>o-Nitrotoluene</b> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub> → (CH <sub>3</sub> ) <sub>2</sub> CO + H <sup>+</sup> + CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub> <sup>-</sup>	4.8 × 10 <sup>8</sup>	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	76A020
301. 301.078	<b>5-Nitrouracil</b> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + NX	(7 ± 3.5) × 10 <sup>8</sup>		p.r.		73A150
302. 302.016 302.078	<b>Norpseudopelletierine-N-oxy</b> ·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + NPPN (CH <sub>3</sub> ) <sub>2</sub> ĊOH + NPPN	(5.1 ± 0.5) × 10 <sup>8</sup> (8.1 ± 0.8) × 10 <sup>8</sup>		p.r. p.r.		71G061 71G061
303. 303.078	<b>Orotate ion</b> (CH <sub>3</sub> ) <sub>2</sub> ĊOH + orotate	(1 ± 0.5) × 10 <sup>8</sup>		p.r.		73A150
<i>Paraquat See 1,1'-Dimethyl-4,4'-bipyridinium ion 183.</i>						
304. 304.075	<b>Penicillamine</b> ·CH <sub>2</sub> OH + RSH → CH <sub>3</sub> OH + RS· (→ RSSR <sup>-</sup> )	(1.1 ± 0.1) × 10 <sup>8</sup>		p.r.	P.b.k. at 450 nm in soln. contg. 0.5 or 1 mol L <sup>-1</sup> MeOH.	73A073
305. 305.001	<b>3-Pentanone</b> ·CH <sub>3</sub> + C <sub>2</sub> H <sub>5</sub> COC <sub>2</sub> H <sub>5</sub> → CH <sub>4</sub> + CH <sub>3</sub> ĊHCOC <sub>2</sub> H <sub>5</sub>	1.4 × 10 <sup>4</sup>	~1	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
306. 306.075	<b>1,10-Phenanthroline</b> ·CH <sub>2</sub> OH + phen ·CH <sub>2</sub> O <sup>-</sup> + phen	<10 <sup>6</sup> <10 <sup>6</sup> <1 × 10 <sup>7</sup>	7 13 ~13	p.r. p.r.	P.b.k. in soln. contg. 0.2 mol L <sup>-1</sup> MeOH. P.b.k. at 490 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH.	79A148 80A115
306.076	CH <sub>3</sub> ĊHO <sup>-</sup> + phen → ·phenH	(8 ± 2) × 10 <sup>8</sup>	~13	p.r.	P.b.k. at 490 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> EtOH.	80A115
306.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + phen → ·phenH	(3.2 ± 0.2) × 10 <sup>9</sup>	1	p.r.	P.b.k. at 520 nm in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	80A115
		1.9 × 10 <sup>9</sup>	3.0	p.r.	P.b.k. in Ar-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	79A148
		1 × 10 <sup>8</sup>	5.5	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	80A115
		<1 × 10 <sup>7</sup>	8			
		1.2 × 10 <sup>7</sup>	7		P.b.k. in soln. contg. 2-PrOH and acetone.	79A148
	(CH <sub>3</sub> ) <sub>2</sub> ĊO <sup>-</sup> + phen → ·phenH	6.0 × 10 <sup>8</sup>	13		P.b.k. in Ar-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	79A148
		5.7 × 10 <sup>8</sup>	13	p.r.	P.b.k.	79A305
		(3.0 ± 0.5) × 10 <sup>9</sup>	~13	p.r.	P.b.k. at 490 nm in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	80A115
307. 307.078	<b>Phenazine</b> (CH <sub>3</sub> ) <sub>2</sub> ĊO <sup>-</sup> + Pz → (CH <sub>3</sub> ) <sub>2</sub> CO + ·PzH	3 × 10 <sup>9</sup>	13	p.r.	p.b.k.	79A305
308. 308.056	<b>Phenosafarine</b> NH <sub>2</sub> ĊHCO <sub>2</sub> <sup>-</sup> + dye	1.9 × 10 <sup>9</sup> (uncor.)	7	p.r.	D.k. at 520 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 66% e-transfer.	73A078
308.075	·CH <sub>2</sub> OH + dye	1.2 × 10 <sup>9</sup> (uncor.)	7	p.r.	D.k. at 520 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 22% e-transfer.	73A078
308.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + dye	3.2 × 10 <sup>9</sup>	7	p.r.	D.k. at 520 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 82% e-transfer.	73A078

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
309.	<b>Phenoxide ion</b>					
309.020	$\cdot\text{CH}_2\text{CHO} + \text{C}_6\text{H}_5\text{O}^- (+ \text{H}_2\text{O}) \rightarrow$ $\text{CH}_3\text{CHO} + \text{C}_6\text{H}_5\text{O}\cdot + \text{OH}^-$	$4.3 \times 10^6$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
310.	<b><i>o</i>-Phenylenediamine</b>					
310.020	$\cdot\text{CH}_2\text{CHO} + \text{H}_2\text{NC}_6\text{H}_4\text{NH}_2 \rightarrow$ $\text{CH}_3\text{CHO} + \text{H}_2\text{NC}_6\text{H}_4\dot{\text{N}}\text{H}$	$7.3 \times 10^7$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
311.	<b><i>p</i>-Phenylenediamine</b>					
311.020	$\cdot\text{CH}_2\text{CHO} + \text{H}_2\text{NC}_6\text{H}_4\text{NH}_2 \rightarrow$ $\text{CH}_3\text{CHO} + \text{H}_2\text{NC}_6\text{H}_4\dot{\text{N}}\text{H}$	$4.0 \times 10^8$	$\sim 11.5$	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol.	79A051
312.	<b>Promethazine</b>					
312.048	$\cdot\text{CCl}_3 + \text{PZ}$	slow	12	p.r.	Suggested that fast reaction is with $\text{CCl}_3\dot{\text{O}}_2$ and not $\cdot\text{CCl}_3$ .	78B128 80A053
313.	<b>1-Propanol</b>					
313.047	$\cdot\text{CF}_3 + \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	$(4.4 \pm 0.7) \times 10^4$ (rel.)	9–10	$\gamma$ -r.	C.k. with addn. of $\text{CF}_3$ to propylene obs. $G(\text{CF}_3\text{H})$ ; rel. to $k(\text{CF}_3 + \text{HCO}_2^-)$ $= 3.4 \times 10^5$ .	70G407
314.	<b>2-Propanol</b>					
314.001	$\cdot\text{CH}_3 + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{CH}_4 + (\text{CH}_3)_2\dot{\text{C}}\text{OH}$	$3.4 \times 10^3$ (rel.)		$\gamma$ -r.	C.k. with O <sub>2</sub> ; radical from MeI; rel. to $k(\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
		$\geq (1.6 \pm 0.3) \times 10^3$ (rel.)	1	phot.	Est. from effect of addn. of 2-PrOH on CH <sub>4</sub> and C <sub>2</sub> H <sub>6</sub> yields in photolysis of $\text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_3^{2+}$ .	71F579
314.015	$\cdot\text{CH}_2\text{CHOHCH}_3 + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $(\text{CH}_3)_2\text{CHOH} + (\text{CH}_3)_2\dot{\text{C}}\text{OH}$	$> 53 \pm 10$		$\gamma$ -r.	Est. from rel. rates and $G(-\text{H}_2\text{O}_2)$ in soln. contg. H <sub>2</sub> O <sub>2</sub> and 2-PrOH.	70G104
314.047	$\cdot\text{CF}_3 + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{CHF}_3 + (\text{CH}_3)_2\dot{\text{C}}\text{OH}$	$(9.2 \pm 0.9) \times 10^4$ (rel.)	9–10	$\gamma$ -r.	C.k. with addn. of $\cdot\text{CF}_3$ to propylene; obs. $G(\text{CHF}_3)$ ; rel. to $k(\cdot\text{CF}_3 + \text{HCO}_2^-)$ $= 3.4 \times 10^5$ .	70G407
314.048	$\cdot\text{CCl}_3 + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{CCl}_3^- + \text{H}^+ + (\text{CH}_3)_2\dot{\text{C}}\text{OH}$	$\sim 10^4$		p.r.	Reaction proposed to explain initial increase of abs. of PNAP <sup>-</sup> in soln. contg. CCl <sub>4</sub> -PNAP-2-PrOH-acetone.	73A140
	$\cdot\text{CCl}_3 + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{CHCl}_3 + (\text{CH}_3)_2\dot{\text{C}}\text{OH}$	79		$\gamma$ -r.	Calcd. from dependence of $G(\text{Cl}^-)$ on dose rate in soln. contg. CCl <sub>4</sub> and 2-PrOH assuming $2k(\cdot\text{CCl}_3 + \cdot\text{CCl}_3) = 10^9$ .	71G778
315.	<b>Propionic acid</b>					
315.001	$\cdot\text{CH}_3 + \text{CH}_3\text{CH}_2\text{CO}_2\text{H} \rightarrow$ $\text{CH}_4 + \text{CH}_3\dot{\text{C}}\text{HCO}_2\text{H}$	$3.0 \times 10^3$	$\sim 1$	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
316.	<b>Propionitrile</b>					
316.001	$\cdot\text{CH}_3 + \text{CH}_3\text{CH}_2\text{CN} \rightarrow$ $\text{CH}_4 + \text{CH}_3\dot{\text{C}}\text{HCN}$	$1.2 \times 10^3$	$\sim 1$	chem.	Radical from dimethyl sulfoxide in Ti <sup>III</sup> -H <sub>2</sub> O <sub>2</sub> soln.; est. from esr meas. and values for competing reactions.	75D188
317.	<b>Propylene</b>					
317.001	$\cdot\text{CH}_3 + \text{CH}_3\text{CH}=\text{CH}_2 \rightarrow \text{addn.}$	$5.3 \times 10^3$ (rel.)		$\gamma$ -r.	C.k. with 2-PrOH in soln. contg. MeI obs. $G(\text{CH}_4)$ ; rel. to $k(\text{CH}_3 + \text{O}_2) = 4.7 \times 10^9$ .	67G041
317.047	$\cdot\text{CF}_3 + \text{CH}_3\text{CH}=\text{CH}_2 \rightarrow \text{addn.}$	$(7.2 \pm 0.4) \times 10^7$ (rel.)	9–10	$\gamma$ -r.	C.k. with 2-PrOH; radical from CF <sub>3</sub> Cl; obs. $G(\text{CF}_3\text{H})$ by H abstr. from formate rel. to $k(\text{CF}_3 + \text{HCO}_2^-) = 3.4 \times 10^5$ .	70G407
<i>1,1'-Propylene-2,2'-bipyridinium ion See Trimethylene-2,2'-bipyridinium ion 354.</i>						
318.	<b>Pteridine</b>					
318.075	$\cdot\text{CH}_2\text{OH} + \text{P}$	$3.6 \times 10^8$	6.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. MeOH.	76A060



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
319.	<b>Pterin</b>					
319.075	$\cdot\text{CH}_2\text{OH} + \text{PtH}_2^+$	$9.0 \times 10^7$ (uncor.)	0.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln.	76A060
	$\cdot\text{CH}_2\text{OH} + \text{PtH}$	$\leq 10^7$	7.0		contg. MeOH; $\sim 40\%$ $e$ -transfer	
	$\cdot\text{CH}_2\text{O}^- + \text{Pt}^-$	$6.0 \times 10^8$	13.0		at pH 0.8, $\sim 100\%$ at pH 13.	
319.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{PtH}$	$3.7 \times 10^7$ (uncor.)	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln.	76A060
	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{Pt}^-$	$1.2 \times 10^9$	13.0		contg. EtOH; $\sim 40\%$ $e$ -transfer at pH 7, $\sim 100\%$ at pH 13.	
319.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PtH}_2^+$	$2.0 \times 10^9$	0.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln.	76A060
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PtH}$	$4.5 \times 10^8$	7.0		contg. 2-PrOH; $\sim 100\%$ $e$ -transfer.	
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pt}^-$	$\leq 10^7$	9.4			
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Pt}^-$	$1.5 \times 10^9$	13.0			
319.147	$\cdot\text{CO}_2^- + \text{PtH}$	$4.6 \times 10^8$	7.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln.	76A060
	$\cdot\text{CO}_2^- + \text{Pt}^-$	$\leq 10^7$	9.5–13.0		contg. formate ion; 100% $e$ -transfer at pH 7.	
320.	<b>Purine</b>					
320.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PH}_2^+$	$2.7 \times 10^9$	$\sim 0$	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg.	75A060
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PH}$	$< 10^7$	6.0		1–2 mol L <sup>-1</sup> 2-PrOH.	
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{P}^-$	$< 10^7$	13.3			
320.147	$\cdot\text{CO}_2^- + \text{PH}$	$< 10^7$	6.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln.	75A060
321.	<b>Pyrazine</b>					
321.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PzH}^+$	$2.8 \times 10^9$	$\sim 0$	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg.	74A127
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pz}$	$\leq 10^7$	5.0, 11.0		2-PrOH; $k \approx 5 \times 10^9$ in 70% HClO <sub>4</sub> soln.	
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Pz}$	$1.7 \times 10^9$	13.6			
321.147	$\cdot\text{CO}_2^- + \text{Pz}$	$\leq 10^7$	5.0, 11.0	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate ion.	74A127
322.	<b>Pyrazinecarboxylic acid</b>					
322.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PzCO}_2\text{H} \rightarrow (\text{CH}_3)_2\dot{\text{C}}\text{O} + \cdot\text{Pz(H)CO}_2\text{H}$	$\sim 8.5 \times 10^8$		p.r.	P.b.k.	78A222
323.	<b>Pyrene</b>					
323.075	$\cdot\text{CH}_2\text{O}^- + \text{Py}$	no reaction	13	p.r.	Buildup of Py <sup>-</sup> in micellar soln. contg. 0.2 mol L <sup>-1</sup> MeOH after phot. attributed to triplet pyrene; $k(\cdot\text{CH}_2\text{O}^- + {}^3\text{Py}) = 1.8 \times 10^{10}$ .	76A062
323.076	$\text{CH}_3\dot{\text{C}}\text{HO}^- + \text{Py} \rightarrow \text{CH}_3\dot{\text{C}}\text{HO} + \text{Py}^-$	$1.7 \times 10^8$	13	p.r.	P.b.k. at 495 nm in soln. contg. 0.2 mol L <sup>-1</sup> EtOH and 10 <sup>-3</sup> mol L <sup>-1</sup> pyrene solubilized by 5 x 10 <sup>-3</sup> mol L <sup>-1</sup> hexadecyl trimethylammonium bromide; after photolysis $k(\text{R} + {}^3\text{Py}) = 8 \times 10^9$ .	76A062
323.078	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Py} \rightarrow (\text{CH}_3)_2\dot{\text{C}}\text{O} + \text{Py}^-$	$2.3 \times 10^9$	13	p.r.	P.b.k. at 495 nm in soln. contg. 2 x 10 <sup>-4</sup> mol L <sup>-1</sup> pyrene, 0.2 mol L <sup>-1</sup> 2-PrOH, 0.1 mol L <sup>-1</sup> acetone and 5 x 10 <sup>-3</sup> mol L <sup>-1</sup> hexadecyl trimethyl-ammonium bromide; after photolysis $k(\text{R} + {}^3\text{Py}) = 2.3 \times 10^9$ .	76A062
323.147	$\cdot\text{CO}_2^- + \text{Py}$	no reaction	13	p.r.	Buildup of 495 nm abs. in micellar soln. (CO <sub>2</sub> -satd.) only after photolysis; $k(\cdot\text{CO}_2^- + {}^3\text{Py}) = 5 \times 10^9$ ; calcd. $k(\cdot\text{CO}_2^- + \text{Py}^-) = 1.8 \times 10^{10}$ from d.k. at 495 nm in micellar CO <sub>2</sub> -free soln. contg. formate ion.	76A062
324.	<b>Pyridazine</b>					
324.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PdH}^+ \rightarrow (\text{CH}_3)_2\dot{\text{C}}\text{O} + \cdot\text{PdH}_2^+$	$2.6 \times 10^9$	$\sim 0$	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH.	74A127
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pd}$	$\leq 10^7$	5.0, 11.0			74A127
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Pd} \rightarrow (\text{CH}_3)_2\dot{\text{C}}\text{O} + \cdot\text{PdH} + \text{OH}^-$	$2.1 \times 10^9$	13.6			74A127

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
324.147	$\cdot\text{CO}_2^- + \text{Pd}$	$<10^7$	5.0, 11.0	p.r.	$<10\%$ $e$ -transfer.	74A127
325. 325.078	<b>4-Pyridinecarboxaldoxime</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{HN}^+\text{C}_5\text{H}_4\text{CH}=\text{NOH}$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{NC}_5\text{H}_4\text{CH}=\text{NO}^-$	$1.7 \times 10^9$ $<10^7$ $1.7 \times 10^8$	acid 7 13.3	p.r.	P.b.k. (pyridinyl radical) in $\text{N}_2\text{O}$ -satd. soln. contg. 2-PrOH; $e$ -transfer in strong acid or base.	76A182
326. 326.078	<b>2-Pyridinecarboxaldoxime methochloride</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CH}_3\text{N}^+\text{C}_5\text{H}_4\text{CH}=\text{NOH}$	$6 \times 10^8$	3,7, 11	p.r.	P.b.k.; $e$ -transfer.	76A182
<i>3-Pyridinecarboxamide See Nicotinamide 275.</i>						
<i>4-Pyridinecarboxamide See Isonicotinamide 244.</i>						
<i>3-Pyridinecarboxylic acid See Nicotinic acid 277.</i>						
<i>4-Pyridinecarboxylic acid See Isonicotinic acid 245.</i>						
327. 327.078	<b>Pyridoxal-5-phosphate</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{P} \rightarrow \cdot\text{P}$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{P} \rightarrow \cdot\text{P}$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{P} \rightarrow \cdot\text{P}$	$5.8 \times 10^8$ $1.3 \times 10^8$ $2.9 \times 10^8$	1.0 5.6 10.0, 13.3	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	75A024
328. 328.078	<b>Pyridoxine</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PH}$	$<10^7$	1,7, 13.5	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 1 mol L <sup>-1</sup> 2-PrOH.	75A024
329. 329.078	<b>Pyrimidine</b> $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{PmH}^+ \rightarrow$ $(\text{CH}_3)_2\dot{\text{C}}\text{O} + \cdot\text{PmH}_2^+$ $(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Pm}$ $(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{Pm}$	$2.2 \times 10^9$ $<10^7$ $<10^7$	$\sim 0$ 5.0, 11.0 13.6	p.r. p.r. p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 2-PrOH. P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 2-PrOH. P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 2-PrOH.	74A127 74A127 74A127
329.147	$\cdot\text{CO}_2^- + \text{Pm}$	$<10^7$	5.0, 11.0	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. formate ion; $<10\%$ $e$ -transfer.	74A127
330. 330.113	<b>Pyruvate ion</b> $\cdot\text{A}^- + \text{CH}_3\text{COCO}_2^-$ ( $\cdot\text{A}^-$ = ascorbate radical)	$<10$	8.6	p.r.	D.k. at 360 nm in $\text{N}_2\text{O}$ -satd. soln. contg. ascorbate ion.	75A240
331. 331.076	<b>Quinoxaline</b> $\text{CH}_3\dot{\text{C}}\text{HOH} + \text{Qx} \rightarrow$ $\text{CH}_3\text{CHO} + \cdot\text{QxH}$	$6.5 \times 10^7$ (uncor.)	6.5	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. EtOH; 70% $e$ -transfer.	74A127
331.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{QxH}^+ \rightarrow$ $(\text{CH}_3)_2\dot{\text{C}}\text{O} + \cdot\text{QxH}_2^+$	$3.7 \times 10^9$	$\sim 0$	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 2 mol L <sup>-1</sup> $\text{HClO}_4$ and 2 mol L <sup>-1</sup> 2-PrOH; $k \approx$ $7.0 \times 10^8$ at 70% $\text{HClO}_4$ .	74A127
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Qx} \rightarrow$ $(\text{CH}_3)_2\dot{\text{C}}\text{O} + \cdot\text{QxH}$	$1.6 \times 10^8$	6.5	p.r.	P.b.k. in $\text{N}_2\text{O}$ -satd. soln. contg. 2-PrOH.	74A127
<i>Resorcinol See 1,3-Dihydroxybenzene 177.</i>						
332. 332.147	<b>Rhodamine B</b> $\cdot\text{CO}_2^- + \text{RhB} \rightarrow \text{CO}_2 + \text{RhB}_{\text{red}}$	$(1.8 \pm 0.5) \times 10^8$		p.r.	D.k. at 510 nm as well as p.b.k. at 410 nm in soln. contg. 0.1 mol L <sup>-1</sup> formate ion.	67E053
333. 333.056	<b>Riboflavin</b> $\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{RF}$	$(2.7 \pm 0.3) \times 10^9$	8.0	p.r.	P.b.k. at 560 nm in $\text{N}_2\text{O}$ -satd. soln. contg. glycine; $e$ -transfer.	73A104

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
333.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + RF	(2.3 ± 0.2) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 560 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; <i>e</i> -transfer.	73A104
333.115	ĊHOHCO <sub>2</sub> <sup>-</sup> + RF	(9.3 ± 0.9) × 10 <sup>8</sup>	7.0	p.r.	P.b.k. at 560 nm in N <sub>2</sub> O-satd. soln. contg. glycolate ion; <i>e</i> -transfer.	73A104
333.147	·CO <sub>2</sub> <sup>-</sup> + RF	1.4 × 10 <sup>9</sup>	alk	p.r.	D.k. at 460 nm as well as p.b.k. at 560 nm (semiquinone) in N <sub>2</sub> O-satd. soln. contg. formate ion.	69G238
		3.6 × 10 <sup>9</sup>	5.9			
		3.6 × 10 <sup>9</sup>	3	p.r.	D.k. at 420 nm in N <sub>2</sub> O-satd. soln. contg. formate ion.	69G238
		3.0 × 10 <sup>9</sup>	H <sub>a</sub> = -2			
		(1.7 ± 0.2) × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 560 nm in N <sub>2</sub> O-satd. soln. contg. formate ion; <i>e</i> -transfer.	73A104
334.	<b>Riboflavine 5'-phosphate (Flavine mononucleotide)</b>					
334.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + FMN	(1 ± 0.5) × 10 <sup>9</sup>		p.r.		73A150
335.	<b>Ribonuclease</b>					
335.045	·CHCl <sub>2</sub> + RNase	6 × 10 <sup>8</sup>	11	p.r.	P.b.k. in soln. contg. CHCl <sub>3</sub> and <i>tert</i> -BuOH.	73A140
336.	<b>Safranin T</b>					
336.056	NH <sub>2</sub> ĊHCO <sub>2</sub> <sup>-</sup> + Dye	1.6 × 10 <sup>9</sup> (uncor.)	7	p.r.	D.k. at 520 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 60% <i>e</i> -transfer.	73A078
336.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + Dye	2.8 × 10 <sup>9</sup>	7	p.r.	D.k. at 520 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 86% <i>e</i> -transfer.	73A078
337.	<b>meso-Tetra(4-carboxyphenyl)porphyrin</b>					
337.078	(CH <sub>3</sub> ) <sub>2</sub> ĊOH + H <sub>2</sub> TCPP → H <sub>2</sub> TCPP <sup>-</sup>	(9 ± 1) × 10 <sup>8</sup>	7-11	p.r.	P.b.k. at 460 and 700 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> 2-PrOH.	79A143
338.	<b>Tetrachlorobenzoquinone (Chloranil)</b>					
338.076	CH <sub>3</sub> ĊHOH + O=C <sub>6</sub> Cl <sub>4</sub> =O → <i>e</i> -transfer	(2.9 to 3.4) × 10 <sup>9</sup>		p.r.	P.b.k. in CTAB or NaLS micelles; soln. contains 0.1 mol L <sup>-1</sup> EtOH; cor. for <i>e</i> <sub>aq</sub> <sup>-</sup> reaction.	76A104
339.	<b>1,2,4,5-Tetracyanobenzene</b>					
339.076	CH <sub>3</sub> ĊHOH + C <sub>6</sub> H <sub>2</sub> (CN) <sub>4</sub> → <i>e</i> -transfer	(3.3 to 4.1) × 10 <sup>9</sup>		p.r.	P.b.k. in CTAB or NaLS micelles; soln. contains 0.1 mol L <sup>-1</sup> EtOH; cor. for <i>e</i> <sub>aq</sub> <sup>-</sup> reaction.	76A104
340.	<b>Tetrahydrofuran</b>					
340.048	·CCl <sub>3</sub> + C <sub>4</sub> H <sub>8</sub> O	57			Calcd. from dependence of <i>G</i> (Cl <sup>-</sup> ) on dose rate in soln. contg. CCl <sub>4</sub> and tetrahydrofuran.	71G778
<i>Tetramethyl-p-benzoquinone See Duroquinone 198.</i>						
<i>N,N,N',N'-Tetramethyldiazenedicarboxamide See Diamide 170.</i>						
341.	<b>1,1'-Tetramethylene-2,2'-bipyridinium ion</b>					
341.147	·CO <sub>2</sub> <sup>-</sup> + BP <sup>2+</sup> → CO <sub>2</sub> + ·BP <sup>-</sup>	9.0 × 10 <sup>9</sup>	7.0	p.r.	P.b.k. at 680 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate.	76A169
		7 × 10 <sup>9</sup>	6.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate ion.	78A321
342.	<b>2,2,6,6-Tetramethyl-4-oxo-1-piperidinyloxy (TAN)</b>					
342.004	<i>c</i> -Ċ <sub>3</sub> H <sub>9</sub> + TAN	(4.0 ± 0.4) × 10 <sup>8</sup>	acid	p.r.	Condy. changes in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> cyclopentane; addn. reaction.	76A067
342.016	·CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> OH + TAN	(2.8 ± 0.3) × 10 <sup>8</sup>	2.4	p.r.	Condy. changes in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH; addn. reaction.	76A067



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
342.075	$\cdot\text{CH}_2\text{OH} + \text{TAN}$	$7.2 \times 10^8$ (rel.)	5-6	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. 1 mol L <sup>-1</sup> MeOH; $k(\cdot\text{CH}_2\text{OH} + \text{ferricyanide}) = 4.0 \times 10^9$ .	71G618
		$(7.2 \pm 0.7) \times 10^8$	acid	p.r.	Condy. changes in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH; addn. reaction.	76A067
342.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{TAN}$	$6.4 \times 10^8$ (rel.)	5-6	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. EtOH assuming $k(\text{R} + \text{ferricyanide}) = 5.3 \times 10^9$ .	71G618
		$(4.0 \pm 0.4) \times 10^8$	acid	p.r.	Condy. changes in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; $e$ -transfer.	76A067
342.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{TAN}$	$3.9 \times 10^8$ (rel.)	5-6	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH, rel. to $k(\text{R} + \text{ferricyanide}) = 4.7 \times 10^9$ ; $k = 4.7 \times 10^8$ rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	71G618
		$(4.3 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; addn. reaction.	76A067
342.105	$\text{R} + \text{TAN}$ (R = Radicals from polyethylene oxide)	$1.9 \times 10^8$ (rel.)	5-6	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. PEO(20000), rel. to $k(\text{R} + \text{ferricyanide}) = 2.1 \times 10^9$ ; similar values were obtained for PEO(600) and PEO(200).	71G618
342.108	$\text{R} + \text{TAN}$ (R = Radicals from deoxy-ribose)	$3.9 \times 10^8$ (rel.)	5-6	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. deoxyribose, rel. to $k(\text{R} + \text{ferricyanide}) = 2.8 \times 10^9$ .	71G618
342.111	$\text{R} + \text{TAN}$ (R = Radicals from glucose)	$<1 \times 10^8$ (rel.)	5-6	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. glucose, rel. to $k(\text{R} + \text{ferricyanide}) = 1.9 \times 10^9$ .	71G618
		$(5.9 \pm 0.6) \times 10^7$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> glucose.	76A067
342.147	$\cdot\text{CO}_2^- + \text{TAN}$	$7.0 \times 10^8$	7-8	p.r.	P.b.k. at 310 nm in N <sub>2</sub> O-satd. soln. contg. formate.	71G618
		$5.4 \times 10^8$ (rel.)	7-8	p.r.	C.k. in N <sub>2</sub> O-satd. soln. contg. formate, rel. to $k(\cdot\text{CO}_2^- + \text{ferricyanide}) = 1.06 \times 10^9$ .	71G618
343.	<b>2,2,6,6-Tetramethyl-4-hydroxy-1-piperidinyloxy (TMPN)</b>					
343.004	$c\text{-}\dot{\text{C}}_5\text{H}_9 + \text{TMPN}$	$(4.3 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> cyclopentane; addn. reaction.	76A067
343.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{TMPN}$	$(1.5 \pm 0.2) \times 10^8$	2.6	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> <i>tert</i> -BuOH; addn. reaction.	76A067
343.075	$\cdot\text{CH}_2\text{OH} + \text{TMPN}$	$(4.4 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH; addn. reaction.	76A067
343.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{TMPN}$	$(4.9 \pm 0.5) \times 10^8$	acid	p.r.	Condy change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; $e$ -transfer.	76A067
343.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{TMPN}$	$(3.6 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; $e$ -transfer.	76A067
343.111	$\text{R} + \text{TMPN}$ (R = Radicals from glucose)	$(4.9 \pm 0.5) \times 10^7$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> glucose; addn. reaction.	76A067
344.	<b>N,N,N',N'-Tetramethyl-p-phenylenediamine</b>					
344.020	$\cdot\text{CH}_2\text{CHO} + \text{TMPD} (+ \text{H}_2\text{O}) \rightarrow \text{CH}_3\text{CHO} + \text{TMPD}\cdot + \text{OH}^-$	$2.0 \times 10^9$	~11.5	p.r.	Soln. contg. N <sub>2</sub> O and ethylene glycol; radical cation ident. by abs. spectra.	70A051
345.	<b>2,2,5,5-Tetramethyl-1-pyrrolidinyloxy-3-carboxamide (NX-s)</b>					
345.001	$\cdot\text{CH}_3 + \text{NX-s}$	$(7.5 \pm 0.8) \times 10^8$	3.0	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> -10 <sup>-1</sup> mol L <sup>-1</sup> dimethyl sulfoxide; addn. reaction.	76A152

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
345.004	$c\text{-}\dot{\text{C}}_5\text{H}_9 + \text{NX-s}$	$(3.5 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> cyclopentane; addn. reaction.	76A152
345.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{NX-s}$	$(4.7 \pm 0.5) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> ethylene; addn. reaction.	76A152
345.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{NX-s}$	$(1.8 \pm 0.2) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> <i>tert</i> -BuOH; addn. reaction.	76A152
345.075	$\cdot\text{CH}_2\text{OH} + \text{NX-s}$	$(4.6 \pm 0.5) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH; addn. reaction.	76A152
345.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{NX-s}$	$(4.3 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; <i>e</i> -transfer.	76A152
345.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX-s}$	$(3.3 \pm 0.3) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; <i>e</i> -transfer.	76A152
345.111	$\text{R} + \text{NX-s}$ (R = Radicals from glucose)	$(5.1 \pm 0.5) \times 10^7$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> glucose; <i>e</i> -transfer.	76A152
346.	<b>2,2,5,5-Tetramethyl-3-pyrrolin-1-yloxy-3-carboxamide (NX-u)</b>					
346.001	$\cdot\text{CH}_3 + \text{NX-u}$	$(7.8 \pm 0.8) \times 10^8$	acid.	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> dimethyl sulfoxide; addn. reaction.	76A152
346.004	$c\text{-}\dot{\text{C}}_5\text{H}_9 + \text{NX-u}$	$(3.6 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> cyclopentane; addn. reaction.	76A152
346.014	$\cdot\text{CH}_2\text{CH}_2\text{OH} + \text{NX-u}$	$(4.8 \pm 0.5) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> ethylene; addn. reaction.	76A152
346.016	$\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \text{NX-u}$	$(2.0 \pm 0.2) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 10 <sup>-2</sup> –10 <sup>-1</sup> mol L <sup>-1</sup> <i>tert</i> -BuOH; addn. reaction.	76A152
346.075	$\cdot\text{CH}_2\text{OH} + \text{NX-u}$	$(3.5 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH; addn. reaction.	76A152
346.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{NX-u}$	$(6.2 \pm 0.6) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; <i>e</i> -transfer.	76A152
346.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{NX-u}$	$(3.6 \pm 0.4) \times 10^8$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; <i>e</i> -transfer.	76A152
346.111	$\text{R} + \text{NX-u}$ (R = Radicals from glucose)	$(4.3 \pm 0.4) \times 10^7$	acid	p.r.	Condy. change in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> glucose; <i>e</i> -transfer.	76A152
347.	<b>Tetranitromethane</b>					
347.075	$\cdot\text{CH}_2\text{OH} + \text{C}(\text{NO}_2)_4 \rightarrow$ $\text{CH}_2\text{O} + \text{NO}_2 + \text{H}^+ + \text{C}(\text{NO}_2)_3^-$	$(5.0 \pm 1) \times 10^9$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> MeOH.	64G133
347.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{C}(\text{NO}_2)_4 \rightarrow$ $\text{CH}_3\text{CHO} + \text{NO}_2 + \text{H}^+ + \text{C}(\text{NO}_2)_3^-$	$5.6 \times 10^9$	~1	p.r.	P.b.k. at 366 nm in soln. contg. 0.5 mol L <sup>-1</sup> EtOH.	65G183
		$\sim 3.5 \times 10^9$		p.r.	P.b.k. in solns. contg. 0.1 mol L <sup>-1</sup> EtOH and either 0.1 mol L <sup>-1</sup> Na dodecyl sulfate or 0.02 mol L <sup>-1</sup> dodecyltrimethylammonium chloride.	76A104
347.077	$\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HOH} + \text{C}(\text{NO}_2)_4 \rightarrow$ $\text{CH}_3\text{CH}_2\text{CHO} + \text{NO}_2 + \text{H}^+ + \text{C}(\text{NO}_2)_3^-$	$(4.7 \pm 1) \times 10^9$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 1-PrOH.	64G133
347.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}(\text{NO}_2)_4 \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ + \text{NO}_2 + \text{C}(\text{NO}_2)_3^-$	$(5.0 \pm 1) \times 10^9$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH.	64G133
347.092a	$\cdot\text{CH}_2\text{OCH}_3 + \text{C}(\text{NO}_2)_4 \rightarrow$ $\text{CH}_3\text{OCH}_2\text{O}\dot{\text{N}}^+(\text{O}^-)\text{C}(\text{NO}_2)_3 \longleftrightarrow$ $\text{CH}_3\text{OCH}_2\text{ON}(\dot{\text{O}})\text{C}(\text{NO}_2)_3$	$6 \times 10^9$		p.r.	P.b.k. at 300 nm in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> methyl ether; adduct decomposes to nitroform anion, $k = 1.1 \times 10^4 \text{ s}^{-1}$ .	80A071
347.093	$\text{CH}_3\text{CH}_2\text{O}\dot{\text{C}}\text{HCH}_3 + \text{C}(\text{NO}_2)_4 \rightarrow$ $\text{C}(\text{NO}_2)_3^-$	$\sim 4 \times 10^9$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. ethyl ether.	80A071
347.093a	$\cdot\text{CH}_2\text{OC}(\text{CH}_3)_3 + \text{C}(\text{NO}_2)_4 \rightarrow$ $\text{C}(\text{NO}_2)_3^-$	$\sim 3 \times 10^9$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. methyl <i>tert</i> -butyl ether.	80A071

TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
347.094a	$\cdot\text{CH}_2\text{OCH}_2\text{OCH}_3 + \text{C}(\text{NO}_2)_4 \rightarrow \text{CH}_3\text{OCH}_2\text{OCH}_2\text{ON}(\dot{\text{O}})\text{C}(\text{NO}_2)_3$	$5 \times 10^9$	4–13	p.r.	P.b.k. at 300 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> dimethoxymethane; adduct decomposes to nitroform anion ( $\lambda$ 350 nm), $k = 8.2 \times 10^2$ s <sup>-1</sup> .	80A071
347.101	$\text{CH}_2\text{OH}\dot{\text{C}}\text{HOH} + \text{C}(\text{NO}_2)_4$	$1.7 \times 10^9$		p.r.	P.b.k. in soln. contg. 0.2 mol L <sup>-1</sup> ethylene glycol.	73G126
347.102	$\text{CH}_3\text{CHOH}\dot{\text{C}}\text{HOH} + \text{C}(\text{NO}_2)_4$	$3.2 \times 10^9$		p.r.	P.b.k. in soln. contg. 0.2 mol L <sup>-1</sup> 1,2-propanediol.	73G126
347.103	$\text{CH}_3\text{CHOH}\dot{\text{C}}\text{OHCH}_3 + \text{C}(\text{NO}_2)_4$	$3.3 \times 10^9$		p.r.	P.b.k. in soln. contg. 0.2 mol L <sup>-1</sup> 2,3-butanediol.	73G126
347.106	$\text{CH}_2\text{OHCHOH}\dot{\text{C}}\text{HOH} + \text{C}(\text{NO}_2)_4$ (+ $\text{CH}_2\text{OH}\dot{\text{C}}\text{OHCH}_2\text{OH}$ )	$(2.4 \pm 0.3) \times 10^9$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> glycerol.	64G133
347.111	$\text{R} + \text{C}(\text{NO}_2)_4$ (R = Radicals from glucose)	$2.6 \times 10^9$		p.r.	P.b.k. at 366 nm in N <sub>2</sub> O-satd. soln. 0.25 mol L <sup>-1</sup> glucose.	65G183
347.112	$\text{R} + \text{C}(\text{NO}_2)_4$ (R = Radicals from sucrose)	$(7.0 \pm 1) \times 10^8$		p.r.	P.b.k. at 350 nm in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> sucrose.	64G133
		$(8.5 \pm 1) \times 10^8$		p.r.	P.b.k. in soln. contg. 0.1 or 0.01 mol L <sup>-1</sup> sucrose.	65G183
347.147	$\cdot\text{CO}_2^- + \text{C}(\text{NO}_2)_4$	$(4 \pm 1) \times 10^9$		p.r.	P.b.k.; independent of pH between 3 and 7.	70G303
348.	<b>Thiamine</b>					
348.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{TmH}^+ \rightarrow (\text{CH}_3)_2\text{CO} + \cdot\text{TmH}_2^+$	$2.2 \times 10^8$	0.5	p.r.	P.b.k. in soln. contg. 2-PrOH.	77A034
	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Tm} \rightarrow (\text{CH}_3)_2\text{CO} + \cdot\text{TmH}^+$	$1.9 \times 10^8$	6.6	p.r.	P.b.k. in soln. contg. 2-PrOH.	77A034
349.	<b>Thiazole</b>					
349.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{C}_3\text{H}_3\text{SNH}^+ \rightarrow (\text{CH}_3)_2\text{CO} + \cdot\text{C}_3\text{H}_3\text{SNH}_2^+$	$6.2 \times 10^8$	0.8	p.r.	P.b.k. in soln. contg. 2-PrOH.	77A034
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{C}_3\text{H}_3\text{SN}$		13.3	p.r.	No electron transfer.	77A034
350.	<b>Thionine</b>					
350.056	$\text{NH}_2\dot{\text{C}}\text{HCO}_2^- + \text{Dye}$	$3.2 \times 10^9$	8	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. glycine; 85% <i>e</i> -transfer.	73A078
350.075	$\cdot\text{CH}_2\text{OH} + \text{Dye}$	$2.6 \times 10^9$	8	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. MeOH; 86% <i>e</i> -transfer.	73A078
350.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Dye}$	$4.2 \times 10^9$	8	p.r.	D.k. at 600 nm in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; 88% <i>e</i> -transfer.	73A078
351.	<b>Trichloroacetate ion</b>					
351.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CCl}_3\text{CO}_2^-$	$5 \times 10^6$ (rel.)		p.r.	C.k. in soln. contg. 2-PrOH and acetone; rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
	$(\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{CCl}_3\text{CO}_2^-$	$3 \times 10^8$ (rel.)	13	p.r.	C.k. in soln. contg. 2-PrOH and acetone; rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
352.	<b>Trifluoroacetate ion</b>					
352.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{CF}_3\text{CO}_2^-$	$< 1 \times 10^5$ (rel.)		p.r.	C.k. in soln. contg. 2-PrOH and acetone; rel. to $k(\text{R} + \text{PNAP}) = 3.8 \times 10^9$ .	73A140
353.	<b>2,3,5-Trimethylbenzoquinone</b>					
353.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow (\text{CH}_3)_2\text{CO} + \text{H}^+ + \cdot\text{Q}^+$	$3.6 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 1 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
354.	<b>1,1'-Trimethylene-2,2'-bipyridinium ion</b>					
354.147	$\cdot\text{CO}_2^- + \text{BP}^{2+} \rightarrow \text{CO}_2 + \cdot\text{BP}^+$	$7.5 \times 10^9$	7.0	p.r.	P.b.k. at 680 nm in N <sub>2</sub> O-satd. soln. contg. 0.2 mol L <sup>-1</sup> formate ion.	76A169
		$1.1 \times 10^{10}$	6.8	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> formate.	78A321



TABLE 2. Rate constants for reactions of aliphatic radicals with organic solutes in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
355.	<b>Trinitrobenzenesulfonate ion</b>					
355.111	$R + (\text{NO}_2)_3\text{C}_6\text{H}_2\text{SO}_3^-$ (R = Radicals from glucose)	$1.1 \times 10^9$		p.r.		77R167
356.	<b>2,4,6-Trinitrobenzoate ion</b>					
356.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + (\text{NO}_2)_3\text{C}_6\text{H}_2\text{CO}_2^- \rightarrow$ $(\text{CH}_3)_2\text{CO} + \text{H}^+ +$ $(\dot{\text{N}}\text{O}_2)(\text{NO}_2)_2\text{C}_6\text{H}_2\text{CO}_2^-$	$3.9 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $k$ at pH 0.8 within 20–30%.	76A111
356.147	$\cdot\text{CO}_2^- + (\text{NO}_2)_3\text{C}_6\text{H}_2\text{CO}_2^-$	$3.4 \times 10^9$	7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. formate; $k$ at pH 0.8 within 20–30%.	76A111
357.	<b>Trypan blue</b>					
357.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Dye}$	$(3 \pm 1.5) \times 10^9$		p.r.		73A150
358.	<b>Tyrosine</b>					
358.048	$\cdot\text{CCl}_3 + \text{ArO}^-$	slow	12	p.r.	Suggested that fast reaction is $\text{CCl}_3\dot{\text{O}}_2 + \text{ArO}^- \rightarrow \text{ArO}\cdot$ ( $k = 1.6 \times 10^8$ ) and $\cdot\text{CCl}_3$ is less reactive.	78B128 80A053
359.	<b>Ubiquinone</b>					
359.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \cdot\text{Q}^- + \text{H}^+$	$1.9 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 7 mol L <sup>-1</sup> 2-PrOH and 1 mol L <sup>-1</sup> acetone.	73G125
	<i>Vitamin B<sub>5</sub> See Nicotinamide 275.</i>					
	<i>Vitamin B<sub>12</sub> See Cyanocobalamin (under Cobalt(III) ions) 53.</i>					
	<i>Vitamin B<sub>12a</sub> See Hydroxocobalamin (under Cobalt(III) ions) 54.</i>					
	<i>Vitamin B<sub>12r</sub> See Cobalamin (under Cobalt(II) ions) 10.</i>					
360.	<b>Vitamin K<sub>1</sub></b>					
360.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{Q} \rightarrow$ $(\text{CH}_3)_2\text{CO} + \cdot\text{Q}^- + \text{H}^+$	$1.7 \times 10^9$	7	p.r.	P.b.k. in soln. contg. 5 mol L <sup>-1</sup> 2-PrOH and 2 mol L <sup>-1</sup> acetone.	73G125
	<i>Vitamin K<sub>3</sub> See 2-Methyl-1,4-naphthoquinone 262.</i>					

TABLE 3. Rate constants for acid- and base-catalyzed proton transfer reactions of aliphatic radicals in aqueous solution

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
361.	<b>Hydronium ion</b>					
361.075	$\cdot\text{CH}_2\text{OH} + \text{H}_3\text{O}^+ \rightleftharpoons \cdot\text{CH}_2\text{OH}_2^+ + \text{H}_2\text{O}$	$k_f = 1.76 \times 10^8$ $k_r = 3.6 \times 10^7$		chem. phot.	Esr; radicals from Ti(III)-H <sub>2</sub> O <sub>2</sub> -MeOH. Esr; soln. contg. 10% acetone and 5% MeOH.	65D040 66D162
		$k_f = (3.19 \pm 0.04) \times 10^7$		chem.	Esr; radicals from Ti(III)-H <sub>2</sub> O <sub>2</sub> -MeOH at 30.4°C; $E_a = 2.8 \pm 5$ kcal mol <sup>-1</sup> (11.7 kJ mol <sup>-1</sup> ) at 5.7-41°C; $A = 9.5$ s <sup>-1</sup> .	68M058
361.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{H}_3\text{O}^+ \rightleftharpoons (\text{CH}_3)_2\dot{\text{C}}\text{OH}_2^+ + \text{H}_2\text{O}$	$k_f = 7.2 \times 10^7$		phot.	Esr; soln. contg. 10% acetone and 5% 2-PrOH.	66D162
		$k_f = (6.44 \pm 0.30) \times 10^7$		chem.	Esr; radicals from Ti(III)-H <sub>2</sub> O <sub>2</sub> -2-PrOH.	68M058
361.148	$\text{CH}_3\dot{\text{C}}(\text{O})\text{OCH}_3 + \text{H}_3\text{O}^+ \rightarrow \text{CH}_3\dot{\text{C}}(\text{OH})\text{OCH}_3 + \text{H}_2\text{O}$	$1.4 \times 10^{10}$		p.r.	D.k. at 360 nm in soln. contg. methyl acetate and HClO <sub>4</sub> (1-5 x 10 <sup>-5</sup> mol L <sup>-1</sup> ).	78A402
362.	<b>Hydroxide ion</b>					
362.022	$\cdot\text{CH}=\text{CHOH} + \text{OH}^- \rightleftharpoons [\cdot\text{CH}=\text{CHO}^-] + \text{H}_2\text{O}$	$k_f = (1.5 \pm 0.3) \times 10^{10}$	9.4-10	p.r.	D.k. at 260 nm in 10 <sup>-3</sup> mol L <sup>-1</sup> acetylene soln.; hydrolysis product is formylmethyl radical.	78A007
362.060	$\text{CH}_3\text{CONH}\dot{\text{C}}\text{HCOO}^- + \text{OH}^- \rightleftharpoons \text{CH}_3\text{CON}^-\dot{\text{C}}\text{HCOO}^- + \text{H}_2\text{O}$	$k_f = (8 \pm 2) \times 10^8$	>13	e-r.	Est. by esr from pH dependence of line width in N <sub>2</sub> O-satd. soln. contg. acetylglycine; $k_f = 1.6 \times 10^7$ ; ( $pK_a = 14.0$ ).	76D198
362.062	$\text{R}(\text{NH}) + \text{OH}^- \rightleftharpoons \text{R}(\text{N}^-) + \text{H}_2\text{O}$ (R = Radicals from glycine anhydride)	$\sim (8 \pm 2) \times 10^9$	$\sim 9.2$ - 10.5	p.r.	P.b.k. at 300 nm in N <sub>2</sub> O-satd. soln. contg. 4 x 10 <sup>-3</sup> mol L <sup>-1</sup> glycine anhydride.	71G554
362.063	$\text{R}(\text{NH}) + \text{OH}^- \rightleftharpoons \text{R}(\text{N}^-) + \text{H}_2\text{O}$ (R = Radicals from alanine anhydride)	$\sim (1.1 \pm 0.2) \times 10^{10}$	$\sim 9.2$ - 10.5	p.r.	P.b.k. at 300 nm in N <sub>2</sub> O-satd. soln. contg. 4 x 10 <sup>-3</sup> mol L <sup>-1</sup> alanine anhydride.	71G554
362.074	$-(\text{CH}_2)_2\text{CONH}\dot{\text{C}}(\text{COO}^-)- + \text{OH}^- \rightleftharpoons -(\text{CH}_2)_2\text{CON}^-\dot{\text{C}}(\text{COO}^-)- + \text{H}_2\text{O}$	$k_f = 1 \times 10^9$		e-r.	Est. by esr from pH dependence of line width in N <sub>2</sub> O-satd. soln. contg. 2-pyrrolidone-5-carboxylic acid; ( $pK_a = 12.7$ ).	77D087
362.076	$\text{CH}_3\dot{\text{C}}\text{HOH} + \text{OH}^- \rightleftharpoons \text{CH}_3\dot{\text{C}}\text{HO}^- + \text{H}_2\text{O}$	$k_f = 7 \times 10^9$	11-12	e-r.	Esr line broadening in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> EtOH; $k_f = 4.1 \times 10^5$ ( $pK_a = 11.51$ ).	73D065
362.078	$(\text{CH}_3)_2\dot{\text{C}}\text{OH} + \text{OH}^- \rightleftharpoons (\text{CH}_3)_2\dot{\text{C}}\text{O}^- + \text{H}_2\text{O}$	$k_f = 9 \times 10^9$	11-12	e-r.	Esr line broadening in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> 2-PrOH; $k_f = 1.8 \times 10^6$ ( $pK_a = 12.03$ ).	73D065
		$k_f = 7.5 \times 10^9$	10.4- 11.0	p.r.	Condy. in soln. contg. 1 mol L <sup>-1</sup> acetone gave $k_f = (1.4 \pm 0.1) \times 10^6$ ; $k_f$ calcd. from $pK_a = 12.00$ .	80A323
362.101	$\cdot\text{CHOHCH}_2\text{OH} + \text{OH}^- \rightarrow [\cdot\text{CHO}\cdot\text{CH}_2\text{OH}] + \text{H}_2\text{O} \rightarrow \cdot\text{CH}_2\text{CHO} + \text{OH}^-$	$\sim 10^{10}$	10	p.r.	Absorption spectrum immediately after pulse in N <sub>2</sub> O-satd. soln. contg. ethylene glycol decays in < 10 μs to formylmethyl spectrum.	73G004
362.114	$\cdot\text{CHOHCONH}_2 + \text{OH}^- \rightleftharpoons \cdot\text{CHO}^-\text{CONH}_2 + \text{H}_2\text{O}$	$k_f = 1.1 \times 10^{10}$	8.3- 9.9	p.r.	Calcd. from effect of [OH <sup>-</sup> ] in buffer soln. on d.k. at 290 nm in N <sub>2</sub> O-satd. soln. contg. glycolamide; cor. to $I = 0$ .	75A053
362.115	$\cdot\text{CHOHCO}_2^- + \text{OH}^- \rightleftharpoons \cdot\text{CHO}^-\text{CO}_2^- + \text{H}_2\text{O}$	$k_f = 3.9 \times 10^9$	9.5- 10.5	p.r.	Calcd. from effect of [OH <sup>-</sup> ] in buffer soln. on d.k. at 270 nm in N <sub>2</sub> O-satd. glycolate soln.; cor. to $I = 0$ ; $k = (3.1 \pm 0.4) \times 10^9$ in unbuffered soln.	75A053
362.116	$\cdot\text{COH}(\text{CH}_3)\text{CONH}_2 + \text{OH}^- \rightleftharpoons \cdot\text{CO}^-(\text{CH}_3)\text{CONH}_2 + \text{H}_2\text{O}$	$k_f = 1.1 \times 10^{10}$	8.0- 10.2	p.r.	Calcd. from effect of [OH <sup>-</sup> ] in buffer soln. on d.k. at 300 nm in N <sub>2</sub> O-satd. lactamide soln.; cor. to $I = 0$ .	75A053

TABLE 3. Rate constants for acid- and base-catalyzed proton transfer reactions of aliphatic radicals in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
362.134	$[\text{CH}_2\text{CHCO}_2\text{H}]^- + \text{OH}^-$	$7.7 \times 10^8$	>7	p.r.	Calcd. from effect of $[\text{OH}^-]$ on d.k. in acrylate soln.; cor. to $I = 0$ ; for mechanism and evidence for protonation at the $\beta$ position see 74G033.	76A113
362.135	$[\text{CH}_3\text{CHCHCO}_2\text{H}]^- + \text{OH}^- \rightarrow \text{CH}_3\text{CH}_2\dot{\text{C}}\text{HCO}_2^- + \text{OH}^-$	$\sim 1.2 \times 10^8$	>7.5	p.r.	Calcd. from effect of $[\text{OH}^-]$ on d.k. in <i>trans</i> -crotonate soln.; cor. to $I = 0$ .	76A113
362.136	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CO}_2\text{H}]^- + \text{OH}^- \rightarrow (\text{CH}_3)_2\dot{\text{C}}\text{CO}_2^- + \text{OH}^-$	$8 \times 10^8$	>5.3	p.r.	Calcd. from effect of $[\text{OH}^-]$ on d.k. in soln. contg. methacrylate; cor. to $I = 0$ .	76A113
362.137	$[(\text{CH}_3)_2\text{CCHCO}_2\text{H}]^- + \text{OH}^- \rightarrow (\text{CH}_3)_2\text{CH}\dot{\text{C}}\text{CO}_2^- + \text{OH}^-$	$3 \times 10^6$	>8.0	p.r.	Calcd. from effect of $[\text{OH}^-]$ on d.k. in $\beta,\beta$ -dimethylacrylate soln.; cor. to $I = 0$ .	76A113
362.139	$[\text{CH}_3\text{CHCHCHCHCO}_2\text{H}]^- + \text{OH}^-$	$\sim 2 \times 10^7$	>6.4	p.r.	Calcd. from effect of $[\text{OH}^-]$ on d.k. in soln. contg. sorbate; cor. to $I = 0$ .	76A113
363.	<b>Water</b>					
363.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{OH}^-$	$1.4 \times 10^5 \text{ s}^{-1}$	9.3–13	p.r.	D.k. in acrylamide soln. contg. $10^{-3}$ mol L <sup>-1</sup> borate buffer; general acid catalysis by buffer components, e.g. $\text{B}(\text{OH})_3$ , $\text{NH}_4^+$ , $(\text{CH}_3)_3\text{NH}^+$ , $\text{HCO}_3^-$ , $\text{HPO}_4^{2-}$ , etc.	75A052
363.129	$[\text{CH}_3\text{CHCHCONH}_2]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{CH}_2\dot{\text{C}}\text{HCONH}_2 + \text{OH}^-$	$2 \times 10^4 \text{ s}^{-1}$	10.2	p.r.	D.k. in <i>trans</i> -crotonamide soln. contg. borate buffer, as above.	75A052
363.130	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CONH}_2]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\dot{\text{C}}(\text{CH}_3)\text{CONH}_2 + \text{OH}^-$	$1.3 \times 10^6 \text{ s}^{-1}$	10–13	p.r.	D.k. in methacrylamide soln. contg. borate buffer, as above.	75A052
363.131	$[(\text{CH}_3)_2\text{CCHCONH}_2]^- + \text{H}_2\text{O} \rightarrow (\text{CH}_3)_2\text{CH}\dot{\text{C}}\text{HCONH}_2 + \text{OH}^-$	$2 \times 10^4 \text{ s}^{-1}$	10.2	p.r.	D.k. in $\beta,\beta$ -dimethylacrylamide soln. cor. to $I = 0$ .	75A052
363.132	$[\text{CH}_2\text{CHCON}(\text{CH}_3)_2]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\dot{\text{C}}\text{HCON}(\text{CH}_3)_2 + \text{OH}^-$	$3.7 \times 10^5 \text{ s}^{-1}$	10	p.r.	D.k. in <i>N,N</i> -dimethylacrylamide soln. contg. borate buffer, as above.	75A052
363.134	$[\text{CH}_2\text{CHCO}_2\text{H}]^- + \text{H}_2\text{O}$	$4.0 \times 10^4 \text{ s}^{-1}$ $3.0 \times 10^4 \text{ s}^{-1}$	6.0 9.5	p.r.	D.k. in acrylate soln.; extrapolated to zero concn. of buffer.	76A113
363.135	$[\text{CH}_3\text{CHCHCO}_2\text{H}]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{CH}_2\dot{\text{C}}\text{HCO}_2^- + \text{H}_2\text{O}$	$1.0 \times 10^4 \text{ s}^{-1}$	10.4	p.r.	D.k. in crotonate soln.; extrapolated to zero concn. of buffer.	76A113
363.136	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CO}_2\text{H}]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\dot{\text{C}}(\text{CH}_3)\text{CO}_2^- + \text{H}_2\text{O}$	$2.0 \times 10^5 \text{ s}^{-1}$	9.0	p.r.	D.k. in methacrylate soln.; extrapolated to zero concn. of buffer.	76A113
363.137	$[(\text{CH}_3)_2\text{CCHCO}_2\text{H}]^- + \text{H}_2\text{O} \rightarrow (\text{CH}_3)_2\text{CH}\dot{\text{C}}\text{HCO}_2^- + \text{H}_2\text{O}$	$9 \times 10^3 \text{ s}^{-1}$	11.0	p.r.	D.k. in $\beta,\beta$ -dimethylacrylate soln.; extrapolated to zero concn. of buffer.	76A113
363.138	$[\text{CH}_2\text{C}(\text{CH}_3)\text{CO}_2\text{CH}_3]^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\dot{\text{C}}(\text{CH}_3)\text{CO}_2\text{CH}_3 + \text{OH}^-$	$4.5 \times 10^5 \text{ s}^{-1}$	9.8–12.8	p.r.	D.k. in methyl methacrylate soln. contg. borate buffer; see 363.128.	75A052
363.139	$[\text{CH}_3\text{CHCHCHCHCO}_2\text{H}]^- + \text{H}_2\text{O}$	$5.0 \times 10^4 \text{ s}^{-1}$	11.1	p.r.	D.k. in sorbate soln.; extrapolated to zero concn. of buffer.	76A113
364.	<b>Ammonium ion</b>					
364.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{NH}_4^+ \rightarrow \text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{NH}_3$	$9.5 \times 10^6$		p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
364.134	$[\text{CH}_2\text{CHCO}_2\text{H}]^- + \text{NH}_4^+ \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{NH}_4^+$	$1.8 \times 10^6$	>7	p.r.	Calcd. from effect of $[\text{NH}_4^+]$ on d.k. in acrylate soln.; cor. to $I = 0$ .	76A113
365.	<b>Ammonia</b>					
365.114	$\cdot\text{CHOHCONH}_2 + \text{NH}_3 \rightleftharpoons \cdot\text{CHO}\cdot\text{CONH}_2 + \text{NH}_4^+$	$k_f = 1.2 \times 10^9$	8.05, 8.25	p.r.	Calcd. from effect of $[\text{NH}_3]$ in buffer soln. on d.k. at 290 nm in $\text{N}_2\text{O}$ -satd. glycolamide soln.; cor. to $I = 0$ .	75A053
365.115	$\cdot\text{CHOHCO}_2^- + \text{NH}_3 \rightleftharpoons \cdot\text{CHO}\cdot\text{CO}_2^- + \text{NH}_4^+$	$k_f = 7.5 \times 10^8$	9.26, 9.75	p.r.	Calcd. from effect of $[\text{NH}_3]$ in buffer soln. on d.k. at 270 nm in $\text{N}_2\text{O}$ -satd. glycolate soln.; cor. to $I = 0$ .	75A053



TABLE 3. Rate constants for acid- and base-catalyzed proton transfer reactions of aliphatic radicals in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
365.116	$\cdot\text{COH}(\text{CH}_3)\text{CONH}_2 + \text{NH}_3 \rightleftharpoons \cdot\text{CO}^-(\text{CH}_3)\text{CONH}_2 + \text{NH}_4^+$	$k_f = 9.7 \times 10^8$	8.05	p.r.	Calcd. from effect of $[\text{NH}_3]$ in buffer soln. on d.k. at 300 nm in $\text{N}_2\text{O}$ -satd. lactamide soln.; cor. to $I = 0$ .	75A053
365.134	$[\text{CH}_2\text{CHCO}_2\text{H}]^- + \text{NH}_3 \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{NH}_3$	$(2.8 \pm 1.0) \times 10^6$	>7	p.r.	Calcd. from effect of $[\text{NH}_3]$ on d.k. in acrylate soln.; cor. to $I = 0$ .	76A113
366.	<b>Tetrahydroborate ion</b>					
366.114	$\cdot\text{CHOHCONH}_2 + \text{B}(\text{OH})_4^- \rightleftharpoons \cdot\text{CHO}^-\text{CONH}_2 + \text{B}(\text{OH})_3 + \text{H}_2\text{O}$	$9.1 \times 10^8$	8.25	p.r.	Calcd. from effect of borate on d.k. at 290 nm in $\text{N}_2\text{O}$ -satd. soln. contg. glycolamide; cor. to $I = 0$ .	75A053
366.115	$\cdot\text{CHOHCOO}^- + \text{B}(\text{OH})_4^- \rightleftharpoons \cdot\text{CHO}^-\text{COO}^- + \text{B}(\text{OH})_3 + \text{H}_2\text{O}$	$1.4 \times 10^7$	9.16, 9.75	p.r.	Calcd. from effect of borate on d.k. at 270 nm in $\text{N}_2\text{O}$ -satd. soln. contg. glycolate; cor. to $I = 0$ .	75A053
366.116	$\cdot\text{COH}(\text{CH}_3)\text{CONH}_2 + \text{B}(\text{OH})_4^- \rightleftharpoons \cdot\text{CO}^-(\text{CH}_3)\text{CONH}_2 + \text{B}(\text{OH})_3 + \text{H}_2\text{O}$	$3.1 \times 10^8$	8.30	p.r.	Calcd. from effect of borate on d.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. lactamide; cor. to $I = 0$ .	75A053
366.134	$[\text{CH}_2\text{CHCO}_2\text{H}]^- + \text{B}(\text{OH})_4^- \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{B}(\text{OH})_4^-$	$2.3 \times 10^5$	>7	p.r.	Calcd. from effect of borate on d.k. in acrylate soln.; cor. to $I = 0$ .	76A113
367.	<b>Boric acid</b>					
367.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{B}(\text{OH})_3 + \text{H}_2\text{O} \rightarrow \text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{B}(\text{OH})_4^-$	$1.6 \times 10^6$	9.8	p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
367.134	$[\text{CH}_2\text{CHCO}_2\text{H}]^- + \text{B}(\text{OH})_3 \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{B}(\text{OH})_3$	$2.8 \times 10^5$	>7	p.r.	Calcd. from effect of borate on d.k. in acrylate soln.	76A113
368.	<b>Phosphate ion</b>					
368.114	$\cdot\text{CHOHCONH}_2 + \text{HPO}_4^{2-} \rightleftharpoons \cdot\text{CHO}^-\text{CONH}_2 + \text{H}_2\text{PO}_4^-$	$k_f = 1.8 \times 10^9$	8.3	p.r.	Calcd. from effects of $[\text{HPO}_4^{2-}]$ on d.k. at 290 nm in $\text{N}_2\text{O}$ -satd. glycolamide soln.; cor. to $I = 0$ .	75A053
368.116	$\cdot\text{COH}(\text{CH}_3)\text{CONH}_2 + \text{HPO}_4^{2-} \rightleftharpoons \cdot\text{CO}^-(\text{CH}_3)\text{CONH}_2 + \text{H}_2\text{PO}_4^-$	$k_f = 8.5 \times 10^8$	6.2	p.r.	Calcd. from effect of $[\text{HPO}_4^{2-}]$ on d.k. at 300 nm in $\text{N}_2\text{O}$ -satd. lactamide soln.; cor. to $I = 0$ .	75A053
368.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{H}_2\text{PO}_4^- \rightarrow \text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{HPO}_4^{2-}$	$3.8 \times 10^6$		p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
	$[\text{CH}_2\text{CHCONH}_2]^- + \text{HPO}_4^{2-} \rightarrow \text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{PO}_4^{3-}$	$8.6 \times 10^4$		p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
368.133	$[(\text{CH}_3)_2\text{NCONNCON}(\text{CH}_3)_2]^- + \text{H}_2\text{PO}_4^{2-} \rightarrow (\text{CH}_3)_2\text{NCONHNCON}(\text{CH}_3)_2 + \text{HPO}_4^{2-}$	$(7 \pm 1) \times 10^7$	6-9	p.r.	Effect of $[\text{H}_2\text{PO}_4^-]$ on d.k. at 300 nm in soln. contg. $10^{-4}$ mol L <sup>-1</sup> Diamide and 0.2 mol L <sup>-1</sup> <i>tert</i> -BuOH.	75A194
368.134	$[\text{CH}_2\text{CHCOOH}]^- + \text{HPO}_4^{2-} \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{HPO}_4^{2-}$	$7 \times 10^3$	>7	p.r.	Calcd. from effect of $[\text{HPO}_4^{2-}]$ on d.k. in acrylate soln.; cor. to $I = 0$ .	76A113
	$[\text{CH}_2\text{CHCOOH}]^- + \text{H}_2\text{PO}_4^- \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{H}_2\text{PO}_4^-$	$5.7 \times 10^6$	>7	p.r.	Calcd. from effect of $[\text{H}_2\text{PO}_4^-]$ on d.k. in acrylate soln.; cor. to $I = 0$ .	76A113
369.	<b>Pyrophosphate ion</b>					
369.114	$\cdot\text{CHOHCONH}_2 + \text{HP}_2\text{O}_7^{3-} \rightarrow \cdot\text{CHO}^-\text{CONH}_2 + \text{H}_2\text{P}_2\text{O}_7^{2-}$	$1.1 \times 10^9$	6.10, 6.40	p.r.	Calcd. from effect of $[\text{HP}_2\text{O}_7^{3-}]$ on d.k. at 290 nm in $\text{N}_2\text{O}$ -satd. glycolamide soln.; cor. to $I = 0$ .	75A053
369.115	$\cdot\text{CHOHCOO}^- + \text{P}_2\text{O}_7^{4-} \rightarrow \cdot\text{CHO}^-\text{COO}^- + \text{HP}_2\text{O}_7^{3-}$	$5.8 \times 10^6$	9.26, 9.75	p.r.	Calcd. from effect of $[\text{P}_2\text{O}_7^{4-}]$ on d.k. at 270 nm in $\text{N}_2\text{O}$ -satd. glycolate soln.; cor. to $I = 0$ .	75A053
369.116	$\cdot\text{COH}(\text{CH}_3)\text{CONH}_2 + \text{HP}_2\text{O}_7^{3-} \rightarrow \cdot\text{CO}^-(\text{CH}_3)\text{CONH}_2 + \text{H}_2\text{P}_2\text{O}_7^{2-}$	$3.7 \times 10^8$	8.1, 9.3	p.r.	Calcd. from effect of $[\text{HP}_2\text{O}_7^{3-}]$ on d.k. at 300 nm in $\text{N}_2\text{O}$ -satd. soln. contg. lactamide; $k = 8.5 \times 10^8$ for $\text{P}_2\text{O}_7^{4-}$ .	75A053
369.134	$[\text{CH}_2\text{CHCOOH}]^- + \text{HP}_2\text{O}_7^{3-} \rightarrow \text{CH}_3\dot{\text{C}}\text{HCO}_2^- + \text{HP}_2\text{O}_7^{3-}$	$9.1 \times 10^4$	>7	p.r.	Calcd. from effect of $[\text{HP}_2\text{O}_7^{3-}]$ on d.k. in acrylate soln.; cor. to $I = 0$ .	76A113

TABLE 3. Rate constants for acid- and base-catalyzed proton transfer reactions of aliphatic radicals in aqueous solution—Continued

No.	Reaction	$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
370.	<b>Bicarbonate ion</b>					
370.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{HCO}_3^- \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{CO}_3^{2-}$	$8.6 \times 10^5$		p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
371.	<b>Trimethylammonium ion</b>					
371.128	$[\text{CH}_2\text{CHCONH}_2]^- + (\text{CH}_3)_3\text{NH}^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + (\text{CH}_3)_3\text{N}$	$2.1 \times 10^9$		p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
372.	<b>Ethylammonium ion</b>					
372.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{C}_2\text{H}_5\text{NH}_3^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{C}_2\text{H}_5\text{NH}_2$	$1.5 \times 10^7$		p.r.	D.k. in soln. contg. acrylamide.	75A052 74A182
373.	<b>Diethylammonium ion</b>					
373.128	$[\text{CH}_2\text{CHCONH}_2]^- + (\text{C}_2\text{H}_5)_2\text{NH}_2^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + (\text{C}_2\text{H}_5)_2\text{NH}$	$7.2 \times 10^7$		p.r.	D.k. in soln. contg. acrylamide.	75A052 74A182
374.	<b>Triethylammonium ion</b>					
374.128	$[\text{CH}_2\text{CHCONH}_2]^- + (\text{C}_2\text{H}_5)_3\text{NH}^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + (\text{C}_2\text{H}_5)_3\text{N}$	$1.3 \times 10^8$	10.3– 11.8	p.r.	D.k. in soln. contg. acrylamide; cor. to $I = 0$ .	75A052 74A182
375.	<b>Cyclohexylammonium ion</b>					
375.128	$[\text{CH}_2\text{CHCONH}_2]^- + c\text{-C}_6\text{H}_{11}\text{NH}_3^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + c\text{-C}_6\text{H}_{11}\text{NH}_2$	$1.4 \times 10^7$		p.r.	D.k. in soln. contg. acrylamide.	75A052 74A182
376.	<b>Pyrrolidinium ion</b>					
376.128	$[\text{CH}_2\text{CHCONH}_2]^- + c\text{-C}_4\text{H}_8\text{NH}_2^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + c\text{-C}_4\text{H}_8\text{NH}$	$2.0 \times 10^8$		p.r.	D.k. in soln. contg. acrylamide.	75A052 74A182
377.	<b>Piperidinium ion</b>					
377.128	$[\text{CH}_2\text{CHCONH}_2]^- + c\text{-C}_5\text{H}_{10}\text{NH}_2^+ \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + c\text{-C}_5\text{H}_{10}\text{NH}$	$6.3 \times 10^7$		p.r.	D.k. in soln. contg. acrylamide.	75A052 74A182
378.	<b>Glycine</b>					
378.128	$[\text{CH}_2\text{CHCONH}_2]^- + \text{H}_3\text{N}^+\text{CH}_2\text{CO}_2^- \rightarrow$ $\text{CH}_3\dot{\text{C}}\text{HCONH}_2 + \text{H}_2\text{NCH}_2\text{CO}_2^-$	$2.5 \times 10^7$		p.r.	D.k. in soln. contg. acrylamide.	75A052 74A182

TABLE 4. Rate constants for second order decay of selected aliphatic radicals in aqueous solution—Continued

No.	Reaction	$k^*$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
.001	<b>Methyl</b> $\cdot\text{CH}_3 + \cdot\text{CH}_3 \rightarrow \text{C}_2\text{H}_6$	$(1.24 \pm 0.2) \times 10^9$	5.5	p.r.	D.k. in soln. contg. $10^{-2}$ mol L <sup>-1</sup> CH <sub>4</sub> and 0.25 mol L <sup>-1</sup> N <sub>2</sub> O; $\epsilon = 850$ L mol <sup>-1</sup> cm <sup>-1</sup> at 220 nm.	72G445
		$(1.6 \pm 0.2) \times 10^9$	4.4	p.r.	D.k. in soln. contg. 0.4 mol L <sup>-1</sup> CH <sub>4</sub> (30 atm) and $2.5 \times 10^{-3}$ mol L <sup>-1</sup> N <sub>2</sub> O; $\epsilon = 1600$ and $975$ L mol <sup>-1</sup> cm <sup>-1</sup> at 213 and 220 nm., resp.; $E_a = 3.9 \pm 0.4$ kcal mol <sup>-1</sup> (16 kJ mol <sup>-1</sup> ).	75A055
.002	<b>Ethyl</b> $\cdot\text{CH}_2\text{CH}_3 + \cdot\text{CH}_2\text{CH}_3$	$(9.6 \pm 2) \times 10^8$	5.5	p.r.	D.k. in soln. contg. $10^{-2}$ mol L <sup>-1</sup> ethane and 0.25 mol L <sup>-1</sup> N <sub>2</sub> O; $\epsilon = 520$ and $330$ L mol <sup>-1</sup> cm <sup>-1</sup> at 220 and 250 nm, resp.	72G445
		$(1.2 \pm 0.2) \times 10^9$	4.4	p.r.	D.k. in soln. contg. $5 \times 10^{-2}$ mol L <sup>-1</sup> ethane (30 atm) and $2.5 \times 10^{-3}$ mol L <sup>-1</sup> N <sub>2</sub> O; $\epsilon = 790, 750, 700, 545$ and $375$ L mol <sup>-1</sup> cm <sup>-1</sup> at 210, 215, 220, 230 and 250 nm, resp.; $E_a = 3.9 \pm 0.4$ kcal mol <sup>-1</sup> (16 kJ mol <sup>-1</sup> ).	75A055
.004	<b>Cyclopentyl</b> $c\text{-}\dot{\text{C}}_5\text{H}_9 + c\text{-}\dot{\text{C}}_5\text{H}_9 \rightarrow$ $\text{C}_{10}\text{H}_{18}$ and $\text{C}_5\text{H}_{10} + \text{C}_5\text{H}_8$	$(1.0 \pm 0.2) \times 10^9$	7	p.r.	D.k. in soln. contg. N <sub>2</sub> O and $1.3 \times 10^{-5}$ mol L <sup>-1</sup> cyclopentane; $\epsilon = 480$ L mol <sup>-1</sup> cm <sup>-1</sup> at 248 nm.	74A051
.016	<b>2-Hydroxy-2,2-dimethylethyl</b> $\cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH} + \cdot\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$	$7 \times 10^8$ $6 \times 10^8$	6 13.5	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. <i>tert</i> -BuOH; $\epsilon = 900, 200$ and $30$ L mol <sup>-1</sup> cm <sup>-1</sup> at 225, 250 and 280 nm, resp. $2k/\epsilon = 1.3 \times 10^7$ and $1.1 \times 10^7$ at pH 6 and 13.5, resp.	69G419
.020	<b>Formylmethyl</b> $\cdot\text{CH}_2\text{CHO} + \cdot\text{CH}_2\text{CHO}$	$4.5 \times 10^8$	10	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.1 mol L <sup>-1</sup> ethylene glycol; $\epsilon = 260$ L mol <sup>-1</sup> cm <sup>-1</sup> at 300 nm.	73G004
		$4.5 \times 10^8$	9.8	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. $10^{-3}$ mol L <sup>-1</sup> acetylene; $2k/\epsilon = 3.4 \times 10^6$ cm s <sup>-1</sup> .	78A007
.021	<b>2-Hydroxyethenyl</b> $\cdot\text{CH}=\text{CHOH} + \cdot\text{CH}=\text{CHOH} \rightarrow$ $\text{HOCH}=\text{CHCH}=\text{CHOH}$	$3.2 \times 10^9$	<7	p.r.	P.b.k. in N <sub>2</sub> O-satd. soln. contg. $10^{-3}$ mol L <sup>-1</sup> acetylene at 240 nm ( $\epsilon = 8.1 \times 10^3$ L mol <sup>-1</sup> cm <sup>-1</sup> ); $2k/\epsilon = 2.0 \times 10^5$ cm s <sup>-1</sup> ; prod. rearranges to succinaldehyde.	78A007
.035	<b>Carboxymethyl</b> $\cdot\text{CH}_2\text{CO}_2^- + \cdot\text{CH}_2\text{CO}_2^-$	$(5.5 \pm 1) \times 10^8$	6.5	p.r.	D.k. at 366 nm in N <sub>2</sub> O-satd. soln. contg. Na acetate; $\epsilon = 780$ L mol <sup>-1</sup> cm <sup>-1</sup> .	76A082
		$5 \times 10^8$	10	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. acetate ion; $\epsilon = 800$ L mol <sup>-1</sup> cm <sup>-1</sup> at 350 nm.	69G446
		$9 \times 10^8$	3	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. acetic acid; $\epsilon = 650$ L mol <sup>-1</sup> cm <sup>-1</sup> at 320 nm; $pK_a = 4.5$ .	69G446
.048	<b>Trichloromethyl</b> $\cdot\text{CCl}_3 + \cdot\text{CCl}_3 \rightarrow \text{C}_2\text{Cl}_6$	$(3.7 \pm 0.9) \times 10^8$	5.5–12.4	p.r.	D.k. at 230 nm in $10^{-4}$ mol L <sup>-1</sup> CCl <sub>4</sub> soln.; $\epsilon = 2300 \pm 450$ L mol <sup>-1</sup> cm <sup>-1</sup> .	74A043



TABLE 4. Rate constants for second order decay of selected aliphatic radicals in aqueous solution—Continued

No.	Reaction	$k^a$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
.075	<b>Hydroxymethyl</b> ·CH <sub>2</sub> OH + ·CH <sub>2</sub> OH ·CH <sub>2</sub> O <sup>-</sup> + ·CH <sub>2</sub> O <sup>-</sup>	1.2 x 10 <sup>9</sup> 4.5 x 10 <sup>8</sup>	6 12	p.r.	D.k. at 280 and 350 nm, resp., at pH 6 and 12 in N <sub>2</sub> O-satd. soln. contg. MeOH; $2k/\epsilon = 6.2 \times 10^6$ and $1.5 \times 10^6$ at pH 6 and 12, resp. ( $\epsilon = 390$ and $600$ ); $pK_a = 10.7$ [66G074].	69G419
	·CH <sub>2</sub> OH + ·CH <sub>2</sub> OH	(1.5 ± 1) x 10 <sup>9</sup>		p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. MeOH; $\epsilon = 270 \pm 15$ L mol <sup>-1</sup> cm <sup>-1</sup> at 310 nm.	77A011
.076	<b>1-Hydroxyethyl</b> CH <sub>3</sub> ·CHOH + CH <sub>3</sub> ·CHOH → CH <sub>3</sub> CHOHCHOHCH <sub>3</sub> (I)	(5.5 ± 1.5) x 10 <sup>8</sup> $k_I = (4.4 \pm 1) \times 10^8$	~7	p.r.	D.k. at 289.4 nm in deaerated soln. contg. EtOH; $\epsilon = 260 \pm 65$ L mol <sup>-1</sup> cm <sup>-1</sup> ; relative disproportionation and combination est. from product yields.	62G140
	→ CH <sub>3</sub> CHO + CH <sub>3</sub> CH <sub>2</sub> OH (II)	$k_{II} = (1.1 \pm 0.3) \times 10^8$ (7 ± 2) x 10 <sup>8</sup>	~1	p.r.	D.k. at 296.7 nm in deaerated soln. contg. EtOH and 0.8 N H <sub>2</sub> SO <sub>4</sub> ; $\epsilon = 240 \pm 45$ L mol <sup>-1</sup> cm <sup>-1</sup> .	63G045
		(1.2 ± 0.3) x 10 <sup>9</sup>	6	p.r.	D.k. at 280 and 300 nm, resp., at pH 6 and 13 in N <sub>2</sub> O-satd. soln. contg. EtOH; $pK_a = 11.6$ [66G074];	69G419
	CH <sub>3</sub> ·CHO <sup>-</sup> + CH <sub>3</sub> ·CHO <sup>-</sup>	(2.5 ± 0.5) x 10 <sup>8</sup>	13		$2k/\epsilon = 4.7 \times 10^6$ and $5.7 \times 10^6$ resp., at pH 6 and 13.	
.078	<b>1-Hydroxy-1-methylethyl</b> (CH <sub>3</sub> ) <sub>2</sub> ·CHOH + (CH <sub>3</sub> ) <sub>2</sub> ·CHOH (CH <sub>3</sub> ) <sub>2</sub> ·C·O <sup>-</sup> + (CH <sub>3</sub> ) <sub>2</sub> ·C·O <sup>-</sup>	(7 ± 1.5) x 10 <sup>8</sup> (2 ± 0.4) x 10 <sup>8</sup>	6 13.3	p.r.	D.k. at 280 and 300 nm, resp. at pH 6 and 13.3 in N <sub>2</sub> O-satd. soln. contg. 2-PrOH; $2k/\epsilon = 2.1 \times 10^6$ and $2.7 \times 10^5$ at pH 6 and 13.3, resp.; $pK_a = 12.2$ [66G074].	69G419
		(6.5 ± 0.5) x 10 <sup>8</sup>		p.r.	D.k. at 310 nm in N <sub>2</sub> O-satd. soln. contg. 0.02 mol L <sup>-1</sup> 2-PrOH; $\epsilon = 353 \pm 15$ L mol <sup>-1</sup> cm <sup>-1</sup> .	77A011
.101	<b>1,2-Dihydroxyethyl</b> ·CHOHCH <sub>2</sub> OH + ·CHOHCH <sub>2</sub> OH	3.4 x 10 <sup>8</sup>	6.0	p.r.	D.k. at 250 nm in N <sub>2</sub> O-satd. soln. contg. ethylene glycol; $2k/\epsilon = 1.1 \times 10^6$ ( $\epsilon = 610$ L mol <sup>-1</sup> cm <sup>-1</sup> ).	73G004
.113	<b>Ascorbate radical</b> ·A <sup>-</sup> + ·A <sup>-</sup> (+ H <sup>+</sup> ) → A + HA <sup>-</sup>	(1.9 ± 0.2) x 10 <sup>5</sup>	8.7	p.r.	D.k. at 360 nm in N <sub>2</sub> O-satd. soln. contg. ascorbate; $\epsilon = 3300$ L mol <sup>-1</sup> cm <sup>-1</sup> [77A036]; $pK_a = -0.45$ [72D181].	75A240
.115	<b>(Carboxy)hydroxymethyl</b> ·CHOHCO <sub>2</sub> H + ·CHOHCO <sub>2</sub> H ·CHOHCO <sub>2</sub> <sup>-</sup> + ·CHOHCO <sub>2</sub> <sup>-</sup> ·CHO <sup>-</sup> CO <sub>2</sub> <sup>-</sup> + ·CHO <sup>-</sup> CO <sub>2</sub> <sup>-</sup>	(6.5 ± 1) x 10 <sup>8</sup> (4.3 ± 0.7) x 10 <sup>8</sup> (7.5 ± 1) x 10 <sup>6</sup>	1 7.2 12	p.r.	D.k. in N <sub>2</sub> O-satd. soln. contg. 0.05 mol L <sup>-1</sup> glycolic acid ( $\epsilon = 6000$ L mol <sup>-1</sup> cm <sup>-1</sup> ); $pK_a = 4.6, 8.8$ .	69G447
.123	<b>Dihydroxymethyl</b> ·CH(OH) <sub>2</sub> + ·CH(OH) <sub>2</sub> → CH <sub>2</sub> (OH) <sub>2</sub> + HCO <sub>2</sub> <sup>-</sup> + H <sup>+</sup>	3.6 x 10 <sup>8</sup> 2.3 x 10 <sup>8</sup>	6.3 12	p.r.	D.k. at 250 nm, as well as condy. change, in N <sub>2</sub> O-satd. soln. contg. formaldehyde, as well as in soln. contg. <i>tert</i> -BuOH and CO; $\epsilon \cong 850$ and $1500$ L mol <sup>-1</sup> cm <sup>-1</sup> at pH 6 and 11.8, resp.; $pK_a = 9.5$ .	71G424

TABLE 4. Rate constants for second order decay of selected aliphatic radicals in aqueous solution—Continued

No.	Reaction	$k^a$ (L mol <sup>-1</sup> s <sup>-1</sup> )	pH	Method	Comment	Ref.
.147	<b>Carboxyl</b> $\cdot\text{CO}_2^- + \cdot\text{CO}_2^- \rightarrow \text{C}_2\text{O}_4^{2-}$	$5 \times 10^8$	5	p.r.	D.k. at 250 nm in CO <sub>2</sub> -satd. soln. contg. 10 <sup>-2</sup> mol L <sup>-1</sup> formate; $\epsilon = 2250$ L mol <sup>-1</sup> cm <sup>-1</sup> ; $k$ cor. to $I = 0$ .	65G384
		$7.5 \times 10^8$	3.1,9	p.r.	D.k. at 255 nm in N <sub>2</sub> O-satd. soln. contg. formate, as well as in CO <sub>2</sub> -satd. soln. contg. formate; $\epsilon_{\text{max}} = 3000$ L mol <sup>-1</sup> cm <sup>-1</sup> at 235 nm.	69G446
		$8.5 \times 10^8$	13			
		$(4.5 \pm 0.15) \times 10^8$	2.8-7	p.r.	D.k. in CO-satd. soln. at 260 nm ( $\epsilon = 2200$ L mol <sup>-1</sup> cm <sup>-1</sup> ) as well as condy. change; same result in N <sub>2</sub> O-satd. formate soln.	70G303
		$(3.8 \pm 0.35) \times 10^8$	2.7-13	p.r.	D.k. at 250 nm in N <sub>2</sub> O-satd. 0.1 mol L <sup>-1</sup> formate soln.; $\epsilon = 2050$ L mol <sup>-1</sup> cm <sup>-1</sup> (1100 at pH 0); $k$ cor. to $I = 0$ ; $\text{p}K_a(\cdot\text{CO}_2\text{H}) = 1.4$ .	73G085
		$(8.5 \pm 1) \times 10^8$	0			

<sup>a</sup>  $k$  is defined by the expression  $-\text{d}[\text{R}]/\text{dt} = 2k[\text{R}]^2$ .

## Index to Radical Reactions

Radicals are listed in number order. Following the name of the radical are the entry numbers for the substrates for which reactions of that radical are included in tables 1-3. For example, the first reaction in the tables for methyl radical is under entry 9 and the reaction is numbered 9.001.

- 001 Methyl 9, 10, 20, 59, 69, 80, 81a, 83, 93, 94, 115, 116, 117, 147, 149, 156, 160, 200, 201, 202, 217, 219, 221a, 222, 223, 230, 241, 242, 243, 248, 254, 255, 267, 305, 314, 315, 316, 317, 345, 346.
- 002 Ethyl 69, 81a, 83, 93.
- 002a 1-Methylethyl 69, 81a.
- 002b 1,1-Dimethylethyl 69, 81a.
- 003 Radicals from pentane 94.
- 004 Cyclopentyl 88, 93, 94, 342, 343, 345, 346.
- 005 2-Chloroethyl 81a.
- 006 2-Carboxy-2,2-dimethylethyl 55, 71, 104.
- 007 Radicals from oleate + OH 93.
- 008 Radicals from linoleate + OH 93.
- 009 Radicals from linolenate + OH 93.
- 010 Radicals from arachidonate + OH 93.
- 011 2-Aminoethyl 262.
- 012 2-Amino-2,2-dimethylethyl 71, 104.
- 013 2-Amino-2-carboxy-2-methylethyl 71, 104.
- 014 2-Hydroxyethyl 57, 58, 59, 69, 81a, 93, 200, 202, 253, 345, 346.
- 015 2-Hydroxypropyl 314.
- 016 2-Hydroxy-2,2-dimethylethyl 1, 2, 4, 5, 6, 9, 10, 10a, 11, 55, 57, 58, 59, 71, 72, 77, 81a, 83, 88, 89, 99, 104, 105, 106, 113, 114, 123, 161, 163, 197, 206, 227, 236, 237, 238, 239, 253, 262, 302, 342, 343, 345, 346.
- 016a Radicals from cyclopentene + OH 94.
- 017 2-Carboxy-2-hydroxy-2-methylethyl 71, 104.
- 018 Radicals from allyl alcohol + OH 161, 278, 279, 283.
- 019 Radicals from crotyl alcohol + OH 278, 279, 283.
- 020 Formylmethyl 9, 55, 81a, 128, 135, 176, 177, 207, 227, 258, 266, 309, 310, 311, 344.
- 021 1-Formylethyl 227.
- 022 2-Hydroxyethenyl 119, 362.
- 023 2-Oxopropyl 227.
- 024 1-Acetyethyl 227.
- 025 2-Oxocyclohexyl 227.
- 026 1-Formyl-2-hydroxyethyl 227.
- 027 4-Hydroxy-2-oxobutyl 227.
- 028 1-Formyl-4-hydroxybutyl 227.
- 029 1-Formyl-5-hydroxypentyl 227.
- 030 Radicals from *meso*-erythritol 227.
- 031 Radicals from xylitol 227.
- 032 2-Oxo-3,4,5-trihydroxycyclohexyl 227.
- 033 Radicals from sorbitol 161, 227.
- 034 Carbamoylmethyl 262.
- 035 Carboxymethyl 9, 12, 55, 59, 81a, 93, 100, 101, 102, 110.
- 036 Methoxycarbonylmethyl 93.
- 037 1-Carboxyethyl 55, 59, 278.
- 038 1-Carboxypropyl 278.
- 039 1-Carboxy-2-hydroxyethyl 278.
- 040 1-Carboxy-2-hydroxypropyl 278.
- 040a 1,2-Dicarboxyethyl 81a.
- 041 1,2-Dicarboxy-2-hydroxyethyl 59, 278.
- 042 Dicarboxymethyl 12, 55, 100, 110, 135, 228, 278.
- 043 Chloromethyl 69, 81a, 83, 223.
- 044 Difluoromethyl
- 045 Dichloromethyl 69, 83, 335.
- 046 Dichlorofluoromethyl 202.
- 047 Trifluoromethyl 130, 147, 149, 200, 202, 212, 214, 255, 292, 313, 314, 317.
- 048 Trichloromethyl 59, 81a, 83, 205, 223, 312, 314, 340, 358.
- 049 1-Aminoethyl 262.
- 050 1-Amino-1-methylethyl 222, 262.
- 051 1-(*N,N*-Diethylamino)ethyl 262.
- 052 Triethylammonioethyl 93.
- 053 (*N*-Acetylamino)methyl 206, 262.
- 054 (*N*-Formyl-*N*-methylamino)methyl 12, 55, 100.
- 055 (*N*-Acetyl-*N*-methylamino)methyl 71, 104, 262.



- 056 Amino(carboxy)methyl 12, 69, 100, 133, 134, 138, 159, 173, 228, 231, 232, 260, 262, 274, 276, 308, 333, 336, 350.
- 057 Amino(carbamoyl)methyl 93, 262.
- 058 Radicals from glycylglycinamide + OH 262.
- 059 Carboxy(*N*-methylamino)methyl 262.
- 060 *N*-Acetylamino(carboxy)methyl 362.
- 061 *N*-Acetyl-*N*-methylamino(carboxy)methyl 262.
- 062 Radical from glycine anhydride + OH 93, 138, 162, 163, 262, 273, 362.
- 063 Radical from alanine anhydride + OH 93, 262, 362.
- 064 Radical from sarcosine anhydride + OH 93.
- 065 Radicals from diglycine + OH 262.
- 066 Radicals from glycylsarcosine + OH 262.
- 067 Radicals from acetyldiglycine + OH 93, 262.
- 068 Radicals from triglycine + OH 133, 134, 138, 262, 274.
- 069 Radicals from acetyltriglycine + OH 262.
- 070 Radicals from acetyltrialanine + OH 262.
- 071 Radicals from acetyltrisarcosine + OH 262.
- 072 Radicals from acetylserineamide + OH 262.
- 073 Radicals from acetylasparagine + OH 262.
- 074 Radical from 2-pyrrolidone-5-carboxylic acid 362.
- 075 Hydroxymethyl 1, 2, 4, 5, 6, 9, 10a, 12, 13, 15, 16, 17, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 45a, 55, 57, 59, 60, 68, 69, 70, 73, 77, 81a, 88, 89, 92, 94, 95, 97, 98, 98a, 99, 99a, 100, 102, 106, 110, 111a, 113, 114, 121, 123, 135, 136, 137, 138, 143, 148, 152, 154, 161, 162, 163, 173, 183, 185a, 197, 204, 206, 211, 214, 220, 226, 231, 232, 233, 236, 239, 253, 260, 262, 268, 270, 271, 276, 278, 279, 283, 293, 297, 304, 306, 308, 318, 319, 323, 342, 343, 345, 346, 347, 350, 361.
- 076 1-Hydroxyethyl 2, 3, 4, 5, 9, 12, 15, 16, 36, 37, 38, 39, 40, 41, 42, 43, 45, 45a, 55, 59, 60, 68, 69, 73, 76, 77, 81, 81a, 86, 87, 88, 93, 94, 95, 98a, 99, 99a, 100, 102, 106, 111, 111a, 118, 123, 137, 138, 148, 154, 161, 163, 164, 165, 185a, 199, 209, 220, 224, 226, 249, 253, 262, 268, 269, 270, 271, 278, 279, 283, 297, 306, 319, 323, 331, 338, 339, 342, 343, 345, 346, 347, 362.
- 077 1-Hydroxypropyl 69, 93, 106, 148, 278, 279, 283, 297, 347.
- 078 1-Hydroxy-1-methylethyl 1, 2, 4, 5, 6, 7, 8, 9, 10, 10a, 12, 15, 16, 24, 25, 26, 27, 28, 29, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 45a, 46, 47, 48, 49, 50, 51, 52, 55, 57, 59, 60, 67, 68, 69, 70, 73, 74, 75, 76, 77, 79, 81a, 83, 84, 85, 87, 88, 89, 93, 94, 95, 96, 97, 98, 98a, 99, 99a, 100, 102, 103, 106, 111, 111a, 113, 114, 118, 120, 121, 123, 124, 127, 129, 131, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 145, 146, 148, 151, 152, 153, 154, 155, 156a, 157, 159, 161, 163, 170, 171, 173, 174, 178, 180, 181, 182, 184, 185, 185a, 186, 187, 188, 189, 190, 191, 192, 193, 197, 198, 206, 210, 215, 216, 220, 221, 221a, 222, 223, 225, 228, 229, 231, 232, 233, 234, 235, 238a, 244, 245, 249, 250, 252, 253, 257, 259, 260, 261, 262, 263, 264, 265, 268, 269, 270, 271, 272, 273, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 306, 307, 308, 319, 320, 321, 322, 323, 324, 325, 326, 327, 329, 329, 331, 333, 334, 336, 337, 342, 343, 345, 346, 347, 348, 349, 350, 351, 352, 353, 356, 357, 359, 360, 361, 362.
- 079 1-Hydroxybutyl 58a, 65a, 161, 262, 278, 279, 283, 297.
- 080 1-Hydroxy-1-methylpropyl 60, 93, 148.
- 081 1-Hydroxy-2-methylpropyl 69, 93, 161, 283, 297.
- 082 1-Hydroxypentyl 69.
- 083 (Cyclohexyl)hydroxymethyl 148.
- 084 1-Hydroxycyclobutyl 69, 283.
- 085 1-Hydroxycyclopentyl 69, 283.
- 086 1-Hydroxycyclohexyl 69, 283.
- 087 1-Hydroxycycloheptyl 69, 283.
- 088 1-Hydroxycyclooctyl 69, 283.
- 089 2,2,2-Trifluoro-1-hydroxyethyl 9.
- 090 Acetoxymethyl 93.
- 091 1-Sulfatoethyl 69.
- 092 Radicals from sodium dodecylsulfate 138.
- 092a Methoxymethyl 69, 81a, 347.
- 093 1-Ethoxyethyl 12, 15, 16, 37, 38, 39, 39, 40, 42, 55, 81a, 94, 100, 101, 102, 152, 283, 347.
- 093a 1,1-Dimethylethoxymethyl 347.
- 094 2-Tetrahydrofuryl 94, 152.
- 094a (Methoxymethoxy)methyl 347.
- 095 1,4-Dihydroxybutyl 161.
- 096 1,3-Dihydroxy-2,2-di(hydroxymethyl)propyl 95, 163, 225.
- 097 2-Carboxy-1-hydroxy-1-methylethyl 69, 83, 131, 133, 134, 181, 228, 262.
- 098 2-Amino-1-hydroxyethyl 9.
- 099 2-Amino-1-phosphoryloxyethyl 278.
- 100 2-Amino-2-carboxy-1-hydroxyethyl 60, 93.
- 101 1,2-Dihydroxyethyl 5, 9, 10, 29, 55, 69, 93, 347, 362.
- 102 1,2-Dihydroxypropyl 347.
- 103 1,2-Dihydroxy-1-methylpropyl 347.

- 104 2,5-Dioxacyclohexyl 12, 15, 16, 42, 55, 81a, 83, 94, 100, 101, 102, 283.
- 105 Radicals from polyethyleneglycol 4, 5, 69, 93.
- 106 1,2,3-Trihydroxypropyl 5, 69, 93, 163, 225, 278, 347.
- 107 1-Hydroxy-1-methyl-2-oxopropyl 12, 100.
- 108 Radicals from deoxyribose + OH 69, 93, 138, 262, 278, 279, 342.
- 109 Radicals from ribose + OH 206, 227, 262, 278.
- 110 Radicals from ribose phosphate + OH 262, 278.
- 111 Radicals from glucose + OH 69, 93, 158, 161, 194, 195, 227, 279, 287, 342, 343, 345, 346, 347, 355.
- 112 Radicals from sucrose + OH 347.
- 113 Radical from ascorbate + OH 59, 93, 163, 179, 214, 246, 247, 255, 262, 330.
- 114 (Carbamoyl)hydroxymethyl 262, 362, 365, 366, 368, 369.
- 115 (Carboxy)hydroxymethyl 55, 69, 81a, 93, 133, 138, 228, 262, 274, 333, 362, 365, 366, 368, 369.
- 116 1-Carbamoyl-1-hydroxyethyl 138, 362, 365, 366, 368, 369.
- 117 1-Carboxy-1-hydroxyethyl 12, 55, 69, 93, 100, 134, 138, 148, 163, 225, 262, 278, 283.
- 118 1-Hydroxy-1-(methoxycarbonyl)ethyl 262.
- 119 (Dicarboxy)hydroxymethyl 262.
- 120 1,2-Dicarboxy-1-hydroxyethyl 163, 262, 278.
- 121 1,2-Dicarboxy-1,2-dihydroxyethyl 163, 165, 225, 262, 270.
- 122 1,3-Dicarboxy-1-hydroxypropyl 262.
- 123 Dihydroxymethyl 93, 283.
- 124 Dimethoxymethyl 94, 347.
- 125 2,5-Dioxacyclopentyl 94.
- 126 2,2,2-Trichloro-1,1-dihydroxyethyl 93.
- 127 Carbamoyl 12, 55, 100.
- 128 Electron adduct of acrylamide 1, 123, 363, 364, 367, 368, 370, 371, 373, 374, 375, 376, 377, 378.
- 129 Electron adduct of crotonamide 363.
- 130 Electron adduct of methacrylamide 363.
- 131 Electron adduct of  $\beta,\beta$ -dimethylacrylamide 363.
- 132 Electron adduct of *N,N*-dimethylacrylamide 363.
- 133 Electron adduct of Diamide 368.
- 134 Electron adduct of acrylate 362, 364, 365, 366, 367, 368, 369.
- 135 Electron adduct of crotonate 362, 363.
- 136 Electron adduct of methacrylate 362, 363.
- 137 Electron adduct of  $\beta,\beta$ -dimethylacrylate 362, 363.
- 138 Electron adduct of methyl methacrylate 363.
- 139 Electron adduct of sorbate 362, 363.
- 141 Electron adduct of dimethyl fumarate 93, 206.
- 142 Electron adduct of glycine anhydride 118, 137, 160a, 162, 163, 215a, 216, 253a, 262.
- 143 Electron adduct of alanine anhydride 118, 137, 160a, 162, 262.
- 144 Electron adduct of sarcosine anhydride 118, 137, 162, 262.
- 146 Electron adduct of acetylglycylglycinamide 262.
- 147 Carboxyl 1, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 44, 46, 49, 50, 53, 54, 55, 56, 59, 60, 61, 62, 63, 64, 65, 66, 69, 70, 74, 76, 78, 81a, 82, 88, 89, 90, 91, 92, 93, 94, 98, 99, 100, 103, 107, 108, 109, 110, 111, 112, 113, 114, 118, 121, 122, 123, 125, 126, 132, 133, 134, 138, 143, 144, 146, 150, 151, 152, 158, 159, 163, 164, 165, 167, 168, 169, 170, 171, 172, 173, 175, 183, 184, 185a, 190, 191, 192, 193, 195, 196, 199, 203, 206, 208, 209, 213, 214, 216, 218, 220, 225, 228, 231, 233, 233a, 240, 249, 251, 252, 256, 260, 262, 270, 271, 276, 278, 279, 283, 284, 285, 286, 287a, 297, 319, 320, 321, 323, 324, 329, 332, 333, 341, 342, 347, 354, 356.
- 148 Electron adduct of methyl acetate 361.



# Radical Formula Index

The radical numbers following the name are contained in the *List of Aliphatic Radicals* preceding table 1 and in the *Index to Radicals* just preceding this index.

CCl <sub>2</sub> F	Dichlorofluoromethyl, 046	C <sub>3</sub> H <sub>8</sub> N	1-Amino-1-methylethyl, 050
CCl <sub>3</sub>	Trichloromethyl, 048	C <sub>4</sub> H <sub>4</sub> O <sub>4</sub> <sup>-</sup>	Electron adduct of maleate, 140
CF <sub>3</sub>	Trifluoromethyl, 047	C <sub>4</sub> H <sub>5</sub> N <sub>2</sub> O <sub>2</sub>	Radical from glycine anhydride + OH, 062
CHCl <sub>2</sub>	Dichloromethyl, 045	C <sub>4</sub> H <sub>5</sub> O <sub>4</sub>	1,2-Dicarboxyethyl, 040a
CHF <sub>2</sub>	Difluoromethyl, 044	C <sub>4</sub> H <sub>5</sub> O <sub>5</sub>	1,2-Dicarboxy-1-hydroxyethyl, 120
CH <sub>2</sub> Cl	Chloromethyl, 043		1,2-Dicarboxy-2-hydroxyethyl, 041
CH <sub>2</sub> NO	Carbamoyl, 127	C <sub>4</sub> H <sub>5</sub> O <sub>6</sub>	1,2-Dicarboxy-1,2-dihydroxyethyl, 121
CH <sub>3</sub>	Methyl, 001	C <sub>4</sub> H <sub>6</sub> NO <sub>3</sub>	<i>N</i> -Acetylamino(carboxy)methyl, 060
CH <sub>3</sub> O	Hydroxymethyl, 075	C <sub>4</sub> H <sub>6</sub> O <sub>2</sub> <sup>-</sup>	Electron adduct of crotonate, 135
CH <sub>3</sub> O <sub>2</sub>	Dihydroxymethyl, 123		Electron adduct of methacrylate, 136
CO <sub>2</sub> <sup>-</sup>	Carboxyl, 147	C <sub>4</sub> H <sub>7</sub> NO <sup>-</sup>	Electron adduct of crotonamide, 129
C <sub>2</sub> H <sub>2</sub> Cl <sub>3</sub> O <sub>2</sub>	2,2,2-Trichloro-1,1-dihydroxyethyl, 126		Electron adduct of methacrylamide, 130
C <sub>2</sub> H <sub>2</sub> F <sub>3</sub> O	2,2,2-Trifluoro-1-hydroxyethyl, 089	C <sub>4</sub> H <sub>7</sub> N <sub>2</sub> O <sub>2</sub> <sup>-</sup>	Electron adduct of glycine anhydride, 142
C <sub>2</sub> H <sub>3</sub> O	Formylmethyl, 020		Radicals from glycyglycinamide + OH, 058
	2-Hydroxyethenyl, 022	C <sub>4</sub> H <sub>7</sub> N <sub>2</sub> O <sub>3</sub>	Radicals from diglycine + OH, 065
C <sub>2</sub> H <sub>3</sub> O <sub>2</sub>	Carboxymethyl, 035	C <sub>4</sub> H <sub>7</sub> O	1-Acetyethyl, 024
C <sub>2</sub> H <sub>3</sub> O <sub>3</sub>	(Carboxy)hydroxymethyl, 115		1-Hydroxycyclobutyl, 084
C <sub>2</sub> H <sub>4</sub> Cl	2-Chloroethyl, 005	C <sub>4</sub> H <sub>7</sub> O <sub>2</sub>	Radicals from crotyl alcohol + OH, 019
C <sub>2</sub> H <sub>4</sub> NO	Carbamoylmethyl, 034		2-Tetrahydrofuryl, 094
C <sub>2</sub> H <sub>4</sub> NO <sub>2</sub>	Amino(carboxy)methyl, 056		1-Carboxypropyl, 038
	(Carbamoyl)hydroxymethyl, 114	C <sub>4</sub> H <sub>7</sub> O <sub>3</sub>	2,5-Dioxacyclohexyl, 104
C <sub>2</sub> H <sub>4</sub> O <sub>4</sub> S <sup>-</sup>	1-Sulfatoethyl, 091		1-Hydroxy-1-methyl-2-oxopropyl, 107
C <sub>2</sub> H <sub>5</sub>	Ethyl, 002		4-Hydroxy-2-oxobutyl, 027
C <sub>2</sub> H <sub>5</sub> N <sub>2</sub> O	Amino(carbamoyl)methyl, 057		2-Carboxy-1-hydroxy-1-methylethyl, 097
C <sub>2</sub> H <sub>5</sub> O	1-Hydroxyethyl, 076		2-Carboxy-2-hydroxy-2-methylethyl, 017
	2-Hydroxyethyl, 014	C <sub>4</sub> H <sub>8</sub> NO	1-Carboxy-2-hydroxypropyl, 040
	Methoxymethyl, 092a	C <sub>4</sub> H <sub>8</sub> NO <sub>2</sub>	1-Hydroxy-1-(methoxycarbonyl)ethyl, 118
C <sub>2</sub> H <sub>5</sub> O <sub>2</sub>	1,2-Dihydroxyethyl, 101	C <sub>4</sub> H <sub>9</sub>	Radicals from <i>meso</i> -erythritol, 030
C <sub>2</sub> H <sub>6</sub> N	1-Aminoethyl, 049	C <sub>4</sub> H <sub>9</sub> O	( <i>N</i> -Acetyl- <i>N</i> -methylamino)methyl, 055
	2-Aminoethyl, 011		2-Amino-2-carboxy-2-methylethyl, 013
C <sub>2</sub> H <sub>6</sub> NO	2-Amino-1-hydroxyethyl, 098		1,1-Dimethylethyl, 002b
C <sub>2</sub> H <sub>7</sub> NO <sub>4</sub> P	2-Amino-1-phosphoryloxyethyl, 099		1-Ethoxyethyl, 093
C <sub>3</sub> H <sub>3</sub> O <sub>4</sub>	Dicarboxymethyl, 042		1-Hydroxybutyl, 079
C <sub>3</sub> H <sub>3</sub> O <sub>5</sub>	(Dicarboxy)hydroxymethyl, 119	C <sub>4</sub> H <sub>9</sub> O <sub>2</sub>	2-Hydroxy-2,2-dimethylethyl, 016
C <sub>3</sub> H <sub>4</sub> O <sub>2</sub> <sup>-</sup>	Electron adduct of acrylate, 134		1-Hydroxy-1-methylpropyl, 080
C <sub>3</sub> H <sub>5</sub> NO <sup>-</sup>	Electron adduct of acrylamide, 128	C <sub>4</sub> H <sub>9</sub> O <sub>3</sub>	1-Hydroxy-2-methylpropyl, 081
C <sub>3</sub> H <sub>5</sub> O	1-Formylethyl, 021		1,4-Dihydroxybutyl, 095
	2-Oxopropyl, 023	C <sub>4</sub> H <sub>9</sub> O <sub>2</sub>	1,2-Dihydroxy-1-methylpropyl, 103
	Radicals from allyl alcohol + OH, 018	C <sub>4</sub> H <sub>10</sub> N	Radicals from crotyl alcohol + OH, 019
C <sub>3</sub> H <sub>5</sub> O <sub>2</sub>	Acetoxymethyl, 090	C <sub>5</sub> H <sub>7</sub> NO <sub>3</sub>	2-Amino-2,2-dimethylethyl, 012
	1-Carboxyethyl, 037		Radical from 2-pyrrolidone-5-carboxylic acid, 074
	2,5-Dioxacyclopentyl, 125	C <sub>5</sub> H <sub>7</sub> O <sub>4</sub>	2-Oxocyclopentyl, 032
	1-Formyl-2-hydroxyethyl, 026	C <sub>5</sub> H <sub>7</sub> O <sub>5</sub>	1,3-Dicarboxy-1-hydroxypropyl, 122
	Methoxycarbonylmethyl, 036	C <sub>5</sub> H <sub>8</sub> NO <sub>3</sub>	<i>N</i> -Acetyl- <i>N</i> -methylamino(carboxy)methyl, 061
C <sub>3</sub> H <sub>5</sub> O <sub>3</sub>	1-Carboxy-1-hydroxyethyl, 117	C <sub>5</sub> H <sub>8</sub> O <sub>2</sub> <sup>-</sup>	Electron adduct of β,β-dimethylacrylate, 137
	1-Carboxy-2-hydroxyethyl, 039		Electron adduct of methyl methacrylate, 138
C <sub>3</sub> H <sub>6</sub> NO	( <i>N</i> -Acetylamino)methyl, 053	C <sub>5</sub> H <sub>9</sub>	Cyclopentyl, 004
	( <i>N</i> -Formyl- <i>N</i> -methylamino)methyl, 054	C <sub>5</sub> H <sub>9</sub> NO <sup>-</sup>	Electron adduct of <i>N,N</i> -dimethylacrylamide, 132
C <sub>3</sub> H <sub>6</sub> NO <sub>2</sub>	1-Carbamoyl-1-hydroxyethyl, 116		Electron adduct of β,β-dimethylacrylamide, 131
	Carboxy( <i>N</i> -methylamino)methyl, 059	C <sub>5</sub> H <sub>9</sub> N <sub>2</sub> O <sub>3</sub>	Radicals from acetylserineamide, 072
C <sub>3</sub> H <sub>6</sub> NO <sub>3</sub>	2-Amino-2-carboxy-1-hydroxyethyl, 100	C <sub>5</sub> H <sub>9</sub> O	1-Hydroxycyclopentyl, 085
C <sub>3</sub> H <sub>6</sub> O <sub>2</sub> <sup>-</sup>	Electron adduct of methyl acetate, 148		Radicals from cyclopentene + OH, 016a
C <sub>3</sub> H <sub>7</sub>	1-Methylethyl, 002a	C <sub>5</sub> H <sub>9</sub> O <sub>2</sub>	2-Carboxy-2,2-dimethylethyl, 006
C <sub>3</sub> H <sub>7</sub> O	1-Hydroxy-1-methylethyl, 078		1-Formyl-4-hydroxybutyl, 028
	1-Hydroxypropyl, 077	C <sub>5</sub> H <sub>9</sub> O <sub>4</sub>	Radicals from ribose + OH, 109
	2-Hydroxypropyl, 015		Radicals from deoxyribose + OH, 108
C <sub>3</sub> H <sub>7</sub> O <sub>2</sub>	1,2-Dihydroxypropyl, 102		Radicals from xylitol, 031
	Dimethoxymethyl, 124	C <sub>5</sub> H <sub>9</sub> O <sub>8</sub> P	Radicals from ribose phosphate + OH, 110
	(Methoxymethoxy)methyl, 094a	C <sub>5</sub> H <sub>11</sub>	Radicals from pentane, 003
C <sub>3</sub> H <sub>7</sub> O <sub>2</sub>	Radicals from allyl alcohol + OH, 018	C <sub>5</sub> H <sub>11</sub> N <sub>2</sub> O <sub>3</sub>	Radicals from glycy sarcosine + OH, 066
C <sub>3</sub> H <sub>7</sub> O <sub>3</sub>	1,2,3-Trihydroxypropyl, 106	C <sub>5</sub> H <sub>11</sub> O	1,1-Dimethylethoxymethyl, 093a
			1-Hydroxypentyl, 082
		C <sub>5</sub> H <sub>11</sub> O <sub>4</sub>	1,3-Dihydroxy-2,2-di(hydroxymethyl)propyl, 096
		C <sub>6</sub> H <sub>6</sub> O <sub>6</sub> <sup>-</sup>	Radical from ascorbate + OH, 113
		C <sub>6</sub> H <sub>8</sub> O <sub>2</sub> <sup>-</sup>	Electron adduct of sorbate, 139



$C_6H_8O_4^-$  Electron adduct of dimethyl fumarate, 141  
 $C_6H_9N_2O_2$  Radical from alanine anhydride + OH, 063  
Radical from sarcosine anhydride + OH, 064  
 $C_6H_9N_2O_4$  Radicals from acetyldiglycine +  $\bar{O}H$ , 067  
 $C_6H_9O$  2-Oxocyclohexyl, 025  
 $C_6H_{10}N_2O_4$  Radicals from acetylasparagine + OH, 073  
 $C_6H_{10}N_3O_4$  Radicals from triglycine + OH, 068  
 $C_6H_{11}N_2O_2^-$  Electron adduct of alanine anhydride, 143  
Electron adduct of sarcosine anhydride, 144  
 $C_6H_{11}O$  1-Hydroxycyclohexyl, 086  
 $C_6H_{11}O_2$  1-Formyl-5-hydroxypentyl, 029  
 $C_6H_{11}O_5$  Radicals from sorbitol, 033  
 $C_6H_{11}O_6$  Radicals from glucose + OH, 111  
 $C_6H_{12}N_2O_3^-$  Electron adduct of acetylglycylglycinamide, 146  
 $C_6H_{12}N_4O_2^-$  Electron adduct of Diamide, 133

$C_6H_{14}N$  1-(*N,N*-Diethylamino)ethyl, 051  
 $C_7H_{13}O$  (Cyclohexyl)hydroxymethyl, 083  
1-Hydroxycycloheptyl, 087  
 $C_8H_{12}N_3O_5$  Radicals from acetyltriglycine + OH, 069  
 $C_8H_{14}N_3O_5^-$  Electron adduct of acetyltriglycine, 145  
 $C_8H_{15}O$  1-Hydroxycyclooctyl, 088  
 $C_8H_{19}N^+$  Triethylammonioethyl, 052  
 $C_{11}H_{18}N_3O_5$  Radicals from acetyltrialanine + OH, 070  
Radicals from acetyltrisarcosine + OH, 071  
 $C_{12}H_{21}O_{12}$  Radicals from sucrose + OH, 112  
 $C_{12}H_{24}O_4S^-$  Radicals from sodium dodecylsulfate, 092  
 $C_{18}H_{28}O_2^-$  Radicals from linolenate + OH, 009  
 $C_{18}H_{30}O_2^-$  Radicals from linoleate + OH, 008  
 $C_{18}H_{32}O_2^-$  Radicals from oleate + OH, 007  
 $C_{20}H_{30}O_2^-$  Radicals from arachidonate + OH, 010

# Substrate Formula Index

Substrates for radical reactions are indexed by molecular formula. Entry numbers for the reactions in tables 1-3 are listed following the name.

Ag <sup>+</sup>	Silver(I) ion, 1	C <sub>3</sub> H <sub>7</sub> NO <sub>2</sub> S	Cysteine, 162
Ag <sub>2</sub> <sup>+</sup>	Silver(I) ion, complex with Ag(0), 2	C <sub>3</sub> H <sub>8</sub> O	1-Propanol, 313
BH <sub>3</sub> O <sub>3</sub>	Boric acid, 367	C <sub>3</sub> H <sub>10</sub> N <sup>+</sup>	2-Propanol, 314
BH <sub>4</sub> O <sub>4</sub> <sup>-</sup>	Tetrahydroborate ion, 366	C <sub>4</sub> H <sub>2</sub> N <sub>2</sub> O <sub>4</sub>	Trimethylammonium ion, 371
BrCoH <sub>15</sub> N <sub>5</sub> <sup>2+</sup>	Pentaamminebromocobalt(III) ion, 15	C <sub>4</sub> H <sub>3</sub> BrN <sub>2</sub> O <sub>2</sub>	Alloxan, 126
BrH <sub>15</sub> N <sub>5</sub> Ru <sup>2+</sup>	Bromopentaammineruthenium(III) ion, 101	C <sub>4</sub> H <sub>3</sub> NO <sub>2</sub> S	5-Bromouracil, 146
BrO <sub>3</sub> <sup>-</sup>	Bromate ion, 3	C <sub>4</sub> H <sub>3</sub> N <sub>3</sub> O <sub>4</sub>	2-Nitrothiophene, 298
Br <sub>2</sub> Hg	Mercuric bromide, 74	C <sub>4</sub> H <sub>3</sub> O <sub>4</sub> <sup>-</sup>	3-Nitrothiophene, 299
CCl <sub>4</sub>	Carbon tetrachloride, 152	C <sub>4</sub> H <sub>4</sub> N <sub>2</sub>	5-Nitouracil, 301
CHCl <sub>3</sub>	Chloroform, 157	C <sub>4</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub>	Fumarate ion, 214
CHO <sub>2</sub> <sup>-</sup>	Formate ion, 212	C <sub>4</sub> H <sub>4</sub> O <sub>4</sub>	Pyrazine, 321
CHO <sub>3</sub> <sup>-</sup>	Bicarbonate ion, 370	C <sub>4</sub> H <sub>5</sub> N <sub>3</sub>	Pyridazine, 324
CH <sub>2</sub> Cl <sub>2</sub>	Dichloromethane, 174	C <sub>4</sub> H <sub>5</sub> N <sub>3</sub> O <sub>2</sub>	Pyrimidine, 329
CH <sub>2</sub> O	Formaldehyde, 211	C <sub>4</sub> H <sub>6</sub>	2-Nitropyrrole, 295
CH <sub>3</sub> NO <sub>2</sub>	Nitromethane, 292	C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub>	3-Nitropyrrole, 296
CH <sub>4</sub> O	Methanol, 255	C <sub>4</sub> H <sub>6</sub> O <sub>2</sub>	Fumaric acid, 215
CH <sub>4</sub> S	Methanethiol, 254	C <sub>4</sub> H <sub>7</sub> N	Maleic acid, 252
CH <sub>5</sub> N	Mercurous cyanide, 77	C <sub>4</sub> H <sub>8</sub>	4-Aminopyrimidine, 129
CN <sub>4</sub> O <sub>8</sub>	Tetranitromethane, 347	C <sub>4</sub> H <sub>8</sub> O	2-Methyl-5-nitroimidazole, 264
C <sub>2</sub> Cl <sub>3</sub> O <sub>2</sub> <sup>-</sup>	Trichloroacetate ion, 351	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	Butadiene, 147
C <sub>2</sub> F <sub>3</sub> O <sub>2</sub> <sup>-</sup>	Trifluoroacetate ion, 352	C <sub>4</sub> H <sub>10</sub> N <sup>+</sup>	Glycine anhydride, 218
C <sub>2</sub> H <sub>2</sub>	Acetylene, 119	C <sub>4</sub> H <sub>10</sub> O	2,3-Butanedione, 148
C <sub>2</sub> H <sub>2</sub> IN	Iodoacetonitrile, 237	C <sub>4</sub> H <sub>10</sub> O <sub>2</sub> S <sub>2</sub>	Isobutyronitrile, 243
C <sub>2</sub> H <sub>2</sub> IO <sub>2</sub> <sup>-</sup>	Iodoacetate ion, 235	C <sub>4</sub> H <sub>12</sub> N <sup>+</sup>	1-Butene, 149
C <sub>2</sub> H <sub>3</sub> ClO <sub>2</sub>	Chloroacetic acid, 156	C <sub>4</sub> H <sub>12</sub> N <sub>2</sub> S <sub>2</sub>	Isobutylene, 241
C <sub>2</sub> H <sub>3</sub> Cl <sub>3</sub> O <sub>2</sub>	Chloral hydrate, 155	C <sub>4</sub> H <sub>16</sub> BrCoFN <sub>4</sub> <sup>+</sup>	Tetrahydrofuran, 340
C <sub>2</sub> H <sub>3</sub> IO <sub>2</sub>	Iodoacetic acid, 236	C <sub>4</sub> H <sub>16</sub> Br <sub>2</sub> CoN <sub>4</sub> <sup>+</sup>	Ethyl acetate, 201
C <sub>2</sub> H <sub>3</sub> N	Acetonitrile, 117	C <sub>4</sub> H <sub>16</sub> Cl <sub>2</sub> CoN <sub>4</sub> <sup>+</sup>	Isobutyric acid, 242
C <sub>2</sub> H <sub>4</sub>	Ethylene, 202	C <sub>4</sub> H <sub>16</sub> Cl <sub>2</sub> N <sub>4</sub> Pt <sup>2+</sup>	Methyl propionate, 267
C <sub>2</sub> H <sub>4</sub> Cu <sup>+</sup>	Copper(I) ion, complex with ethylene, 58	C <sub>4</sub> H <sub>16</sub> CoF <sub>2</sub> N <sub>4</sub> <sup>+</sup>	Pyrrrolidinium ion, 376
C <sub>2</sub> H <sub>4</sub> INO	Iodoacetamide, 234	C <sub>4</sub> H <sub>18</sub> ClCoN <sub>4</sub> O <sup>2+</sup>	Ethyl ether, 205
C <sub>2</sub> H <sub>4</sub> O <sub>2</sub>	Acetic acid, 115	C <sub>4</sub> H <sub>19</sub> ClCoN <sub>5</sub> <sup>2+</sup>	Dithiothreitol, 197
C <sub>2</sub> H <sub>4</sub> O <sub>3</sub>	Glycolic acid, 219	C <sub>4</sub> N <sub>4</sub> Ni <sup>2-</sup>	Diethylammonium ion, 373
C <sub>2</sub> H <sub>5</sub> IO	Iodoethanol, 238	C <sub>5</sub> FeN <sub>6</sub> O <sup>3-</sup>	Cystamine, 160a
C <sub>2</sub> H <sub>5</sub> NO <sub>2</sub>	Glycine, 217, 378	C <sub>5</sub> FeN <sub>6</sub> O <sup>4-</sup>	<i>cis</i> -Bromobis(ethylenediamine)
C <sub>2</sub> H <sub>6</sub> O	Ethanol, 200	C <sub>5</sub> H <sub>2</sub> NO <sub>5</sub> <sup>-</sup>	fluorocobalt(III) ion, 40
C <sub>2</sub> H <sub>6</sub> OS	2-Mercaptoethanol, 253	C <sub>5</sub> H <sub>3</sub> N <sub>2</sub> O <sub>4</sub> <sup>-</sup>	<i>trans</i> -Dibromobis(ethylenediamine)
C <sub>2</sub> H <sub>6</sub> O <sub>2</sub>	Ethylene glycol, 204	C <sub>5</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub>	cobalt(III) ion, 39
C <sub>2</sub> H <sub>7</sub> NS	Cysteamine, 161	C <sub>5</sub> H <sub>4</sub> N <sub>2</sub> O <sub>4</sub>	<i>cis</i> -Dichlorobis(ethylenediamine)
C <sub>2</sub> H <sub>8</sub> N <sup>+</sup>	Ethylammonium ion, 372	C <sub>5</sub> H <sub>4</sub> N <sub>4</sub>	cobalt(III) ion, 41
C <sub>2</sub> H <sub>15</sub> CoN <sub>5</sub> O <sub>2</sub> <sup>+</sup>	(Acetato)pentaamminecobalt(III) ion, 20	C <sub>5</sub> H <sub>4</sub> O <sub>5</sub> <sup>2-</sup>	<i>trans</i> -Dichlorobis(ethylenediamine)
C <sub>2</sub> HgN <sub>2</sub>	Mercuric cyanide, 78	C <sub>5</sub> H <sub>10</sub> O	cobalt(III) ion, 42
C <sub>2</sub> HgN <sub>2</sub> S <sub>2</sub>	Mercuric thiocyanate, 79	C <sub>5</sub> H <sub>11</sub> NO <sub>2</sub> S	<i>trans</i> -Dichlorobis(ethylenediamine)-
C <sub>3</sub> HBrNO <sub>2</sub> S	2-Bromo-5-nitrothiazole, 145	C <sub>5</sub> H <sub>12</sub> N <sup>+</sup>	platinum(IV) ion, 97
C <sub>3</sub> H <sub>3</sub> NO <sub>2</sub>	Cyanoacetic acid, 160	C <sub>5</sub> H <sub>15</sub> CoN <sub>6</sub> O <sub>2</sub> <sup>2+</sup>	<i>trans</i> -Bis(ethylenediamine)
C <sub>3</sub> H <sub>3</sub> NS	Thiazole, 349	C <sub>5</sub> H <sub>18</sub> CoN <sub>7</sub> O <sub>2</sub> <sup>2+</sup>	difluorocobalt(III) ion, 43
C <sub>3</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	2-Nitroimidazole, 289		<i>cis</i> -Aqua chlorobis(ethylene-
	4-Nitroimidazole, 290		diamine)cobalt(III) ion, 38
	2-Amino-5-nitrothiazole, 127		<i>cis</i> -Amminechlorobis(ethylene-
C <sub>3</sub> H <sub>3</sub> O <sub>3</sub> <sup>-</sup>	Pyruvate ion, 330		diamine)cobalt(III) ion, 37
C <sub>3</sub> H <sub>3</sub> IO <sub>2</sub>	3-Iodopropionic acid, 239		Tetracyanonickelate(II) ion, 90
C <sub>3</sub> H <sub>5</sub> N	Propionitrile, 316		Pentacyanonitrosylferrate(III) ion, 70
C <sub>3</sub> H <sub>5</sub> NO	Acrylamide, 123		Pentacyanonitrosylferrate(III) radical ion
	3-Hydroxypropionitrile, 230		(electron adduct), 71
C <sub>3</sub> H <sub>5</sub> O <sub>2</sub> S <sup>-</sup>	3-Mercaptopropionate ion, 253a		5-Nitrofuroate ion, 288
C <sub>3</sub> H <sub>5</sub> O <sub>3</sub> <sup>-</sup>	Lactate ion, 247		Orotate ion, 303
C <sub>3</sub> H <sub>6</sub>	Propylene, 317		Pyrazinecarboxylic acid, 322
C <sub>3</sub> H <sub>6</sub> O	Acetone, 116		Nifuroxime, 278
C <sub>3</sub> H <sub>6</sub> O <sub>2</sub>	Propionic acid, 315		Purine, 320
C <sub>3</sub> H <sub>6</sub> O <sub>3</sub>	Lactic acid, 248		α-Ketoglutarate ion, 246

$C_5H_{20}CoN_6^{3+}$	cobalt(III) ion, 33	$C_7H_8NO^+$	3-Carboxy-1-methylpyridinium ion, 153
$C_5H_{23}Co_2N_8O_4^{3+}$	Pentaammine(pyridine)cobalt(III) ion, 31	$C_7H_9ClN_2O$	4-Carboxy-1-methylpyridinium ion, 154
$C_6Cl_4O_2$	Hexaamminebis( $\mu$ -hydroxy)[ $\mu$ -(pyrazine-carboxylato)]dicobalt(III) ion, 35	$C_7H_9N_2O^+$	2-Pyridinecarboxaldoxime methochloride, 326
$C_6FeN_6^{3-}$	Tetrachlorobenzoquinone, 338	$C_7H_{11}N_3O_3$	3-Carbamoyl-1-methylpyridinium ion, 151
$C_6H_2N_3O_9S^-$	Ferricyanide ion, 69	$C_7H_{18}CoN_7O_6^{2+}$	Misonidazole, 272
$C_6H_4ClNO_2$	Trinitrobenzenesulfonate ion, 355		Pentaammine(2,4-dinitrobenzoato)cobalt(III) ion, 27
$C_6H_4N_2O_4$	<i>p</i> -Chloronitrobenzene, 158		Pentaammine(3,5-dinitrobenzoato)cobalt(III) ion, 28
	1,2-Dinitrobenzene, 187	$C_7H_{19}CoN_6O_4^{2+}$	Pentaammine(4-nitrobenzoato)cobalt(III) ion, 26
	1,3-Dinitrobenzene, 188		Pentaammine(2-nitrobenzoato)cobalt(III) ion, 24
	1,4-Dinitrobenzene, 189		Pentaammine(3-nitrobenzoato)cobalt(III) ion, 25
$C_6H_4N_4$	Pteridine, 318	$C_7H_{20}CoN_5O_2^{2+}$	Pentaammine(benzoato)cobalt(III) ion, 23
$C_6H_4N_4O_2$	Lumazine, 250	$C_7H_{24}Co_2N_7O_6^{3+}$	Hexaamminebis( $\mu$ -hydroxy)[ $\mu$ -(4-nitrobenzoato)]dicobalt(III) ion, 29
$C_6H_4O_2$	Benzoquinone, 138	$C_8H_3NO_6^{2-}$	2-Nitroisophthalate ion, 291
$C_6H_5NO$	Nitrosobenzene, 297	$C_8H_6NO_4^-$	3-Methyl-2-nitrobenzoate ion, 263
$C_6H_5NO_2$	Isonicotinic acid, 245	$C_8H_6N_2$	Quinoxaline, 331
	Nicotinic acid, 277	$C_8H_7NO_3$	<i>p</i> -Nitroacetophenone, 279
	Nitrobenzene, 283	$C_8H_8O$	Acetophenone, 118
$C_6H_5NO_3$	<i>o</i> -Nitrophenol, 294	$C_8H_8O_2$	2,3-Dimethylbenzoquinone, 180
$C_6H_5NO_4$	1,3-Dihydroxy-2-nitrobenzene, 178		2,5-Dimethylbenzoquinone, 181
$C_6H_5N_5O$	Pterin, 319	$C_8H_9NO_3$	2,6-Dimethylbenzoquinone, 182
$C_6H_5O^-$	Phenoxide ion, 309		3-Methoxy-2-nitrotoluene, 257
$C_6H_6CoN_3O_6^-$	Nitrioltriacetatocobalt(II) ion, 11	$C_8H_9O^-$	4-Ethylphenoxide ion, 207
$C_6H_6NO^-$	4-Aminophenoxide ion, 128	$C_8H_{10}NO_6P$	Pyridoxal-5-phosphate, 327
$C_6H_6N_2O$	Isonicotinamide, 244	$C_8H_{10}N_2O$	<i>N,N</i> -Dimethyl-4-nitrosoaniline, 185a
	Nicotinamide, 275	$C_8H_{11}NO_2$	2-(3,4-Dihydroxyphenyl)ethylamine, 179
$C_6H_6N_2O_2$	4-Pyridinecarboxaldoxime, 325	$C_8H_{11}NO_3$	Pyridoxine, 328
	<i>o</i> -Nitroaniline, 280	$C_8H_{12}CuN_4O_3$	Glycylhistidincopper(II) complex, 64
	<i>m</i> -Nitroaniline, 281	$C_8H_{12}NO_2$	Norpseudopelletierine- <i>N</i> -oxyl, 302
	<i>p</i> -Nitroaniline, 282	$C_8H_{13}O_2S_2^-$	Lipoate ion, 249
$C_6H_6N_4$	9-Methylpurine, 269	$C_8H_{19}N_6O_2^{2+}$	Pentaammine(4-cyanobenzoato)cobalt(III) ion, 30
$C_6H_6O_2$	1,2-Dihydroxybenzene, 176	$C_8H_{21}CoN_6O_4^{2+}$	Pentaammine(4-nitrophenylacetato)cobalt(III) ion, 22
	1,3-Dihydroxybenzene, 177		Pentaammine(phenylacetato)cobalt(III) ion, 21
	Hydroquinone, 227	$C_8H_{22}CoN_5O_2^{2+}$	3-Iodotyrosine, 240
$C_6H_7N$	Aniline, 130	$C_9H_{10}INO_3$	2,3,5-Trimethylbenzoquinone, 353
$C_6H_7NO_2$	<i>N</i> -Ethylmaleimide, 206	$C_9H_{10}O_2$	Tyrosine, 358
$C_6H_7O_6^-$	Ascorbate ion, 135	$C_9H_{11}NO_3$	2,2,5,5-Tetramethyl-3-pyrrolin-1-yloxy-3-carboxamide, 346
$C_6H_8N_2$	<i>o</i> -Phenylenediamine, 310	$C_9H_{15}N_2O_2$	2,2,6,6-Tetramethyl-4-oxo-1-piperidin-yloxy, 342
	<i>p</i> -Phenylenediamine, 311		2,2,5,5-Tetramethyl-1-pyrrolidin-yloxy-3-carboxamide, 345
$C_6H_8O_4$	Dimethyl fumarate, 184	$C_9H_{16}NO_2$	2,2,6,6-Tetramethyl-4-hydroxy-1-piperidin-yloxy, 343
$C_6H_9N_3O_3$	Metronidazole, 271		1,2,4,5-Tetracyanobenzene, 339
$C_6H_{11}CuN_3O_4^{2+}$	Glycylglycylglycinatocopper(II) complex, 61	$C_{10}H_2N_4$	1,4-Naphthoquinone-2-sulfonate ion, 274
	Diamide, 170	$C_{10}H_5O_5S^-$	1,4-Naphthoquinone, 273
$C_6H_{12}N_4O_2$	Cyclohexylammonium ion, 375	$C_{10}H_6O_2$	2-Hydroxy-1,4-naphthoquinone, 228
$C_6H_{14}N^+$	Triethylammonium ion, 374	$C_{10}H_6O_3$	2,2'-Bipyridine, 143
$C_6H_{16}N^+$	Pentaammine(pyridinecarboxylato)cobalt(III) ion, 32	$C_{10}H_8N_2$	Ethylenediaminetetraacetatocuprate(I) ion, 58a
$C_6H_{19}CoN_6O_2^{2+}$	Tris(ethylenediamine)cobalt(III) ion, 36	$C_{10}H_{12}CuN_2O_8^{3-}$	Ethylenediaminetetraacetatocuprate(II) ion, 65a
	2,4,6-Trinitrobenzoate ion, 356	$C_{10}H_{12}CuN_2O_8^{2-}$	Duroquinone, 198
$C_6H_{24}CoN_6^{3+}$	2,4-Dinitrobenzoate ion, 190	$C_{10}H_{12}O_2$	Adenosine, 124
$C_7H_2N_3O_8^-$	2,5-Dinitrobenzoate ion, 191	$C_{10}H_{13}N_5O_4$	$\beta$ -Alanylhistidincopper(II) complex, 65
$C_7H_3N_2O_6^-$	3,4-Dinitrobenzoate ion, 192	$C_{10}H_{14}CuN_4O_3$	Adenosine monophosphate, 125
	3,5-Dinitrobenzoate ion, 193	$C_{10}H_{14}N_5O_7P$	<i>N,N,N',N'</i> -Tetramethyl- <i>p</i> -phenylenediamine, 344
$C_7H_3N_3O_4$	3,5-Dinitrobenzonitrile, 194	$C_{10}H_{16}N_2$	Camphor, 150
$C_7H_4ClN$	<i>m</i> -Chlorobenzonitrile, 156a		Glutathione, 215a
$C_7H_4NO_4^-$	<i>o</i> -Nitrobenzoate ion, 284	$C_{10}H_{17}N_3O_6S$	1-(2,4-Dinitrophenyl)pyridinium ion, 195
	<i>m</i> -Nitrobenzoate ion, 285	$C_{11}H_8N_3O_4^+$	2-Methyl-1,4-naphthoquinone, 262
	<i>p</i> -Nitrobenzoate ion, 286	$C_{11}H_8O_2$	<i>N</i> -Formylkynurenine, 213
$C_7H_4NO_5^-$	3-Hydroxy-2-nitrobenzoate ion, 229	$C_{11}H_{12}N_2O_4$	1-Methylguanosine, 261
$C_7H_4N_2O_2$	<i>p</i> -Nitrobenzonitrile, 287	$C_{11}H_{15}N_5O_5$	Dichloroindophenol, 173
$C_7H_5NO_5$	<i>p</i> -Nitroperoxybenzoic acid, 293	$C_{12}H_7ClNO$	
$C_7H_6N_2O_5$	3,5-Dinitroanisole, 186		
$C_7H_6O_2$	2-Methylbenzoquinone, 259		
$C_7H_7NO$	3-Acetylpyridine, 120		
$C_7H_7NO_2$	<i>o</i> -Nitrotoluene, 300		
$C_7H_7NO_3$	3-Methyl-2-nitrophenol, 265		
$C_7H_7N_5O$	3-Methylpterin, 268		
$C_7H_7O^-$	4-Methylphenoxide ion, 266		
$C_7H_7O_2^-$	4-Methoxyphenoxide ion, 258		



$C_{12}H_8N_2$	1,10-Phenanthroline, 306 Phenazine, 307	$C_{19}H_{19}N_7O_6$	Folic acid, 210
$C_{12}H_9NO$	3-Benzoylpyridine, 141 4-Benzoylpyridine, 142 Indophenol, 233 Thionine, 350	$C_{20}H_6Br_4O_5^{2-}$	Eosin, 199
$C_{12}H_9N_3S$	Azobenzene, 136	$C_{20}H_{10}O_5^{2-}$	Fluorescein, 209
$C_{12}H_{10}N_2$	2,3-Dimethylnaphthoquinone, 185	$C_{20}H_{19}N_4^+$	Safranine T, 336
$C_{12}H_{10}O_2$	1,1'-Ethylene-2,2'-bipyridinium ion, 203	$C_{21}H_{28}N_7O_{10}P_2$	Nicotinamide-adenine dinucleotide, 276
$C_{12}H_{12}N_2^{2+}$	1,1'-Dimethyl-4,4'-bipyridinium ion, 183	$C_{22}H_{18}N_2^{2+}$	1,1'-Diphenyl-4,4'-bipyridinium ion, 196
$C_{12}H_{14}N_2^{2+}$	Thiamine, 348	$C_{24}H_{16}N_4^{2+}$	1,1'-Di(4-cyanophenyl)-4,4'-bipyridinium ion, 175
$C_{12}H_{17}ClN_4OS$	Histidinecopper(II) complex, 63	$C_{24}H_{22}N_2^{2+}$	1,1'-Dibenzyl-4,4'-bipyridinium ion, 171
$C_{12}H_{18}CuN_6O_4$	2-Benzoylpyridine, 140	$C_{25}H_{30}ClN_3^+$	Crystal Violet, 159
$C_{12}N_9NO$	Acridine, 121	$C_{28}H_{31}N_2O_3^+$	Rhodamine B, 332
$C_{13}H_9N$	Benzophenone, 137	$C_{30}H_{24}CoN_6^{3+}$	Tris(2,2'-bipyridine)cobalt(III) ion, 44
$C_{13}H_{10}O$	3-Benzoyl-1-methylpyridinium ion, 139	$C_{30}H_{24}N_6Rh^{3+}$	Tris(2,2'-bipyridine)rhodium(III) ion, 98
$C_{13}H_{12}NO^+$	1,1-Trimethylene-2,2'-bipyridinium ion, 354	$C_{30}H_{24}N_6Ru^{2+}$	Tris(2,2'-bipyridine)ruthenium(II) ion, 99
$C_{13}H_{14}N_2^{2+}$	9,10-Antraquinone-2,6-disulfonate ion, 134	$C_{30}H_{24}N_6Ru^{3+}$	Tris(2,2'-bipyridine)ruthenium(III) ion, 105
$C_{14}H_6O_7S_2^{2-}$	9,10-Antraquinone-1-sulfonate ion, 132	$C_{30}H_{28}FeN_4O_6$	Deuteroheme, 223
$C_{14}H_7O_5S^-$	9,10-Antraquinone-2-sulfonate ion, 133	$C_{30}H_{28}FeN_4O_6^+$	Deuterohemin, 222
$C_{14}H_8O_2$	9,10-Antraquinone, 131	$C_{31}H_{46}O_2$	Vitamin K <sub>1</sub> , 360
$C_{14}H_{14}ClN_3$	Acriflavin, 122	$C_{32}H_{34}N_4O_6$	Deuteroporphyrin dimethyl ester, 221a
$C_{14}H_{16}N_2^{2+}$	1,1'-Tetramethylene-2,2'-bipyridinium ion, 341	$C_{34}H_{24}N_6Na_4O_{14}S_4$	Trypan blue, 357
$C_{14}H_{18}N_2O_2^{2+}$	1,1'-Bis(2-hydroxyethyl)-4,4'-bipyridinium ion, 144	$C_{34}H_{32}ClFeN_4O_4$	Hemin, 224
$C_{14}H_{26}CoN_4O_2^+$	Dihydroxy(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(III) ion, 52	$C_{34}H_{38}N_4O_6$	Hematoporphyrin, 220
$C_{14}H_{27}CoN_4O_2^{2+}$	Aquahydroxy(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(III) ion, 51	$C_{36}H_{24}CoN_6^{3+}$	Tris(1,10-phenanthroline)cobalt(III) ion, 45
$C_{14}H_{28}CoN_4O_2^{2+}$	Diaqua(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(II) ion, 7	$C_{36}H_{24}N_6Rh^{3+}$	Tris(1,10-phenanthroline)iron(III) ion, 72
$C_{14}H_{28}CoN_4O_2^{3+}$	Diaqua(2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene)cobalt(III) ion, 50	$C_{36}H_{24}N_6Ru^{2+}$	Tris(1,10-phenanthroline)rhodium(III) ion, 98a
$C_{16}H_6N_2O_{14}S_4^{4-}$	Indigo tetrasulfonate, 232	$C_{40}H_{30}N_{10}O_6^{2+}$	Tris(1,10-phenanthroline)ruthenium(II) ion, 99a
$C_{16}H_8N_2O_8S_2^{2-}$	Indigo disulfonate, 231	$C_{40}H_{46}ClFeN_6O_8S_2$	Nitro blue tetrazolium, 287a
$C_{16}H_{10}$	Pyrene, 323	$C_{42}H_{58}CoN_6^{3+}$	Hemin c, 225
$C_{16}H_{16}N_2O_4$	1,1'-Dicarboxyethyl-4,4'-bipyridinium, 172	$C_{48}H_{30}N_4O_8$	Tris(5,6-dimethyl-1,10-phenanthroline)cobalt(III) ion, 45a
$C_{16}H_{18}N_3S^+$	Methylene blue, 260	$C_{62}H_{90}CoN_{13}O_{14}P$	<i>meso</i> -5-Tetra(4-carboxyphenyl)porphyrin, 337
$C_{16}H_{32}Cl_2CoN_4^+$	Dichloro(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(III) ion, 49	$C_{62}H_{91}CoN_{13}O_{15}P$	Cobal(II)amin, 10
$C_{16}H_{32}CuN_4^{2+}$	(5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(II) ion, 60	$C_{63}H_{90}CoN_{14}O_{14}P$	Hydroxocob(III)alamin(Vitamin B12a), 54
$C_{16}H_{32}N_4Ni^{2+}$	(5,7,7,12,12,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)nickel(II) ion, 92	$C_{64}H_{24}CoN_{16}O_{24}S_8^{8-}$	Cyanocob(III)alamin(Vitamin B12), 53
$C_{16}H_{34}CoN_4O_2^+$	(5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)dihydroxycobalt(III) ion, 48	$Cd^+$	3,10,17,24-Tetrasulfophthalocyaninecobalt(II) ion dimer, 10a
$C_{16}H_{35}CoN_4O_2^{2+}$	Aqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)hydroxycobalt(III) ion, 47	$Cd^{2+}$	Cadmium(I) ions, 4
$C_{16}H_{36}CoN_4O_2^{2+}$	Diaqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(II) ion, 9	$ClCoH_{15}N_5S^{2+}$	Cadmium(II) ion, 5
$C_{16}H_{36}CoN_4O_2^{3+}$	Diaqua(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(III) ion, 46	$ClH_{15}N_5Ru^{2+}$	Pentaamminechlorocobalt(III) ion, 16
$C_{16}H_{36}N_4Ni^{2+}$	(5,7,7,12,12,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)nickel(II) ion, 91	$Cl_2Hg$	Chloropentaammineruthenium(III) ion, 102
$C_{17}H_{20}N_2S$	Promethazine, 312	$Cl_2Ir^{2-}$	Mercuric chloride, 75
$C_{17}H_{20}N_4O_6$	Riboflavin, 333	$Co^{2+}$	Hexachloroiridate(II) ion, 81a
$C_{17}H_{20}N_4O_9P^-$	Riboflavine 5'-phosphate, 334	$CoFH_{15}N_5S^{2+}$	Cobalt(II) ions, 6
$C_{18}H_{15}N_4^+$	Phenosafarine, 308	$CoH_{15}N_6O_2^{2+}$	Pentaamminefluorocobalt(III) ion, 17
		$CoH_{15}N_6O_3^{2+}$	Pentaammine(nitrito- <i>N</i> )cobalt(III) ion, 18
		$CoH_{16}N_5O^{2+}$	Pentaammine(nitrato- <i>O</i> )cobalt(III) ion, 19
		$CoH_{17}N_5O^{3+}$	Pentaamminehydroxycobalt(III) ion, 14
		$Cr^{2+}$	Pentaammineaquacobalt(III) ion, 13
		$Cr^{3+}$	Hexaamminecobalt(III) ion, 12
		$Cu^+$	Chromium(II) ion, 55
		$Cu^{2+}$	Chromium(III) ion, 56
		$Eu^{3+}$	Copper(I) ions, 57
		$Fe^{2+}$	Copper(II) ions, 59
		$Fe^{3+}$	Europium(III) ion, 66
		$Ga^{2+}$	Iron(II) ion, 67
		$HO^-$	Iron(III) ions, 68
		$HO_2P_2^{3-}$	Gallium(II) ions, 73
		$H_2O$	Hydroxide ion, 362
		$H_2O_2$	Pyrophosphate ion, 369
		$H_2O_4P^-$	Water, 363
		$H_2S$	Hydrogen peroxide, 94
		$H_3N$	Phosphate ion, 368
		$H_3NO$	Hydrogen sulfide, 106
		$H_3O^+$	Ammonia, 365
		$H_4N^+$	Hydroxylamine, 84
		$H_{15}N_6ORu^{3+}$	Hydronium ion, 361
			Ammonium ion, 364
			Nitrosopentaammineruthenium(III) ion, 103

$\text{H}_{15}\text{N}_6\text{ORu}^{2+}$	Nitrosopentaammineruthenium(III) radical ion (electron adduct), 104
$\text{H}_{18}\text{N}_6\text{Ru}^{3+}$	Hexaammineruthenium(III) ion, 100
$\text{HgI}_2$	Mercuric iodide, 76
$\text{IO}_3^-$	Iodate ion, 81
$\text{I}_2$	Iodine, 80
$\text{Mn}^{2+}$	Manganese(II) ions, 82
$\text{MnO}_4^-$	Permanganate ion, 83
$\text{NO}_2^-$	Nitrite ion, 86
$\text{NO}_3^-$	Nitrate ion, 87
$\text{N}_2\text{O}$	Nitrous oxide, 85
$\text{Ni}^+$	Nickel(I) ion, 88
$\text{Ni}^{2+}$	Nickel(II) ions, 89

$\text{O}_2$	Oxygen, 93
$\text{O}_2\text{S}$	Sulfur dioxide, 107
$\text{O}_6\text{S}_4^{2-}$	Tetrathionate ion, 108
$\text{Pb}^+$	Lead(I) ions, 95
$\text{Pb}^{2+}$	Lead(II) ions, 96
$\text{Sc}^{2+}$	Scandium(II) ions, 109
$\text{Ti}^{3+}$	Titanium(III) ions, 110
$\text{Ti}^+$	Thallium(I) ion, 111
$\text{Ti}_2^+$	Thallium(I) ion, complex with $\text{Tl}(0)$ , 111a
$\text{Yb}^{3+}$	Ytterbium(III) ions, 112
$\text{Zn}^+$	Zinc(I) ion, 113
$\text{Zn}^{2+}$	Zinc(II) ion, 114



## References

- 61G019 Kelly, P., Smith, M., The radiation chemistry of azide solutions. Part II. The action of x-rays (200 kv) on deaerated aqueous solutions of sodium azide in the presence of different organic solutes, *J. Chem. Soc.* 1487-94 (1961).
- 62G140 Taub, I.A., Dorfman, L.M., Pulse radiolysis studies. II. Transient spectra and rate processes in irradiated ethanol and aqueous ethanol solution, *J. Am. Chem. Soc.* **84**: 4053-9 (1962).
- 63F005 Buxton, G., Wilmarth, W.K., Aqueous chemistry of inorganic free radicals. V. The use of carbon monoxide as a scavenger for hydroxyl radicals generated by the photolysis of hydrogen peroxide, *J. Phys. Chem.* **67**: 2835-41 (1963).
- 63G045 Dorfman, L.M., Taub, I.A., Pulse radiolysis studies. III. Elementary reactions in aqueous ethanol solution, *J. Am. Chem. Soc.* **85**: 2370-4 (1963).
- 64G133 Asmus, K.-D., Henglein, A., Ebert, M., Keene, J.P., Pulsradiolytische Untersuchung schneller Reaktionen von hydratisierten Elektronen, freien Radikalen und Ionen mit Tetranitromethan in waessriger Loesung, *Ber. Bunsenges. Phys. Chem.* **68**: 657-63 (1964).
- 65D040 Fischer, H., Rapid proton exchange of the free radical  $\cdot\text{CH}_2\text{OH}$  as studied by E.S.R., *Mol. Phys.* **9**: 149-52 (1965).
- 65F006 Dainton, F.S., Fowles, P., The photolysis of aqueous systems at 1849 Å. I. Solutions containing nitrous oxide, *Proc. Roy. Soc. (London) Ser. A* **287**: 295-311 (1965).
- 65G183 Rabani, J., Mulac, W.A., Matheson, M.S., The pulse radiolysis of aqueous tetranitromethane. I. Rate constants and the extinction coefficient of  $e_{aq}^-$ . II. Oxygenated solutions, *J. Phys. Chem.* **69**: 53-70 (1965).
- 65G375 Guetlbauer, F., Getoff, N., Kinetic studies on the radiation-chemistry carboxylation of methanol, *Z. Phys. Chem. (Frankfurt Am Main)* **47**: 299-305 (1965).
- 65G384 Keene, J.P., Raef, Y., Swallow, A.J., Pulse radiolysis studies of carboxyl and related radicals, *Pulse Radiolysis*, Ebert, M.; Keene, J.P.; Swallow, A.J.; Baxendale, J.H. (eds.), Academic Press, New York, 1965, p.99-106.
- 65G396 Keene, J.P., Land, E.J., Swallow, A.J., A pulse radiolysis study of methylene blue, *Pulse Radiolysis*, Ebert, M.; Keene, J.P.; Swallow, A.J.; Baxendale, J.H. (eds.), Academic Press, New York, 1965, p.227-45.
- 66D162 Zeldes, H., Livingston, R., Paramagnetic resonance study of liquids during photolysis. II. Acetone and solutions containing acetone, *J. Chem. Phys.* **45**(6): 1946-54; (1966).
- 66G074 Asmus, K.-D., Henglein, A., Wigger, A., Beck, G., Pulsradiolytische Versuche zur elektrolytischen Dissoziation von aliphatischen Alkoholradikalen, *Ber. Bunsenges. Phys. Chem.* **70**(7): 756-8 (1966).
- 66G316 Guetlbauer, F., Getoff, N., Bestimmung des  $g(e_{aq}^-)$ -Wertes und der Dissoziationskonstante des  $\text{COOH}$ -Radikals in waessriger Methanolloesung, *Z. Phys. Chem. (Frankfurt Am Main)* **51**(5-6): 255-63 (1966).
- 66G432 Asmus, K.-D., Wigger, A., Henglein, A., Pulsradiolytische Untersuchung einiger Elementarprozesse der Nitrobenzol reduktion, *Ber. Bunsenges. Phys. Chem.* **70**(6): 862-8 (1966).
- 66G433 Asmus, K.-D., Beck, G., Henglein, A., Wigger, A., Pulsradiolytische Untersuchung der Oxydation und Reduktion des Nitrosobenzols in waessriger Loesung, *Ber. Bunsenges. Phys. Chem.* **70**(6): 869-74 (1966).
- 67E053 Pruetz, W., Land, E.J., Lichtemission von waessrigen Farbstoffloesungen nach Bestrahlung mit Elektronenimpulsen, *Biophysik* **3**: 349-60 (1967).
- 67G038 Chrysochoos, J., Ovadia, J., Grossweiner, L.I., Pulse radiolysis of aqueous eosin, *J. Phys. Chem.* **71**(6): 1629-36 (1967).
- 67G041 Thomas, J. K., Pulse radiolysis of aqueous solutions of methyl iodide and methyl bromide. The reactions of iodine atoms and methyl radicals in water, *J. Phys. Chem.* **71**(6): 1919-1925 (1967).
- 67G094 Seddon, W.A., Allen, A.O., Radiation chemistry of aqueous solutions of ethanol, *J. Phys. Chem.* **71**: 1914-8 (1967).
- 67G102 Basson, R.A., van der Linde, H.J., Radical yield of propionaldehyde in the radiolysis of n-propanol, *Chem. Commun.* (2): 91-2 (1967).
- 67G262 Karmann, W., Henglein, A., Pulse-radiolytic investigations of the reactions of organic radicals with hydrogen sulfide, *Ber. Bunsenges. Phys. Chem.* **71**(4): 421-4 (1967).
- 67G269 Cullis, C.F., Francis, J.M., Raef, Y., Swallow, A.J., Studies of radiation-induced reactions of ethylene in aqueous solution. II. Reactions in the presence of oxygen as studied by pulse radiolysis and  $\gamma$ -irradiation techniques, *Proc. Roy. Soc. (London) Ser. A* **300**: 443-54 (1967).
- 67G333 Sibirskaya, G.K., Pikaev, A.K., The radiolysis of aqueous sulfuric acid solutions of methyl alcohol, *Bull. Acad. Sci. USSR, Div. Chem. Sci. (Engl. Transl.)* (1): 183-5 (1967), *Transl. from Izv. Akad. Nauk SSSR, Ser. Khim.* (1): 190-2 (1967).
- 67G729 Adams, G.E., Michael, B.D., Richards, J.T., Three stage electron transfer in aqueous and alcoholic solutions, *Nature (London)* **215**: 1248-50 (1967).
- 68G132 Adams, G.E., McNaughton, G.S., Michael, B.D., Pulse radiolysis of sulphur compounds. Part 2.- Free radical repair by hydrogen transfer from sulphydryl compounds, *Trans. Faraday Soc.* **64**(4): 902-10 (1968).
- 68G172 Cordier, P., Grossweiner, L.I., Pulse radiolysis of aqueous fluorescein, *J. Phys. Chem.* **72**(6): 2018-26 (1968).
- 68G249 Lilie, J., Beck, G., Henglein, A., Pulsradiolytische Untersuchung des Acetoinradikals und des Diacetylanions in waessriger Loesung, *Ber. Bunsenges. Phys. Chem.* **72**(4): 529-33 (1968).
- 68G308 Adams, G.E., Michael, B.D., Willson, R.L., Electron transfer studies by pulse radiolysis, *Adv. Chem. Ser.* **81**: 289-308 (1968).
- 68G441 Land, E.J., Swallow, A.J., One-electron reactions in biochemical systems as studied by pulse radiolysis I. Nicotinamide-adenine dinucleotide and related compounds, *Biochim. Biophys. Acta* **162**: 327-37 (1968).
- 68G539 Volkert, W.A., Some radical reactions with amino acids in aqueous solutions, Ph.D. Thesis, Univ. of Missouri, 1968, 164p.
- 68M058 Poupo, R., Loewenstein, A., Kinetic study of proton exchange in radicals derived from simple alcohols, *J. Chem. Soc. B* 949-51 (1968).
- 69F176 Taha, I.A.I., Kuntz, R.R., Methyl radical reactions in aqueous solutions. I. Hydrogen abstraction from acetone, *J. Phys. Chem.* **73**(12): 4406-9 (1969).
- 69G019 Shankar, J., Rama Rao, K.V.S., Shastri, L.V., Peroxide formation in the  $\gamma$  radiolysis of aerated aqueous solutions of methyl iodide, *J. Phys. Chem.* **73**(1): 52-7 (1969).
- 69G052 Buxton, G.V., Dainton, F.S., Kalecinski, J., The radiation chemistry of aqueous solutions of sodium nitroprusside, *Int. J. Radiat. Phys. Chem.* **1**: 87-98 (1969).
- 69G156 Baxendale, J.H., Khan, A.A., The pulse radiolysis of p-nitrosodimethylaniline in aqueous solution, *Int. J. Radiat. Phys. Chem.* **1**(1): 11-24 (1969).
- 69G238 Land, E.J., Swallow, A.J., One-electron reactions in biochemical systems as studied by pulse radiolysis. II. Riboflavin, *Biochemistry* **8**: 2117-25 (1969).
- 69G270 Gruenbein, W., Henglein, A., Pulsradiolytische Untersuchung zweibasischer Radikale der Reduktion von Nitrophenolen in waessriger Loesung, *Ber. Bunsenges. Phys. Chem.* **73**(4): 376-82 (1969).
- 69G419 Simic, M., Neta, P., Hayon, E., Pulse radiolysis of alcohols in aqueous solution, *J. Phys. Chem.* **73**(11): 3794-800 (1969).
- 69G446 Neta, P., Simic, M., Hayon, E., Pulse radiolysis of aliphatic acids in aqueous solutions. I. Simple monocarboxylic acids, *J. Phys. Chem.* **73**(12): 4207-13 (1969).



- 69G447 Simic, M., Neta, P., Hayon, E., Pulse radiolysis of aliphatic acids in aqueous solutions. II. Hydroxy and polycarboxylic acids, *J. Phys. Chem.* **73**(12): 4214-9 (1969).
- 69G522 Adams, G.E., Willson, R.L., Pulse radiolysis studies on the oxidation of organic radicals in aqueous solution, *Trans. Faraday Soc.* **65**: 2981-7 (1969).
- 69G524 Morita, M., Tajima, M., Fujimaki, M., Radiation chemistry of foods. Part II. Reactivity of some food constituents with a model secondary radical,  $\text{CH}_2\text{OH}$ , *Agric. Biol. Chem.* **33**(2): 250-5 (1969).
- 69G553 Karmann, W., Granzow, A., Meissner, G., Henglein, A., Die Pulsradiolyse einfacher Mercaptane in Luft freier waessriger Loesung, *Int. J. Radiat. Phys. Chem.* **1**: 395-405 (1969).
- 69G826 Zimbrick, J.D., Ward, J.F., Myers, L.S.Jr., Studies on the chemical bases of cellular radiosensitization by 5-bromouracil substitution in DNA. I. Pulse and steady-state radiolysis of 5-bromouracil and thymine, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **16**(6): 505-23 (1969).
- 70G052 Chambers, K.W., Collinson, E., Dainton, F.S., Addition of  $e^-_{aq}$ , H- and  $\cdot\text{OH}$  to acrylamide in aqueous solution and reactions of the adducts, *Trans. Faraday Soc.* **66**: 142-62 (1970).
- 70G080 Wander, R., Gall, B.L., Dorfman, L.M., The absolute reactivity of the oxide radical ion with methanol and ethanol in water, *J. Phys. Chem.* **74**(8): 1819-21 (1970).
- 70G104 Burchill, C.E., Ginns, I.S., Radiation-induced oxidation of 2-propanol by hydrogen peroxide in aqueous solutions, *Can. J. Chem.* **48**: 1232-8 (1970).
- 70G241 Pruetz, W.A., Land, E.J., Chemiluminescent reactions after pulse radiolysis of aqueous solutions of acriflavin. Effects of halides and pseudo halides, *J. Phys. Chem.* **74**(10): 2107-14 (1970).
- 70G254 Barker, G.C., Fowles, P., Stringer, B., Pulse radiolytic induced transient electrical conductance in liquid solutions. Part 2. Radiolysis of aqueous solutions of  $\text{NO}_3^-$ ,  $\text{NO}_2^-$  and  $\text{Fe}(\text{CN})_6^{3-}$ , *Trans. Faraday Soc.* **66**(6): 1509-19 (1970).
- 70G303 Fojtik, A., Czapski, G., Henglein, A., Pulse radiolytic investigation of the carboxyl radical in aqueous solution, *J. Phys. Chem.* **74**(17): 3204-8 (1970).
- 70G338 Burchill, C.E., Ginns, I.S., The radiation-induced oxidation of ethanol and methanol by hydrogen peroxide in aqueous solution, *Can. J. Chem.* **48**(16): 2628-32 (1970).
- 70G407 Bullock, G., Cooper, R., Reactions of aqueous trifluoromethyl radicals, *Trans. Faraday Soc.* **66**: 2055-64 (1970).
- 70G560 Willson, R.L., Pulse radiolysis studies of electron transfer in aqueous disulphide solutions, *Chem. Commun.* (21): 1425-6 (1970).
- 70R013 Willson, R.L., Pulse radiolysis studies of electron transfer reactions in aerobic solution, *Chem. Commun.* (16): 1005 (1970).
- 71D227 Burchill, C.E., Jones, P.W., Electron spin resonance studies of reactions of photo-generated organic radicals with hydrogen peroxide in aqueous solution, *Can. J. Chem.* **49**(24): 4005-16 (1971).
- 71F579 Kantrowitz, E.R., Hoffman, M.Z., Endicott, J.F., Ultraviolet photochemistry of acetatopentaammine cobalt(III) in aqueous solution, *J. Phys. Chem.* **75**(12): 1914-20 (1971).
- 71G026 Balkas, T.I., Fendler, J.H., Schuler, R.H., The radiation chemistry of aqueous solutions of  $\text{CFCl}_3$ ,  $\text{CF}_2\text{Cl}_2$ , and  $\text{CF}_3\text{Cl}$ , *J. Phys. Chem.* **75**(4): 455-66 (1971).
- 71G041 Getoff, N., Schwoerer, F., Markovic, V.M., Sehested, K., Nielsen, S.O., Pulse radiolysis of oxalic acid and oxalates, *J. Phys. Chem.* **75**(6): 749-55 (1971).
- 71G061 Fielden, E.M., Roberts, P.B., Pulse radiolysis studies of the radiosensitizer nor-pseudo-pelletierine-N-oxyl (NPPN). I. Radiation chemistry, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **20**(4): 355-62 (1971).
- 71G284 Neta, P., Electron spin resonance study of irradiated aqueous solutions of fumarate ion. The use of fumarate for radical trapping, *J. Phys. Chem.* **75**(17): 2570-4 (1971).
- 71G327 Land, E.J., Swallow, A.J., One-electron reactions in biochemical systems as studied by pulse radiolysis. V. Cytochrome c, *Arch. Biochem. Biophys.* **145**: 365-72 (1971).
- 71G424 Stockhausen, K., Henglein, A., Pulsradiolytische Untersuchung des Mechanismus der Oxidation und Autoxidation von Formaldehyd in waessriger Loesung, *Ber. Bunsenges. Phys. Chem.* **75**(8): 833-40 (1971).
- 71G554 Hayon, E., Simic, M., Pulse radiolysis study of cyclic peptides in aqueous solution. Absorption spectrum of the peptide radical  $-\text{NHCHCO}-$ , *J. Am. Chem. Soc.* **93**(25): 6781-6 (1971).
- 71G618 Willson, R.L., Pulse radiolysis studies on reaction of triacetoneamine-N-oxyl with radiation-induced free radicals, *Trans. Faraday Soc.* **67**(10): 3008-19 (1971).
- 71G619 Willson, R.L., Pulse radiolysis studies of electron transfer in aqueous quinone solutions, *Trans. Faraday Soc.* **67**(10): 3020-9 (1971).
- 71G778 Koester, R., Asmus, K.-D., Die Reduktion von Tetrachlorkohlenstoff durch hydratisierte Elektronen, H-Atome und reduzierende Radikale, *Z. Naturforsch. Teil B* **26**(11): 1104-8 (1971).
- 71G929 Byakov, V.M., Zhiltsova, L.V., Kalyazin, E.P., Kovalyova, E.P., Petryaev, E.P., On the mechanism of radiolysis of aqueous methanol solutions, *Proc. of the Third Tihany Symposium on Radiation Chemistry*, Vol. 2, J. Dobo and P. Hedvig (eds.), Akademiai Kiado, Budapest, Hungary, 1972, p.1301-9.
- 72A002 Nilsson, K., The reduction of ferricytochrome c studied by pulse radiolysis, *Isr. J. Chem.* **10**(6): 1011-9 (1972).
- 72A006 Adams, G.E., Redpath, J.L., Bisby, R.H., Cundall, R.B., The use of free radical probes in the study of mechanisms of enzyme inactivation, *Isr. J. Chem.* **10**(6): 1079-93 (1972).
- 72A018 Cohen, H., Meyerstein, D., On the mechanism of reduction of cobalt(III) and ruthenium(III) hexaammine complexes by several aliphatic radicals, *J. Am. Chem. Soc.* **94**(20): 6944-8 (1972).
- 72D181 Laroff, G.P., Fessenden, R.W., Schuler, R.H., The electron spin resonance spectra of radical intermediates in the oxidation of ascorbic acid and related substances, *J. Am. Chem. Soc.* **94**(26): 9062-73 (1972).
- 72G070 Cohen, H., Meyerstein, D., Formation of chromium-carbon bonds in aqueous solution: A pulse radiolytic study, *J. Chem. Soc., Chem. Commun.* (6): 320-1 (1972).
- 72G144 Hayon, E., Simic, M., Radiation sensitization reactions of N-ethylmaleimide with model compounds, *Radiat. Res.* **50**(3): 464-78 (1972).
- 72G167 Burchill, C.E., Wollner, G.P., Radiation-induced oxidation of 2-propanol by nitrous oxide in alkaline aqueous solution, *Can. J. Chem.* **50**(11): 1751-6 (1972).
- 72G244 Lilie, J., Behar, D., Sujdak, R.J., Schuler, R.H., On the lifetime of trifluoromethyl radical in aqueous solution, *J. Phys. Chem.* **76**(18): 2517-20 (1972).
- 72G340 Hoffman, M.Z., Simic, M., Ligand-to-metal intramolecular electron transfer in the reduction of p-nitrobenzoato pentaamminecobalt(III) ion, *J. Am. Chem. Soc.* **94**(5): 1757-9 (1972).
- 72G359 Nelson, D.A., Hayon, E., Ketyl radicals of benzoylpyridines, *J. Phys. Chem.* **76**(22): 3200-7 (1972).
- 72G388 Hoffman, M.Z., Hayon, E., One-electron reduction of the disulfide linkage in aqueous solution. Formation, protonation, and decay kinetics of the  $\text{RSSR}^-$  radical, *J. Am. Chem. Soc.* **94**(23): 7950-7 (1972).
- 72G391 Hulme, B.E., Land, E.J., Phillips, G.O., Pulse radiolysis of 9,10-anthraquinones. Part I. Radicals, *J. Chem. Soc., Faraday Trans. 1* **68**(10): 1992-2002 (1972).
- 72G445 Stevens, G.C., Clarke, R.M., Hart, E.J., Radiolysis of aqueous methane solutions, *J. Phys. Chem.* **76**(25): 3863-7 (1972).
- 72G462 Martin, J.E., Hart, E.J., Adamson, A.W., Gafney, H., Halpern, Jack, Chemiluminescence from the reaction of the hydrated electron with tris(bipyridyl)ruthenium(III), *J. Am. Chem. Soc.* **94**(26): 9238-40 (1972).
- 72G610 Stelter, L.H., Chemical effects of direct and indirect interactions of X-radiation with iodine containing amino acids, Ph.D., Thesis, Univ. of Toledo, Ohio, 1972, 115p.
- 72R003 Nucifora, G., Smaller, B., Remko, R., Avery, E.C., Transient radicals of DNA bases by pulse radiolysis. Effects of cysteine and cysteamine as radioprotectors, *Radiat. Res.* **49**: 96-111 (1972).



- 72R046 Blackburn, R., Cox, D.L., Phillips, G.O., Effects of gamma radiation on vitamin B12 systems, *J. Chem. Soc., Faraday Trans. 1* **68**(9): 1687-96 (1972).
- 72R057 Simic, M., Hayon, E., A model of radiation sensitization by quinones, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **22**(5): 507-11 (1972).
- 73A020 Redpath, J.L., Pulse radiolysis of dithiothreitol, *Radiat. Res.* **54**(3): 364-74 (1973).
- 73A027 Schoeneshoefer, M., Pulsradiolytische Untersuchung ueber das  $S_4O_6^{3-}$ -Radikal als Zwischenprodukt der Thiosulfatoxidation und Tetrathionatreduktion, *Int. J. Radiat. Phys. Chem.* **5**(4): 375-86 (1973).
- 73A047 Rao, P.S., Hayon, E., One-electron redox reactions of free radicals in solution. Rate of electron transfer processes to quinones, *Biochim. Biophys. Acta* **292**: 516-33 (1973).
- 73A052 Hayon, E., Simic, M., Acid-base properties of organic peroxy radicals,  $\cdot OORH$ , in aqueous solution, *J. Am. Chem. Soc.* **95**(20): 6681-4 (1973).
- 73A057 Ellis, J.D., Green, M., Sykes, A.G., Buxton, G.V., Sellers, R.M., Pulse radiolysis of titanium(III) and other metal(III) ions in the presence of formic acid, *J. Chem. Soc., Dalton Trans.* (16): 1724-8 (1973).
- 73A062 Greenstock, C.L., Dunlop, I., Kinetics of competing free radical reactions with nitroaromatic compounds, *J. Am. Chem. Soc.* **95**(21): 6917-9 (1973).
- 73A064 Pecht, I., Pulse radiolysis study of the reduction of spinach ferredoxin, *FEBS (Fed. Eur. Biochem. Soc.) Lett.* **33**(2): 259-61 (1973).
- 73A070 Simic, M., Neta, P., Hayon, E., Reactions of hydroxyl radicals with unsaturated aliphatic alcohols in aqueous solution. A spectroscopic and electron spin resonance radiolysis study, *J. Phys. Chem.* **77**(22): 2662-7 (1973).
- 73A073 Purdie, J.W., Gillis, H.A., Klassen, N.V., The pulse radiolysis of penicillamine and penicillamine disulfide in aqueous solution, *Can. J. Chem.* **51**(18): 3132-42 (1973).
- 73A074 Farrington, J.A., Ebert, M., Land, E.J., Fletcher, K., Bipyridylum quaternary salts and related compounds. V. Pulse radiolysis studies of the reaction of paraquat radical with oxygen. Implications for the mode of action of bipyridyl herbicides, *Biochim. Biophys. Acta* **314**(22): 372-81 (1973).
- 73A075 Hoffman, M.Z., Simic, M., Rate of reduction of cobalt(III) complexes by  $CO_2^{\cdot -}$  radicals in aqueous solution, *Inorg. Chem.* **12**(10): 2471-2 (1973).
- 73A078 Rao, P.S., Hayon, E., Reduction of dyes by free radicals in solution. Correlation between reaction rate constants and redox potentials, *J. Phys. Chem.* **77**(23): 2753-6 (1973).
- 73A099 Greenstock, C.L., Dunlop, I., Pulse radiolysis studies of nitrofurans: Chemical radiosensitization, *Radiat. Res.* **56**(3): 428-40 (1973).
- 73A102 Bansal, K.M., Henglein, A., Janata, E., Sellers, R.M., Pulse radiolysis and polarography. Influence of the double layer capacity on electrode reactions of short lived radicals in aqueous solutions, *Ber. Bunsenges. Phys. Chem.* **77**(12): 1139-44 (1973).
- 73A104 Rao, P.S., Hayon, E., Rate constants of electron transfer processes in solution: Dependence on the redox potential of the acceptor, *Nature (London)* **243**: 344-6 (1973).
- 73A140 Willson, R.L., Slater, T.F., Carbon tetrachloride and biological damage: pulse radiolysis studies of associated free radical reactions, *Fast Processes in Radiation Chemistry and Biology*, G.E. Adams, E.M. Fielden and B.D. Michael (eds.), John Wiley & Sons, 1975, p.147-61 (Proceedings of the Fifth L.H. Gray Conference, University of Sussex, Sept. 10-14, 1973).
- 73A146 Faraggi, M., Electron transfer reactions in structural units of copper proteins, *Ibid.*, p.285-308.
- 73A150 Willson, R.L., Free radical chain oxidation by carbon tetrachloride and related compounds: Model pulse radiolysis studies, *Trans. Biochem. Soc.* **1**: 929-31 (1973).
- 73D065 Laroff, G.P., Fessenden, R.W., Equilibrium and kinetics of the acid dissociation of several hydroxyalkyl radicals, *J. Phys. Chem.* **77**(10): 1283-8 (1973).
- 73F161 Mittal, L.J., Mittal, J.P., Hayon, E., Photo-induced decarboxylation of aliphatic acids and esters in solution. Dependence upon state of protonation of the carboxyl group, *J. Phys. Chem.* **77**(12): 1482-7 (1973).
- 73F285 Mittal, L.J., Mittal, J.P., Hayon, E., Primary processes in the photochemistry of phenylalanine and derivatives in aqueous solution. Biphotonic photoionization and photodissociation reactions, *J. Am. Chem. Soc.* **95**(19): 6203-10 (1973).
- 73G004 Bansal, K.M., Graetzel, M., Henglein, A., Janata, E., Polarographic and optical absorption studies of radicals produced in the pulse radiolysis of aqueous solutions of ethylene glycol, *J. Phys. Chem.* **77**(1): 16-9 (1973).
- 73G026 Hayon, E., Simic, M., Addition of hydroxyl radicals to pyrimidine bases and electron transfer reactions of intermediates to quinones, *J. Am. Chem. Soc.* **95**(4): 1029-35 (1973).
- 73G030 Anbar, M., Bambenek, M., Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. I. Hydrated electron, *NSRDS-NBS* **43**, 1973, 67 p.
- 73G039 Buxton, G.V., Dainton, F., McCracken, D.R., Radiation chemical study of the reaction of  $Ni^+$ ,  $Co^+$  and  $Cd^+$  with  $N_2O$ . Evidence for the formation of a hyperoxidised state by oxygen atom transfer, *J. Chem. Soc., Faraday Trans. 1* **69**(1): 243-54 (1973).
- 73G043 Nazhat, N.B., Asmus, K.-D., Reduction of mercuric chloride by hydrated electrons and reducing radicals in aqueous solutions. Formation and reactions of  $HgCl$ , *J. Phys. Chem.* **77**(5): 614-20 (1973).
- 73G049 Simic, M., Hayon, E., Comparison between the electron transfer reactions from free radicals and their corresponding peroxy radicals to quinones, *Biochem. Biophys. Res. Commun.* **50**(2): 364-9 (1973).
- 73G062 Eriksen, T., Henglein, A., Stockhausen, K., Pulse radiolytic oxidation of chloral hydrate in oxygenated and deoxygenated aqueous solutions, *J. Chem. Soc., Faraday Trans. 1* **69**(2): 337-45 (1973).
- 73G068 Rao, P.S., Hayon, E., Experimental determination of the redox potential of the superoxide radical  $\cdot O_2^-$ , *Biochem. Biophys. Res. Commun.* **51**(2): 468-73 (1973).
- 73G085 Buxton, G.V., Sellers, R.M., Acid dissociation constant of the carboxyl radical. Pulse radiolysis studies of aqueous solutions of formic acid and sodium formate, *J. Chem. Soc., Faraday Trans. 1* **69**(3): 555-9 (1973).
- 73G097 Hayon, E., Simic, M., Acid-base properties of radical anions of cis and trans isomers. I. Fumarates and maleates, *J. Am. Chem. Soc.* **95**(8): 2433-9 (1973).
- 73G112 Rabani, J., Klug-Roth, D., Lilie, J., Pulse radiolytic investigations of the catalyzed disproportionation of peroxy radicals. Aqueous cupric ions, *J. Phys. Chem.* **77**(9): 1169-75 (1973).
- 73G114 Greenstock, C.L., Dunlop, I., Neta, P., Radiation chemical studies of the oxidation and reduction of nitrofurans. Oxidative denitration by OH radicals, *J. Phys. Chem.* **77**(9): 1187-90 (1973).
- 73G116 Faraggi, M., Leopold, J.G., The reduction of cobalamin. A pulse radiolysis study, *Biochem. Biophys. Res. Commun.* **50**(2): 413-20 (1973).
- 73G122 Adams, G.E., Willson, R.L., Ketyl radicals in aqueous solution. Pulse radiolysis study, *J. Chem. Soc., Faraday Trans. 1* **69**(4): 719-29 (1973).
- 73G125 Patel, K.B., Willson, R.L., Semiquinone free radicals and oxygen. Pulse radiolysis study of one electron transfer equilibria, *J. Chem. Soc., Faraday Trans. 1* **69**(4): 814-25 (1973).
- 73G126 Asmus, K.-D., Moekkel, H., Henglein, A., Pulse radiolytic study of the site of OH $\cdot$  radical attack on aliphatic alcohols in aqueous solution, *J. Phys. Chem.* **77**(10): 1218-21 (1973).
- 73G249 Behar, D., Samuni, A., Fessenden, R.W., Reactions of iron(II) and titanium(III) with organic radicals, *J. Phys. Chem.* **77**(17): 2055-9 (1973).
- 73R150 Bensasson, R., Land, E.J., Optical and kinetic properties of semireduced plastoquinone and ubiquinone: Electron acceptors in photosynthesis, *Biochim. Biophys. Acta* **325**: 175-81 (1973).



- 74A002 Simic, M., Lilie, J., Kinetics of ammonia detachment from reduced cobalt(III) complexes based on conductometric pulse radiolysis, *J. Am. Chem. Soc.* **96**(1): 291-2 (1974).
- 74A007 Shafferman, A., Stein, G., Reduction of ferricytochrome c by some free radical agents, *Science* **183**: 428-30 (1974).
- 74A010 Micic, O.I., Cercek, B., Solvent participation in electron transfer reactions, *J. Phys. Chem.* **78**(3): 285-9 (1974).
- 74A037 Kelm, M., Lilie, J., Henglein, A., Janata, E., Pulse radiolytic study of  $\text{Ni}^+$ . Nickel-carbon bond formation, *J. Phys. Chem.* **78**(9): 882-7 (1974).
- 74A040 Harel, Y., Meyerstein, D., On the mechanism of reduction of porphyrins. A pulse radiolytic study, *J. Am. Chem. Soc.* **96**(9): 2720-7 (1974).
- 74A043 Lesigne, B., Gilles, L., Woods, R.J., Spectra and decay of trichloromethyl radicals in aqueous solution, *Can. J. Chem.* **52**(7): 1135-9 (1974).
- 74A051 Rabani, J., Pick, M., Simic, M., Pulse radiolysis of cyclopentane in aqueous solutions, *J. Phys. Chem.* **78**(11): 1049-51 (1974).
- 74A072 Mulazzani, G.G., Ward, M.D., Semerano, G., Emmi, S.S., Giordani, P., Gamma and pulse radiolysis of tetracyanonickelate(II) anion in aqueous solution, *Int. J. Radiat. Phys. Chem.* **6**(3): 187-201 (1974).
- 74A074 Butler, J., Jayson, G.G., Swallow, A.J., Oxidation of ferrous ions by an aliphatic dioxy radical, *J. Chem. Soc., Faraday Trans. 1* **70**(7): 1394-401 (1974).
- 74A078 Lilie, J., Heckel, E., Lamb, R.C., The pulse radiolytic decomposition of p-nitroperoxybenzoic acid, *J. Am. Chem. Soc.* **96**(17): 5543-7 (1974).
- 74A089 Bruehlmann, U., Hayon, E., One-electron redox reactions of water-soluble vitamins. I. Nicotinamide (vitamin  $\text{B}_3$ ) and related compounds, *J. Am. Chem. Soc.* **96**(19): 6169-74 (1974).
- 74A099 Rabani, J., Klug-Roth, D., Henglein, A., Pulse radiolytic investigations of  $\text{OHCH}_2\text{O}_2$  radicals, *J. Phys. Chem.* **78**(21): 2089-3 (1974).
- 74A105 Blackburn, R., Erkol, A.Y., Phillips, G.O., One-electron reactions in some cobalamins, *J. Chem. Soc., Faraday Trans. 1* (9): 1693-1701 (1974).
- 74A106 Neta, P., Patterson, L.K., Substituted pyridinyl radicals in aqueous solutions. Formation, reactivity, and acid-base equilibria, *J. Phys. Chem.* **78**(22): 2211-7 (1974).
- 74A127 Moorthy, P.N., Hayon, E., Intermediates produced from the one-electron reduction of nitrogen heterocyclic compounds in solution, *J. Phys. Chem.* **78**(25): 2615-20 (1974).
- 74A135 Willson, R.L., Gilbert, B.C., Marshall, P.D.R., Norman, R.O.C., Metronidazole ('Flagyl'): A pulse radiolysis and e.s.r. study, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **26**(5): 427-34 (1974).
- 74A146 Cohen, H., Meyerstein, D., Chromium-carbon bonds in aqueous solutions. A pulse radiolytic study, *Inorg. Chem.* **13**(10): 2434-43 (1974).
- 74A167 Mulazzani, G.G., Emmi, S., Roffi, G., Hoffman, M.Z., Radiation chemically induced reduction of  $\text{Rh}(\text{dipyr})_3^{3+}$  in aqueous solution, *Quaderni dell'Area di Ricerca dell'Emilia-Romagna, C.N.R., Vol. 5, Lab. di Fotochimica e Radiazioni d'Alta Energia, Rapporto annuale 1973, Bologna, Consiglio Nazionale delle Ricerche, 1974, p.111-3.*
- 74A182 Madhavan, V., Spectra and dynamics of transients formed by the action of pulse radiolytically generated hydrated electrons and hydroxyl radicals on acrylic acid derivatives, *Ph.D. Thesis, Boston University, 1974, p. 156.*
- 74D144 Gilbert, B.C., Norman, R.O.C., Sealy, R.C., Electron spin resonance studies. Part XL. A kinetic investigation of the oxidation of oxygen-substituted carbon radicals by hydrogen peroxide, *J. Chem. Soc., Perkin Trans. 2* (7): 824-9 (1974).
- 74D286 Gilbert, B.C., Norman, R.O.C., Sealy, R.C., Electron spin resonance studies. Part XLII. The reactions of iodine-containing compounds with hydroxyl and carbon radicals in aqueous solution, *J. Chem. Soc., Perkin Trans. 2* (12): 1435-41 (1974).
- 74F246 Rowan, N.S., Hoffman, M.Z., Milburn, R.M., Intermediates in the photochemistry of tris(oxalato)cobaltate(III) ion in aqueous solution. Free and coordinated radicals, *J. Am. Chem. Soc.* **96**(19): 6060-7 (1974).
- 74F528 Moger, G., Mink, G., Nemes, I., Cserhegyi, A., Moger, D., Kinetic isotope study of the reaction of methyl radicals with amino acids and aliphatic acids in aqueous solutions. I. Kinetic isotope effect of deuterium in the abstraction of hydrogen from an  $\alpha\text{-C-H}$  bond, *Kinetics Catalysis* **15**(3): 512-7 (1974), *Transl. from Kinet. Katal.* **15**(3): 573-80 (1974).
- 74F529 Moger, G., Nemes, I., Cherkhedii, A., Kinetic-isotope study of the reactions of amino acids and aliphatic acids with methyl radicals in aqueous solutions. II. Relative rate constants for hydrogen abstraction from methine and methylene groups, *Kinetics Catalysis* **15**(4): 762-6 (1974), *Transl. from Kinet. Katal.* **15**(4): 851-6 (1974).
- 74F644 Roche, T.S., Endicott, J.F., Saturated macrocyclic ( $\text{N}_4$ ) complexes of cobalt(III) containing cobalt-alkyl bonds. Preparation and properties, *Inorg. Chem.* **13**(7): 1575-80 (1974).
- 74G033 Fessenden, R.W., Chawla, O.P., A proposed mechanism for the base catalyzed protonation of the  $\text{e}_{\text{aq}}^-$  adduct to acrylate, *J. Am. Chem. Soc.* **96**: 2262-3 (1974).
- 74G207 Pribush, A.G., Brusentseva, S.A., Shubin, V.N., Dolin, P.I., Gamma-radiolysis of air-saturated aqueous solutions of propyl alcohols, *High Energy Chem.* **8**(2): 137-9 (1974), *Transl. from Khim. Vys. Energ.* **8**(2): 167-70 (1974).
- 74G275 Olson, K.R., Hoffman, M.Z., One-electron reduction of cobalt(III) hexa-ammine ion by hydroxymethyl radicals in aqueous solution, *J. Chem. Soc., Chem. Commun.* (22): 938-9 (1974).
- 74M382 Kimura, M., Kinetics and mechanism of the reaction between formate and peroxodisulfate in aqueous solution, *Inorg. Chem.* **13**(4): 841-6 (1974).
- 75A012 Simic, M.G., Taub, I.A., Tocci, J., Hurwitz, P.A., Free radical reduction of ferricytochrome-C, *Biochem. Biophys. Res. Commun.* **62**(2): 161-7 (1975).
- 75A017 Graetzel, M., Kozak, J.J., Thomas, J.K., Electron reactions and electron transfer reactions catalyzed by micellar systems, *J. Chem. Phys.* **62**(5): 1632-40 (1975).
- 75A024 Moorthy, P.N., Hayon, E., One-electron redox reactions of water-soluble vitamins. III. Pyridoxine and pyridoxal phosphate (vitamin  $\text{B}_6$ ), *J. Am. Chem. Soc.* **97**(8): 2048-52 (1975).
- 75A027 Buxton, G.V., Sellers, R.M., Pulse radiolysis study of monovalent cadmium, cobalt, nickel and zinc in aqueous solution. Pt. 1. Formation and decay of the monovalent ions, *J. Chem. Soc., Faraday Trans. 1* **71**(3): 558-68 (1975).
- 75A049 Armor, J.N., Hoffman, M.Z., Reactivity of coordinated nitrosyls. IV. One-electron reduction of ruthenium nitrosylpentaammine(3+) ion in aqueous solution, *Inorg. Chem.* **14**(2): 444-6 (1975).
- 75A051 Rao, P.S., Hayon, E., Redox potentials of free radicals. III. Reevaluation of the method, *J. Am. Chem. Soc.* **97**(11): 2986-9 (1975).
- 75A052 Madhavan, V., Lichtin, N.N., Hayon, E., Protonation reactions of electron adducts of acrylamide derivatives. A pulse radiolytic-kinetic spectrophotometric study, *J. Am. Chem. Soc.* **97**(11): 2989-95 (1975).
- 75A053 Bell, J.A., Grunwald, E., Hayon, E., Kinetics of deprotonation of organic free radicals in water. Reaction of  $\text{HOCHCO}_2^-$ ,  $\text{HOCHCONH}_2$ , and  $\text{HOCH}_2\text{CONH}_2$  with various bases, *J. Am. Chem. Soc.* **97**(11): 2995-3000 (1975).
- 75A055 Hickel, B., Absorption spectra and kinetics of methyl and ethyl radicals in water, *J. Phys. Chem.* **79**(11): 1054-9 (1975).
- 75A056 Moorthy, P.N., Hayon, E., Radical intermediates produced from the one-electron reduction of lumazine in water, *J. Phys. Chem.* **79**(11): 1059-62 (1975).
- 75A060 Moorthy, P.N., Hayon, E., Free radical intermediates produced from the one-electron reduction of purine, adenine, and guanine derivatives in water, *J. Am. Chem. Soc.* **97**(12): 3345-50 (1975).



- 75A064 Kelm, M., Lilie, J., Henglein, A., Pulse radiolytic investigation of the reduction of cadmium(II) ions, *J. Chem. Soc., Faraday Trans. 1* **5**(9): 1132-42 (1975).
- 75A079 Barkatt, A., Rabani, J., Metal precipitation from pulse irradiated solutions of cadmium(II) and similar cations, *J. Phys. Chem.* **79**(14): 1359-62 (1975).
- 75A080 Hoffman, M.Z., Hayon, E., Solute environmental effects in the one-electron reduction of lysozyme in aqueous solution, *J. Phys. Chem.* **79**(14): 1362-9 (1975).
- 75A099 Semerano, G., Flash photolysis and pulse radiolysis in coordination chemistry, *Coord. Chem. Rev.* **16**(1/2): 185-94 (1975).
- 75A113 Wiltling, J., Van Buuren, K.J.H., Braams, R., Van Gelder, B.F., The mechanism of reduction of cytochrome c as studied by pulse radiolysis, *Biochim. Biophys. Acta* **376**(2): 285-97 (1975).
- 75A118 Eriksen, T.E., Pulse radiolytic investigation of the  $\text{SO}_2^-$  radical ion, *Radiochem. Radioanal. Lett.* **22**(1): 33-40 (1975).
- 75A125 Brede, O., Helmstreit, W., Mehnert, R., Pulradiolyse von Benzophenon in Waessriger Loesung, *Z. Phys. Chem. (Leipzig)* **256**(3): 513-21 (1975).
- 75A153 Barkatt, A., Rabani, J., Kinetics of spur reactions of electrons in ethylene glycol-water glassy ice. A pulse radiolytic study, *J. Phys. Chem.* **79**(24): 2592-7 (1975).
- 75A166 Pribush, A.G., Brusentseva, S.A., Shubin, V.N., Dolin, P.I., Mechanism of the reaction of  $\alpha$ -hydroxyisopropyl radicals with ferrous ions, *High Energy Chem.* **9**(3): 206-8 (1975), *Trans. from Khim. Vys. Energ.* **9**(3): 235-7 (1975).
- 75A169 Blackburn, R., Kyaw, M., Phillips, G.O., Swallow, A.J., Free radical reactions in the coenzyme B12 system, *J. Chem. Soc., Faraday Trans. 1* **71**(11): 2277-87 (1975).
- 75A188 Storer, D.K., Waltz, W.L., Brodovitch, J.C., Eager, R.L., A pulse radiolysis study of some platinum(II) and platinum(IV) complex ions in aqueous solutions. The formation and characterization of platinum(I) and platinum(III) transients, *Int. J. Radiat. Phys. Chem.* **7**(6): 693-704 (1975).
- 75A194 Whillans, D.W., Neta, P., Radiation chemical studies of the sensitizer Diamide, *Radiat. Res.* **64**(3): 416-30 (1975).
- 75A195 Faraggi, M., Redpath, J.L., Tal, Y., Pulse radiolysis studies of electron transfer reaction in molecules of biological interest. I. The reduction of a disulfide bridge by peptide radicals, *Radiat. Res.* **64**(3): 452-66 (1975).
- 75A203 Fujita, S., Horii, H., Mori, T., Taniguchi, S., Pulse radiolysis of  $\text{Hg}(\text{CN})_2$  in aqueous solutions, *Bull. Chem. Soc. Jpn.* **48**(11): 3067-72 (1975).
- 75A207 Whillans, D.W., Adams, G.E., Electron transfer oxidation of DNA radicals by paranitroacetophenone, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **28**(6): 501-10 (1975).
- 75A240 Bielski, B.H.J., Richter, H.W., Chan, P.C., Some properties of the ascorbate free radical, *Ann. N.Y. Acad. Sci.* **258**: 231-7 (1975).
- 75A241 Goff, H., Simic, M.G., Free radical reduction of hemin c, *Biochim. Biophys. Acta* **392**: 201-6 (1975).
- 75D188 Gilbert, B.C., Norman, R.O.C., Placucci, G., Sealy, R.C., Electron spin resonance studies. Part XLV. Reactions of the methyl radical with some aliphatic compounds in aqueous solution, *J. Chem. Soc., Perkin Trans. 2* (9): 885-91 (1975).
- 75F361 Walrant, P., Santus, R., Grossweiner, L.I., Photosensitizing properties of N-formylkynurenine, *Photochem. Photobiol.* **22**(1/2): 63-5 (1975).
- 75G001 Anbar, M., Farhataziz, Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. II. Hydrogen atom, *NSRDS-NBS 51*, 1975, 56 p.
- 75G002 Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. I. Hydrated electron, supplemental data, *NSRDS-NBS 43-Supp*, 1975, 43 p.
- 75G106 Packer, J.E., Richardson, R.K., Free-radical reductions of arenediazonium ions in aqueous solution. Part III. Kinetics of reactions of toluene-p-diazonium ions with ethanol, propan-2-ol, and acetaldehyde, *J. Chem. Soc., Perkin Trans. 2* (7): 751-5 (1975).
- 75M087 Gilbert, B.C., Norman, R.O.C., Sealy, R.C., Electron spin resonance studies. Part XLIII. Reaction of dimethyl sulphoxide with the hydroxyl radical, *J. Chem. Soc., Perkin Trans. 2* (4): 303-8 (1975).
- 75M297 Brooks, H.B., Sicillo, F., Electron spin resonance study of the  $\text{TiF}_2^+-\text{H}_2\text{O}_2$  reaction system, *J. Phys. Chem.* **79**(23): 2544-7 (1975).
- 75M298 Brooks, H.B., Sicillo, F., Electron spin resonance study of effects of sulfate and chloride ions on kinetics of the titanium(III)-hydrogen peroxide reaction system, *J. Phys. Chem.* **79**(23): 2547-50 (1975).
- 76A001 Tait, A.M., Hoffman, M.Z., Hayon, E., The reactivity of cobalt(II) complexes containing unsaturated macrocyclic ligands in aqueous solution, *J. Am. Chem. Soc.* **98**(1): 86-93 (1976).
- 76A016 Faraggi, M., Leopold, J.G., Pulse radiolysis studies of electron-transfer reaction in molecules of biological interest. II. The reduction of  $\text{Cu}(\text{II})$ -peptide complexes, *Radiat. Res.* **65**(2): 238-49 (1976).
- 76A020 Neta, P., Meisel, D., Substituent effects on nitroaromatic radical anions in aqueous solution, *J. Phys. Chem.* **80**(5): 519-24 (1976).
- 76A039 Tait, A.M., Hoffman, M.Z., Hayon, E., Reactivity of nickel(II) and copper(II) complexes containing 14-membered macrocyclic ligands in aqueous solution, *Inorg. Chem.* **15**(4): 934-9 (1976).
- 76A042 Jungbluth, H., Beyrich, J., Asmus, K.-D., Reduction of mercuric halides and pseudohalides in aqueous solution. Formation and some physicochemical properties of  $\text{HgCl}$ ,  $\text{HgBr}$ ,  $\text{HgI}$ ,  $\text{HgSCN}$ , and  $\text{HgCN}$  radical molecules, *J. Phys. Chem.* **80**(10): 1049-53 (1976).
- 76A060 Moorthy, P.N., Hayon, E., One-electron redox reactions of water-soluble vitamins. II. Pterin and folic acid, *J. Org. Chem.* **41**(9): 1607-13 (1976).
- 76A062 Frank, A.J., Graetzel, M., Henglein, A., Janata, E., Electron transfer reactions of singlet and triplet pyrene in micelles with various radical anions in aqueous solution, *Ber. Bunsenges. Phys. Chem.* **80**(4): 294-300 (1976).
- 76A067 Asmus, K.-D., Nigam, S., Willson, R.L., Kinetics of nitroxyl radical reactions. A pulse-radiolysis conductivity study, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **29**(3): 211-9 (1976).
- 76A068 Schuler, R.H., Neta, P., Zemel, H., Fessenden, R.W., Conversion of hydroxyphenyl to phenoxyl radicals. A radiolytic study of the reduction of bromophenols in aqueous solution, *J. Am. Chem. Soc.* **98**(13): 3825-31 (1976).
- 76A070 Wardman, P., Clarke, E.D., One-electron reduction potentials of substituted nitroimidazoles measured by pulse radiolysis, *J. Chem. Soc., Faraday Trans. 1* **72**(6): 1377-90 (1976).
- 76A072 Buxton, G.V., Sellers, R.M., McCracken, D.R., Pulse radiolysis study of monovalent cadmium, cobalt, nickel and zinc in aqueous solution. Part 2. Reactions of the monovalent ions, *J. Chem. Soc. Faraday Trans 1* **72**(6): 1464-76 (1976).
- 76A075 Greenstock, C.L., Ruddock, G.W., Neta, P., Pulse radiolysis and ESR studies of the electron-affinic properties of nitroheterocyclic radiosensitizers, *Radiat. Res.* **66**(3): 472-84 (1976).
- 76A082 Abramovitch, S., Rabani, J., Pulse radiolytic investigations of peroxy radicals in aqueous solutions of acetate and glycine, *J. Phys. Chem.* **8**(5): 609-11 (1976).
- 76A087 Fujita, S., Horii, H., Mori, T., Taniguchi, S., Pulse radiolysis of  $\text{HgBr}_2$  in aqueous solutions, *Bull. Chem. Soc. Jpn.* **49**(5): 1250-3 (1976).
- 76A103 Toffel, P., Henglein, A., The temperature dependence of the electrolytic dissociation of some 1-hydroxy-alkyl radicals in aqueous solution (a pulse radiolysis study), *Ber. Bunsenges. Phys. Chem.* **80**(6): 525-8 (1976).
- 76A104 Frank, A.J., Graetzel, M., Henglein, A., Janata, E., The influence of the interface potential on the reactions of hydrated electrons and neutral radicals with acceptors in micelles, *Ber. Bunsenges. Phys. Chem.* **80**(6): 547-51 (1976).
- 76A109 Pick-Kaplan, M., Rabani, J., Pulse radiolytic studies of aqueous  $\text{Mn}(\text{ClO}_4)_2$  solutions, *J. Phys. Chem.* **80**(17): 1840-3 (1976).



- 76A111 Neta, P., Simic, M.G., Hoffman, M.Z., Pulse radiolysis and electron spin resonance studies of nitroaromatic radical anions. Optical absorption spectra, kinetics, and one-electron redox potentials, *J. Phys. Chem.* **80**(18): 2018-23 (1976).
- 76A113 Madhavan, V., Lichtin, N.N., Hayon, E., Electron adducts of acrylic acid and homologues. Spectra, kinetics, and protonation reactions. A pulse radiolysis study, *J. Org. Chem.* **41**(13): 2320-6 (1976).
- 76A127 Seki, H., Ilan, Y.A., Ilan, Y., Stein, G., Reactions of the ferri-ferrocyclochrome-c system with superoxide/oxygen and  $\text{CO}_2^-/\text{CO}_2$  studied by fast pulse radiolysis, *Biochim. Biophys. Acta* **440**(3): 573-86 (1976).
- 76A132 Ilan, Y., Rabani, J., On some fundamental reactions in radiation chemistry: Nanosecond pulse radiolysis, *Int. J. Radiat. Phys. Chem.* **8**(5): 609-11 (1976).
- 76A137 Ilan, Y.A., Rabani, J., Czapski, G., One electron reduction of metmyoglobin and methemoglobin and the reaction of the reduced molecule with oxygen, *Biochim. Biophys. Acta* **446**(1): 277-86 (1976).
- 76A152 Nigam, S., Asmus, K.-D., Willson, R.L., Electron transfer and addition reactions of free nitroxyl radicals with radiation induced radicals, *J. Chem. Soc., Faraday Trans. 1* **72**(10): 2324-40 (1976).
- 76A169 Anderson, R.F., Electron transfer and equilibria between pyridinyl radicals and FAD, *Ber. Bunsenges. Phys. Chem.* **80**(10): 969-72 (1976).
- 76A170 Breitenkamp, M., Henglein, A., Lilie, J., Mechanism of the reduction of lead ions in aqueous solution (a pulse radiolysis study), *Ber. Bunsenges. Phys. Chem.* **80**(10): 973-9 (1976).
- 76A182 Neta, P., Intermediates in the reduction of oximes in aqueous solutions, *Radiat. Res.* **68**(3): 422-32 (1976).
- 76A201 Venturi, M., Breccia, A., Busi, F., Mulazzani, Q.G., Gamma and pulse radiolysis of methanolic solutions of triphenylphosphine, *Int. J. Radiat. Phys. Chem.* **8**(6): 673-7 (1976).
- 76A203 Tait, A.M., Hoffman, M.Z., Hayon, E., The reaction of radiation-generated free radicals with cobalt(II) and cobalt(III) complexes containing macrocyclic ligands, *Int. J. Radiat. Phys. Chem.* **8**(6): 691-6 (1976).
- 76A241 Frank, A.J., Radiation induced redox reactions in micellar solutions, Micellization, Solubilization, Microemulsions, [Proc. Int. Symp.], 1976, K.L. Mittal (ed.), Plenum, New York, N.Y., Pub. 1977, Vol. 2, p.549-68.
- 76A254 Soylemez, T., Balkas, T.I., Studies on the reactions between aliphatic hydrocarbon radicals and radiolytically formed hydrogen peroxide, Proc. Fourth Tihany Symposium on Radiation Chemistry (Keszthely, Hungary, 1-6 June 1976), P. Hedvig and R. Schiller (eds.), Akad. Kiado, Budapest, 1977, p.853-9.
- 76D198 Kirino, Y., Taniguchi, H., An esr study of the acid dissociation of NH protons. 1. Linear peptide radicals and related radicals, *J. Am. Chem. Soc.* **98**(17): 5089-96 (1976).
- 76D199 Trifunac, A.D., Chemically induced dynamic electron polarization. Ionic strength and radical concentration dependence, *J. Am. Chem. Soc.* **98**(17): 5202-6 (1976).
- 76D419 Shuvalov, V.F., Moravskii, A.P., Radical generation kinetics in the system  $\text{NH}_2\text{OH} + \text{Ti(III)} + \text{alcohols}$ , *Kinet. Catal.* **17**(4): 760-6 (1976), Transl. from *Kinet. Katal.* **17**(4): 870-6 (1976).
- 76G021 Buxton, G.V., Green, J.C., Higgins, R., Kanji, S., Formation of epoxides in the oxidation of  $\beta$ -hydroxyalkyl radicals by copper(II) in aqueous solution, *J. Chem. Soc., Chem. Commun.* (5): 158-9 (1976).
- 77A002 Buechler, Hch., Bruehlmann, U., Buehler, R.E., A transient equilibrium of outer-inner charge transfer complexes from OH-radical reactions with aqueous alkyl iodide, *Chem. Phys.* **19**(1): 43-50 (1977).
- 77A011 Rabani, J., Mulac, W.A., Matheson, M.S., Pulse radiolysis studies of  $\text{Zn}^+$  reactions, *J. Phys. Chem.* **81**(2): 99-104 (1977).
- 77A025 Freiberg, M., Meyerstein, D., An intermediate with a copper-carbon bond formed by the reaction of copper ions with  $\cdot\text{CH}_2\text{CO}_2\text{H}$  radicals in aqueous solutions, *J. Chem. Soc., Chem. Commun.* (4): 127-8 (1977).
- 77A027 Simic, M.G., Hoffman, M.Z., Brezniak, N.V., Kinetics of ligand-to-metal intramolecular electron transfer in cobalt(III)-amine complexes containing a coordinated radical, *J. Am. Chem. Soc.* **99**(7): 2166-72 (1977).
- 77A034 Moorthy, P.N., Hayon, E., One-electron redox reactions of water-soluble vitamins. 4. Thiamin (vitamin  $\text{B}_1$ ), biotin, and pantothenic acid, *J. Org. Chem.* **42**(5): 879-85 (1977).
- 77A036 Schuler, R.H., Oxidation of ascorbate anion by electron transfer to phenoxyl radicals, *Radiat. Res.* **69**(3): 417-33 (1977).
- 77A100 Cohen, H., Meyerstein, D., Aliphatic radicals as reducing agents of cobalt(III) and ruthenium(III) complexes: A pulse radiolytic study, *J. Chem. Soc., Dalton Trans.* (11): 1056-62 (1977).
- 77A118 van der Linde, H.J., A pulse radiolytic study of aqueous nitroaniline solutions, *Radiat. Phys. Chem.* **10**(3): 199-206 (1977).
- 77A120 Cheney, R.P., Simic, M.G., Hoffman, M.Z., Taub, I.A., Asmus, K.-D., One-electron reduction of pentacyanonitrosyl ferrate(II) ion in aqueous solution, *Inorg. Chem.* **16**(9): 2187-92 (1977).
- 77A128 Shafferman, A., Stein, G., Conformation dependent participation of the protein in electron equivalent transfer to cytochrome c, *Biochim. Biophys. Acta* **462**(1): 161-70 (1977).
- 77A138 Bettelheim, A., Faraggi, M., The reduction of Cu(II) complexes of histidine and histidyl peptides. A pulse radiolysis study, *Radiat. Res.* **72**(1): 71-80 (1977).
- 77A169 Neta, P., Levanon, H., Spectrophotometric study of the radicals produced by the reduction of syn- and anti-azobenzene, *J. Phys. Chem.* **81**(24): 2288-92 (1977).
- 77A177 Patterson, L.K., Small, R.D., Jr., Scaiano, J.C., Reaction of paraquat radical cations with oxygen. A pulse radiolysis and laser photolysis study, *Radiat. Res.* **72**(2): 218-25 (1977).
- 77A191 Ashworth, B., Gilbert, B.C., Norman, R.O.C., Electron spin resonance studies. Part 53. An investigation of the generation of aryl radicals and of their abstraction reactions in aqueous solution, *J. Chem. Res. (S)* (4): 94-5 (1977).
- 77D087 Taniguchi, H., Kirino, Y., An esr study of the acid dissociation of NH protons. 2. Cyclic peptide radicals and related radicals, *J. Am. Chem. Soc.* **99**(11): 3625-31 (1977).
- 77F005 Endicott, J.F., Ferraudi, G.J., A flash photolytic investigation of low energy homolytic processes in methylcobalamin, *J. Am. Chem. Soc.* **99**(1): 243-5 (1977).
- 77F639 Ferraudi, G.J., Endicott, J.F., Photostimulated solvolytic reactions of a macrocyclic copper complex, *Inorg. Chem.* **16**(11): 2762-6 (1977).
- 77G011 Farhataziz, Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. III. Hydroxyl radical and perhydroxyl radical and their radical ions, *NSRDS-NBS* 59, 1977, 122p.
- 77G411 Berdnikov, V.M., Zhuravleva, O.S., Terent'eva, L.A., Kinetics and mechanism of elementary electron transfer in the oxidation of alcohol radicals by hydrated Fe(III) ions, *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **26**(10): 2050-6 (1977), Transl. from *Izv. Akad. Nauk SSSR, Ser. Khim.* (10): 2214-20 (1977).
- 77R091 Seki, H., Imamura, M., Illian, X.A., Ilan, Y., Stein, G., Oxidation and reduction of cytochrome C with  $\text{Br}_2^-$ ,  $\text{O}_2^-$  and  $\text{CO}_2^-$  studied by pulse radiolysis, *J. Radiat. Res.* **18**(1): 6 (1977).
- 77R167 Biaglow, J.E., Jacobson, B., Greenstock, C.L., Raleigh, J., Effect of nitrobenzene derivatives on electron transfer in cellular and chemical models, *Mol. Pharmacol.* **13**: 269-82 (1977).
- 78A007 Anderson, R.F., Schulte-Frohlinde, D., Reactions induced by hydroxyl radical attack on acetylene in aqueous solution. A pulse radiolysis study, *J. Phys. Chem.* **82**(1): 22-6 (1978).
- 78A033 Evers, E.L., Jayson, G.G., Swallow, A.J., Pulse radiolysis study of protoferrin IX intercalated in sodium dodecyl sulphate micelles, *J. Chem. Soc., Faraday Trans. 1* **74**(2): 418-26 (1978).
- 78A068 Mulazzani, Q.G., Emmi, S., Fucchi, P.G., Hoffman, M.Z., Venturi, M., On the nature of  $\text{Ru}(\text{bpy})_3^+$  in aqueous solution, *J. Am. Chem. Soc.* **100**(3): 981-3 (1978).
- 78A070 Jonah, C.D., Matheson, M.S., Meisel, D., Reaction of  $\text{e}_{\text{aq}}^-$  into excited states of  $\text{Ru}(\text{bpy})_3^{2+}$ , *J. Am. Chem. Soc.* **100**(5): 1449-56 (1978).



- 78A095 Bobrowski, K., Suwalski, J.P., Zagorski Z.P., Pulse radiolysis of tetraalkylammonium hydroxides in alkaline solution containing oxygen, *Radiat. Phys. Chem.* **11**(2): 71-6 (1978).
- 78A108 Wieghardt, K., Cohen, H., Meyerstein, D., Intramolecular electron transfer through a bridging carboxylate group coordinated to two cobalt(III)-ions: A pulse radiolytic study, *Ber. Bunsenges. Phys. Chem.* **82**(4): 388-92 (1978).
- 78A110 Cheney, R.P., Hoffman, M.Z., Lust, J.A., Redox-catalyzed aquation of nitrosylpentaammine ruthenium(3+) ion, *Inorg. Chem.* **17**(5): 1177-80 (1978).
- 78A165 Fujita, S., Horii, H., Mori, T., Taniguchi, S., Studies of emission species observed on pulse radiolysis of mercuric halides in aqueous systems, *J. Phys. Chem.* **82**(15): 1693-6 (1978).
- 78A175 Parsons, B.J., Schulte-Frohlinde, D., von Sonntag, C., Radiation chemistry of DNA model compounds. XIII. Reaction of  $\text{Br}_3^{2-}$  with 2-deoxy-D-ribose. A preferred attack at C-1, *Z. Naturforsch., Teil B* **33B**(6): 666-8 (1978).
- 78A176 Ilan, Y., Ilan, Y.A., Czapski, G., Do copper ions influence the reduction of ferricytochrome C by  $\text{O}_2^-$ , *Biochim. Biophys. Acta* **503**(2): 399-401 (1978).
- 78A177 Sutton, H.C., Seddon, W.A., Sopchysyn, F.C., Pulse radiolysis: evidence for the reaction of  $\text{HO}_2$  with  $\text{NO}_2$  in aqueous solution, *Can. J. Chem.* **56**(14): 1961-4 (1978).
- 78A200 Elroi, H., Meyerstein, D., Kinetics of the formation and decomposition of carbon-cobalt(III) bonds in aqueous solutions by the reaction of aliphatic free radicals with a coenzyme B-12r model cobalt(II) complex, *J. Am. Chem. Soc.* **100**(17): 5540-8 (1978).
- 78A222 Wieghardt, K., Cohen, H., Meyerstein, D., Intramolecular electron transfer of coordinated pyrazinecarboxylate radicals to cobalt(III): A pulse radiolytic study, *Angew. Chem., Int. Ed. Engl.* **17**(8): 608-9 (1978).
- 78A232 Favaudon, V., Ferradini, C., Pucheault, J., Gilles, L., Le Gall, J., Kinetics of reduction of the cytochrome  $\text{C}_3$  from *Desulfovibrio vulgaris*, *Biochem. Biophys. Res. Commun.* **84**(2): 435-40 (1978).
- 78A256 Lilie, J., Hanrahan, R.J., Henglein, A.,  $\text{O}^-$  transfer reactions of the carbonate radical anion, *Radiat. Phys. Chem.* **11**(5): 225-7 (1978).
- 78A261 Chen, T.S.-S., Radiation induced reactions in micellar and polymer systems, Ph.D., Thesis, Univ. of Notre Dame, Notre Dame, IN, 1978, 182p.
- 78A288 Simic, M.G., Taub, I.A., Fast electron transfer processes in cytochrome C and related metalloproteins, *Biophys. J.* **24**(1): 285-94 (1978).
- 78A321 Farrington, J.A., Ebert, M., Land, E.J., Bipyridylum quaternary salts and related compounds. Part 6. Pulse radiolysis studies of the reaction of paraquat radical analogues with oxygen, *J. Chem. Soc., Faraday Trans. 1* **74**(3): 665-75 (1978).
- 78A322 Buxton, G.V., Green, J.C., Reactions of some simple  $\alpha$ - and  $\beta$ -hydroxyalkyl radicals with  $\text{Cu}^{2+}$  and  $\text{Cu}^+$  ions in aqueous solution. A radiation chemical study, *J. Chem. Soc., Faraday Trans. 1* **74**(3): 697-714 (1978).
- 78A365 Hasegawa, K., Patterson, L.K., Pulse radiolysis studies in model lipid systems: Formation and behavior of peroxy radicals in fatty acids, *Photochem. Photobiol.* **28**(4-5): 817-23 (1978).
- 78A394 Shtamm, E.V., Purmal, A.P., Skurlatov, Yu.I., The photochemical and photocatalytic oxidation of ascorbic acid in the photolysis of hydrogen peroxide, *Russ. J. Phys. Chem.* **52**(1): 61-3 (1978), *Trans. from Zh. Fiz. Khim.* **52**(1): 117-20 (1978).
- 78A402 Nenadovic, M.T., Micic, O.I., Pulse radiolysis of methyl acetate in aqueous solutions, *Radiat. Phys. Chem.* **12**(3-4): 85-9 (1978).
- 78A403 Hoffman, M.Z., Simic, M.G., Mulazzani, Q.G., Emmi, S., Fuochi, P.G., Venturi, M., One-electron reduction of 2,2'-bipyridine in aqueous solution, *Radiat. Phys. Chem.* **12**(3-4): 111-3 (1978).
- 78A410 Tausch-Treml, R., Henglein, A., Lilie, J., Reactivity of silver atoms in aqueous solution. II. A pulse radiolysis study, *Ber. Bunsenges. Phys. Chem.* **82**(12): 1335-43 (1978).
- 78A417 Fedorova, O.S., Berdnikov, V.M., Radiation-chemical oxidation of ascorbic acid in aqueous solution in the presence of oxygen, *High Energy Chem.* **12**(5): 387-9 (1978), *Trans. from Khim. Vys. Energ.* **12**(5): 463-5 (1978).
- 78B128 Packer, J.E., Slater, T.F., Willson, R.L., Reactions of the carbon tetrachloride-related peroxy free radical ( $\text{CCl}_3\text{O}_2^-$ ) with amino acids: Pulse radiolysis evidence, *Life Sci.* **23**(26): 2617-20 (1978).
- 78F301 Ferraudi, G., Photochemical generation of metastable methylcopper complexes. Oxidation-reduction of methyl radicals by copper complexes, *Inorg. Chem.* **17**(9): 2506-8 (1978).
- 79A034 Simic, M.G., Hoffman, M.Z., Cheney, R.P., Mulazzani, Q.G., One-electron reduction of tris(2,2'-bipyridine) and tris(1,10-phenanthroline) complexes of cobalt(III) in aqueous solution, *J. Phys. Chem.* **83**(4): 439-43 (1979).
- 79A036 Debey, P., Land, E.J., Santus, R., Swallow, A.J., Electron transfer from pyridinyl radicals, hydrated electrons,  $\text{CO}_2^-$  and  $\text{O}_2^-$  to bacterial cytochrome P450, *Biochem. Biophys. Res. Commun.* **86**(4): 953-60 (1979).
- 79A051 Steenzen, S., Oxidation of phenolates and phenylenediamines by 2-alkanonyl radicals produced from 1,2-dihydroxy and 1-hydroxy-2-alkoxyalkyl radicals, *J. Phys. Chem.* **83**(5): 595-9 (1979).
- 79A134 Cheney, R.P., Pell, S.D., Hoffman, M.Z., Radiation synthesis of iron(II) and ruthenium(II) alkyl nitroso complexes, *J. Inorg. Nucl. Chem.* **41**(4): 489-93 (1979).
- 79A143 Neta, P., Scherz, A., Levanon, H., Electron transfer reactions involving porphyrins and chlorophyll a, *J. Am. Chem. Soc.*, **101**:3624-9 (1979).
- 79A148 Mulazzani, Q.G., Emmi, S., Fuochi, P.G., Venturi, M., Hoffman, M.Z., Simic, M.G., One-electron reduction of aromatic nitrogen heterocycles in aqueous solution. 2,2'-Bipyridine and 1,10-phenanthroline, *J. Phys. Chem.* **83**(12): 1582-90 (1979).
- 79A153 van Leeuwen, J.W., Tromp, J., Nauta, H., Reduction of ferricytochrome c, methemoglobin and metmyoglobin by hydroxyl and alcohol radicals, *Biochim. Biophys. Acta* **577**(2): 394-9 (1979).
- 79A158 Endicott, J.F., Netzel, T.L., Early events and transient chemistry in the photolysis of alkylcobalamins, *J. Am. Chem. Soc.* **101**(14): 4000-2 (1979).
- 79A174 Janovsky, I., On the radiolytic reduction of tris(1,10-phenanthroline)/Fe(III) complex in acid aqueous solution, *Radiochem. Radioanal. Lett.* **39**(4-5): 337-41 (1979).
- 79A176 Tamminga, J.J., van den Ende, C.A.M., Warman, J.M., Hummel, A., Absorption spectra and reaction kinetics of the radical anion, radical cation and OH adduct(s) of 3,5-dinitroanisole studied by pulse radiolysis, *Recl. Trav. Chim. Pays-Bas* **98**(5): 305-15 (1979).
- 79A190 Sukhov, N.L., Ershov, B.G., Pulse radiolysis of aqueous Ga(III) solutions, *High Energy Chem.* **13**(1): 45-9 (1979), *Trans. from Khim. Vys. Energ.* **13**(1): 55-60 (1979).
- 79A191 Land, E.J., Swallow, A.J., Some free radical reactions of camphor in relation to the action of cytochrome P450, *J. Chem. Soc., Faraday Trans. 1* **75**: 1849-56 (1979).
- 79A204 Harrington, P.C., Wilkins, R.G., The reduction of methemerythrin by  $\text{e}_{\text{aq}}^-$  and  $\text{CO}_2^-$  from pulse radiolysis studies, *J. Biol. Chem.* **254**(16): 7505-8 (1979).
- 79A213 Hoffman, M.Z., Kimmel, D.W., Simic, M.G., Intramolecular electron transfer in the reaction of hydroxyl radicals with (pyridine)pentaamminecobalt(III) ion in aqueous solution, *Inorg. Chem.* **18**(9): 2479-85 (1979).
- 79A255 Bhattacharyya, S.N., Srisankar, E.V., Radiation chemistry of cobalt(II) nitilotriacetate in aqueous solutions, *J. Chem. Soc., Faraday Trans. 1* **75**(9): 2089-99 (1979).
- 79A305 Neta, P., Electron transfer reactions involving acridine and related compounds, *J. Phys. Chem.* **83**(24): 3096-101 (1979).



- 79A312 Ilan, Y., Shafferman, A., Feinberg, B.A., Lau, Y.-K., Partitioning of electrostatic and conformational contributions in the redox reactions of modified cytochromes c, *Biochim. Biophys. Acta* **548**(3): 565-78 (1979).
- 79A327 Houee-Levin, C., Gardes-Albert, M., Ferradini, C., Pucheault, J., Pulse radiolytic investigations of the redox system alloxan-dialuric acid: Evidence for a radical intermediate, *Biochem. Biophys. Res. Commun.* **91**(3): 1196-200 (1979).
- 79A341 Micic, O.I., Nenadovic, M.T., Pulse-radiolytic investigation of the reduction of titanium(III) ions in aqueous solutions, *J. Chem. Soc., Dalton Trans.* (12): 2011-4 (1979).
- 79A371 Shieh, J.J., Sellers, R.M., Hoffman, M.Z., Taub, I.A., Radiation chemistry of myoglobin, its derivatives, and model compounds in aqueous solution, *Radiation Biology and Chemistry. Research Developments*, H.E. Edwards, S. Navaratnam, B.J. Parsons and G.O. Phillips (eds.), Elsevier, N.Y., 1979, p. 179-88.
- 79E282 Land, E.J., Truscott, T.G., Triplet excited state of coumarin and 4',5'-dihydropsoalene: Reaction with nucleic acid bases and amino acids, *Photochem. Photobiol.* **29**(5): 861-6 (1979).
- 79G191 Henglein, A., Catalysis of the reduction of  $\text{Ti}^+$  and of  $\text{CH}_2\text{Cl}_2$  by colloidal silver in aqueous solution, *J. Phys. Chem.* **83**(22): 2858-62 (1979).
- 79N061 Almgren, M., Grieser, F., Thomas, J.K., Rate of exchange of surfactant monomer radicals and long chain alcohols between micelles and aqueous solutions. A pulse radiolysis study, *J. Chem. Soc., Faraday Trans. 1* **75**: 1674-87 (1979).
- 80A011 Brault, D., Bizet, C., Morliere, P., Rougee, M., Land, E.J., Santus, R., Swallow, A.J., One-electron reduction of ferrideuteroporphyrin IX and reaction of the oxidized and reduced forms with chlorinated methyl radicals, *J. Am. Chem. Soc.* **102**(3): 1015-20 (1980).
- 80A053 Packer, J.E., Willson, R.L., Bahnmann, D., Asmus, K.-D., Electron transfer reactions of halogenated aliphatic peroxy radicals: measurement of absolute rate constants by pulse radiolysis, *J. Chem. Soc., Perkin Trans. 2* (2): 296-9 (1980).
- 80A054 Soylemez, T., von Sonntag, C., Hydroxyl radical-induced oligomerization of ethylene in deoxygenated aqueous solution, *J. Chem. Soc., Perkin Trans. 2* (2): 391-4 (1980).
- 80A066 Beitz, J.V., Miller, J.R., Cohen, H., Wiegardt, K., Meyerstein, D., Activation energies for an intramolecular electron transfer reaction, *Inorg. Chem.* **19**(4): 966-8 (1980).
- 80A071 Eibenberger, J., Schulte-Frohlinde, D., Steenken, S., One-electron oxidation of  $\alpha$ -monoalkoxyalkyl radicals by tetranitromethane via an intermediate adduct. Influence of radical structure on rate of decomposition of the adduct, *J. Phys. Chem.* **84**(7): 704-10 (1980).
- 80A085 Bielski, B.H.J., Shiue, G.G., Bajuk, S., Reduction of nitro blue tetrazolium by  $\text{CO}_2^-$  and  $\text{O}_2^-$  radicals, *J. Phys. Chem.* **84**(8): 830-33 (1980).
- 80A115 Teply, J., Janovsky, I., Mehnert, R., Brede, O., Pulse radiolytic study of unstable intermediates in the radiolysis of aqueous solutions of 1,10-phenanthroline, *Radiat. Phys. Chem.* **15**(2-3): 169-75 (1980).
- 80A123 Butler, J., Henglein, A., Elementary reactions of the reduction of  $\text{Ti}^+$  in aqueous solution, *Radiat. Phys. Chem.* **15**(5): 603-12 (1980).
- 80A146 Ferraudi, G., Patterson, L.K., Free radical reactions with dimeric cobalt(II) sulphophthalocyanine. A pulse radiolytic study, *J. Chem. Soc., Dalton Trans.* 476-80 (1980).
- 80A153 Buitenhuis, R., Bakker, C.N.N., Stock, F.R., Louwrier, P.W.F., Pulse radiolysis study of  $\text{Cu}(\text{EDTA})^{2-}$  in aqueous solution in the presence of alcohols, *Radiat. Phys. Chem.* **16**(1): 6-10 (1980).
- 80A189 Freiberg, M., Lilie, J., Meyerstein, D., Nature and mechanism of decomposition of the complex of copper(I) with 5,7,7,12,14,14-hexamethyl 1,4,8,11-tetraazacyclotetradeca-4,11-diene in aqueous solutions. A pulse radiolysis study, *Inorg. Chem.* **19**(7): 1908-12 (1980).
- 80A197 Houee-Levin, C., Gardes-Albert, M., Ferradini, C., Pucheault, J., Radiolysis study of the alloxan-dialuric acid couple, *Radiat. Res.* **83**: 270-8 (1980).
- 80A204 Elliot, A.J., Wilkinson, F., Armstrong, D.A., A pulse radiolysis study of zinc(II)-insulin, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* **38**(1): 1-10 (1980).
- 80A227 Venturi, M., Emmi, S., Fuochi, P.G., Mulazzani, Q.G., Nucleophilic addition of  $\alpha$ -hydroxyalkyl radicals to polypyridyl complexes of chromium(III), cobalt(III), rhodium(III), and ruthenium(II) in aqueous solution, *J. Phys. Chem.* **84**(17): 2160-6 (1980).
- 80A277 Freiberg, M., Meyerstein, D., Reactions of aliphatic free radicals with copper cations in aqueous solution. Part 2. Reactions with cupric ions: A pulse radiolysis study, *J. Chem. Soc. Faraday Trans. 1* **76**: 1825-37 (1980).
- 80A278 Freiberg, M., Mulac, W.A., Schmidt, K.H., Meyerstein, D., Reactions of aliphatic free radicals with copper cations in aqueous solutions. Part 3. Reactions with cuprous ions: A pulse radiolysis study, *J. Chem. Soc. Faraday Trans 1* **76**: 1838-48 (1980).
- 80A282 Bothe, E., Schulte-Frohlinde, D., Reaction of dihydroxymethyl radical with molecular oxygen in aqueous solution, *Z. Naturforsch. Teil B* **35B**: 1035-9 (1980).
- 80A323 Janata, E., Schuler, R.H., Direct observation of the protonation of acetone ketyl radical by conductometric pulse radiolysis, *J. Phys. Chem.* **84**(25): 3351-3 (1980).
- 80N062 Henglein, A., Colloidal silver catalyzed multi-electron transfer processes in aqueous solution, *Ber. Bunsenges. Phys. Chem.* **84**: 253-9 (1980).
- 81A001 Neta, P., Behar, D., Intramolecular electron transfer and dehalogenation of anion radicals. 3. Halobenzonitriles and cyano-benzyl halides, *J. Am. Chem. Soc.* **103**: 103-6 (1981).
- 81A002 Brault, D., Neta, P., Reactions of iron porphyrins with methyl radicals, *J. Am. Chem. Soc.* **103**(10): 2705-10 (1981).
- 81A003 Steenken, S., Neta, P., Oxidation of substituted alkyl radicals by  $\text{IrCl}_6^{2-}$ ,  $\text{Fe}(\text{CN})_6^{3-}$  and  $\text{MnO}_4^-$  in aqueous solution. Electron transfer versus chlorine transfer from  $\text{IrCl}_6^{2-}$ , *J. Am. Chem. Soc.* **104**(5): 1244-8 (1982).

U.S. DEPT. OF COMM. <b>BIBLIOGRAPHIC DATA SHEET</b> (See instructions)		1. PUBLICATION OR REPORT NO. NSRDS-NBS 70	2. Performing Organ. Report No.	3. Publication Date October 1982
4. TITLE AND SUBTITLE Rate Constants for Reactions of Aliphatic Carbon-Centered Radicals in Aqueous Solution				
5. AUTHOR(S) Alberta B. Ross and Pedatsur Neta				
6. PERFORMING ORGANIZATION (If joint or other than NBS, see instructions) Radiation Chemistry Data Center Radiation Laboratory University of Notre Dame Notre Dame, Indiana 46556			7. Contract/Grant No.	8. Type of Report & Period Covered N/A
9. SPONSORING ORGANIZATION NAME AND COMPLETE ADDRESS (Street, City, State, ZIP) National Bureau of Standards Department of Commerce Washington, D. C. 20234				
10. SUPPLEMENTARY NOTES Library of Congress Catalog Card Number: 82-2096  <input type="checkbox"/> Document describes a computer program; SF-185, FIPS Software Summary, is attached.				
11. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here)  Rate constants have been compiled for reactions of various transient aliphatic radicals produced mostly by radiolysis in aqueous solutions. In certain cases the radicals have been produced by photolysis or by a thermal chemical reaction. Data are included for aliphatic carbon-centered radicals only, i.e., substituted alkyl radicals in the broad sense, but not for oxyl or aminyl radicals. Reactions of the aliphatic radicals with inorganic and organic compounds are included.				
12. KEY WORDS (Six to twelve entries; alphabetical order; capitalize only proper names; and separate key words by semicolons) alkyl radicals; aminoalkyl radicals; aqueous solution; carboxyalkyl radicals; chemical kinetics; electron transfer; haloalkyl radicals; hydroxyalkyl radicals; photolysis; radical anions; radiolysis; rates.				
13. AVAILABILITY <input checked="" type="checkbox"/> Unlimited <input type="checkbox"/> For Official Distribution. Do Not Release to NTIS <input checked="" type="checkbox"/> Order From Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.  <input type="checkbox"/> Order From National Technical Information Service (NTIS), Springfield, VA. 22161			14. NO. OF PRINTED PAGES 103  15. Price \$5.50	





**Announcement of New Publications in  
National Standard Reference Data Series**

Superintendent of Documents,  
Government Printing Office,  
Washington, D.C. 20402

Dear Sir:

Please add my name to the announcement list of new publications to be issued  
in the series: National Standard Reference Data Series—National Bureau of  
Standards.

Name\_\_\_\_\_

Company\_\_\_\_\_

Address\_\_\_\_\_

City\_\_\_\_\_ State\_\_\_\_\_ Zip Code\_\_\_\_\_

(Notification Key N-519)



## NBS TECHNICAL PUBLICATIONS

### PERIODICALS

**JOURNAL OF RESEARCH**—The Journal of Research of the National Bureau of Standards reports NBS research and development in those disciplines of the physical and engineering sciences in which the Bureau is active. These include physics, chemistry, engineering, mathematics, and computer sciences. Papers cover a broad range of subjects, with major emphasis on measurement methodology and the basic technology underlying standardization. Also included from time to time are survey articles on topics closely related to the Bureau's technical and scientific programs. As a special service to subscribers each issue contains complete citations to all recent Bureau publications in both NBS and non-NBS media. Issued six times a year. Annual subscription: domestic \$18; foreign \$22.50. Single copy, \$4.25 domestic; \$5.35 foreign.

### NONPERIODICALS

**Monographs**—Major contributions to the technical literature on various subjects related to the Bureau's scientific and technical activities.

**Handbooks**—Recommended codes of engineering and industrial practice (including safety codes) developed in cooperation with interested industries, professional organizations, and regulatory bodies.

**Special Publications**—Include proceedings of conferences sponsored by NBS, NBS annual reports, and other special publications appropriate to this grouping such as wall charts, pocket cards, and bibliographies.

**Applied Mathematics Series**—Mathematical tables, manuals, and studies of special interest to physicists, engineers, chemists, biologists, mathematicians, computer programmers, and others engaged in scientific and technical work.

**National Standard Reference Data Series**—Provides quantitative data on the physical and chemical properties of materials, compiled from the world's literature and critically evaluated. Developed under a worldwide program coordinated by NBS under the authority of the National Standard Data Act (Public Law 90-396).

NOTE: The principal publication outlet for the foregoing data is the Journal of Physical and Chemical Reference Data (JPCRD) published quarterly for NBS by the American Chemical Society (ACS) and the American Institute of Physics (AIP). Subscriptions, reprints, and supplements available from ACS, 1155 Sixteenth St., NW, Washington, DC 20056.

**Building Science Series**—Disseminates technical information developed at the Bureau on building materials, components, systems, and whole structures. The series presents research results, test methods, and performance criteria related to the structural and environmental functions and the durability and safety characteristics of building elements and systems.

**Technical Notes**—Studies or reports which are complete in themselves but restrictive in their treatment of a subject. Analogous to monographs but not so comprehensive in scope or definitive in treatment of the subject area. Often serve as a vehicle for final reports of work performed at NBS under the sponsorship of other government agencies.

**Voluntary Product Standards**—Developed under procedures published by the Department of Commerce in Part 10, Title 15, of the Code of Federal Regulations. The standards establish nationally recognized requirements for products, and provide all concerned interests with a basis for common understanding of the characteristics of the products. NBS administers this program as a supplement to the activities of the private sector standardizing organizations.

**Consumer Information Series**—Practical information, based on NBS research and experience, covering areas of interest to the consumer. Easily understandable language and illustrations provide useful background knowledge for shopping in today's technological marketplace.

*Order the above NBS publications from: Superintendent of Documents, Government Printing Office, Washington, DC 20402.*

*Order the following NBS publications—FIPS and NBSIR's—from the National Technical Information Services, Springfield, VA 22161.*

**Federal Information Processing Standards Publications (FIPS PUB)**—Publications in this series collectively constitute the Federal Information Processing Standards Register. The Register serves as the official source of information in the Federal Government regarding standards issued by NBS pursuant to the Federal Property and Administrative Services Act of 1949 as amended, Public Law 89-306 (79 Stat. 1127), and as implemented by Executive Order 11717 (38 FR 12315, dated May 11, 1973) and Part 6 of Title 15 CFR (Code of Federal Regulations).

**NBS Interagency Reports (NBSIR)**—A special series of interim or final reports on work performed by NBS for outside sponsors (both government and non-government). In general, initial distribution is handled by the sponsor; public distribution is by the National Technical Information Services, Springfield, VA 22161, in paper copy or microfiche form.



U.S. Department of Commerce  
National Bureau of Standards

Washington, D.C. 20234  
Official Business

Penalty for Private Use \$300



POSTAGE AND FEES PAID  
U.S. DEPARTMENT OF COMMERCE  
COM-215

THIRD CLASS  
BULK RATE