



Technical Note No. 31

Boulder Laboratories

AN ATLAS OF OBLIQUE-INCIDENCE IONOGRAMS

BY VAUGHN AGY, KENNETH DAVIES
AND ROGER SALAMAN



U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS

THE NATIONAL BUREAU OF STANDARDS

Functions and Activities

The functions of the National Bureau of Standards are set forth in the Act of Congress, March 3, 1901, as amended by Congress in Public Law 619, 1950. These include the development and maintenance of the national standards of measurement and the provision of means and methods for making measurements consistent with these standards; the determination of physical constants and properties of materials; the development of methods and instruments for testing materials, devices, and structures; advisory services to government agencies on scientific and technical problems; invention and development of devices to serve special needs of the Government; and the development of standard practices, codes, and specifications. The work includes basic and applied research, development, engineering, instrumentation, testing, evaluation, calibration services, and various consultation and information services. Research projects are also performed for other government agencies when the work relates to and supplements the basic program of the Bureau or when the Bureau's unique competence is required. The scope of activities is suggested by the listing of divisions and sections on the inside of the back cover.

Publications

The results of the Bureau's work take the form of either actual equipment and devices or published papers. These papers appear either in the Bureau's own series of publications or in the journals of professional and scientific societies. The Bureau itself publishes three periodicals available from the Government Printing Office: The Journal of Research, published in four separate editions, presents complete scientific and technical papers; the Technical News Bulletin presents summary and preliminary reports on work in progress; and Basic Radio Propagation Predictions provide data for determining the best frequencies to use for radio communications throughout the world. There are also five series of nonperiodical publications: Monographs, Applied Mathematics Series, Handbooks, Miscellaneous Publications, and Technical Notes.

Information on the Bureau's publications can be found in NBS Circular 460, Publications of the National Bureau of Standards (\$1.25) and its Supplement (\$1.50), available from the Superintendent of Documents, Government Printing Office, Washington 25, D.C.

NATIONAL BUREAU OF STANDARDS

Technical Note

31

November 1959

AN ATLAS OF OBLIQUE-INCIDENCE IONOGRAMS

by

Vaughn Agy, Kenneth Davies
and Roger Salaman

NBS Technical Notes are designed to supplement the Bureau's regular publications program. They provide a means for making available scientific data that are of transient or limited interest. Technical Notes may be listed or referred to in the open literature. They are for sale by the Office of Technical Services, U. S. Department of Commerce, Washington 25, D. C.

DISTRIBUTED BY

UNITED STATES DEPARTMENT OF COMMERCE
OFFICE OF TECHNICAL SERVICES
WASHINGTON 25, D. C.

Price \$ 2.25



AN ATLAS OF OBLIQUE-INCIDENCE IONOGrams

by

Vaughn Agy, Kenneth Davies
and Roger Salaman

INTRODUCTION

This atlas is intended to serve a twofold purpose: first, to provide a compilation of records, of a type becoming standard in the field of ionospheric research, for those workers who are not now familiar with them, and second, to present records which are characteristic of the specific paths used by the National Bureau of Standards for consideration by those using other paths.

To date, NBS sweep-frequency experiments have been carried out over two approximately east-west paths: Sterling, Va. - St. Louis, Mo. (about 1150 km), and Sterling, Va. - Boulder, Colo. (about 2370 km). The experimental data on these paths were taken under the supervision of R. Silberstein and P. G. Sulzer. For each path a midpoint vertical-incidence station was established within about one mile of the geographical midpoint of the great circle path. For the Sterling - St. Louis path the midpoint station was near Batavia, Ohio and for the Sterling - Boulder path it was near Carthage, Illinois. The shorter path was used from August 1951 to December 1952 when the St. Louis equipment was moved west to Boulder. From September 1953 to May 1955, routine records were made over the Sterling-Boulder path, after which time various minor equipment changes were made to improve the time-delay records and to obtain records giving more propagation information for the path.

The data concerning the end-point stations are tabulated below:

<u>Equipment</u>	<u>Power</u>	<u>Time</u>	<u>Sweep Frequency</u>	<u>Pulse Length</u>	<u>Pulse Rep Rate</u>	<u>Antennas</u>
C-3 (modified)	10 kw	12 min*	2-25 Mc	50 μ s**	25/sec	Horizontal Rhombic

*Changed in 1955 to $7\frac{1}{2}$ minutes.

**Unless otherwise noted in the body of the atlas.

For the Sterling-St. Louis path the time at which each record was made (indicated below and to the left of each record shown) is 75° West Meridian Time. For the Sterling-Boulder path, 90° West Meridian Time was used. The time given for each oblique-incidence ionogram is that at which the frequency passed upward through 23 Mc (changed to 24 Mc during the first months of 1958).

The relatively slow sweep time allowed for manual adjustment, when necessary, to synchronize the two equipments. The midpoint vertical-incidence equipment operated in normal fashion, sweeping from 1 Mc to 25 Mc in 15 seconds once every 6 minutes and later once every 3 minutes. The 3 minute spacing between sweeps allowed selection of the vertical-incidence ionogram made within $1\frac{1}{2}$ minutes of the time the slower oblique-incidence sweep passed through the frequency of interest, e.g., the MUF.

The oblique-incidence ionograms are plots of time-delay against frequency (on a linear scale). The ordinate (time-delay) scale, however, just as for vertical-incidence ionograms, is calibrated in terms of the apparent virtual half-path. In order to determine the absolute time delay (or virtual half-path) between the two stations at a given frequency, it is necessary to take the arithmetic mean of the ordinates on the two end-point records. For given virtual heights* of reflection, the virtual half-paths are plotted in Figure 1 for the Sterling-St. Louis path and in Figure 2 for the Sterling-Boulder path. In addition to the 1-, 2-, 3-hop modes, the M and N modes are presented involving both F-layer reflections and E-layer reflections (with the E-layer virtual height assumed to be 100 km).

The ionograms for the shorter path usually show vertical-incidence as well as oblique-incidence traces; for the longer path, however, in order to avoid undue compression of the virtual half-path scale, the first 1000 km of half-path are not recorded. On some of the later records, one or both scales are expanded in order to allow examination of some of the detailed structure of the recorded traces.

In some cases (e.g. Section I-3) the midpoint vertical-incidence records are included for comparison. Although not always labeled, these records are easily identified by the logarithmic frequency scale. The end-point records, unless otherwise noted, were those made at the western terminus of the path, i.e. St. Louis for the shorter path and Boulder for the longer path. The primary reason for this choice is the fact that these records usually are freer of interference than those made at Sterling.

*Virtual height in this sense, for the 1-hop mode, is the height above a spherical earth of the apex of an isosceles triangle whose base is the chord between the two stations and whose legs are equal in length to the absolute virtual half-path between the stations.




FIGURE 1




FIGURE 2

The body of the atlas is divided into three parts, giving examples of records made: (1) over the Sterling-St. Louis (1150 km) path, (2) over the Sterling-Boulder (2370 km) path during the routine phase of this part of the experiment, and (3) over the Sterling-Boulder path during the more recent phase of the experiment in which equipment changes were made to increase the observable detail in the ionograms and to make measurements of recorded pulse amplitudes. Some sequences of ionograms are included to show the variations during a day, or during shorter periods when the development of a specific phenomenon can be seen. Other single records are included simply to show the occurrence rather than the development of a particular phenomenon. Many of the records for which interpretations are not readily evident are included because of the interest they may arouse. However, it is not the purpose of the atlas to give detailed interpretations of the ionograms shown.

For the most part, the analysis of the records has consisted of a comparison of the MUF determined directly from the oblique-incidence records with that obtained by applying the appropriate Smith transmission curves (see references) to the midpoint vertical-incidence ionograms. It is worth noting that although, on the average, the scaling of MUF from the vertical-incidence midpoint records agrees well with the observed MUF, there are frequent discrepancies in detail. For an eventual understanding of oblique-incidence propagation, the discrepancies are just as important as the cases of close agreement.

The use of various descriptive terms has developed to aid in the description of certain aspects of the oblique-incidence ionograms. One of these terms is "nose", which is suggested by the shape of the trace on an oblique-incidence ionogram indicating the merging of a high-angle and a low-angle ray. It has become customary occasionally to speak of the "junction frequency" or simply the "nose frequency" for a particular layer when referring to the classical MUF for that layer, but when complications arise, as shown in Sections II-3 and III-3, "nose" may also be used to refer to a junction frequency different from the MUF.

REFERENCES
(NBS work of special interest)

1. Ionospheric Radio Propagation, Chap. 6, NBS Circular 462 (1948).
2. P. G. Sulzer and E. E. Ferguson, Proc. I.R.E. 40, 1124 (1952).
3. B. Wieder, J. Geophys. Research, 60, 395 (1955).
4. P. G. Sulzer, J. Geophys. Research, 60, 411 (1955).
5. V. Agy and K. Davies, J. Research NBS, 63D, No. 2 (Sept-Oct. 1959).

INDEX OF SELECTIONS

I. Sterling-St. Louis - 1150 km

1. Diurnal Variations
2. Post Sunrise Development of the F Layers
3. Spread Echo
4. MUF Extensions
5. Equipment Sensitivity
6. Importance of the High-Angle Ray
7. Sporadic E
8. Moving Irregularities
9. Unusual and Complex Traces

II. Sterling-Boulder - 2370 km

(Routine)

1. Diurnal Variations
2. Post Sunrise Development
3. Miscellaneous Sequences

Apparent Stratifications
Complex Traces
MUF Extension
Sporadic E

4. Miscellaneous Ionograms


Multi-hop Traces
F1 MUF
E2 Trace
O-X Frequency Separation

III. Sterling-Boulder - 2370 km

(Experimental)

1. Magnetic Storm
 - A. 3-day Sequence
 - B. Detailed Sequence During Storm
2. Morning and Afternoon Sequences
3. Inner and Outer "Nose"
4. "Disturbed F2 Nose"
5. Disturbed Conditions
6. Rarely Observed Ionograms
7. Equipment Effects
8. Fixed Frequency vs. Sweep Frequency
9. Field Strength Variations near the MUF

I. Sterling-St. Louis - 1150 km



Sterling-St. Louis

Sequences Showing Diurnal Variations

Winter Day
January 9-10, 1952

$$\Sigma K_p = 28^\circ$$

Equinoctial Day (Spring)
March 5-6, 1952

$$\Sigma K_p = 47^\circ$$

Equinoctial Day (Spring)
April 2, 1952

$$\Sigma K_p = 48-$$

Summer Day
May 28, 1952

$$\Sigma K_p = 38+$$


Summer Day
June 25-26, 1952

$$\Sigma K_p = 30^\circ$$


Equinoctial Day (Fall)
September 18-19, 1952

$$\Sigma K_p = 8-$$


JANUARY 9, 1952




1204



1504





1804





2052

JANUARY 10, 1952




MARCH 5 - 6, 1952



1817
10 15 20
Mc


0013
20 15 10 5
Mc




1505
10 15 20
Mc

2053
20 15 10 5
Mc


MARCH 6, 1952



0901




0701




1201


APRIL 2, 1952




1510 Mc



1127




2058




1810


APRIL 3, 1952




0010




0258



0858




20




0558

MAY 28, 1952




↓
100 Km
↓

20
15
10 Mc
5
1508




↓
100 Km
↓

20
15
10 Mc
5
2108



↓
100 Km
↓


20
15
10 Mc
5
1205



↓
100 Km
↓


20
15
10 Mc
5
1808

MAY 29, 1952




0256

100 Km




0908

100 Km



0008


100 Km




0556

100 Km


JUNE 25, 1952




JUNE 26, 1952



0256 5 10 15 20
Mc







0904 5 10 15 20
Mc




25
0256 5 10 15 20
Mc


SEPTEMBER 18, 1952




SEPTEMBER 19, 1952




0007




0255



0631



0907



Sterling-St. Louis

Sequences Showing Development of the F Layers
After Sunrise


April 24, 1952

The F1 trace is seen to develop out of the low angle F2 trace and the night F layer is continuous with the day F2 layer.


May 15, 1952

Here the F2-layer trace develops out of the Pedersen ray trace of the night F layer. The night F layer is continuous with the day F1 layer.


APRIL 24, 1952




0757




0821



0857




0933




1009


MAY 15, 1952




0701




0725



0649



0713



Sterling-St. Louis

Ionograms Showing Spread Echo

May 1, 1952

The midpoint vertical-incidence record is not only spread but weak, indicating high absorption. The oblique-incidence record is somewhat spread but appears to be relatively strong.

May 29, 1952

On these records spread echoes are evident at vertical incidence but the oblique-incidence traces are relatively sharp.

May 29, 1952


Moderate spread on the oblique-incidence records accompanies severe spread seen at vertical incidence.

The great amount of spread apparent on the midpoint vertical-incidence ionograms represents a complicated ionospheric structure which shows itself in the complex traces seen on the oblique-incidence ionograms.

July 25, October 21, 1952


The July 25 ionograms show more spread at oblique incidence than at vertical incidence; the October 21 records show comparable spread.

MAY 1, 1952




↓
100 Km
↑

MAY 29, 1952
0400




↓
100 Km
↑

MAY 1, 1952



↓
100 Km
↑

MAY 29, 1952
0409



↓
100 Km
↑

0242

2 3 4 5 6 8 10
Mc

20

15

5


10
Mc

20

15
Mc

5
Mc


MAY 29, 1952




0400



0501




0408




0508


MAY 29, 1952




0532




0512



0544




0530



0608


OBLIQUE




0548

VERTICAL


JULY 25, 1952




JULY 25, 1952




OCTOBER 21, 1952
0648



OCTOBER 21, 1952
0636



OCTOBER 21, 1952
2103



Sterling-St. Louis


Ionograms Showing MUF Extensions

September 11, 1952; October 3, September 3, 1951; September 12, July 18, September 26, 1952

Note the following:


- (1) The echoes at frequencies above that which the high-angle and low-angle rays merge. This has been named the MUF extension.
- (2) On July 18, September 12 and September 26, it is difficult to separate the 2-hop E echo from the 1-hop F1 echo.
- (3) On September 26 there is no extension on the F2 trace, perhaps because of the presence of the underlying F1 layer.
- (4) The presence of the MUF extension does not appear to be correlated with the presence of sporadic E.

SEPTEMBER 11, 1952




1935

SEPTEMBER 11, 1952




2211

OCTOBER 3, 1951




2124

SEPTEMBER 3, 1951






2224

SEPTEMBER 12, 1952



JULY 18, 1952



SEPTEMBER 26, 1952



Sterling-St. Louis

Ionograms Showing Effects of Equipment
Sensitivity



May 28, July 18, September 5, 1952

- (1) Note in the May 28 records the relative strength of the high-angle F1 ray.
- (2) In the ionograms for July 18 the number of multiples is markedly reduced as the gain is reduced.
- (3) In the September 5 records the disappearance of the Es trace is to be noted.



MAY 28, 1952




JULY 18, 1952



SEPTEMBER 5, 1952





Sterling-St. Louis

Ionograms Showing Relative Importance of High-Angle Ray

May 14, 1952

The one-hop low-angle ray is cut off by sporadic E which is evident on the oblique-incidence record (1945) but not on the vertical-incidence ionogram (1942). The high-angle ray may, therefore, be relatively important (note the presence of M and N-type echoes).

May 15, 1952

Here again the low-angle ray has been cut off by intense sporadic E.


October 3, November 18, December 2, 1952

The high-angle ray here is relatively unimportant. This is especially typical of winter day conditions.


February 13, July 17, August 28, September 11, 1952

An example showing the relatively rare high-angle E-layer ray.


MAY 14, 1952




1909




1906




1945



1942




1957




2000


MAY 15, 1952



0833




0845




0857

OCTOBER 3, 1951




1230

OCTOBER 3, 1951




1205

NOVEMBER 18, 1952




1411

DECEMBER 2, 1952



1218

FEBRUARY 13, 1952




1525

20
15
10 Mc

1826
5
10 Mc

AUGUST 28, 1952




1533

20
15
10 Mc

1613
5
10 Mc

SEPTEMBER 11, 1952




100 Km

100 Km

20
15
10 Mc

1513
5
10 Mc

JULY 17, 1952



100 Km

20
15
10 Mc

1826
5
10 Mc


Sterling-St. Louis

Ionograms Showing Sporadic E Reflections


May 15, July 17, 1952


On both records Es extends out past 20 Mc and the record for May 15 shows blanketing of the F-layer echoes. The record for July 17 gives an example of complex echo structure.

MAY 15, 1952



JULY 17, 1952





Sterling-St. Louis

Sequences Showing Moving Irregularities


April 30, 1952

Moving irregularity seen on oblique-incidence ionograms
not apparent on the vertical-incidence records.


May 28, 1952

Apparent downward motion of irregularity indicated.


APRIL 30, 1952




2122




2118



2134




2130



2146


OBLIQUE



2136

VERTICAL

MAY 28, 1952




10

5

2220
Mc

↓
100 Km
↓




10

5

2208
Mc

↓
100 Km
↓




10

5

2156
Mc

↓
100 Km
↓




10

5

2256
Mc

↓
100 Km
↓




10

5

2244
Mc

↓
100 Km
↓



10

5

2230
Mc

↓
100 Km
↓

MOVING IRREGULARITIES

Sterling-St. Louis

Ionograms Showing Unusual and Complex Traces

February 6, March 19, December 3, 1952

Apparent stratifications and complex traces.

May 22, 1952



Here the complexities show up in the 2-hop echo trace.

August 8, May 8, 1952

The Pedersen ray ordinary and extraordinary traces are differently shaped.



FEBRUARY 6, 1952

MARCH 19, 1952




MARCH 19, 1952


DECEMBER 3, 1952




DECEMBER 3, 1952




MAY 22, 1952



0841




0853




0905

AUGUST 8, 1951



1113

MAY 8, 1952



2332

II. Sterling-Boulder - 2370 km
(Routine)

Sterling-Boulder
(Routine)

Sequences Showing Diurnal Variation

Winter Day
February 3-4, 1954

$$\Sigma K_p = 18-$$

Winter Day
February 10-11, 1954

$$\Sigma K_p = 18+$$

Equinoctial Day
March 31-April 1, 1954

$$\Sigma K_p = 15-$$


Summer Day
May 11-12, 1954

$$\Sigma K_p = 15-$$


Summer Day
August 4-5, 1954

$$\Sigma K_p = 10$$


FEBRUARY 3-4, 1954




20
15
10
Mc
5
2103



20
15
10
Mc
5
1815




20
15
10
Mc
5
0303




20
15
10
Mc
5
0015

FEBRUARY 4, 1954



0603




1203



0903




1503




2003


FEBRUARY 10, 1954




1200



1512



1812




2100

20
15
10
Mc

20
15
10
Mc

20
15
10
Mc

FEBRUARY 11, 1954




0012



0300




0600




0900


MARCH 31, 1954



1158




1258




1510



1810




2058




2358


APRIL 1, 1954




0258




0558



0858





1158




1458


MAY 11, 1954





MAY 12, 1954




MAY 12, 1954




MIDPOINT VERTICAL INCIDENCE




AUGUST 4, 1954




1157



1509




2057



1757




2057




1509


AUGUST 4-5, 1954




0257



0857



2357



0545

Sterling-Boulder
(Routine)

Sequences Showing Morning Bifurcation of the F layer

April 1, 1954

"Splitting at the bottom": F1 layer develops under
the nighttime F layer.


August 11, 1954

Continuity between nighttime F layer and daytime F1 layer.


April 14, 1954

"Splitting at the top": Nighttime F layer is continuous
with the daytime F1 layer.


APRIL 1, 1954




0610



0634




0658



0722


APRIL 1, 1954




0746



0810







0834



100 Km

AUGUST 11, 1954



AUGUST 11, 1954



0735

20
15
10 Mc
5
0748



0748

20
15
10 Mc
5









0801

20
15
10 Mc
5

0813

20
15
10 Mc
5

APRIL 14, 1954



Sterling-Boulder
(Routine)

Miscellaneous Sequences

November 5, 1953

Oblique-incidence records showing development of apparent stratifications, possibly produced by "off-path" reflections. Some ionospheric roughness indicated by slight spread and presence of Z-trace on vertical-incidence records.

November 12, 1953

Complexities in oblique-incidence records, perhaps associated with "oblique" reflection and forked trace evident on the midpoint vertical-incidence ionograms shown.

June 23, 1954

Sporadic E reflections out to 24 Mc; pronounced MUF extension.

July 21, 1954

Complex nose structure; Es reflections.










August 25, 1954

Es evident from traces between those for 1XF and 2XF. Complex nose structure.


November 12, 1953

Apparent stratification and inner "nose". (See footnote in Section III-3)


NOVEMBER 5, 1953




NOVEMBER 5, 1953




0203



0318




0327




0339
MIDPOINT VERTICAL INCIDENCE


NOVEMBER 12, 1953




1853



1905






1917




1929


NOVEMBER 12, 1953



JUNE 23, 1954




1803




1815

JUNE 23, 1954




1827




1839

JULY 21, 1954




1922




1934

JULY 21, 1954




1946




1958


AUGUST 25, 1954




2035



2047




2059




2111


NOVEMBER 12, 1953



1805



1817



1829

Sterling-Boulder
(Routine)

Miscellaneous Ionograms

January 7, 1954

Five F-layer modes present

July 22, 1954

The record indicates the difficulty encountered in specifying exactly the F1 MUF for this path.


August 25, 1954

E2, F1, and F2 traces evident.

April 5, 1955


The Pedersen ray gives a clear indication of the increase in O-X frequency separation as the angle of incidence increases.

JANUARY 7, 1954




0517

JULY 22, 1954




0934

AUGUST 25, 1954



1059

APRIL 5, 1955



1647

III. Sterling-Boulder - 2370 km
(Experimental)

(All ionograms with expanded frequency scales
were made using 20 μ s pulses)

III-1A

Sterling-Boulder
(Experimental)

June 28-29, 1958 Magnetic Storm





Ionogram Series from Quiet Day before Storm
to Quiet Day after Storm

Date	Midpoint Time	K _p	Date	Midpoint Time	K _p
6/27/59	0759	1+	6/29/59	0759	8-
	1059	1+		1059	6+
	1359	1+		1359	4+
	1659	3o		1659	3-
	1959	2+		1959	2o
	2259	3+		2259	3-
6/28/59	0159	4o	6/30/59	0152	2-
	0459	4-		0459	1+
	0759	3o		0759	2-
	1059	5-		1059	3-
	1414	7o		1407	3o
	1659	8-		1559	2+
	1959	7o			
	2259	8o			
6/29/59	0159	7-			
	0429	7o			

BOULDER

JUNE 27, 1958

WASHINGTON




OBlique

VERTICAL


BOULDER

JUNE 27-28, 1958


WASHINGTON




1959



2259




0159




0459

OBlique


VERTICAL



2000



0200




0500


BOULDER

JUNE 28, 1958


WASHINGTON




0759



1059




1414




1659

OBLIQUE


VERTICAL




0800



1000



1415




1700

BOULDER

JUNE 28-29, 1958


WASHINGTON



1959

2000


MC



0200

2259

MC




0429

0500

MC


OBLIQUE

VERTICAL




2300

MC



0200

MC









0500

MC

BOULDER

JUNE 29, 1958

WASHINGTON




OBlique

VERTICAL


BOULDER

JUNE 29-30, 1958


WASHINGTON




1959




2259




2000



0152



0200



0459


OBLIQUE

VERTICAL


BOULDER

JUNE 30, 1958


WASHINGTON




0759




1059




0800



1100



1400



1407

1559

OBLIQUE

VERTICAL

III-1B

**Sterling-Boulder
(Experimental)**


Detailed Sequence During Magnetic Storm

June 28-29, 1958

BOULDER

JUNE 28, 1958


WASHINGTON




1822




1815




1944




1945




1959



2000



2144




OBlique

VERTICAL


BOULDER

JUNE 28 - 29, 1958


WASHINGTON



2237




2259



0037


OBLIQUE

BOULDER




0030


VERTICAL



2236



2300




0030


BOULDER

JUNE 29, 1958

WASHINGTON



0152




0159




0200

OBLIQUE

BOULDER




0245



VERTICAL


JUNE 29, 1958

WASHINGTON



0337

BOULDER




0330

WASHINGTON




0445

OBLIQUE



BOULDER



VERTICAL


BOULDER

0430

BOULDER

JUNE 29, 1958


WASHINGTON



0514


BOULDER

WASHINGTON



0530

WASHINGTON




0759

OBLIQUE

BOULDER

WASHINGTON



0800

VERTICAL

Sterling-Boulder
(Experimental)

Morning and Afternoon Sequences

June 20, 1958

Unusual morning sequence showing the development of an F1 layer and then an F2 layer.

June 9, 1958

Normal afternoon sequence. The trace of a possible off-path signal may be seen from 1813 to 1835 CST.


August 6, 1958

Normal afternoon sequence. A disturbance in the F2 layer can be seen on the oblique- and vertical-incidence ionograms at 1454 CST.

WASHINGTON

JUNE 20, 1958

BOULDER




VERTICAL

OBLIQUE

WASHINGTON

JUNE 20, 1958







VERTICAL

OBLIQUE





BOULDER

JUNE 9, 1958



BOULDER







JUNE 9, 1958



CARTHAGE

AUGUST 6, 1957

BOULDER









OBLIQUE

VERTICAL

CARTHAGE

AUGUST 6, 1957

BOULDER









OBLIQUE

VERTICAL

CARTHAGE

AUGUST 6, 1957

BOULDER



OBlique

VERTICAL

Sterling-Boulder
(Experimental)

"Inner"* and "Outer" Nose

August 9, 1957
August 23, 1957


Sequences showing development of "inner" nose on oblique-incidence ionograms and simultaneous midpoint vertical-incidence disturbances.

June 19, 1958

Sequence showing development of "outer" nose on oblique-incidence ionograms and blanketing by Es on end-point vertical-incidence ionograms.






*Occasionally, as shown in the following records, in addition to the well defined "nose" representing the classical F2 MUF, a second nose appears at frequencies below (or above) the classical MUF. The additional nose, for the sake of brevity, is referred to as an "inner" (or "outer") nose.

STERLING



CARTHAGE

AUGUST 9, 1957









OBLIQUE

VERTICAL

STERLING

CARTHAGE

AUGUST 9, 1957






OBLIQUE

VERTICAL

STERLING







AUGUST 23, 1957



OBLIQUE

CARTHAGE

AUGUST 23, 1957




VERTICAL

WASHINGTON


JUNE 19, 1958

BOULDER




2014

2015



2022

2020



2029

2030

OBLIQUE

VERTICAL

WASHINGTON

JUNE 19, 1958


BOULDER




2037




2035



2044
Mc



2045
Mc



2050
Mc

OBLIQUE

VERTICAL

Sterling-Boulder
(Experimental)

"Disturbed F2 Nose"

April 30, 1957

Some spread can be seen on midpoint vertical-incidence ionograms at time of "disturbed nose" on oblique-incidence ionograms.

April 20, 1958

May 12, 1958

Sequences of "disturbed-nose" oblique-incidence ionograms.




June 24, 1958

Spread nose on oblique-incidence ionogram and spread F on end-point vertical-incidence ionograms.

STERLING

CARTHAGE

APRIL 30, 1957




OBLIQUE


VERTICAL

BOULDER


APRIL 20, 1958




1620



1710




1732




1740

BOULDER


MAY 12, 1958




1611




1618




1626



1633




1656




1703

CARTHAGE

APRIL 30, 1957



2208




2209


STERLING

OBLIQUE

VERTICAL




2245



2246

OLUDER


APRIL 20, 1958



1547

Mc


25



1555

Mc


25



1602

Mc

25




1617

Mc


25

BOULDER


JUNE 24, 1958



WASHINGTON



0210
BOULDER



0330
VERTICAL

III-5

Sterling-Boulder
(Experimental)

Disturbed Conditions

February 5, 1957

Sequence showing traveling disturbance in oblique- and vertical-incidence ionograms.







June 9, 1958

Disturbed oblique-incidence and vertical-incidence night ionograms. K_p remained between 5+ and 5-. (See also the June 9, 1958 example of Morning and Afternoon Sequences in Section III-2)

CARTHAGE

FEBRUARY 5, 1957

STERLING






OBLIQUE

VERTICAL




CARTHAGE

FEBRUARY 5, 1957

STERLING




OBLIQUE




VERTICAL

BOULDER


JUNE 9, 1958




2056



2104




2111




2119

BOULDER


JUNE 9, 1958




2126



2156







2204



2211


BOULDER

JUNE 9, 1958




BOULDER


JUNE 9, 1958




2326



2334



2341











2349

BOULDER

WASHINGTON

JUNE 9, 1958



VERTICAL

III-6

Sterling-Boulder (Experimental)

Rarely Observed Ionograms

May 5, 1958

The appearance of the extraordinary component as a nose extension.

April 9, 1957

The rounded F2 nose may be attributed to the similarity between the vertical-incidence ionogram and the transmission curve.




August 2, 1956

August 30, 1957

The high-angle ray of the E mode is seldom seen on the 2400 km path.


BOULDER

MAY 5, 1958



BOULDER


APRIL 9, 1957



2109

OBLIQUE

CARTHAGE




2109

VERTICAL


BOULDER

AUGUST 2, 1956




STERLING

AUGUST 30, 1957



BOULDER



Sterling-Boulder
(Experimental)

Equipment Effects


The June 24, 1958, 0659 and 0707 CST ionograms were received using 100 and 20 μ s transmitted pulses respectively. The June 26, 1958, 1944 CST ionogram was received with every 10th transmitter pulse changed to 100 or 20 μ s. The loss of detail with the longer pulse can be seen.

The top records on August 1, 1958 are those ordinarily received. The bottom records were taken simultaneously with large differentiation in the receiving system.

The 1323 CST ionogram of July 23, 1957 illustrates the effect of transmitter double pulsing. The parallel traces on the March 12, 1957 and April 23, 1957 ionograms may also be attributed to a deformed transmitter pulse.

BOULDER

JUNE 24, 1958




0659



0707



JUNE 26, 1958






1944

BOULDER

AUGUST 1, 1958




100 Km




BOULDER

JULY 23, 1957




STERLING


MARCH 12, 1957



APRIL 23, 1957



APRIL 23, 1957



1323

2032

1855

Sterling-Boulder
(Experimental)

Fixed Frequency vs. Sweep Frequency


May 7, 1958

The first ionogram shows the pulse received on 20.084 Mc, from 1920 to 2100 CST. The following sweep-frequency ionograms were obtained during the same period. Both transmitters used a 25 μ s pulse at a 25 pps repetition rate. The output of the fixed-frequency transmitter was about 200 kw.


The hesitation of the F2 MUF at about 20 Mc appears as a nose extension on the fixed-frequency record.

BOULDER


MAY 7, 1958




20.084 Mc



1941




1956




2011

BOULDER


MAY 7, 1958




2026



2041



2056



2111

Sterling-Boulder
(Experimental)A Sequence of A-scan Records Showing Field Strength
Variations Near the MUF

In addition to the time-delay studies, some attention has been given to received field strengths. Here is shown a series of A-scan records made at Boulder and the corresponding oblique-incidence ionograms. The frequency at which the A-scan records were made increases upward in each column and from left to right. These were made at the rate of $2\frac{1}{2}$ per second (or every .025 Mc). On each record, time delay increases toward the right. The pulse amplitudes — the scale is approximately linear — vary somewhat due to normal fading.

At frequencies greater than 14.8 Mc (near the top of the first column) the Pedersen ray first makes its appearance. At about 15.1 Mc (at the top of the second column) it shows two component pulses, the ordinary and the extraordinary modes. At about 15.3 Mc (near the top of the third column) similar slight separation of the O and X modes in the low-angle ray can be seen. Following the progression upward in the fourth column, as the low-angle and high-angle rays begin to overlap (just above 15.4 Mc) the pulse structure becomes complex; at 15.475 Mc the ordinary rays have combined and at 15.55 Mc only the extraordinary ray pulses are left. These have combined at about 15.625 Mc and have disappeared at 15.7 Mc.

OBLIQUE INCIDENCE AMPLITUDES
NEAR MUF

BOULDER - STERLING PATH (2370 KM)
JULY 16, 1957, 2054 CST

