

CRPL-F 111

REMOVED FROM THE LIBRARY.

FOR OFFICIAL USE

100-100000  
Merry, R. W. Big  
7 (LC/1953)

## IONOSPHERIC DATA

ISSUED  
NOVEMBER 1953

U. S. DEPARTMENT OF COMMERCE  
NATIONAL BUREAU OF STANDARDS  
CENTRAL RADIO PROPAGATION LABORATORY  
WASHINGTON, D. C.



## IONOSPHERIC DATA

## CONTENTS

	<u>Page</u>
Symbols, Terminology, Conventions . . . . .	2
World-Wide Sources of Ionospheric Data . . . .	5
Hourly Ionospheric Data at Washington, D. C. .	7, 12, 20, 46
Ionospheric Storminess at Washington, D. C. .	7, 32
Relative Sunspot Numbers . . . . .	7, 33
Radio Propagation Quality Figures . . . . .	8, 34
Observations of the Solar Corona . . . . .	9, 36
Sudden Ionosphere Disturbances . . . . .	10, 42
Observations of Solar Flares . . . . .	10, 44
Indices of Geomagnetic Activity . . . . .	11, 45
Errata . . . . .	11
Tables of Ionospheric Data . . . . .	12
Graphs of Ionospheric Data . . . . .	46
Index of Tables and Graphs of Ionospheric Data in CRPL-Fill . . . . .	70

## SYMBOLS, TERMINOLOGY, CONVENTIONS

Beginning with data reported for January 1952, the symbols, terminology, and conventions for the determination of median values used in this report (CRPL-F series) conform as far as practicable to those adopted at the Sixth Meeting of the International Radio Consultative Committee (C.C.I.R.) in Geneva, 1951. Excerpts concerning symbols and terminology from Document No. 626-E of this Meeting are given on pages 2-7 of the report CRPL-F89, "Ionospheric Data," issued January 1952. Reprints of these pages are available upon request.

Beginning with data for January 1945, median values are published wherever possible. Where averages are reported, they are, at any hour, the average for all the days during the month for which numerical data exist.

The following conventions are used in determining the medians for hours when no measured values are given because of equipment limitations and ionospheric irregularities. Symbols used are those given in Document No. 626-E referred to above.

### a. For all ionospheric characteristics:

Values missing because of A, C, F, L, M, N, Q, S, or T are omitted from the median count.

### b. For critical frequencies and virtual heights:

Values of  $f_{oF2}$  (and  $f_{oE}$  near sunrise and sunset) missing because of E are counted as equal to or less than the lower limit of the recorder. Values of  $h'F2$  (and  $h'E$  near sunrise and sunset) missing for this reason are counted usually as equal to or greater than the median. Other characteristics missing because of E are omitted from the median count.

Values missing because of D are counted as equal to or greater than the upper limit of the recorder.

Values missing because of G are counted:

1. For  $f_{oF2}$ , as equal to or less than  $f_{oF1}$ .
2. For  $h'F2$ , as equal to or greater than the median.

The symbol W is included in the median count only when it replaces a height characteristic. This practice represents a change from that listed in issues previous to CRPL-F78.

Values missing for any other reason are omitted from the median count.

c. For MUF factor (M-factors):

Values missing because of G or W are counted as equal to or less than the median.

Values missing for any other reason are omitted from the median count.

d. For sporadic E ( $E_s$ ):

Values of f $E_s$  missing because of E or G (and B when applied to the daytime E region only) are counted as equal to or less than the median foE, or equal to or less than the lower frequency limit of the recorder.

Values of f $E_s$  missing for any other reason, and values of h $E_s$  missing for any reason at all are omitted from the median count.

Beginning with data for November 1945, doubtful monthly median values for ionospheric observations at Washington, D. C., are indicated by parentheses, in accordance with the practice already in use for doubtful hourly values. The following are the conventions used to determine whether or not a median value is doubtful:

1. If only four values or less are available, the data are considered insufficient and no median value is computed.

2. For the F2 layer, if only five to nine values are available, the median is considered doubtful. The E and F1 layers are so regular in their characteristics that, as long as there are at least five values, the median is not considered doubtful.

3. For all layers, if more than half of the values used to compute the median are doubtful (either doubtful or interpolated), the median is considered doubtful.

The same conventions are used by the CRPL in computing the medians from tabulations of daily and hourly data for stations other than Washington, beginning with the tables in IRPL-F18.

The tables and graphs of ionospheric data are correct for the values reported to the CRPL, but, because of variations in practice in the interpretation of records and scaling and manner of reporting of values, may at times give an erroneous conception of typical ionospheric characteristics at the station. Some of the errors are due to:

- a. Differences in scaling records when spread echoes are present.
- b. Omission of values when  $f_{oF2}$  is less than or equal to  $f_{oFl}$ , leading to erroneously high values of monthly averages or median values.
- c. Omission of values when critical frequencies are less than the lower frequency limit of the recorder, also leading to erroneously high values of monthly average or median values.

These effects were discussed on pages 6 and 7 of the previous F-series report IRPL-F5.

Ordinarily, a blank space in the fEs column of a table is the result of the fact that a majority of the readings for the month are below the lower limit of the recorder or less than the corresponding values of  $f_{oE}$ . Blank spaces at the beginning and end of columns of  $h'Fl$ ,  $f_{oFl}$ ,  $h'E$ , and  $f_{oE}$  are usually the result of diurnal variation in these characteristics. Complete absence of medians of  $h'Fl$  and  $f_{oFl}$  is usually the result of seasonal effects.

The dashed-line prediction curves of the graphs of ionospheric data are obtained from the predicted zero-muf contour charts of the CRPL-D series publications. The following points are worthy of note:

- a. Predictions for individual stations used to construct the charts may be more accurate than the values read from the charts since some smoothing of the contours is necessary to allow for the longitude effect within a zone. Thus, inasmuch as the predicted contours are for the center of each zone, part of the discrepancy between the predicted and observed values as given in the F series may be caused by the fact that the station is not centrally located within the zone.
- b. The final presentation of the predictions is dependent upon the latest available ionospheric and radio propagation data, as well as upon predicted sunspot number.

c. There is no indication on the graphs of the relative reliability of the data; it is necessary to consult the tables for such information.

The following predicted smoothed 12-month running-average Zürich sunspot numbers were used in constructing the contour charts:

Month	Predicted Sunspot Number								
	1953	1952	1951	1950	1949	1948	1947	1946	1945
December	33	53	86	108	114	126	85	38	
November	38	52	87	112	115	124	83	36	
October	17	43	52	90	114	116	119	81	23
September	18	46	54	91	115	117	121	79	22
August	18	49	57	96	111	123	122	77	20
July	20	51	60	101	108	125	116	73	
June	21	52	63	103	108	129	112	67	
May	22	52	68	102	108	130	109	67	
April	24	52	74	101	109	133	107	62	
March	27	52	78	103	111	133	105	51	
February	29	51	82	103	113	133	90	46	
January	30	53	85	105	112	130	88	42	

### WORLD - WIDE SOURCES OF IONOSPHERIC DATA

The ionospheric data given here in tables 1 to 48 and figures 1 to 96 were assembled by the Central Radio Propagation Laboratory for analysis and correlation, incidental to CRPL prediction of radio propagation conditions. The data are median values unless otherwise indicated. The following are the sources of the data in this issue:

República Argentina, Ministerio de Marina:  
 Buenos Aires, Argentina  
 Decepcion I.

Australian Department of Supply and Shipping, Bureau of Mineral Resources, Geology and Geophysics:  
 Watheroo, Western Australia

Meteorological Service of the Belgian Congo and Ruanda-Urundi:  
 Leopoldville, Belgian Congo

Defence Research Board, Canada:

Baker Lake, Canada  
Churchill, Canada  
Fort Chimo, Canada  
Ottawa, Canada  
Prince Rupert, Canada  
Resolute Bay, Canada  
St. John's, Newfoundland  
Winnipeg, Canada

The Royal Netherlands Meteorological Institute:

De Bilt, Holland

Indian Council of Scientific and Industrial Research, Radio Research Committee:  
Calcutta, India

Ministry of Postal Services, Radio Research Laboratories, Tokyo, Japan:

Akita, Japan  
Tokyo (Kokubunji), Japan  
Wakkanai, Japan  
Yamagawa, Japan

Christchurch Geophysical Observatory, New Zealand Department of Scientific  
and Industrial Research:

Christchurch, New Zealand  
Barotonga, Cook Is.

Norwegian Defence Research Establishment, Kjeller per Lillestrom, Norway:

Oslo, Norway  
Tronso, Norway

Manila Observatory:

Baguio, P. I.

Research Institute of National Defence, Stockholm, Sweden:

Upsala, Sweden

Post, Telephone and Telegraph Administration, Berne, Switzerland:

Schwarzenburg, Switzerland

United States Army Signal Corps:

Okinawa I.  
White Sands, New Mexico

National Bureau of Standards (Central Radio Propagation Laboratory);  
Anchorage, Alaska  
Baton Rouge, Louisiana (Louisiana State University)  
Fairbanks, Alaska (Geophysical Institute of the University of Alaska)  
Guam I.  
Maui, Hawaii  
Narsarsuaq, Greenland  
Panama Canal Zone  
Point Barrow, Alaska  
Puerto Rico, W. I.  
San Francisco, California (Stanford University)  
Washington, D. C.

#### HOURLY IONOSPHERIC DATA AT WASHINGTON, D. C.

The data given in tables 49 through 60 follow the scaling practices given in the report IRPL-C61, "Report of International Radio Propagation Conference," pages 36 to 39, and the median values are determined by the conventions given above under "Symbols, Terminology, Conventions." Beginning with September 1949, the data are taken at Ft. Belvoir, Virginia.

#### IONOSPHERIC STORMINESS AT WASHINGTON, D.C.

Table 61 presents ionosphere character figures for Washington, D. C., during October 1953, as determined by the criteria given in the report IRPL-R5, "Criteria for Ionospheric Storminess," together with Cheltenham, Maryland, geomagnetic K-figures, which are usually covariant with them.

#### RELATIVE SUNSPOT NUMBERS

Table 62 lists the daily provisional Zürich relative sunspot number,  $R_Z$ , as communicated by the Swiss Federal Observatory. Publication of the American relative sunspot numbers,  $R_A$ , which usually appear monthly in these pages, is temporarily suspended until new arrangements are made for the reduction of the observations made by the Solar Division of the AAVSO.

Tables 63a and 63b give for September 1953 the radio propagation quality figures for the North Atlantic area, CRPL advance and short-term forecasts, a summary geomagnetic activity index and sundry comparisons, specifically as follows:

- (a) radio propagation quality figures, separately for each 6-hour interval of the Greenwich day, viz., 00-06, 06-12, 12-18, 18-24 hours UT (Universal Time or GCT).
- (b) whole-day radio quality indices (beginning October 1952). Each index is a weighted average of the four quarter-day Q-figures, before rounding off, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which designate whenever possible the days when significant disturbance or unusually quiet conditions will occur.
- (c) short-term forecasts, issued by CRPL every six hours (nominally one hour before 00<sup>h</sup>, 06<sup>h</sup>, 12<sup>h</sup>, 18<sup>h</sup> UT) and applicable to the period 1 to 13 (especially 1 to 7) hours ahead. Note that new scoring rules have been adopted beginning with October 1952 data.
- (d) advance forecasts, issued semiweekly (CRPL-J reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole-day quality indices.
- (e) half-day averages of the geomagnetic K indices measured by the Cheltenham Magnetic Observatory of the U. S. Coast and Geodetic Survey.
- (f) illustration of the comparison of short-term forecasts with Q-figures and also with estimates of radio quality based on CRPL observations only.
- (g) illustration of the outcome of advance forecasts (1 to 3 or 4 days ahead) and, for comparison, the outcome of a type of "blind" forecast. For the latter the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

The radio propagation quality figures are prepared from radio traffic data reported to CRPL by American Telephone and Telegraph Company, Mackay Radio and Telegraph Company, ECA Communications, Inc., Marconi Company, British Admiralty Signal and Radar Establishment, and the following agencies of the U. S. Government:--Coast Guard, Navy, Army Signal Corps, and State Department. The method of calculation, summarized below, is similar to that described in a 1946 report, IRPL-R31, now out of print. Beginning with recalculated figures for January 1952, only reports of radio transmission on North Atlantic paths closely approximating New York-London are included in the estimation of quality. Observations of selected ionospheric characteristics, even though strongly correlated with radio transmission quality, and traffic reports for paths such as New York-Stockholm or New York-Tangier, previously included in the quality-figure determination with low weight, have been left out of the present calculations inasmuch as a sufficient number of homogeneous reports are now available.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the quality scale of the original reports. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year,

with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. The 6-hourly quality figures are (subjectively) weighted means of the reports received for that period. These 6-hourly quality figures replace, beginning January 1953, the half-daily quality figures which formerly appeared in this table.

These quality figures are, in effect, a consensus of reported radio propagation conditions in the North Atlantic area. The reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality because of multipath, interference, etc. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

Note. The North Pacific quality figures, which were published through October 1951, have been temporarily discontinued. Since the establishment of the North Pacific Radio Warning Service at Anchorage, Alaska, a larger number of reports are being received than were previously available in Washington. The preparation of the quality figures will be resumed when sufficient data have been accumulated for determination of conversion tables for these new reports.

## OBSERVATIONS OF THE SOLAR CORONA

Tables 64 through 66 give the observations of the solar corona during October 1953, obtained at Climax, Colorado, by the High Altitude Observatory of Harvard University and the University of Colorado. Tables 67 through 69 list the coronal observations obtained at Sacramento Peak, New Mexico, during October 1953, derived by Harvard College Observatory as a part of its performance of a research contract with the Upper Air Research Observatory, Geophysical Research Directorate, Air Force Cambridge Research Center. The data are listed separately for east and west limbs at 5-degree intervals of position angle north and south of the Solar Equator at the limb. The time of observation is given to the nearest tenth of a day, GCT.

Table 64 gives the intensities of the green (5303A) line of the emission spectrum of the solar corona; table 65 gives similarly the intensities of the first red (6374A) coronal line; and table 66, the intensities of the second red (6702A) coronal line; all observed at Climax in October 1953.

Table 67 gives the intensities of the green (5303 Å) coronal line; table 68, the intensities of the first red (6374 Å) coronal line; and table 69, the intensities of the second red (6702 Å) coronal line; all observed at Sacramento Peak in October 1953.

The following symbols are used in tables 64 through 69: a, observation of low weight; -, corona not visible; and X, position angle not included in plate estimates.

## SUDDEN IONOSPHERE DISTURBANCES

Tables 70, 71, and 72 list respectively the sudden ionosphere disturbances observed at Ft. Belvoir, Virginia, October 1953; in England, October 1953; and in the Netherlands, July and November 1952 and March, May, and August 1953.

## OBSERVATIONS OF SOLAR FLARES

Table 73 gives the preliminary record of solar flares reported to the CRPL. These reports are communicated on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete records are published later in the Quarterly Bulletin of Solar Activity, I.A.U., in various observatory publications, and elsewhere. The present listing serves to identify and roughly describe the phenomena observed. Details should be sought from the reporting observatory.

Reporting directly to the CRPL are the following observatories; Mt. Wilson, McMath-Hulbert, U. S. Naval, Wendelstein, Kanzel and High Altitude at Sacramento Peak, New Mexico. The remainder report to Meudon (Paris) and the data are taken from the Paris-URSIGram broadcast, monitored fairly regularly by the CRPL. The data on solar flares reported from Sacramento Peak, New Mexico, communicated by the High Altitude Observatory at Boulder, Colorado, are provided by Harvard University as the result of work undertaken on an Air Materiel Command Research and Development Contract administered by the Air Force Cambridge Research Laboratories.

The table lists for each flare the reporting observatory, date, times of beginning and ending of observation, duration (when known), total area (corrected for foreshortening), and heliographic coordinates. For the maximum phase of the flare is given the time, intensity, area relative to the total area, and the importance. The column "SID observed" is to indicate when a sudden ionosphere disturbance, noted elsewhere in these reports, occurred at the time of a flare. Times are in Universal Time (GCT).

## INDICES OF GEOMAGNETIC ACTIVITY

Table 74 lists various indices of geomagnetic activity based on data from magnetic observatories widely distributed throughout the world. The indices are: (1) preliminary international character-figures, C; (2) geomagnetic planetary three-hour-range indices, K<sub>p</sub>; (3) magnetically selected quiet and disturbed days.

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm). The magnetically quiet and disturbed days are selected by the international scheme outlined on pages 219-227 in the December 1943 issue of Terrestrial Magnetism and Atmospheric Electricity. The details of the currently used method follow. For each day of a month, its geomagnetic activity is assigned by weighting equally the following four criteria: (1) C; (2) the sum of the eight K<sub>p</sub>'s; (3) the greatest K<sub>p</sub>; and (4) the sum of the squares of the eight K<sub>p</sub>'s.

K<sub>p</sub> is the mean standardized K-index from 11 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g., 5 is  $4\frac{2}{3}$ , 5<sub>o</sub> is  $5\frac{0}{3}$ , and 5<sub>+</sub> is  $5\frac{1}{3}$ . This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of K<sub>p</sub> has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948," published in Washington, D. C., 1949, by the Association of Terrestrial Magnetism and Electricity, International Union of Geodesy and Geophysics. K<sub>p</sub> is available from 1937 to date as noted in F108.

The Committee on Characterization of Magnetic Disturbance, ATME, IUGG, has kindly supplied this table. The Meteorological Office, De Bilt, Holland, collects the data and compiles C and selected days. The Chairman of the Committee computes the planetary index. Current tables are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

### ERRATA

1. CEPL-F 110, p. 50, fig. 8: h'F2 at 02 should read "<320."
2. CRPL-F 110, p. 52, fig. 13: fcJ2 at 05 should read "<3.3."

## TABLES OF IONOSPHERIC DATA

Table 1

Time	Washington, D. C. (38.7°N, 77.1°W)							October 1953	
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	(270)	2.8					3.1		
01	(260)	2.7					3.1		
02	250	2.6					3.1		
03	260	2.6					3.1		
04	(250)	2.5					3.2		
05	(250)	2.4					3.3		
06	(250)	2.8					3.3		
07	240	4.3	230	---	(120)	(1.8)	1.8	3.5	
08	250	5.0	220	---	110	2.4	3.2	3.5	
09	260	5.6	210	3.8	110	2.6	2.9	3.5	
10	270	5.8	200	4.0	100	2.8	2.7	3.4	
11	280	6.2	200	4.1	100	2.9		3.3	
12	290	6.2	200	4.2	100	3.0		3.3	
13	280	6.3	210	4.1	100	3.0		3.3	
14	270	6.2	210	4.0	100	2.9		3.3	
15	270	6.1	220	3.6	100	2.6	2.2	3.3	
16	250	6.0	230	---	110	2.3	2.3	3.4	
17	230	5.7	---	---	(120)	1.8	2.4	3.5	
18	220	4.7					1.3	3.4	
19	240	3.8					2.1	3.3	
20	(250)	3.3					2.5	3.2	
21	(270)	2.8					3.1		
22	(270)	2.7					3.0		
23	(220)	2.7					3.1		

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3

Time	Fairbanks, Alaska (64.9°N, 147.8°W)							September 1953	
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	(340)	---					5.3		
01	(360)	(2.4)					4.8	(3.2)	
02	(380)	(2.1)					5.0	(3.2)	
03	(360)	(2.2)					6.0	(2.8)	
04	(350)	---					5.6	---	
05	350	2.7					4.2	(3.3)	
06	260	3.1	250	3.2	120	2.0	3.4	3.3	
07	320	3.4	220	3.2	110	2.1		3.3	
08	380	3.6	220	3.3	110	2.3	2.4	2.8	
09	350	3.8	210	3.6	110	2.5		2.8	
10	380	4.0	210	3.6	110	2.6		3.0	
11	370	4.2	210	3.7	110	2.6		3.0	
12	350	4.2	220	3.7	110	2.5		3.2	
13	360	4.2	220	3.7	120	2.4		3.2	
14	350	4.2	220	3.7	120	2.3		3.3	
15	320	4.4	220	3.6	120	2.2		3.4	
16	300	4.0	230	---	120	2.0		3.5	
17	260	3.8	240	---	120	1.9		3.4	
18	250	3.8	---	---	---	2.0		3.5	
19	260	3.5						3.4	
20	260	3.1					3.5	3.3	
21	270	2.9					4.8	3.4	
22	300	(2.4)					4.0	(3.2)	
23	300	---					4.8	---	

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 5

Time	Nuuk, Greenland (61.2°N, 45.4°W)							September 1953	
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	(330)	(3.0)					5.6	(2.8)	
01	---	(3.4)					4.8	---	
02	---	---					5.0	---	
03	---	(3.4)					4.8	---	
04	---	(3.4)					5.3	---	
05	(340)	(2.5)					5.2	(3.1)	
06	(270)	(3.4)					4.7	(3.4)	
07	(240)	3.9	---	---	(2.2)	(3.5)		(3.4)	
08	260	4.2	230	---	100	(2.3)	2.6	3.8	
09	310	4.4	220	3.8	100	(2.6)		3.3	
10	330	(4.7)	200	3.8	100	2.7		3.2	
11	390	4.4	220	3.9	100	2.8		3.0	
12	380	4.6	210	3.9	100	2.8		3.0	
13	360	(4.8)	200	3.9	100	2.8		3.1	
14	330	4.9	220	3.8	100	2.6	3.0	3.1	
15	350	4.8	230	3.8	100	2.4	3.2	3.0	
16	(340)	(4.3)	230	(3.6)	100	2.2	3.8	(3.0)	
17	300	(4.1)	230	---	110	2.0	4.2	(3.1)	
18	300	(3.8)					4.6	(3.1)	
19	(300)	(3.6)					5.2	(3.0)	
20	300	(3.6)					7.1	(2.9)	
21	(280)	(3.1)					6.0	(3.0)	
22	(280)	(3.2)					6.2	(2.9)	
23	(320)	(3.0)					6.6	(2.8)	

Time: 45.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

## TABLES OF IONOSPHERIC DATA

Table 1

Time	Washington, D. C. (38.7°N, 77.1°W)							October 1953	
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	(270)	2.8					3.1		
01	(260)	2.7					3.1		
02	250	2.6					3.1		
03	260	2.6					3.1		
04	(250)	2.5					3.2		
05	(250)	2.4					3.3		
06	(250)	2.8					3.3		
07	240	4.3	230	---	(120)	(1.8)	1.8	3.5	
08	250	5.0	220	---	110	2.4	3.2	3.5	
09	260	5.6	210	3.8	110	2.6	2.9	3.5	
10	270	5.8	200	4.0	100	2.8	2.7	3.4	
11	280	6.2	200	4.1	100	2.9		3.3	
12	290	6.2	200	4.2	100	3.0		3.3	
13	280	6.3	210	4.1	100	3.0		3.3	
14	270	6.2	210	4.0	100	2.9		3.3	
15	270	6.1	220	3.6	100	2.6	2.2	3.3	
16	250	6.0	230	---	110	2.3	2.3	3.4	
17	230	5.7	---	---	(120)	1.8	2.4	3.5	
18	220	4.7					1.3	3.4	
19	240	3.8					2.1	3.3	
20	260	3.8					2.1	3.3	
21	270	3.3					2.5	3.2	
22	270	2.8					3.1	3.0	
23	270	2.7					3.0	(3.0)	

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3

Time	Fairbanks, Alaska (64.9°N, 147.8°W)							September 1953	
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	(340)	---					5.3		
01	(360)	(2.4)					4.8	(3.2)	
02	(380)	(2.1)					5.0	(3.2)	
03	(360)	(2.2)					6.0	(2.8)	
04	(350)	---					5.6	---	
05	350	2.7					4.2	(3.3)	
06	260	3.1	250	3.2	120	2.0	3.4	3.3	
07	320	3.4	220	3.2	110	2.1		3.3	
08	380	3.6	220	3.3	110	2.3	2.4	2.8	
09	350	3.8	210	3.6	110	2.5		2.8	
10	380	4.0	210	3.6	110	2.6		3.0	
11	370	4.2	210	3.7	110	2.5		3.2	
12	350	4.2	220	3.7	110	2.5		3.2	
13	360	4.2	220	3.9	100	2.8		3.1	
14	330	4.9	220	3.8	100	2.6	3.0	3.1	
15	350	4.8	230	3.8	100	2.4	3.2	3.0	
16	(340)	(4.3)	230	(3.6)	100	2.2	3.8	(3.0)	
17	300	(4.1)	230	---	110	2.0	4.2	(3.1)	
18	300	(3.8)					4.6	(3.1)	
19	(300)	(3.6)					5.2	(3.0)	
20	300	(3.6)					7.1	(2.9)	
21	(280)	(3.1)					6.0	(3.0)	
22	(280)	(3.2)					6.2	(2.9)	
23	(320)	(3.0)					6.6	(2.8)	

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3

Time	Fairbanks, Alaska (64.9°N, 147.8°W)							September 1953	
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	</th

Table 7

Time	September 1953							
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	2.1						3.2	2.7
01	2.0						2.5	2.7
02	2.0						2.9	2.7
03	2.0						3.2	(2.7)
04	1.9						2.7	2.7
05	2.2	—	—	—	—	—	2.4	2.9
06	2.6	3.2	235	2.8	—	—	2.1	3.1
07	(300)	3.7	230	(3.2)	120	2.0	2.3	3.2
08	2.6	4.2	225	3.4	115	2.2	2.4	3.1
09	340	4.6	215	3.7	110	2.4	2.4	3.0
10	315	4.8	215	3.8	110	2.6	2.7	3.2
11	320	4.8	210	4.0	110	2.6	2.8	3.1
12	320	4.9	210	4.0	115	2.7	3.0	3.1
13	316	4.9	210	3.9	116	2.6	2.4	3.1
14	296	4.9	215	3.8	116	2.6	2.4	3.2
15	275	4.8	220	3.6	115	2.4	1.7	3.2
16	250	4.7	230	(3.3)	115	2.1		3.2
17	260	4.7	240	(3.0)	120	1.8	2.3	3.1
18	240	4.7	255	—	—	—	2.4	3.1
19	245	4.6	—	—	—	—	2.7	3.0
20	255	4.1	—	—	—	—	2.3	3.0
21	260	3.4	—	—	—	—	2.2	3.0
22	300	2.6	—	—	—	—	2.2	2.9
23	400	2.2	—	—	—	—	2.3	2.8

Time: 15.0°W.

Sweep: 1.4 Mc to 17.0 Mc in 6 minutes, automatic operation.

Table 9

Time	September 1953							
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	3.4						3.1
01	290	3.2						3.0
02	280	3.2						3.0
03	270	3.2						3.0
04	280	3.1						3.0
05	280	3.0						3.1
06	240	3.6	—	—	120	1.7		3.2
07	260	5.0	220	3.6	110	2.1	3.0	3.3
08	230	6.4	200	3.9	110	2.6	3.1	3.3
09	200	5.8	200	4.1	110	2.8	3.6	3.2
10	300	6.9	190	4.3	110	2.9	3.5	3.1
11	310	6.1	190	4.4	110	3.1	3.8	3.1
12	320	6.6	200	4.4	110	3.2	3.0	3.0
13	310	6.8	200	4.4	110	3.1	3.2	3.0
14	300	6.6	200	4.3	110	3.1	3.2	3.1
15	280	6.5	220	4.2	100	3.0	2.7	3.2
16	270	6.2	220	3.9	100	2.6	2.4	3.3
17	250	6.0	230	—	110	2.1	3.0	3.3
18	220	5.6	—	—	—	—	2.3	3.4
19	230	4.8	—	—	—	—		3.2
20	250	4.0	—	—	—	—		3.1
21	270	3.6	—	—	—	—		3.1
22	270	3.7	—	—	—	—		3.0
23	280	3.6	—	—	—	—		3.0

Time: 105.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 11

Time	September 1953							
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	3.6					3.2	2.9
01	280	3.8					2.6	3.1
02	260	3.2					2.4	3.2
03	240	3.2					2.4	3.3
04	250	2.4					1.8	3.2
05	270	2.2					1.8	3.0
06	280	2.8					2.1	3.0
07	260	4.9	240	—	110	1.9	3.9	3.3
08	280	5.6	220	4.0	110	2.5	5.0	3.2
09	320	6.2	220	4.3	110	2.8	5.2	3.0
10	380	6.6	210	4.6	110	3.1	6.4	2.7
11	330	7.5	210	4.5	110	3.2	6.2	2.7
12	360	8.5	220	4.5	110	3.4	6.2	2.7
13	340	9.6	220	4.6	110	3.4	5.5	2.9
14	320	9.9	220	4.5	110	3.3	4.9	3.0
15	310	9.4	220	4.3	110	3.1	5.2	3.0
16	290	9.9	220	4.2	110	2.8	4.4	3.2
17	280	10.2	230	3.7	110	2.3	4.8	3.4
18	230	8.3	—	—	—	—	4.0	3.5
19	220	6.6	—	—	—	—	3.4	3.4
20	220	4.4	—	—	—	—	3.0	3.2
21	260	3.8	—	—	—	—	2.8	3.0
22	300	3.3	—	—	—	—	3.2	2.8
23	300	3.4	—	—	—	—	3.6	3.0

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 12

San Francisco, California (37.4°N, 122.3°W)

Time	September 1953							
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	(280)	(3.1)						2.5
01	(270)	(3.0)						2.2
02	(280)	(3.0)						2.1
03	(280)	(2.8)						2.2
04	(280)	(2.8)						2.1
05	(280)	(2.8)						2.7
06	270	(3.2)						2.8
07	300	(4.0)	220	3.4	110	(2.0)	3.3	(3.2)
08	340	4.6	210	(3.7)	110	(2.4)	3.9	3.1
09	340	4.9	200	(4.0)	110	(2.6)	4.1	3.1
10	340	5.2	190	(4.2)	100	(2.9)	3.6	3.1
11	340	5.3	200	4.2	100	(3.0)	4.0	3.1
12	330	5.5	200	4.2	100	(3.1)	4.0	3.0
13	330	5.6	210	(4.2)	100	(3.1)	4.0	3.0
14	320	5.6	210	4.1	110	(3.0)	3.6	3.2
15	300	5.2	220	4.0	110	(2.8)	3.4	3.2
16	280	5.3	230	(3.8)	110	(2.5)	2.0	3.2
17	250	6.4	230	—	120	2.0	3.2	3.4
18	240	4.9	250	—	—	—	2.7	3.4
19	230	(4.3)	—	—	—	—	2.4	3.2
20	(280)	4.0	—	—	—	—	2.6	3.2
21	(280)	(3.7)	—	—	—	—	3.8	(3.1)
22	(280)	(3.3)	—	—	—	—	3.2	(3.1)
23	(280)	(3.2)	—	—	—	—	3.1	(3.1)

Time: 120.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 10

Time	September 1953							
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	310	4.0						2.2
01	280	3.8						3.0
02	260	3.5						3.2
03	250	3.2						3.2
04	280	2.8						(3.1)
05	270	3.0						(3.1)
06	240	4.6	—	—	—	—		3.5
07	230	6.4	220	—	110	2.1	3.5	3.7
08	350	6.6	210	4.0	110	2.5	4.2	3.6
09	260	6.6	200	4.3	110	2.8	4.4	3.6
10	280	6.8	200	4.5	110	3.0	4.8	3.2
11	310	8.1	200	4.6	110	3.1	4.4	3.1
12	300	8.9	200	4.6	110	3.1	4.4	3.1
13	300	8.9	200	4.6	110	3.2	4.4	3.2
14	280	8.7	210	4.5	110	3.1	3.9	3.1
15	290	9.6	220	4.3	110	3.0	3.8	3.2
16	260	10.0	220	4.0	110	2.7	4.0	3.4
17	250	9.2	230	—	110	2.1	4.8	3.4
18	230	9.2	—	—	—	—		3.6
19	210	7.3	—	—	—	—		3.5
20	220	5.0	—	—	—	—		3.6
21	260	3.8	—	—	—	—		3.6
22	300	3.9	—	—	—	—		2.9
23	300	4.0	—	—	—	—		2.9

Time: 125.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 12

Time	September 1963							
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	3.8						2.1
01	260	3.9						2.8
02	230	3.8						3.3
03	220	3.6						2.3
04	220	3.0						2.0
05	240	2.6						3.1
06	220	4.8	210	—	110	1.8	3.7	
07	240	6.4	220	—	100	2.5	4.0	3.5
08	270	5.7	200	4.3	100	2.9	4.0	3.4
09	280	6.0						

Table 13

Time	September 1953						
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	280	3.8					3.0
01	270	3.8					3.2
02	260	3.2					3.3
03	250	3.0					3.4
04	260	3.2					2.2 (3.3)
05	250	2.3					3.4
06	260	2.7					3.2
07	240	6.8	250	—	120	2.1	2.6 3.4
08	250	7.1	210	—	110	2.6	3.3 3.1
09	200	7.8	200	—	110	2.9	4.0 2.9
10	330	8.6	190	4.4	110	3.1	4.4 2.6
11	340	8.4	200	4.4	(110)	(3.3)	4.1 2.6
12	350	8.6	200	4.6	110	(3.4)	4.5 2.6
13	340	8.8	210	4.6	110	(3.4)	4.3 2.6
14	340	9.1	210	4.4	110	(3.2)	4.7 2.8
15	320	9.9	220	4.3	110	3.1	5.2 2.9
16	300	10.7	220	4.1	110	2.8	6.6 3.1
17	270	11.0	230	—	110	(2.3)	4.0 3.2
18	260	10.6	240	—			3.4 3.2
19	240	10.0					2.6 3.2
20	240	8.4					2.2 3.3
21	240	6.6					3.2 3.0
22	260	5.4					3.2 2.9
23	280	4.6					3.0 2.9

Time: 150.0°E.

Sweep: 1.0 Mc to 25.0 Mc in 16 seconds.

Table 15

Time	August 1953						
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	250	3.7	—	—	—	1.5	3.0
01	250	3.4	—	—	140	1.4	3.0
02	260	3.3	—	—	120	1.5	3.1
03	270	3.3	230	2.8	120	1.6	3.0
04	280	3.6	250	3.1	120	1.7	3.0
05	300	3.6	240	3.0	120	1.8	2.9
06	430	2.4	240	3.2	110	2.1	2.8
07	460	3.6	230	3.3	110	2.3	2.6
08	530	3.6	220	3.4	110	2.4	2.6
09	460	4.0	220	3.6	110	2.6	(2.5)
10	400	4.0	220	3.5	100	2.6	2.8
11	430	4.0	210	3.6	100	2.6	(2.7)
12	460	4.0	210	3.6	100	2.6	(2.6)
13	600	4.0	210	3.6	100	2.6	(2.6)
14	500	3.9	210	3.6	100	2.6	(2.6)
15	420	4.0	210	3.4	110	2.6	(2.7)
16	400	4.0	220	3.4	110	2.4	2.9
17	400	4.0	220	3.3	110	2.3	2.8
18	390	4.0	220	3.3	110	2.1	2.8
19	310	3.8	230	3.2	120	2.0	3.0
20	260	3.8	240	3.0	120	1.9	3.0
21	260	3.9	250	—	120	1.7	3.0
22	260	3.9	240	—	110	1.6	3.1
23	260	3.7	240	—	130	1.4	3.0

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 16 seconds.

Table 17

Time	August 1953						
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	290	3.6				8.0	(2.9)
01	300	3.2				8.0	(3.1)
02	300	3.4				7.0	(3.0)
03	300	2.0	—	—	120	2.6	7.0 (3.0)
04	300	3.0	—	—	110	2.3	6.8 —
05	300	3.4	—	—	100	2.6	4.6 (2.9)
06	300	3.7	300	3.6	100	3.1	4.0 3.0
07	6	< 4.0	240	3.7	110	3.6	G
08	400	4.0	210	3.8	100	3.2	2.6
09	(500)	< 4.0	210	3.9	100	3.1	G
10	620	< 4.1	200	< 4.0	100	3.1	G
11	650	4.2	200	4.0	100	3.0	2.4
12	500	4.2	210	4.0	100	3.1	2.4
13	630	4.2	220	4.0	100	3.1	2.4
14	420	4.4	210	4.0	110	3.0	2.7
15	420	4.5	210	4.0	110	3.0	2.7
16	370	4.6	230	3.5	110	2.9	2.8
17	370	4.6	220	3.8	110	2.8	2.8
18	320	4.2	240	3.5	110	2.8	2.9
19	300	4.0	260	—	110	2.8	3.0
20	300	4.0	—	—	110	3.0	7.1 3.0
21	290	3.8			120	2.0	9.0 (3.0)
22	300	3.5	—	—		9.3	(3.0)
23	280	3.3	—	—		> 10.0	(3.0)

Time: 90.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 14

Time	Panama Canal Zone (9.4°N, 79.9°W)						
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	270	3.8					3.1
01	240	4.0					1.7 3.2
02	230	3.7					2.0 3.4
03	220	3.2					3.3
04	250	3.0					2.0 3.2
05	240	2.7					2.8 3.2
06	260	2.9					3.0 3.2
07	240	4.9	220	—	110	2.0	4.1 3.4
08	280	5.6	220	4.3	110	2.6	4.2 3.3
09	350	6.1	220	4.4	110	3.0	4.9 2.9
10	370	7.3	210	4.5	100	3.3	4.3 2.8
11	380	8.8	210	4.6	100	3.4	5.1 2.7
12	340	10.0	200	4.6	100	3.4	4.9 2.9
13	330	11.0	220	4.5	110	3.4	5.0 3.0
14	320	11.8	220	4.6	100	3.3	6.0 3.0
15	290	12.0	220	4.3	110	3.2	5.4 3.1
16	260	12.6	220	4.2	110	2.9	4.4 3.3
17	250	11.6	220	3.8	110	2.4	4.3 3.4
18	220	9.6					3.4 3.4
19	220	7.4					3.1 3.2
20	230	6.8					2.2 3.2
21	250	4.6					2.1 3.0
22	280	4.2					2.0 2.9
23	290	4.1					1.8 2.9

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 16

Time	Baker Lake, Canada (64.3°N, 96.0°W)						
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	220	3.5				E	3.1 3.0
01	230	3.2				E	6.0 3.0
02	250	3.0				E	4.0 3.0
03	250	3.0				—	4.0 3.0
04	250	2.8				—	2.8 3.0
05	260	3.2	220	3.0	100	1.9	4.1 3.0
06	310	3.4	210	3.0	100	2.2	5.2 (2.9)
07	450	3.7	200	3.4	100	2.4	2.9 2.8
08	500	(3.9)	200	3.6	100	2.7	4.0 (2.4)
09	< 3.9	200	3.8	100	2.8	3.8	G
10	560	(4.0)	240	3.9	100	3.2	3.7 (2.5)
11	G	(4.0)	200	3.9	100	3.1	3.7 G
12	G	< 4.0	200	3.9	100	3.1	3.7 G
13	410	4.1	200	3.9	100	3.1	2.4
14	410	4.3	200	3.8	100	3.0	2.7
15	380	4.6	200	3.8	100	3.0	2.8
16	390	4.4	200	3.8	100	2.9	2.8
17	360	4.4	220	3.7	100	2.7	5.8 2.9
18	350	4.2	220	3.6	100	2.3	4.0 2.9
19	250	4.2	220	3.0	100	2.0	6.0 3.0
20	400	4.3	200	4.0	100	3.0	3.5 2.9
21	420	4.4	200	4.0	100	3.0	3.3 2.8
22	400	4.4	200	4.0	100	3.0	3.0 2.8
23	380	3.3	200	4.0	100	3.0	3.7 2.9

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 18

Time	Fort Chimo, Canada (58.1°N, 68.8°W)						
	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	290	2.9					5.0 3.0
01	300	2.9					6.0 (3.0)
02	300	2.8					5.2 (3.0)
03	(290)	(2.8)					4.8 (3.0)
04	—	(3.0)					4.9 —
05	(300)	3.8	—	—	100	3.3	6.2 (3.0)
06	(300)	3.9	—	—	100	3.1	6.0 3.1
07	350	4.1	230	3.7	100	3.0	4.5 3.0
08	390	4.4	210	3.8	100	3.0	4.2 3.0
09	480	4.1	200	4.0	100	3.0	3.5 2.8
10	400	4.3	200	4.0	100	3.0	3.3 2.9
11	420	4.4	200	4.0	100	3.0	2.8 2.8
12	400	4.6	200	4.0	100	3.0	2.8 2.8
13	400	4.4	200	4.0	100	3.0	2.8 2.8
14	410	4.6	200	4.0	100	3.0	

Table 19

Prince Rupert, Canada (54.3°N, 130.3°W)								August 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	290	2.2				3.7	—		
01	310	2.0				3.8	—		
02	300	2.0				3.7	—		
03	320	2.0				4.0	—		
04	340	2.0				3.6	—		
05	280	2.4				3.4	—		
06	260	2.9	230	3.0	120	1.8	3.2	G	
07	G	< 3.3	210	3.3	110	2.2	2.5	G	
08	G	< 3.6	210	3.6	110	2.4	3.6	G	
09	G	< 3.8	200	3.8	100	2.6	4.8	G	
10	G	< 4.0	200	3.9	100	2.8	4.3	G	
11	480	4.2	200	4.0	100	3.0	4.0	G	
12	440	4.4	200	4.0	100	3.0	4.2	2.8	
13	440	4.4	200	4.0	100	3.0	3.6	2.9	
14	480	4.3	200	4.0	100	3.0	3.7	2.7	
15	490	4.3	210	4.0	100	3.0	3.8	2.5	
16	440	4.3	210	3.9	110	2.8	3.0		
17	400	4.1	220	3.7	110	2.6	3.0		
18	340	4.2	230	3.7	110	2.3	3.2	3.0	
19	250	4.1	240	—	120	1.9	2.4	3.2	
20	250	4.0				1.8	3.0	3.2	
21	250	4.0					4.0	3.2	
22	260	3.5					3.4	—	
23	260	2.8					3.8	—	

Time: 120.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 15 seconds.

Table 20

De Bilt, Holland (52.1°N, 5.2°E)								August 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	260	3.0							2.2
01	< 270	2.6							5.1
02	275	2.5							3.1
03	275	2.4							3.1
04	270	2.6							2.2
05	255	3.3	220	—	—	—	1.6	> 2.0	3.2
06	310	3.8	210	3.8	105	2.1	2.5	5.2	
07	350	4.3	200	3.7	100	2.4	2.8	3.1	
08	320	4.4	200	3.8	100	2.6	3.2	3.2	
09	330	4.8	200	4.0	100	2.8	3.6	3.2	
10	335	4.8	200	4.1	100	2.9	4.2	3.2	
11	340	4.8	200	4.2	100	3.0	3.6	3.2	
12	350	5.0	200	4.2	100	3.0	3.6	3.2	
13	340	4.8	200	4.2	100	3.0	3.0	3.2	
14	360	4.7	200	4.2	100	3.0	3.2	3.2	
15	340	4.7	200	4.0	100	2.9	2.9	3.2	
16	350	4.6	210	3.8	100	2.7	2.9	3.1	
17	295	4.8	220	3.6	100	2.8	3.1	3.2	
18	280	4.9	220	3.2	—	1.7	3.6	3.2	
19	250	3.4						3.0	3.2
20	220	5.4						2.9	3.3
21	220	4.8						3.3	
22	235	4.0						3.3	
23	230	3.4						3.2	

Time: 0.0°.

Sweep: 1.4 Mc to 11.2 Mc in 6 minutes, automatic operation.

Table 21

Winnipeg, Canada (49.9°N, 97.4°W)								August 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	300	2.4				3.2	(3.0)		
01	370	2.3				2.8	—		
02	370	2.5				3.8	—		
03	370	(2.5)				4.0	—		
04	360	(2.3)				3.1	—		
05	290	2.4	—	—		3.1	(3.0)		
06	G	3.1	230	3.1	120	1.8	2.9	(3.0)	
07	G	< 3.5	220	3.4	120	2.3	0		
08	G	< 3.7	210	3.6	110	2.6	G		
09	G	4.0	200	3.8	110	2.8	G		
10	480	4.2	200	4.0	110	3.0	4.2	(2.7)	
11	460	4.4	200	4.0	110	3.1	2.8		
12	450	4.4	200	4.0	110	3.1	2.8		
13	490	4.4	200	4.1	110	3.1	2.7		
14	480	4.3	210	4.0	110	3.1	2.7		
15	430	4.4	210	4.0	110	3.0	2.8		
16	410	4.4	210	3.9	110	2.9	2.9		
17	370	4.5	210	3.8	110	2.7	2.9		
18	330	4.5	220	3.5	120	2.3	3.1		
19	270	4.3	240	—	120	2.1	3.2		
20	260	4.2				3.0	3.2		
21	250	4.2					(3.2)		
22	260	3.2					3.2		
23	300	2.7				2.8	(2.9)		

Time: 90.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 22

St. John's, Newfoundland (47.6°E, 52.7°W)								August 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	320	2.0						2.0	2.9
01	310	2.2							2.9
02	300	2.0						3.0	2.9
03	300	2.0						3.0	3.0
04	280	2.0						3.0	3.0
05	260	3.3	230	3.3	120	E	1.8	3.4	
06	340	3.7	230	3.4	120	2.3	2.7	3.2	
07	410	4.2	230	3.7	120	2.6	3.0	2.9	
08	380	4.3	210	3.9	120	2.9	3.8	3.2	
09	370	4.4	200	4.1	110	3.1	5.7	3.1	
10	390	4.4	200	4.1	120	3.2	2.9	2.9	
11	430	4.6	200	4.2	120	3.3	3.1	2.8	
12	390	4.6	210	4.2	120	3.3	2.7	2.9	
13	460	4.5	210	4.2	110	3.2	3.0	2.7	
14	400	4.6	210	4.1	110	3.1	2.9	2.9	
15	350	4.5	220	4.0	110	2.9	2.9	2.9	
16	350	4.7	230	3.8	120	2.7	2.7	3.0	
17	330	4.9	240	3.4	120	2.3	3.0	3.0	
18	270	5.0	240	3.0	130	E	2.9	3.2	
19	250	5.2	200	3.0	100	2.0	3.4	3.1	
20	250	5.7	—	—	—	3.6	3.5		
21	215	5.5				3.5	3.5		
22	210	5.0					3.5		
23	230	4.0					3.5		

Time: 60.0°W.

Sweep: 0.8 Mc to 10.0 Mc in 18 seconds.

Table 23

Schwarzenburg, Switzerland (46.8°N, 7.3°E)								August 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	250	3.4				3.3			
01	290	3.2				3.3			
02	280	3.0				3.3			
03	300	2.8				3.2			
04	300	2.8				3.3			
05	300	2.7				3.3			
06	225	3.4	—	—	100	2.0	3.0	3.6	
07	280	3.4	200	3.4	100	2.2	3.6	3.6	
08	300	4.2	200	3.6	100	2.5	4.5	3.5	
09	300	4.8	200	3.9	100	2.7	4.0	3.4	
10	300	4.9	200	4.0	100	2.8	5.0	3.5	
11	300	4.8	200	4.0	100	3.0	4.2	3.4	
12	330	5.0	200	4.1	100	3.0	4.4	3.2	
13	330	5.0	200	4.1	100	3.0	4.8	3.2	
14	380	4.8	200	4.1	100	3.0	3.2	3.2	
15	315	5.0	200	4.0	100	2.9	3.9	3.2	
16	310	4.7	200	3.9	100	2.8	3.3	3.3	
17	300	4.7	200	3.8	100	2.6	3.4	3.4	
18	300	4.8	200	3.5	100	2.2	4.0	3.4	
19	270	5.2	200	3.0	100	2.0	4.1	3.4	
20	235	5.7	—	—	—	3.6	3.5		
21	215	5.5				3.5	3.5		
22	210	5.0					3.5		
23	230	4.0					3.5		

Time: 15.0°E.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

Table 24

Ottawa, Canada (45.4°N, 75.9°W)								August 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	





<tbl\_r cells="10" ix="5" maxcspan="

Table 25

Time	August 1953						
	h'F2	foF2	h'F1	foF1	h'X	foE	fEs (M3000)F2
00	300	3.2				3.2	3.0
01	300	3.2				3.5	3.1
02	300	3.0				3.1	
03	200	2.8				2.2	3.1
04	200	2.6				3.1	
05	300	2.5				3.1	
06	280	3.4	250	—	120	—	2.9
07	300	4.1	240	3.5	120	2.1	3.7
08	350	4.7	220	3.8	120	(2.5)	4.1
09	350	5.0	210	4.0	110	(2.8)	4.0
10	430	5.1	200	4.1	110	(3.1)	4.0
11	380	5.1	210	4.2	110	3.1	4.0
12	420	5.2	220	4.2	110	3.2	4.2
13	400	5.2	220	4.2	110	3.2	4.2
14	380	5.2	220	4.2	110	3.2	3.5
15	330	5.3	230	4.0	110	3.0	3.7
16	350	5.3	220	3.9	120	2.8	4.2
17	320	5.4	230	3.6	120	3.4	3.8
18	280	5.6	240	—	120	(2.1)	3.6
19	260	5.1				3.1	3.2
20	250	4.8				2.9	3.2
21	270	4.1				3.1	3.1
22	300	2.4				3.0	3.0
23	200	3.3				2.4	3.0

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

Table 27

Time	August 1953						
	h'F2	foF2	h'F1	foF1	h'X	foE	fEs (M3000)F2
00	820	2.6					2.8
01	300	2.6					2.8
02	280	2.6					2.9
03	250	2.9					3.3
04	210	2.5					3.4
05	260	1.8					3.2
06	270	2.0					3.1
07	230	4.0					2.5
08	240	5.0	220	—	—	—	2.8
09	260	5.3	220	—	120	2.8	3.4
10	270	5.9	220	—	110	2.9	3.8
11	270	5.6	200	3.8	110	3.0	3.9
12	270	6.5	200	4.1	110	3.1	3.9
13	260	7.2	200	4.1	110	3.1	3.9
14	250	7.0	200	4.1	110	2.9	3.8
15	240	6.6	210	3.6	110	2.8	3.3
16	230	5.6	210	—	—	—	2.7
17	220	5.1					3.5
18	220	4.8					3.3
19	240	3.6					3.2
20	270	3.6					3.0
21	260	3.1					3.1
22	270	3.0					3.1
23	300	2.7					2.9

Time: 60.0°W.

Sweep: 1.0 Mc to 26.0 Mc in 30 seconds.

Table 29

Time	July 1953						
	h'F2	foF2	h'F1	foF1	h'X	foE	fEs (M3000)F2
00	280	3.5	220	2.7	110	1.8	3.0
01	260	3.6	220	2.7	110	1.7	3.0
02	270	3.5	220	2.9	110	1.8	
03	310	3.5	220	3.0	110	1.9	3.0
04	400	3.5	210	3.1	100	2.0	3.0
05	410	3.6	210	3.2	100	2.1	2.7
06	430	3.4	210	3.3	100	2.3	2.8
07	6	< 3.5	200	3.4	100	2.4	0
08	0	< 3.8	210	3.4	100	2.5	0
09	0	3.7	200	3.6	100	2.6	0
10	0	< 3.7	200	3.6	100	2.7	0
11	0	< 3.7	200	3.7	100	2.8	0
12	0	< 3.8	200	3.7	100	2.7	0
13	0	< 4.0	200	3.7	100	2.7	0
14	0	< 3.7	200	3.7	100	2.7	0
15	440	4.2	200	3.7	100	2.6	2.7
16	0	(3.9)	200	3.6	100	3.5	0
17	440	4.0	200	3.5	100	2.5	2.8
18	440	3.8	200	3.4	100	2.3	2.7
19	380	3.9	200	3.3	100	2.2	2.8
20	360	3.8	210	3.3	100	2.1	2.9
21	320	3.8	220	3.1	110	2.0	3.0
22	310	3.8	220	3.0	110	1.9	3.0
23	270	3.8	220	3.2	110	1.8	3.1

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 26

Time	August 1953						
	h'F2	foF2	h'F1	foF1	h'X	foE	fEs (M2000)F2
00	220	3.7					2.4
01	220	3.6					2.8
02	(240)	(2.9)					3.1
03	—	(2.3)					2.9
04	(250)	2.4					3.0
05	250	3.3					2.4
06	250	5.7	240	—	125	2.2	3.3
07	270	8.6	230	4.0	120	2.8	3.2
08	280	7.6	225	4.2	120	3.0	2.3
09	290	8.0	220	4.4	115	3.2	4.4
10	300	8.6	210	4.4	115	3.3	5.0
11	300	8.9	200	4.4	115	3.5	5.1
12	310	8.9	200	4.4	115	3.4	4.3
13	320	9.9	200	4.2	115	3.2	3.8
14	300	9.8	200	4.3	120	3.0	2.0
15	285	9.8	225	4.0	120	2.8	3.5
16	280	9.5	260	—	120	2.2	3.5
17	240	9.2					3.2
18	240	9.0					3.0
19	220	8.0					2.6
20	210	6.5					2.4
21	210	3.9					2.2
22	215	3.6					2.4
23	230	3.4					2.3

Time: 0.0°W.

Sweep: 1.0 Mc to 16.0 Mc in 7 seconds.

Table 28

Time	August 1953						
	h'F2	foF2	h'F1	foF1	h'X	foE	fEs (M3000)F2
00	310	2.5					(3.0)
01	310	2.6					(3.0)
02	310	2.8					3.0
03	300	2.7					(3.0)
04	300	2.7					(3.1)
05	300	2.8					(3.2)
06	270	2.5					(3.3)
07	250	3.0					(3.4)
08	230	3.3					(3.5)
09	210	4.3					2.5
10	210	4.6					3.0
11	230	4.6					(3.6)
12	210	4.6					(3.6)
13	210	5.2					(3.6)
14	210	4.6					(3.5)
15	220	4.6					(3.5)
16	220	4.3					(3.6)
17	220	4.2					(3.5)
18	230	3.6					(3.5)
19	270	3.0					(3.3)
20	300	2.6					(3.2)
21	300	2.6					(3.1)
22	310	2.4					(3.1)
23	310	2.5					(3.1)

Time: 60.0°W.

Sweep: 1.5 Mc to 16.0 Mc in 15 minutes, manual operation.

Table 30

Time	July 1953						
	h'F2	foF2	h'F1	foF1	h'X	foE	fEs (M3000)F2
00	240	3.4				1.2	4.0
01	240	3.3				1.4	3.0
02	240	3.2				1.5	4.0
03	250	3.2				1.7	2.0
04	250	3.3	230	2.8	100	1.8	3.8
05	400	3.4	220	3.0	100	1.9	3.9
06	(530)	3.6	200	3.3	100	2.2	2.5
07	(480)	3.8	200	3.4	100	2.5	3.9
08	520	4.0	200	3.7	100	2.7	4.3
09	530	< 4.1	210	3.8	100	3.0	5.1
10	0	< 4.0	200	3.9	100	3.0	3.4
11	600	< 4.0	220	4.0	100	3.3	3.4
12	590	4.1	200	3.9	100	3.1	
13	500	4.2	210	3.9	100	3.1	(2.7)
14	440	4.4	200	3.8	100	3.0	2.7
15	390	4.5	200	3.8	100	3.0	2.9
16	400	4.4	200	3.9	100	2.9	6.0
17	400	4.3	200	3.8	100	2.9	4.0
18	380	4.2	210	3.5	100	2.6	2.8
19	320	4.1	200	3.3	100	2.2	6.0
20	300	4.0	220	3.0	110	2.0	6.0
21	240	3.7	—	—	110	1.8	6.0
22	260	3.7	—	—	110	1.7	4.0
23	250	3.6	—	—	110	1.6	4.4

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 31

Fort Chimo, Canada (58.1°N, 68.5°W)								July 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	3.0	—	—	5.0	5.0	3.0	
01	260	3.0	—	—	5.8	3.2	3.0	
02	290	3.0	—	100	2.8	6.0	(3.1)	
03	300	3.0	—	100	2.8	5.0	3.0	
04	300	3.4	—	100	3.0	5.6	3.0	
05	290	3.8	—	—	100	3.2	5.0	3.2
06	430	< 3.9	220	3.8	100	3.0	5.0	3.0
07	G	< 3.8	210	3.8	100	2.9	4.4	G
08	G	< 4.0	200	3.8	100	3.0	4.2	G
09	G	< 4.0	200	3.9	100	3.0	4.0	G
10	500	4.2	200	4.0	100	3.0	—	2.6
11	G	< 4.1	200	4.0	100	3.0	—	—
12	450	4.3	200	4.0	100	3.0	—	2.8
13	420	4.5	200	4.0	100	3.0	—	2.8
14	420	4.6	200	4.0	100	3.0	—	2.8
15	400	4.5	210	3.9	100	3.0	—	2.8
16	400	4.4	220	3.8	100	3.0	5.0	2.8
17	380	4.3	220	3.7	100	2.8	5.0	3.0
18	390	4.0	220	3.5	100	2.8	4.5	2.9
19	300	3.9	250	—	100	2.5	4.8	3.0
20	260	3.8	—	100	2.2	7.0	3.0	
21	250	3.6	—	100	2.5	7.0	3.0	
22	280	3.2	—	—	—	7.0	3.0	
23	280	3.2	—	—	—	6.5	3.0	

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 33

Winnipeg, Canada (49.9°N, 97.4°W)								July 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	320	2.6	—	—	—	3.8	(3.0)	
01	320	2.8	—	—	—	5.0	—	
02	330	(2.8)	—	—	—	5.0	—	
03	330	2.8	—	—	—	5.0	—	
04	320	2.7	—	—	—	4.5	—	
05	240	2.9	—	—	—	4.0	(3.2)	
06	G	< 3.2	220	3.2	120	2.0	3.5	G
07	G	< 3.5	210	3.4	120	2.4	—	G
08	G	< 3.7	200	3.7	110	2.6	—	G
09	G	< 3.8	210	3.8	110	2.9	—	G
10	G	< 4.0	200	3.9	110	3.0	—	G
11	510	< 4.2	200	4.0	110	3.0	—	G
12	480	4.2	200	4.0	110	3.1	—	2.6
13	600	4.2	200	4.0	110	3.1	—	G
14	465	4.3	200	4.0	110	3.0	—	2.6
15	430	4.3	200	4.0	110	3.0	—	2.8
16	400	4.3	210	3.9	110	2.9	—	2.9
17	370	4.6	210	3.8	110	2.7	—	3.0
18	340	4.6	220	3.6	120	2.4	—	3.0
19	300	4.4	220	3.3	130	2.1	2.8	3.0
20	280	4.5	240	—	—	—	3.2	
21	250	3.8	—	—	—	—	3.0	
22	270	2.2	—	—	—	—	(3.0)	
23	290	2.8	—	—	—	2.4	(3.0)	

Time: 90.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 35

Ottawa, Canada (45.4°N, 75.9°W)								July 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	2.4	—	—	—	3.0	3.0	
01	290	2.4	—	—	—	3.0	3.0	
02	300	2.0	—	—	—	3.3	3.0	
03	(290)	2.0	—	—	—	3.9	(3.0)	
04	300	2.0	—	—	—	2.8	(3.0)	
05	270	2.8	230	2.9	120	1.8	2.3	3.2
06	G	< 3.4	220	3.4	110	2.2	—	G
07	G	< 3.7	220	3.6	110	2.6	—	G
08	G	< 4.0	210	3.8	110	2.8	—	G
09	G	< 4.0	210	4.0	110	3.0	3.0	G
10	540	(4.2)	200	4.0	100	3.1	3.7	G
11	G	(4.1)	200	4.1	110	3.2	4.1	G
12	590	(4.3)	200	4.1	110	3.3	4.0	G
13	G	(4.2)	200	4.0	110	3.3	3.2	G
14	560	(4.5)	210	4.0	110	3.2	3.0	G
15	420	4.6	210	4.0	110	3.1	2.7	
16	410	4.7	220	3.9	110	2.9	2.8	2.8
17	360	4.8	220	3.8	110	2.7	—	2.9
18	510	4.8	230	3.5	110	2.3	3.6	3.1
19	270	4.9	230	—	130	1.9	2.5	3.0
20	250	4.8	—	—	—	—	3.2	
21	250	4.1	—	—	—	—	3.1	
22	260	3.8	—	—	—	—	3.0	
23	280	2.7	—	—	—	2.8	3.0	

Time: 75.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 15 seconds.

Table 32

Prince Rupert, Canada (54.3°N, 130.5°W)								July 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	2.0	—	—	—	—	—	2.2
01	320	2.2	—	—	—	—	—	3.0
02	300	2.0	—	—	—	—	—	3.7
03	300	2.0	—	—	—	—	—	4.0
04	290	2.2	—	—	—	—	—	4.0
05	230	2.8	230	2.8	120	1.7	2.9	G
06	430	3.2	220	3.0	110	2.2	3.1	2.7
07	G	3.5	210	3.4	110	2.2	—	G
08	G	< 3.8	200	3.6	100	2.5	3.1	G
09	G	< 3.9	200	3.8	100	2.8	4.0	G
10	G	< 4.0	200	3.9	100	2.9	4.4	G
11	G	4.1	200	4.0	100	3.0	4.4	G
12	490	4.3	200	4.0	100	3.0	3.7	G
13	470	4.4	200	4.0	100	3.0	4.7	2.7
14	460	4.4	200	4.0	100	3.1	4.5	2.7
15	450	4.2	200	4.0	100	3.0	—	G
16	450	4.2	210	4.0	100	2.9	5.2	2.8
17	420	4.2	210	3.8	100	2.7	3.4	2.9
18	390	4.2	220	3.7	110	2.4	—	3.0
19	340	4.2	230	3.4	110	2.1	3.2	3.0
20	230	4.1	240	—	120	1.8	3.0	3.1
21	250	4.0	—	—	—	—	2.5	(3.1)
22	260	3.8	—	—	—	—	2.6	—
23	280	3.0	—	—	—	—	2.9	—

Time: 120.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 15 seconds.

Table 36

Calcutta, India (22.6°N, 88.4°E)								July 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	4.7	—	—	—	—	—	2.8
01	270	4.7	—	—	—	—	—	3.1
02	(270)	(4.4)	—	—	—	—	—	3.0
03	270	4.1	—	—	—	—	—	3.4
04	265	4.0	—	—	—	—	—	3.4
05	250	(4.2)	—	—	—	—	—	3.4
06	240	4.7	—	—	—	—	—	2.9
07	225	5.8	—	—	—	—	—	3.0
08	240	6.4	—	—	—	—	—	2.7
09	240	6.5	—	—	—	—	—	3.8
10	240	7.2	—	—	—	—	—	3.5
11	250	8.0	—	—	—	—	—	3.7
12	255	9.2	—	—	—	—	—	3.5
13	(270)	9.4	—	—	—	—	—	6.4
14	270	10.5	—	—	—	—	—	4.8
15	270	10.5	—	—	—	—	—	3.6
16	270	10.5	—	—	—	—	—	4.3
17	270	9.9	—	—	—	—	—	3.8
18	240	10.6	—	—	—	—	—	4.1
19	240	9.2	—	—	—	—	—	3.5
20	240	8.8	—	—	—	—	—	3.6
21	240	5.8	—	—	—	—	—	3.2
22	240	6.4	—	—	—	—	—	3.0
23	270	6.3	—	—	—	—	—	2.8

Time: 90.0°E.

Sweep: 0.5 Mc to 18.0 Mc in 10 minutes, semi-automatic operation.

Baguio, P.I. (16.4°N, 120.6°E)

Table 37

July 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	320	3.2					2.8	2.8
01	290	3.4					3.0	(2.9)
02	280	3.1					2.8	(2.9)
03	200	2.0					2.8	(3.0)
04	270	(2.8)						(3.0)
05	250	2.7					3.2	3.2
06	250	4.2					3.8	3.3
07	250	5.3	220	—	110	2.3	5.5	3.2
08	320	5.4	220	4.0	110	2.8	6.1	3.0
09	370	5.7	210	4.0	110	(3.0)	7.0	2.8
10	420	6.2	200	4.2	110	3.2	7.2	2.6
11	440	6.7	200	4.2	110	3.4	6.4	2.4
12	430	7.4	210	4.2	110	3.4	6.9	2.5
13	440	7.6	200	4.1	110	3.8	6.6	2.5
14	430	7.8	200	4.1	110	3.2	6.4	2.5
15	420	7.9	210	4.0	110	3.0	5.8	2.6
16	380	8.4	220	4.0	110	(2.8)	4.8	2.7
17	220	8.4	220	—	110	2.3	4.4	2.9
18	250	8.8					4.7	3.0
19	240	7.9					4.0	3.1
20	260	6.1					3.0	3.1
21	280	8.0					3.1	2.9
22	300	3.9					1.8	2.8
23	220	3.8					2.3	(2.8)

Time: 120.0°E.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 39

June 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	310	3.6	250	—	—	—	6.4	3.1
01	310	3.7	240	—	—	—	5.0	3.1
02	310	3.7	230	—	—	—	5.6	3.1
03	300	3.6	220	3.2	120	—	6.0	3.1
04	280	3.7	230	3.2	110	—	4.4	3.0
05	350	3.8	220	3.4	110	2.0	4.9	2.9
06	440	3.8	230	3.6	100	—	4.7	2.7
07	440	4.1	220	3.7	100	—	5.3	2.8
08	430	4.1	240	3.7	100	—	4.8	2.7
09	410	4.2	220	3.8	100	2.6	4.7	2.8
10	500	4.0	220	3.8	100	2.7	4.8	2.6
11	480	4.1	220	3.8	100	2.8	4.4	2.6
12	480	4.2	210	3.9	100	2.9	3.8	2.6
13	500	4.1	220	3.9	100	2.8	3.5	2.6
14	440	4.2	220	3.8	100	2.8	3.5	2.6
15	450	4.3	210	3.8	100	2.6	2.6	2.7
16	410	4.4	220	3.8	100	2.5	2.5	2.8
17	390	4.4	220	3.7	100	2.4	2.8	2.9
18	370	4.3	230	3.6	110	2.2	3.0	3.0
19	350	4.2	230	3.5	110	2.2	4.0	3.0
20	320	3.9	240	3.3	110	2.0	3.9	3.1
21	330	3.9	250	3.3	110	—	4.4	3.1
22	340	3.7	240	—	—	—	4.5	3.0
23	340	3.8	230	—	—	—	4.9	3.1

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 41

June 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	4.7					5.2	2.8
01	280	4.7					4.3	2.8
02	280	4.5					4.4	3.0
03	270	4.2					3.8	3.0
04	260	4.0					3.5	3.0
05	260	4.8	250	3.0	120	1.8	3.5	3.1
06	300	4.9	250	3.5	110	2.4	4.3	3.2
07	300	8.2	230	3.8	110	2.7	5.2	3.1
08	320	5.6	240	4.0	110	2.9	6.2	3.2
09	360	5.2	230	4.2	110	3.1	6.5	3.0
10	330	5.3	220	4.3	110	3.2	6.5	2.9
11	360	5.4	210	4.2	110	3.2	6.2	3.0
12	400	5.2	230	4.2	110	3.3	6.2	2.9
13	430	5.0	230	4.2	110	3.2	5.8	2.8
14	370	5.3	240	4.1	110	3.0	5.6	2.9
15	350	5.5	240	4.0	110	2.9	5.5	3.0
16	330	5.7	240	3.9	110	2.7	4.8	3.0
17	310	5.8	240	3.6	110	2.5	4.8	3.0
18	300	5.9	250	3.2	120	2.0	5.0	3.0
19	270	5.6	—	—			5.7	3.1
20	250	5.2					4.4	3.1
21	270	5.6					4.5	3.0
22	280	5.0					4.6	2.9
23	260	4.9					4.8	2.9

Time: 125.0°E.

Sweep: 0.85 Mc to 22.0 Mc in 2 minutes.

Table 38

Watheroo, W. Australia (30.3°S, 118.9°E)

July 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	250	3.0						2.0
01	250	3.2						2.0
02	260	3.2						2.0
03	250	3.2						2.0
04	250	3.2						1.9
05	240	3.0						2.0
06	240	2.5						1.9
07	240	3.1						3.3
08	240	4.7	220	2.8	2.8		2.0	3.5
09	250	4.9	220	3.5	2.9		2.5	3.5
10	280	5.4	210	3.9	2.9		2.7	3.3
11	280	8.5	220	4.2	2.9		2.9	3.4
12	300	5.6	210	4.2	4.2		3.0	3.3
13	290	5.7	220	4.2	4.2		3.0	3.3
14	300	5.8	200	4.1	200	4.1	2.9	3.6
15	280	5.6	220	3.9	220	3.9	2.7	3.6
16	260	5.3	230	3.4	230	3.4	2.4	3.4
17	240	4.9	240	2.4	240	2.4	2.0	3.1
18	230	3.8	—	—	2.8			3.4
19	220	3.0	—	—	2.0			3.3
20	220	2.7	—	—	1.8			3.3
21	250	2.7	—	—	1.8			3.1
22	250	3.0	—	—	2.0			3.1
23	250	3.1	—	—	2.0			3.0

Time: 120.0°E.

Sweep: 1.0 Mc to 16.0 Mc in 2 minutes.

Table 42

June 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	4.6					4.5	2.9
01	280	5.2					4.8	(2.9)
02	280	4.4					4.5	2.9
03	260	4.4					4.2	3.0
04	270	4.0					4.2	3.0
05	250	4.1	240	—	120	1.7	3.2	3.2
06	300	5.0	240	3.6	120	2.3	4.7	3.2
07	300	5.4	250	3.9	110	2.7	5.0	3.1
08	300	5.9	230	4.0	110	3.0	6.0	3.2
09	230	5.8	240	4.2	110	3.2	7.0	3.1
10	370	5.1	250	4.3	110	3.2	6.5	2.9
11	380	5.5	220	4.4	110	3.2	6.6	2.8
12	410	5.6	240	4.4	110	3.3	6.6	2.8
13	370	5.8	240	4.2	110	3.2	6.8	2.8
14	360	8.0	230	4.2	110	3.2	6.7	2.9
15	340	6.5	250	4.2	110	3.0	6.0	2.9
16	320	6.6	250	3.9	110	2.7	5.0	2.9
17	310	6.5	220	3.7	110	2.4	5.5	3.0
18	300	6.6	240	3.3	120	1.9	5.3	3.0
19	260	7.0					4.8	3.1
20	260	6.3					4.5	3.0
21	300	5.2					4.6	2.9
22	290	4.8					4.5	2.9
23	290	4.8					5.1	2.9

Time: 125.0°E.

Sweep: 1.0 Mc to 17.2 Mc in 2 minutes.

Table 43

Yamagawa, Japan (31.2°N, 130.6°E)								
Time	b'F2	foF2	b'F1	foF1	b'E	foE	fEs	(M3000)F2
00	280	5.2					4.4	3.0
01	280	4.9					4.2	(3.0)
02	260	4.7					4.0	3.1
03	240	4.7					4.0	(3.2)
04	240	3.9					3.0	2.2
05	240	3.9					3.2	3.4
06	230	4.4	200	—	100	1.8	3.5	3.5
07	240	5.4	220	3.7	100	2.4	5.0	3.4
08	250	5.6	210	4.1	100	2.8	5.0	3.4
09	260	5.7	210	4.2	100	3.0	6.0	3.4
10	300	5.7	230	4.3	100	3.2	7.4	2.2
11	330	5.7	220	4.5	100	3.3	6.9	3.1
12	340	5.8	220	4.4	100	3.5	6.8	3.0
13	340	6.0	200	4.4	100	3.3	6.5	3.1
14	310	7.0	200	4.4	100	3.4	6.3	3.1
15	300	7.2	200	4.2	100	3.1	6.2	3.1
16	300	7.4	230	4.1	100	3.0	6.0	3.1
17	280	7.5	200	5.0	100	2.7	5.2	3.2
18	260	7.2	250	3.6	100	2.8	5.7	3.1
19	230	5.7	—	—	—	—	4.6	3.4
20	220	6.1	—	—	—	—	3.8	3.2
21	240	5.5	—	—	—	—	4.6	3.2
22	260	5.2	—	—	—	—	4.6	3.1
23	270	5.1	—	—	—	—	4.3	3.0

Time: 135.6°E.

Sweep: 0.8 Mc to 20.0 Mc in 15 minutes, manual operation.

Table 45

Hartonga I. (21.3°S, 159.8°W)								
Time	b'F2	foF2	b'F1	foF1	b'E	foE	fEs	(M3000)F2
00	270	3.2						2.9
01	300	3.1						2.9
02	300	3.0						3.0
03	250	3.1						3.0
04	230	3.0						3.2
05	250	2.7						3.0
06	260	2.6						3.0
07	250	4.2	—	1.8				3.4
08	250	6.8	200	2.9	120	3.2	3.2	3.5
09	260	6.3	200	4.9	110	2.6	3.8	3.4
10	270	6.4	200	4.2	110	2.8	4.0	3.5
11	270	6.3	210	4.3	110	3.1	4.2	3.4
12	270	6.2	210	4.3	110	3.1	4.2	3.4
13	270	6.4	200	4.2	110	3.0	4.2	3.4
14	270	6.2	210	4.2	110	2.0	4.1	3.5
15	270	6.0	200	4.0	110	2.8	4.0	3.3
16	260	6.4	240	3.6	110	2.5	3.6	3.3
17	250	6.1	—	2.2	—	1.9	3.5	3.4
18	220	5.8	—	—	—	—	3.0	3.3
19	220	4.6	—	—	—	—	2.9	3.3
20	< 260	3.8	—	—	—	—	2.6	3.1
21	230	3.2	—	—	—	—	2.4	3.1
22	250	3.4	—	—	—	—	3.0	3.1
23	280	3.1	—	—	—	—	3.0	3.1

Time: 157.5°W.

Sweep: 2.0 Mc to 16.0 Mc, manual operation.

Table 47

Hartonga I. (21.3°S, 159.8°W)								
Time	b'F2	foF2	b'F1	foF1	b'E	foE	fEs	May 1953
00	< 270	3.4						3.0
01	260	3.4						3.0
02	270	3.3						3.0
03	260	3.5						3.1
04	250	3.2						2.8
05	< 260	3.0						3.0
06	260	2.7						2.9
07	250	5.2	210	2.0	120	2.3		3.3
08	250	6.8	220	3.4	110	2.4	3.2	3.4
09	250	7.2	210	4.0	110	2.8	4.0	3.5
10	260	7.8	210	4.2	110	3.0	4.2	3.4
11	260	7.0	210	4.3	110	3.1	4.3	3.4
12	280	6.9	220	4.4	110	3.1	4.4	3.3
13	260	7.4	210	4.3	110	3.1	4.4	3.3
14	270	6.8	210	4.2	110	3.0	4.3	3.3
15	260	6.9	210	4.0	110	2.9	4.2	3.3
16	260	7.2	240	3.8	110	2.5	3.7	3.3
17	250	7.2	—	2.8	—	2.2	3.9	2.4
18	240	6.5	—	—	—	—	3.9	3.3
19	240	5.4	—	—	—	—	3.6	3.3
20	250	4.1	—	—	—	—	3.0	3.2
21	250	3.7	—	—	—	—	2.5	3.1
22	260	3.7	—	—	—	—	2.5	3.0
23	250	6.6	—	—	—	—	3.0	3.0

Time: 157.5°W.

Sweep: 2.0 Mc to 16.0 Mc, manual operation.

Calcutta, India (22.6°N, 80.2°E)

Time	b'F2	foF2	b'F1	foF1	b'E	foE	fEs	June 1953
00	(265)	(5.1)						(2.8)
01	(270)	(5.2)						
02	(240)	(4.7)						
03	240	4.6						
04	(210)	(4.3)						
05	(225)	(3.6)						
06	(240)	(4.8)						
07	220	5.8						
08	240	6.4						
09	240	6.4						
10	240	10.5						
11	(250)	8.2						
12	(240)	(10.0)						
13	(225)	(10.1)						
14	(225)	9.9						
15	260	10.2						
16	240	10.1						
17	240	10.5						
18	(225)	(10.0)						
19	(225)	(10.1)						
20	(225)	(8.8)						
21	(220)	(7.0)						
22	(255)	(5.2)						
23	270	5.3						

Time: 90.6°E.

Sweep: 6.5 Mc to 18.0 Mc in 10 minutes, semi-automatic operation.

Table 48

Christchurch, New Zealand (43.6°S, 172.7°E)								
Time	b'F2	foF2	b'F1	foF1	b'E	foE	fEs	May 1953
00	260	2.7						3.0
01	280	2.7						3.0
02	270	2.7						3.0
03	280	2.4						3.0
04	280	2.2						3.1
05	270	2.0						3.1
06	250	1.9						3.2
07	250	2.8						3.2
08	240	4.2	240	2.6		1.8	3.7	3.5
09	240	4.7	230	3.2		2.2	4.3	3.5
10	250	4.9	220	3.6		2.4	4.3	3.5
11	260	5.0	230	3.8		2.6	4.3	3.5
12	270	5.1	220	3.8		2.6	4.3	3.4
13	270	5.5	230	3.7		2.6	4.3	3.4
14	270	5.4	240	3.7		2.4	4.3	3.3
15	250	5.2	280	3.3		2.2	4.3	3.5
16	260	4.9	260	2.6	1.8	3.6	3.4	
17	240	4.6	—	—		—	2.7	3.3
18	240	3.7	—	—		—	2.1	3.1
19	250	3.2	—	—		—	—	3.1
20	260	3.1	—	—		—	—	3.1
21	270	2.7	—	—		—	—	3.1
22	270	2.8	—	—		—	—	3.0
23	270	2.6	—	—		—	—	3.0

Time: 172.5°E.

Sweep: 1.0 Mc to 18.0 Mc in 1 minute 55 seconds.

**HF2** — Km — Observed at **Washington, D.C.**

Km — (Unit) Lat **38.7°N**, Long **77.1°W**

# IONOSPHERIC DATA

Day	Moon Time												Calculated by:
	00	01	02	03	04	05	06	07	08	09	10	11	
1 (27.0) <sup>5</sup> (26.0) <sup>3</sup>	d50	260	240	230	(27.0) <sup>5</sup>	240	230	250	260	270	270	270	270 (27.0) <sup>5</sup>
2 (26.0) <sup>5</sup> (25.0) <sup>3</sup>	440	470	(28.0) <sup>5</sup> (25.0) <sup>3</sup>	480	460	470	300	260	280	270	270	270	270 (25.0) <sup>5</sup>
3 260 (28.0) <sup>5</sup>	250	260	(25.0) <sup>3</sup>	250	250	250	250	260	260	260	260	260	260 (27.0) <sup>5</sup>
4 (27.0) <sup>5</sup>	260	260	230	220	230	240	250	270	280	280	280	280	270 (27.0) <sup>5</sup>
5 260 260	250	240	230	250	220	230	250	(25.0) <sup>4</sup>	260	280	280	280	(25.0) <sup>5</sup> (26.0) <sup>5</sup>
6 (26.0) <sup>5</sup> (26.0) <sup>3</sup>	230	240	230	(27.0) <sup>5</sup>	230	230	230	250	270	270	270	270	270 (26.0) <sup>5</sup>
7 250 360	250	<300	E (31.0) <sup>5</sup>	(27.0) <sup>5</sup>	260	250	250	290	280	260	270	260	260 (25.0) <sup>5</sup>
8 250 (25.0) <sup>5</sup>	270	270	250	260	240	230	250	250	(27.0) <sup>4</sup>	300	260	270	270 (26.0) <sup>5</sup> (27.0) <sup>5</sup>
9 (28.0) <sup>5</sup>	A	A	A	(27.0) <sup>5</sup>	230	230	250	250	270	270	270	270	(27.0) <sup>5</sup> (28.0) <sup>5</sup>
10 (27.0) <sup>5</sup> (26.0) <sup>5</sup>	(27.0) <sup>4</sup>	(26.0) <sup>4</sup>	(27.0) <sup>5</sup>	(27.0) <sup>5</sup>	240	250	260	270	270	270	270	270	(27.0) <sup>5</sup> A A
11 (29.0) <sup>5</sup> (28.0) <sup>5</sup>	(27.0) <sup>5</sup>	(28.0) <sup>5</sup>	(25.0) <sup>5</sup>	250	(25.0) <sup>5</sup>	260	270	300	280	280	270	270	(27.0) <sup>5</sup> (28.0) <sup>5</sup>
12 (31.0) <sup>4</sup> (28.0) <sup>5</sup>	270	(26.0) <sup>5</sup>	250	(25.0) <sup>5</sup>	240	(26.0) <sup>4</sup>	260	270	300	290	290	290	290 (27.0) <sup>5</sup>
13 (29.0) <sup>5</sup> (28.0) <sup>4</sup>	(28.0) <sup>5</sup>	260	240	240	230	260	260	250	280	290	280	270	270 (27.0) <sup>5</sup>
14 270 (29.0) <sup>5</sup>	(28.0) <sup>5</sup>	270	230	(24.0) <sup>4</sup>	210	(24.0) <sup>4</sup>	250	260	270	290	280	270	270 (27.0) <sup>5</sup> (28.0) <sup>5</sup>
15 (26.0) <sup>5</sup> (25.0) <sup>5</sup>	230	240	240	<270	250	230	240	280	(29.0) <sup>5</sup>	300	260	270	270 (27.0) <sup>5</sup>
16 (28.0) <sup>5</sup> (23.0) <sup>5</sup>	E	A	E	E	(33.0) <sup>5</sup>	(26.0) <sup>5</sup>	(24.0) <sup>5</sup>	260	250	260	(30.0) <sup>5</sup>	230	220 (30.0) <sup>5</sup> >70
17 (27.0) <sup>4</sup> (28.0) <sup>5</sup>	280	(28.0) <sup>5</sup>	E	E	(28.0) <sup>5</sup>	240	230	260	260	270	270	270	(27.0) <sup>5</sup> (28.0) <sup>5</sup>
18 (31.0) <sup>5</sup> 290	(27.0) <sup>5</sup>	(27.0) <sup>5</sup>	(24.0) <sup>5</sup>	E	E	(31.0) <sup>5</sup>	260	310	310	330	300	300	280 S K
19 E E	E	E	E	E	E	E	E	E	E	E	E	E	E E K
20 E E	E	E	E	E	E	E	E	E	E	E	E	E	E E K
21 5 5	A	A	S	K	A	S	K	E	E	S	K	A	A E K
22 A A	E	E	E	E	E	E	E	E	E	E	E	A	A E
23 (29.0) <sup>5</sup> (30.0) <sup>5</sup>	250	240	230	230	230	230	230	230	230	230	230	230	230 S K
24 (27.0) <sup>5</sup> (26.0) <sup>5</sup>	(25.0) <sup>5</sup>	(26.0) <sup>5</sup>	(25.0) <sup>5</sup>	(22.0) <sup>5</sup>	(26.0) <sup>5</sup>	220	250	250	260	280	280	280	(27.0) <sup>5</sup> (28.0) <sup>5</sup>
25 (29.0) <sup>5</sup>	220	260	240	240	240	240	240	250	260	260	260	260	(27.0) <sup>5</sup> (28.0) <sup>5</sup>
26 260	250	250	250	250	240	230	230	260	260	250	250	250	250 (27.0) <sup>5</sup>
27 280	250	250	250	(30.0) <sup>4</sup>	E	(29.0) <sup>5</sup>	240	230	270	270	260	270	(29.0) <sup>5</sup> <300 E 270
28 270	280	250	240	240	240	240	240	250	250	260	260	260	260 (27.0) <sup>5</sup> (28.0) <sup>5</sup>
29 (27.0) <sup>5</sup> (26.0) <sup>5</sup>	260	(27.0) <sup>5</sup>	(26.0) <sup>4</sup>	(26.0) <sup>5</sup>	(26.0) <sup>5</sup>	(26.0) <sup>5</sup>	(26.0) <sup>5</sup>	230	220	250	250	240	240 (27.0) <sup>5</sup>
30 (26.0) <sup>5</sup>	270	250	270	260	260	260	260	260	270	270	270	270	(29.0) <sup>5</sup> (28.0) <sup>5</sup> (29.0) <sup>5</sup> A
31 (26.0) <sup>4</sup> (27.0) <sup>5</sup>	270	270	260	260	270	270	270	270	270	270	270	270	(27.0) <sup>5</sup> (27.0) <sup>5</sup>
Median (27.0)	250	260	(25.0)	(25.0)	240	250	260	270	280	290	280	270	270 (27.0)
Count	28	26	27	26	27	26	27	31	31	31	31	31	27 (27.0)

Sweep 1.0 Mc to 25.0 Mc in Q.25 min  
Manual □ Automatic ■

TABLE 50  
 IONOSPHERIC DATA

 Mc 1953  
 (Characteristic) (Month)

Observed at Washington, D.C.

Lat. 38°7'N., Long. 77°10'W

Day	75°W Mean Time												75°W Mean Time														
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23			
1	3.1 F	3.1 F	3.3 F	3.1 F	2.8	2.5	3.2	4.6 F	5.2	5.6	6.2	6.6	6.8	6.6	6.4	5.9	5.4	5.1	5.0	5.0	4.2	3.7	3.5	3.3			
2	2.9	2.8	2.8	2.7	2.6 F	2.6 F	2.7	(2.0) S	4.7	5.2	5.8	5.8	5.6	5.8	6.2	5.9	6.0	5.7	5.4	4.8	4.3	3.5	3.5	3.4			
3	3.1	3.0	3.1	3.0	2.7	3.3	4.9	(5.9) H	5.6	5.4 H	5.6	6.0	6.1	6.2	5.4	5.9	6.0	5.5	4.6	3.8	3.1	3.1	3.0	3.0			
4	3.0	3.1 F	3.2 F	3.2 F	3.0	3.7	3.5	5.3	5.3	5.9	6.2	5.9	6.2	5.7	5.6	5.6	(5.6) S	(5.6) S	(5.6) S	(5.6) S	(3.4) S	(3.4) S	(3.4) S	(3.4) S			
5	(3.1) F	(3.0) F	3.2 F	(3.1) F	3.3	3.2 F	(3.5) F	5.4	5.6	5.4	5.6	5.6	6.2	6.0	6.0	6.1	6.4	6.0	5.4	3.8 S	3.5	3.2	3.2	(3.3) S			
6	3.2	3.1	3.0	3.0	2.9	2.6	3.1 F	4.8	5.9	6.1 H	6.8	6.3	6.2	6.0	6.4	6.4	6.0	5.9	5.5	4.2	3.8	3.7	3.6	3.8			
7	(3.1) S	3.4	3.0	1.8 F	1.7	2.3 F	3.5	4.9	5.0	5.6	5.8	6.4	6.7	6.3	7.1	6.8	6.5	6.0	4.7	4.6	4.1 S	3.7	3.7	3.7	3.7		
8	3.7	3.4 S	3.0 F	3.2 F	3.1 F	3.0 F	3.2	4.3	5.5 H	6.1	6.0	(5.8) H	6.9	7.1	6.1	6.2	6.6	6.4	6.0	4.9	3.8	3.8	3.3	3.0			
9	2.7 S	(2.5) A	A	A	(2.5) H	(2.9) S	4.9	5.8	5.3	6.2	6.8	6.5	6.9	7.3	6.7	6.0	6.1	5.1	4.2	3.7	(3.3) H	3.2	3.0	3.0	3.0		
10	3.1	2.9	(2.8) A	2.5	2.2	(1.9) A	2.7	4.3	4.7	5.4	5.8 H	6.3	6.2	6.6	6.8	6.8	6.9	6.3	5.0	4.2	3.3	2.7	(2.5) H	(2.4) A	(2.5) A		
11	2.3	2.3 F	2.2	2.2	2.4	2.2	2.6	4.3	4.3	5.6	5.5	6.2	6.4	6.8	6.1	6.3	5.9	5.1	4.5	4.3	3.4	2.8	(2.5) A	(2.4) A	(2.5) A		
12	(2.4) H	2.5	2.5	2.6	2.5	2.4	2.8	4.2	4.8	5.2	5.8	5.7	6.0	6.3	6.0	5.9	5.9	5.7	5.0	4.4	3.1	3.5	2.5	(2.5) A	(2.5) A	(2.5) A	
13	2.5	2.6	2.6	2.7	2.7 F	2.5	2.8	4.6	5.9	6.7	6.0	6.0	5.8	6.1	6.8	6.8	6.7	6.0	5.2	4.1	2.9	2.7	2.6	2.7	2.7		
14	(2.6) S	2.5	2.6	2.7	(2.6) F	(2.4) S	(3.0) S	4.8	(5.5) H	6.0	6.0	6.0	6.8	6.1	6.0	6.2	6.1	6.4	5.7	4.7	3.8	3.0	2.6	2.7	2.7		
15	2.8	2.8	2.7	2.6 F	2.5	2.5	(2.4) F	2.6	4.8	5.0	5.7	5.7 K	7.5 K	7.6 K	9.4 K	8.9 K	6.9 K	6.2 K	5.8 S	5.2 K	4.7 K	3.9 K	4.1 K	4.0 K	(3.3) S		
16	(1.1) S	(2.0) P	(2.0) S	<1.0 E	(1.1) S	(2.3) F	(3.1) F	5.2 S	5.2 S	5.4	5.1 S	5.8	5.1 S	5.4	6.5	6.8	6.9	(7.3) S	7.4	6.8	6.4	4.6	3.4	2.9	3.2		
17	2.8 S	2.4 F	2.3	2.4 F	2.3	2.3	2.3	4.9	5.8	5.6	5.6	6.0 H	7.1	8.0	8.2	6.8	7.2	8.2	7.6	6.4	5.4	4.1	3.3	3.0	2.6		
18	2.6 S	(2.5) S	(2.5) F	2.6	2.3	2.2 K	1.0 E	2.2 K	3.6 K	4.6 K	4.6 K	5.6 S	5.0 H	6.0 K	5.6 K	5.8 K	5.2 K	5.7 K	4.7 K	3.5 K	3.2 K	2.7 S	F K	F S	F K		
19	F S	F K	F S	F K	F S	<1.0 E	(1.0 E)	(1.0 E)	(1.8) S	2.8 K	2.8 K	<3.5 K	<3.7 K	4.1 K	4.4 K	4.3 K	4.9 K	4.7 K	4.2 K	4.0 K	3.1 K	2.3 K	(1.9) S	K 17 S	K (1.5) S	K (1.5) S	
20	<1.0 E	<1.0 E	<1.0 E	<1.0 E	<1.0 E	(1.0 E)	(1.0 E)	(1.0 E)	(1.8) S	3.1 K	3.1 K	(3.7 K	4.1 K	4.4 K	4.4 K	4.4 K	5.2 K	5.0 K	5.4 K	5.4 K	4.9 K	4.0 K	3.6 K	2.8 K	2.4 K	(1.7) K	
21	(1.6) S	(1.5) S	(1.5) S	A X	F X	<1.0 E	(1.0 E)	(1.0 E)	2.3	4.9	5.8	5.6	6.0 H	7.1	8.0	8.2	6.8	7.2	8.2	7.6	6.4	5.4	4.1	3.3	3.0	2.6	2.6
22	[1.6] A	[1.6] S	[1.5] S	F S	F K	<1.0 E	F K	(3.6) F	(4.1) F	4.8 K	5.0 K	5.2 K	5.6 S	5.0 H	6.0 K	5.6 K	5.8 K	5.2 K	5.7 K	4.7 K	3.5 K	3.2 K	2.7 S	F K	F S	F K	
23	(2.3) F	(2.4) F	(2.4) F	(2.4) F	(2.3) F	(2.2) S	(2.2) S	4.1	5.0	5.6	6.0	6.3 K	6.4	6.6	6.2	6.1	5.7	5.0	3.6	2.6	2.3	2.0 F	2.0 F	(2.1) S	(2.1) F	(2.1) F	
24	(2.1) F	(2.2) F	(2.4) F	(2.5) F	(2.3) F	(2.4) F	(2.4) F	4.1 F	5.3	5.8	5.6	-6.7	7.5	7.5	6.8	6.8	6.2	5.4	4.3	(3.4) F	2.9 S	(2.5) F	2.6 F	(2.7) E	(2.7) E	(2.7) E	
25	(2.8) F	(2.8) F	(3.0) F	3.1 S	2.8	2.3	2.2	4.0	4.8 V	5.2	5.4 H	6.2	6.2	5.9	5.8	6.1	5.9	5.0	3.7 F	(3.1) S	2.9	2.7	2.6	2.7	2.7	2.7	
26	2.9	2.7	2.9	2.7	2.8	2.7	2.7	3.8	4.7	5.0	5.4	5.5	6.2	6.2	6.4	6.6	6.1	6.2	5.4	3.9	3.3	2.7	2.4	2.6	2.7	2.7	
27	2.8	2.9	2.5	2.2	2.1	2.1	4.1	4.8	5.6	6.3	6.4	6.8	7.6	7.0	7.0	6.4	5.7	4.9	3.2	[3.0] A	2.7	2.8 J	3.0 F	3.0 F	3.0 F	3.0 F	
28	(3.3) F	(3.4) S	(3.5) S	(3.2) S	2.8 F	2.5	4.5	5.2	6.4	6.0	6.2	6.7	7.0	6.5	6.6	6.4	5.1	4.1	3.7	3.3	2.9	2.7	2.7	2.7	2.7	2.7	
29	2.7	2.5 F	2.5 F	2.4 F	2.4 F	2.4 F	2.4 F	4.1	5.8 S	6.0	6.4	6.5	7.5	7.5	7.4	6.2	6.6	6.8	6.5	6.1	5.7	3.7	3.3	2.7	2.2	2.2	2.2
30	2.4	(2.7) F	2.6 F	2.7 F	3.1 F	2.9 F	3.1 F	2.7	3.7	4.8	4.9	5.6 K	6.3	6.7	6.5	5.8	6.2	6.0	5.0	3.9	3.4	3.0	2.8	2.7 F	2.8	2.8	2.8
31	2.9 F	2.8 F	2.8 F	2.7	2.5	2.3	3.8	4.7	4.9	5.1	5.5	5.8	6.0	5.9	5.6	5.6	5.0	(3.5) A	(3.0) A	(2.5) A	2.3	2.4	2.4	2.4	2.4	2.4	
Median	2.8	2.7	2.6	2.6	2.5	2.4	2.8	4.3	5.0	5.6	5.6	6.2	6.2	6.2	6.1	6.0	5.7	4.7	3.8	3.3	2.8	2.7	2.7	2.7	2.7	2.7	
Count	30	30	28	28	28	29	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31

 National Bureau of Standards  
 (Institution)  
 MCC., E.J.W.  
 Scaled by:  
 Calculated by:  
 MCC., E.J.W.

 Sweep-10 Mc to 250 Mc in 0.23 min  
 Manual □ Automatic □

**TABLE 51**  
 Ionospheric Data  
 National Bureau of Standards

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

TABLE 52  
IONOSPHERIC DATA  
Lat. 38°7'N., Long. 77.1°W.  
Observed at Washington, D. C.  
(Characteristic) Km (Unit) October (Month)  
Day 00 01 02 03 04 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23

$h'F_1$

Km

(Unit)

National Bureau of Standards  
(Institution) MCC, E.J.W.

Observed at Lat. 38°7'N., Long. 77.1°W.

Scaled by: Calculated by: MCC, E.J.W.

Day	75°W Mean Time																							
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	Q	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0
2	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
3	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
4	Q	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0
5	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
6	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
7	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
8	Q	230	240	250	260	270	280	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290
9	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
10	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
11	Q	230	240	250	260	270	280	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290
12	Q	230	240	250	260	270	280	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290
13	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
14	Q	200	200	210	220	230	240	250	260	270	280	290	290	290	290	290	290	290	290	290	290	290	290	290
15	Q	210	220	230	240	250	260	270	280	290	290	290	290	290	290	290	290	290	290	290	290	290	290	290
16	Q	240	230	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20
17	Q	230	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10
18	Q	240	230	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20
19	Q	240	230	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20
20	Q	230	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10
21	Q	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0
22	Q	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0
23	Q	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0
24	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
25	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
26	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
27	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
28	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
29	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
30	Q	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0
31	Q	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10	0	0
Median		230	220	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	40	30	20	10
Count		"	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30

Sweep 1.0 Mc to 25.0 Mc in 0.25-min  
Manual  Automatic

**TABLE 53**  
**IONOSPHERIC DATA**

**f<sub>0</sub>F<sub>1</sub>**      **Nc**      **October**, 1953  
(Characteristic)      (Unit)      (Month)

**Observed at Washington, D. C.**

**Lat 38.7°N**

**Long 77.1°W**

Day	75°W												Mean Time												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
2	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
3	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
4	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
5	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
6	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
7	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
8	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
9	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
10	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
11	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
12	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
13	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
14	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
15	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
16	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
17	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
18	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
19	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
20	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
21	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
22	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
23	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
24	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
25	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
26	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
27	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
28	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
29	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
30	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
31	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
Median	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Count	3	20	20	9	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31

Sweep 1.0 Mc to 25.0 Mc in 0.25-min  
Manual  Automatic

National Bureau of Standards

(Institution)

MCC., E.J.W.

Scaled by:

Calculated by:

TABLE 54  
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

Form adopted June 1946  
National Bureau of Standards  
(Institution)  
MCC, E.J.W.

$h^{\prime}E$ , Km (Units)  
(Characteristic) Observed at Lat 38.7°N, Long 77.1°W  
Km (Month)  
October, 1953  
D. C.

IONOSPHERIC DATA  
**75°W Mean Time**

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1																								
2																								
3																								
4																								
5																								
6																								
7																								
8																								
9																								
10																								
11																								
12																								
13																								
14																								
15																								
16																								
17																								
18																								
19																								
20																								
21																								
22																								
23																								
24																								
25																								
26																								
27																								
28																								
29																								
30																								
31																								
Median Count																								
	17	23	27	26	29	30	28	20	21	22	23	20	19	18	17	16	15	14	13	12	11	10	9	8

Sweep 1.0 Mc to 25.0 Mc in 2.5 min  
Manual  Automatic

TABLE 55  
 IONOSPHERIC DATA

fo E Mc October 1953

(Characteristic) (Unit)

Washington, D.C.

(Month)

Lot 38.7°N Long 77.1°W

Observed at \_\_\_\_\_

National Bureau of Standards

(Institution)

MCC., E.J.W.

Calculated by:

MCC., E.J.W.

Doy	75°W												Mean Time												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1																									
2																									
3																									
4																									
5																									
6																									
7																									
8																									
9																									
10																									
11																									
12																									
13																									
14																									
15																									
16																									
17																									
18																									
19																									
20																									
21																									
22																									
23																									
24																									
25																									
26																									
27																									
28																									
29																									
30																									
31																									
Median	(1.8)	24	26	28	28	29	29	30	30	30	30	31	31	31	31	31	31	31	31	31	31	31	31	31	
Count	16	20	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	

 Sweep 1.0 Mc to 25.0 Mc in 0.25 min  
 Manual  Automatic

TABLE 56  
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.  
IONOSPHERIC DATA

Form adopted June 1946

Characteristic	Mc, Km (Unit)	Observed at Washington, D.C.	Lat 38°7'N, Long 77°10'W	75°W Mean Time												Calculated by: MCC, E, J, W	Bureau of Standards (Institution)											
				Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	4.2°V/0	E	E	E	E	E	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	G	G	
2	2.3°V/0	2.4°H/0	E	2.9°/0.0	3.0°/0.0	2.6°/0.0	2.6°/0.0	2.5°/0.0	3.0°/0.0	3.2°/0.0	3.7°/0.0	3.8°/0.0	3.8°/0.0	3.8°/0.0	3.8°/0.0	3.8°/0.0	3.8°/0.0	3.4°/0.0	3.0°/0.0	2.7°/0.0	2.7°/0.0	2.5°/0.0	2.5°/0.0	2.5°/0.0	2.5°/0.0	2.5°/0.0	2.5°/0.0	
3	2.4°/0.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E		
4	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E		
5	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E		
6	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E		
7	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E		
8	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E		
9	2.6°/0.0	3.3°/0.0	4.6°/1.0	5°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	4.9°/1.0	
10	E	2.6°V/1.0	3.7°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	3.4°/1.0	
11	4.8°/1.0	E	3.1°/1.0	2.9°/1.0	3.0°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	3.2°/1.0	
12	4.7°/0.0	2.6°V/1.0	2.2°/1.0	2.4°/1.0	3.1°/1.0	3.1°/1.0	3.3°/1.0	3.3°/1.0	3.5°/1.0	3.5°/1.0	4.0°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	4.6°/1.0	
13	2.5°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	3.2°/0.0	
14	4.9°/0.0	2.3°/0.0	4.0°/0.0	4.0°/0.0	4.3°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	
15	E	3.2°V/0.0	2.4°/0.0	2.4°/0.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
16	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	
17	2.0°V/4.0	3.9°/1.0	3.5°/1.0	2.9°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	3.6°/0.0	
18	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	
19	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	
20	E	E	E	E	4.4°/3.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
21	E	2.4°/3.0	3.0°/1.0	2.6°/1.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
22	2.6°/0.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
23	E	E	E	E	E	3.2°V/1.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
24	E	E	E	E	E	1.6°/1.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
25	E	E	E	E	E	2.3°V/1.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
26	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	
27	E	E	E	E	E	3.8°V/0.0	4.4°V/0.0	4.4°V/0.0	4.4°V/0.0	2.5°V/0.0	2.5°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	2.4°V/0.0	
28	2.3°V/1.0	4.2°/0.0	4.4°V/0.0	4.4°V/0.0	4.4°V/0.0	2.4°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0	2.5°V/0.0										
29	2.6°V/1.0	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
30	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E
31	4.2°/1.0	2.1°/2.0	E	3.7°V/2.0	2.4°/1.0	4.4°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0	3.5°/2.0
Median	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**
Count	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31

Sweep  $\Omega$ , Mc to 25. Mc in 25-min  
Manual  Automatic

TABLE 57  
IONOSPHERIC DATA  
Lat 38°7' N, Long 77°10' W

(M1500)F2, October, 1953  
(Characteristic) (Unit)  
Observed at Washington, D. C.

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

(Month)

75°W

Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	.20 F	.20	.21	.21 F	.22	.23	.24	.23	.22	.23	.23	.22	.23	.23	.24	.23	.23	.22	.23	.23	.22	.21	.21	.22	
2.	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.22	
3	.20	.20	.20	.20	.20	.20	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.22	
4	.21 J	.21 J	.20 S	.20 F	.20 S	.20 J	.20 S	.20 F	.20 J	.20 P															
5	.20 J																								
6	.22 S	.21	.20 J																						
7	.20 J	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21
8	.21	.21 J	.20 S	.20 F	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21
9	.20 J	.21 J	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	
10	.21	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20
11	.20	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21
12	.20 J	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20
13	.20	.20	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21
14	.21 S	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20
15	.21	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22	.22
16	.21 J	.20 S	.20 F	.20 J																					
17	.21 J	.21	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20	.20
18	.20 J																								
19	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	
20	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	E K	
21	.19 J	S	A	F	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	
22	A	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	
23	E	K	F	S	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
24	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
25	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
26	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
27	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
28	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
29	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
30	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
31	E	K	F	E	E	J	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	E	K	
Median	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21	.21
Count	28	27	26	25	26	26	25	25	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26

Manual □ Automatic □  
Sweep LO Mc 25.0 Mc in 0.25 min

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

TABLE 58  
IONOSPHERIC DATA

(M3000)F2, - October, 1953

(Characteristic) (Unit) Washington, D. C.

Observed at Lat 38.7°N, Long 77.0°W

Day	75°W												Mean Time														
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23			
1	3.0	F	3.0	F	3.1	P	3.3	3.5	3.1	3.4	(3.5)	3	3	3.3	3.2	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.2	3.1	3.2		
2	3.1	3.1	3.1	3.1	3.1	(3.1)	3	(3.4)	3.5	3.0	3.5	3.4	3.5	3.2	3.3	3.3	3.3	3.4	2.9	2.9	3.4	3.4	3.3	3.1	3.2		
3	3.0	3.0	3.2	3.1	3.1	3.1	3.1	3.2	3.4	3.4	(3.4)	3.5	(3.5)	H	3.2	3.4	3.2	3.3	3.5	3.5	3.5	3.3	3.3	3.1	3.0		
4	3.1	(3.1)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	3.3	3.4	3.4	3.6	3.5	3.4	3.4	3.3	3.5	3.4	3.2	3.3	3.3	3.3	3.1	3.1	..		
5	(3.1)	(2.9)	(3.1)	(2.9)	(3.1)	(2.9)	(3.1)	(2.9)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)	(3.2)		
6	3.2	3.1	3.2	3.2	3.2	3.3	3.3	3.4	3.4	3.4	3.5	3.5	3.5	3.5	3.5	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4		
7	(3.1)	3	3.1	3.1	3.1	3.1	3.1	3.0	3.0	3.0	(3.0)	F	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0		
8	3.1	(3.1)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0		
9	(3.1)	(3.1)	A	A	A	A	A	(3.1)	H	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)	(3.4)		
10	3.1	3.0	(3.1)	A	3.1	(3.4)	A	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	(3.4)	3.1	
11	3.0	3.0	3.2	F	3.2	3.1	3.1	3.3	3.3	3.2	3.5	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.4	3.4	3.4	3.4	3.4	3.4	3.4	A	
12	(3.0)	A	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	
13	3.0	3.0	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	
14	(3.1)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	
15	3.1	3.2	3.2	3.3	3.3	3.3	3.3	3.4	3.4	3.4	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	
16	(3.1)	P	(3.0)	P	E	K	E	K	E	K	(K)(2.8)	P	N	(3.2)	F	(3.5)	E	K	E	K	E	K	E	K	E	K	
17	(3.1)	J	3.1	F	3.0	(3.0)	F	E	E	E	2.9	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	
18	(2.8)	J	(2.8)	J	(3.1)	J	(2.9)	J	E	K	2.8	K	3.2	K	(3.6)	J	(3.6)	K	3.2	K	3.2	K	3.2	K	3.2	K	A
19	F	S	F	S	E	K	E	K	E	K	K	K	G	K	G	K	G	K	2.4	K	3.1	K	3.1	K	3.1	K	
20	E	K	E	K	E	K	E	K	E	K	K(2.5)	J	3.4	K	2.8	K	2.9	K	2.4	K	3.0	K	2.8	K	2.8	K	
21	A	K	(2.8)	J	S	K	E	K	E	K	(3.3)	K	3.4	K	(3.6)	K	3.4	K	3.2	K	3.2	K	3.2	K	3.2	K	
22	A	K	(2.8)	J	F	S	F	K	E	K	F	K	K(3.4)	J	K(3.2)	F	3.4	K	3.4	K	3.4	K	3.4	K	3.4	K	
23	(3.0)	J	(3.3)	J	3.3	F	(3.4)	J	F	S	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5		
24	(3.3)	J	(3.1)	P	(3.2)	P	(3.2)	P	F	(3.2)	J	3.6	3.5	3.4	3.4	3.3	3.5	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4		
25	(3.2)	F	(3.2)	J	(3.2)	J	(3.2)	J	3.4	3.4	3.2	3.6	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5		
26	3.2	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1		
27	3.0	3.2	3.3	3.3	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0		
28	(3.1)	J	(3.4)	S	(3.4)	J	3.3	F	3.4	3.4	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5		
29	3.0	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1		
30	3.3	(3.1)	J	(3.1)	J	3.0	F	3.2	F	3.1	5	3.2	5	3.3	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	
31	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.2	3.3	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4		
- Median	3.1	3.1	3.1	3.1	3.1	3.2	3.3	3.3	3.5	3.5	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4		
Count	28	27	26	26	26	25	25	25	28	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	

Sweep 10 Mc to 250 Mc in 0.25 min  
Manual □ Automatic ■

C

Form adopted June 1946

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

**TABLE 59**  
**IONOSPHERIC DATA**

(M3000)F1, (Unit)  
 Observed at Washington, D.C.  
 (Month) October, 1953  
 Lat. 38.7°N Long. 77.1°W

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	L	L	3.9	3.7	3.8	3.9	H	3.9	H	3.9	H	3.9
2	L	L	(3.9)	L	3.7	3.8	H	3.8	H	3.7	H	L
3	L	L	L	L	3.7	3.9	H	3.7	H	3.7	H	L
4	L	L	3.8	3.6	3.7	3.9	L	3.7	L	3.7	L	L
5	L	L	(4.0)	L	(3.9)P	(3.9)H	L	3.7	L	(3.9)	L	L
6	L	L	3.8	H	3.6	P	3.7	3.7	L	L	L	L
7	L	L	L	4.0	H	A	3.7	H	3.6	(3.7)H	L	Q
8	G	L	L	L	4.0	H	4.0	3.7	3.6	H	(3.6)H	L
9	G	L	L	L	(3.8)H	A	4.0	(3.8)P	(3.6)P	(3.6)H	L	Q
10	L	L	L	L	3.9	3.7	3.7	3.6	H	3.6	L	Q
11	L	L	L	L	3.7	3.7	3.7	3.6	H	(3.6)P	3.7	L
12	Q	L	L	L	3.7	3.7	3.9	3.9	3.9	3.6	3.7	L
13	L	L	L	L	3.9	3.9	3.9	3.7	3.7	3.6	3.6	L
14	L	L	L	L	3.9	3.8	3.7	3.7	3.8	H	(3.8)L	L
15	Q	L	L	L	(3.6)H	L	K	(3.6)K	L	3.5	H	L
16	L	K	L	X	3.8	K	(3.9)H	3.7	K	3.6	K	L
17	L	K	L	L	(3.9)	K	3.8	L	3.4	P	3.7	H
18	G	K	3.5	K	(3.6)H	K	3.6	K	3.4	K	3.7	K
19	Q	K	3.3	K	3.7	H	3.9	H	3.2	K	3.5	K
20	G	K	3.5	K	3.9	K	3.6	H	4.2	H	3.7	K
21	Q	K	L	K	3.9	K	(3.8)H	3.7	K	3.8	K	3.8
22	Q	K	3.6	K	3.9	K	3.9	K	3.6	K	3.6	K
23	Q	L	L	L	(3.9)H	L	L	(3.5)H	L	3.7	K	L
24	Q	L	L	L	(3.9)H	(3.9)P	(3.7)H	L	3.8	H	3.7	K
25	L	L	L	L	(3.9)H	L	(3.9)H	3.6	H	3.7	H	L
26	G	L	L	L	4.0	H	3.6	H	3.9	H	4.0	K
27	Q	L	L	L	(3.7)P	P	3.7	L	(3.7)H	L	K	Q
28	G	K	L	K	3.9	H	(3.9)H	L	3.9	L	3.7	H
29	L	L	A	A	(4.0)P	(3.7)P	(3.8)H	L	L	L	A	B
30	Q	L	L	L	(3.9)H	L	(3.7)H	L	L	L	B	Q
31	L	L	(3.9)H	(4.0)A	A	3.8	3.6	L	L	L	A	-
	-	-	3.8	3.8	3.9	3.9	3.7	3.7	3.9	-	-	-
3	3	6.0	27	27	31	31	28	28	22	22	2	2

Mean  
Count

Manual □ Automatic □  
 Sweed 10 Mc 10.350 Mc in 0.25 min

TABLE 60  
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.  
IONOSPHERIC DATA

Form adopted June 1946

(M) (500) E , (Unit) (Month)  
Observed at Washington, D. C.  
Lat. 38°7'N, Long 77.1°W

October, 1953

(Month)

W

National Bureau of Standards

(Institution)

McC. E. J. W.

Day	75°W												Mean Time													
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23		
1										A	(4.3) A	4.4	4.4	4.4	4.3	4.2	4.2	4.3	4.4	4.4	4.3	4.2	4.2	4.1		
2										A	(4.2) A	4.4	4.3	4.3	4.3	4.4	4.4	4.3	4.2	4.2	4.1					
3										A	(4.3) A	4.4	4.3	4.4	4.3	A	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1		
4										A	(4.3) A	4.4	4.3	4.3	4.3	4.4	4.3	4.0	4.4	4.4	4.3					
5										A	(4.4) S	4.3	4.3	4.3	4.1	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.3		
6										A	(4.4) A	4.5	4.5	4.5	4.5	(4.2) P										
7										A	4.3	4.5	A	(4.3) A	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
8										A	(4.5) A	4.4	4.4	4.4	4.3	4.4	4.0	4.1	4.2	4.2	4.2	4.2	4.2	4.2		
9										A	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4		
10										A	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4		
11										A	(4.1) A	4.3	4	A	A	4.2	4.2	A	A	A	A	A	A	A		
12										A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
13										A	A	A	A	A	(4.3) P	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3		
14										A	(4.2) P	4.2	4.2	4.2	4.2	(4.1) P	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1		
15										A	(4.4) A	4.4	4.4	4.4	4.4	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2		
16										A	(4.3) P	4.4	4.4	4.4	4.4	(4.3) A	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3		
17										A	(4.2) P	4.2	4.2	4.2	4.2	(4.2) P	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1		
18										A	(4.2) A	4.3	4.3	4.3	4.3	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2		
19										A	(4.3) K	4.1	4.1	4.1	4.1	(4.0) K	B	K	K	K	K	K	K	K		
20										A	(4.3) K	4.1	4.1	4.1	4.1	(4.3) K	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1		
21										A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
22										A	K	K	K	K	K	A	A	A	A	A	A	A	A	A		
23										A	A	A	A	A	A	(4.5) H	A	A	A	A	A	A	A	A		
24										A	(4.2) H	4.3	4.3	4.1	4.1	(4.1) S	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1		
25										S	4.4	4.0	(4.2) H	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	
26										(4.3) A	4.3	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0		
27										(4.3) P	4.3	(4.2) P	A	A	4.1	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
28										S	A	4.3	A	A	A	(4.3) P	(4.3) A	A	A	A	A	A	A	A		
29										S	A	4.3	4.3	A	(4.4) P	(4.4) A	A	(4.4) P	(4.4) A	(4.4) P	(4.4) A	(4.4) P	(4.4) A	(4.4) P		
30										A	A	A	A	A	A	(4.2) A	4.0	A	A	A	A	A	A	A		
31										(4.2) A	4.3	4.3	4.3	4.3	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2		
Median Count	15	19	21	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	

Manual  Automatic

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

3

Table 61

Ionospheric Storminess at Washington, D. C.October 1953

Day	Ionospheric character*		Principal storms		Geomagnetic character**	
	00-12 GCT	12-24 GCT	Beginning GCT	End GCT	00-12 GCT	12-24 GCT
1	1	1			2	2
2	1	2			2	1
3	1	2			2	2
4	1	2			1	2
5	1	2			2	1
6	0	2			1	1
7	1	1			3	2
8	.	1			2	3
9	2	0			3	2
10	1	1			2	2
11	2	1			1	2
12	2	2			1	1
13	2	2			1	1
14	2	2			1	1
15	1	4	1500	----	1	5
16	4	3	----	1100	4	4
17	3	3			4	4
18	3	4	0900	----	4	5
19	7	6	-----	-----	6	4
20	7	5	-----	-----	4	4
21	5	4	-----	-----	4	2
22	5	4	-----	2100	3	3
23	2	1			3	2
24	2	1			1	2
25	1	2			3	2
26	0	2			2	1
27	1	1			4	2
28	1	1			2	2
29	1	1			2	3
30	1	2			2	1
31	2	3			3	1

\*Ionosphere character figure (I-figure) for ionospheric storminess at Washington, D. C., during 12-hour period, on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

\*\*Average for 12 hours of Coltenham, Maryland, geomagnetic K-figures on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.  
----Dashes indicate continuing storm.

Table 62Zürich Provisional Relative Sunspot NumbersOctober 1953

Date	R <sub>Z</sub> *	Date	R <sub>Z</sub> *
1	0	17	9
2	0	18	0
3	7	19	0
4	7	20	0
5	0	21	7
6	13	22	0
7	13	23	0
8	14	24	7
9	11	25	7
10	10	26	2
11	9	27	7
12	9	28	7
13	8	29	0
14	29	30	0
15	22	31	8
16	13	Mean:	7.4

\*Dependent on observations at Zürich Observatory and its stations at Locarno and Arosa.

Table 53a

Radio Propagation Quality Figures  
(Including Comparisons with Short-Term and Advance Forecasts)

September 1952

Day	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day quality index	Advance forecasts (J-reports) for whole day; issued in advance by:			Geomag- netic K Ch	
	00	06	12	18	00	06	12	18		1-4 days	4-7 days	8-25 days	Half day (1)	Day (2)
06.	to 06.	to 12	to 18	to 24										
1	5	(4)	5	6	(4)	(4)	5	5	5	6	7	X	3	3
2	5	(3)	6	5	5	(4)	5	5	(4)	5	7	X	(4)	2
3	(3)	(4)	6	5	(4)	(2)	6	6	(4)	5	7	X	2	(4)
4	(2)	(2)	(4)	(4)	(3)	(2)	(4)	5	(3)	6	7	X	(6)	(4)
5	(2)	(2)	6	6	(2)	(2)	(4)	(4)	(3)	(3)	6	X	(4)	3
6	(4)	(4)	6	6	(3)	(3)	5	6	5	(3)	(4)	X	3	2
7	5	(4)	7	6	5	(4)	6	7	6	(4)	(4)	X	3	3
8	5	(4)	6	6	6	(4)	7	7	5	(4)	(4)	X	3	1
9	5	(4)	7	7	5	(4)	6	7	6	5	5	X	3	1
10	6	6	7	6	5	5	7	7	6	6	6	X	2	2
11	6	5	7	7	5	(4)	6	7	6	6	6	X	3	2
12	6	5	7	7	6	5	7	7	6	7	7	X	2	3
13	7	5	7	7	6	5	6	7	7	7	7	X	3	2
14	7	6	7	7	5	5	7	7	7	7	7	X	1	1
15	7	6	7	7	7	6	7	7	7	7	7	X	2	(4)
16	6	5	7	7	(4)	(4)	6	7	6	7	7	X	3	3
17	5	5	7	6	7	5	6	7	6	7	7	X	3	2
18	6	5	7	7	6	5	6	6	6	6	6	X	3	3
19	(3)	(2)	(4)	(3)	5	(2)	(4)	(4)	(3)	5	5	X	(6)	(4)
20	(2)	(2)	5	5	(3)	(2)	(3)	(4)	(3)	(3)	(3)	X	(5)	(4)
21	(4)	(2)	6	5	(3)	(2)	(4)	5	(4)	(3)	(3)	X	(4)	(4)
22	(4)	(3)	6	5	5	(2)	(4)	5	(4)	(4)	(4)	X	(5)	3
23	(3)	(2)	5	5	(3)	(2)	(4)	(4)	(3)	(4)	(4)	X	(5)	(4)
24	(3)	(3)	5	6	(3)	(3)	5	5	(4)	(4)	(4)	X	(5)	3
25	(4)	5	6	6	(4)	(4)	6	6	5	(4)	(4)	X	3	2
26	6	5	6	6	5	(4)	6	6	6	(4)	(4)	X	2	2
27	5	(4)	6	6	5	(4)	5	5	5	5	(4)	X	(4)	1
28	(3)	(4)	7	6	(4)	(4)	6	6	5	5	5	X	3	1
29	5	6	7	7	5	5	6	7	6	5	5	X	1	1
30	5	5	7	6	5	5	7	7	6	6	5	X	2	2
<b>Score:</b>														
Quiet periods				P	9	6	10	15		9	7			
				S	6	7	14	12		8	10			
				U	2	0	1	0		1	1			
				F	1	0	3	1		2	2			
Disturbed periods				P	4	13	2	0		4	3			
				S	7	3	0	2		4	2			
				U	1	1	0	0		1	1			
				F	0	0	0	0		1	4			

**Scales:****Q-scale of Radio Propagation Quality**

- (1) - useless
- (2) - very poor
- (3) - poor
- (4) - poor to fair
- 5 - fair
- 6 - fair to good
- 7 - good
- 8 - very good
- 9 - excellent

**K-scale of Geomagnetic Activity**

0 to 9, 9 representing the greatest disturbance;  $K_{Ch} \geq 4$  indicates significant disturbance, enclosed in ( ) for emphasis

**Scoring:** (beginning October 1952)

P - Perfect: forecast quality equal to observed

S - Satisfactory: (beginning October 1952)  
forecast quality one grade different from observed

U - Unsatisfactory: forecast quality two or more

grades different from observed when both forecast and observed were  $\geq 5$ , or both  $\leq 5$ 

F - Failure: other times when forecast quality two or more grades different from observed

**Symbols:**

X - probable disturbed date

**Note:** All times are UT (Universal Time or GCT)


Table 63b

## Short-Term Forecasts---September 1953

— forecast

x CRPL observation (not in consensus)

- individual reports of quality  
(adjusted to CRPL scale)



## Outcome of Advance Forecasts (1 to 4 days ahead) --- September 1953




Table 64a

### Coronal observations at Climax, Colorado (5303A), east limb

Date GCT	Degrees north of the solar equator															0°	Degrees south of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	
1953	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-			
Oct	4.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-		
	5.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	1	1	1	1	2	1	1	1	-	-	-	-	-	-	-		
	6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
	7.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
	8.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	9.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	6	8	7	-	-	-	-	-	-	-	-	-	-	-	-	
	10.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	3	4	1	-	-	-	-	-	-	-	-	-	-	-	-	
	11.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	15.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	2	1	1	1	-	-	-	-	-	-	-	-	-	-	-	
	16.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	17.8	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	18.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	20.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	4	3	2	3	3	6	13	8	3	3	-	-	-	-	-	-	-
	22.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	1	1	4	2	5	4	2	2	2	1	1	1	1	1	-	
	23.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	2	1	2	3	1	-	-	-	-	-	-	-	-	-	-	
	24.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	25.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	26.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	3	2	2	2	3	4	1	1	1	2	-	-	-	-	-	
	27.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	28.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	3	3	2	1	2	2	1	1	1	3	1	2	-	-	-	
	29.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	2	2	1	3	2	1	1	1	2	1	1	2	1	-	-	
	30.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	3	2	1	1	2	2	3	3	3	1	1	1	2	2	1	
	31.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	2	1	1	3	2	2	1	1	2	1	1	2	2	1	

Table 65a

### Coronal observations at Climax, Colorado (6374A), east limb

Table 64b

Coronal observations at Climax, Colorado (5303A), west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator																				
	90	85	80	75	70	65	50	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	55	70	75	80	85	90			
1953	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-	-		
Oct	4.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	5	6	3	2	2	1	1	-	-	-	-	-	-	-	-	-	-		
	5.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	6	3	2	2	1	1	-	-	-	-	-	-	-	-	-	-	-		
	6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	1	2	4	9	15	16	3	3	1	1	1	-	-	-	-	-	-		
	7.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	5	12	17	3	3	2	2	1	-	-	-	-	-	-	-	-	-		
	8.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	7	6	2	2	2	1	-	-	-	-	-	-	-	-	-	-		
	*9.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	4	4	1	1	1	1	-	-	-	-	-	-	-	-	-	-	
	10.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-	-	-		
	11.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	15.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	2	1	1	2	10	2	-	-	5	4	1	1	1	-	-	-		
	16.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	2	2	2	4	6	4	4	2	2	1	1	1	-	-	-		
	17.8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
	18.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	6	3	1	1	-	-	-	-	-	-	-	-	-	-	-	-		
	20.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-	-	-	
	22.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	23.8a	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
	24.7	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
	25.7	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
	26.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	27.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	28.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	29.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	30.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	31.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	2	2	1	1	1	1	1	-	-	-	-	-	-	-	-	-	-	-

\* Trace of yellow line 5694A at 30°N.

Table 65b

Coronal observations at Climax, Colorado (6374A), west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90			
1953	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	6	6	4	2	1	1	1	-	-	-	-	1	1	1	1	1	1	2		
Oct	4.6	1	1	1	1	1	-	-	1	1	1	2	2	3	3	3	5	6	5	5	6	6	4	2	1	1	1	1	1	1	1	1	2				
	5.6	1	1	2	1	1	1	1	1	2	2	3	4	4	5	4	5	5	4	5	3	12	3	1	-	-	-	-	1	1	1	1	1				
	6.7	1	1	1	1	1	1	1	1	1	3	3	6	10	12	4	4	3	2	14	1	1	1	1	1	1	1	1	1	1	2	1	1				
	7.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	3	4	3	2	1	1	7	1	-	-	-	-	-	-	-	2		
	8.7	-	-	-	-	-	-	-	1	1	1	1	1	1	1	1	2	3	3	3	1	1	4	1	1	-	-	-	-	-	-	-	-	1			
	9.7	2	1	1	-	-	1	1	1	1	1	1	1	1	1	1	2	2	2	3	3	4	2	2	3	2	1	2	1	1	1	1	1				
	10.6	1	2	2	1	-	-	-	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
	11.7a	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
	15.7	3	3	2	2	2	2	1	1	1	2	2	2	2	3	5	9	17	4	2	3	2	1	1	1	1	1	1	1	1	1	1	1	1			
	16.6	2	2	1	1	1	1	1	1	1	1	2	5	5	12	12	12	5	2	2	2	1	1	1	1	1	1	1	1	1	1	1	-	-			
	17.8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
	18.6	1	1	1	-	2	1	1	1	1	1	1	1	1	1	1	1	3	5	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	20.8	2	1	1	1	1	1	1	1	1	1	4	3	1	1	2	3	2	3	2	3	3	3	2	2	2	2	X	X	-	-	-	-	2			
	22.8a	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	3	3	3	2	3	2	3	3	3	3	-	-	-	-	-	-	1			
	23.8a	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-
	24.7	2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-
	25.7	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-
	26.7a	-	-	-	-	-	-	-	-	-	-	3	3	2	2	2	3	2	2	2	2	3	2	2	2	4	2	1	-	-	-	1	1	1	1	1	
	27.8a	1	1	1	1	1	1	1	1	1	2	2	2	2	2	3	2	2	2	2	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-	-	-
	28.7	1	1	1	1	1	1	1	1	1	2	2	3	3	4	4	5	4	4	5	4	5	4	3	3	2	2	2	2	2	2	2	1	1	1		
	29.6	1	1	1	1	1	1	1	1	1	4	5	4	4	5	5	6	5	4	5	5	5	4	4	4	-	-	-	-	1	1	1	2	1	1		
	30.7	2	2	1	1	1	1	1	1	2	2	3	3	4	4	5	6	5	5	6	6	5	3	3	2	1	1	1	1	1	1	1	2	2			
	31.6	2	1	1	1	1	1	1	2	2	3	4	3	3	4	5	4	4	4	4	5	3	1	1	1	1	1	1	-	-	-	-	-	-			

Table 66a

Coronal observations at Climax, Colorado (6702A), east limb

Date GCT	Degrees north of the solar equator															0°	Degrees south of the solar equator																			
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1953	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Oct	4.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	5.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	7.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	8.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	9.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	10.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	11.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	15.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	16.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	17.8	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	X	X	X	X	X	X	X		
	18.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	20.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	3	2	2	1	-	-	-	-	-	-	-	-	-	-	-		
	22.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	23.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	24.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	25.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	26.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	27.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	28.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	29.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	30.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	31.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

Table 67a

Coronal observations at Sacramento Peak, New Mexico (5303A), east limb

Date GCT	Degrees north of the solar equator															0°	Degrees south of the solar equator																			
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1953	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Oct	1.7	-	2	2	-	2	2	3	4	5	6	5	4	4	4	4	5	7	6	5	4	4	4	5	4	3	3	2	2	2	-	-	-			
	2.7	-	-	-	3	2	2	3	5	6	5	4	3	4	4	4	5	12	13	14	7	7	5	4	3	3	3	2	-	-	2	2	-			
	3.7	-	-	-	-	2	2	2	3	3	3	3	2	2	4	5	7	8	6	3	3	3	4	2	3	2	2	3	-	-	-	-	-			
	4.7a	-	-	-	-	2	2	2	2	3	2	3	2	2	3	4	5	4	5	4	3	3	2	3	3	2	3	3	2	3	2	3	-			
	5.7a	-	-	-	-	2	2	2	2	3	3	3	3	3	2	2	3	3	3	2	3	3	3	2	2	3	3	3	3	2	3	2	3			
	6.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	7.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	8.7a	-	3	2	3	3	3	3	2	2	3	2	3	3	3	3	4	4	4	4	3	4	7	5	3	2	2	3	2	-	-	-	-			
	9.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	4	3	3	4	5	4	4	3	2	3	2	3	2	3	-	
	10.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	11.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	12.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	13.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	14.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	4	3	3	4	5	4	4	3	2	3	2	3	2	3	-		
	15.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	4	3	2	3	2	2	3	2	3	2	3	2	3	-		
	16.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	3	2	3	2	2	3	2	2	3	2	3	2	3	-		
	23.9	-	-	-	2	2	2	2	3	5	6	7	8	6	5	8	8	6	5	4	4	4	3	3	2	2	2	3	2	4	4	3	2			
	25.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	4	4	4	3	2	2	2	3	2	3	3	2	2	-	-		
	26.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	4	5	4	4	4	3	2	2	3	2	3	2	2	-		
	27.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	4	5	4	3	2	2	3	2	3	2	2	3	2	-		
	28.7	-	-	-	2	3	2	4	4	5	5	6	5	5	4	5	3	2	8	11	10	5	3	2	2	2	3	3	3	2	2	-				
	31.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	14	20	19	10	4	3	3	4	3	5	5	3	2	2	-			

Table 65b

### Coronal observations at Climax, Colorado (6702A), west limb

Table 67b

Coronal observations at Sacramento Peak, New Mexico (5303A), west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator																						
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90					
1953																																							
Oct	1.7	-	-	-	2	3	2	3	3	3	2	2	2	3	3	3	3	4	4	3	4	3	3	3	3	3	3	6	3	2	2	2	-	-	-				
	2.7	-	-	-	-	-	2	2	2	3	3	3	4	3	3	2	3	2	2	3	3	2	3	2	2	2	2	2	2	2	-	-	-	-	-				
	3.7a	-	-	-	-	-	-	-	-	2	2	2	2	2	3	3	4	3	2	2	2	2	4	4	3	3	2	3	3	3	2	2	-	-	-				
	4.7a	-	-	-	-	-	-	-	-	2	2	3	3	3	2	2	2	3	3	2	3	3	3	2	3	3	3	2	2	2	3	-	-	-	-				
	5.7a	-	-	-	-	-	-	-	-	2	2	3	3	2	3	3	2	2	2	-	3	8	13	12	5	4	4	3	2	-	-	-	-	-	-	-			
	6.7	-	-	-	-	-	-	-	-	2	2	3	3	2	2	2	3	3	4	2	2	5	10	16	17	10	6	4	3	3	2	-	-	-	-	-			
	7.7	-	-	-	-	-	-	-	-	2	2	3	3	2	3	2	3	3	3	8	4	3	11	14	22	32	23	11	8	6	5	4	4	4	3	3	2	-	
	8.7	-	-	-	-	-	-	-	-	2	2	3	3	3	2	3	2	3	2	3	3	4	5	11	12	5	4	3	3	4	3	2	2	2	-	-			
	9.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	3	4	5	6	5	4	3	4	5	4	3	-	-	-			
	10.7a	-	-	-	-	-	-	-	-	2	2	3	2	2	-	2	2	3	3	2	2	3	3	3	4	4	3	3	2	2	2	3	-	-	-				
	11.7a	-	-	-	-	-	-	-	-	-	2	3	3	3	2	2	2	3	3	2	2	2	2	3	3	3	3	3	3	3	2	4	4	3	-				
	12.8a	X	X	X	X	X	X	X	X	2	2	2	3	2	2	2	2	3	2	-	2	3	3	3	3	2	3	2	-	-	-	-	-	-	-				
	13.7a	-	-	-	-	-	-	-	-	3	3	3	3	4	5	3	3	2	3	2	2	4	5	3	4	3	4	3	4	3	2	2	2	3	-				
	14.7	-	-	-	-	-	-	-	-	2	2	2	3	2	2	3	4	4	4	5	5	7	13	14	6	5	5	8	7	6	6	5	5	8	8	7	3	2	-
	15.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	4	4	5	4	4	4	4	3	4	5	3	2	-	-	-		
	16.7	-	-	-	-	-	-	-	-	-	2	2	2	3	3	2	3	3	3	4	4	5	4	3	3	5	8	6	5	3	4	3	4	3	-	-	-		
	23.9	-	-	-	-	-	-	-	-	-	2	2	2	4	3	4	3	2	2	5	8	7	3	11	14	13	11	4	3	3	4	3	4	2	2	-	-	-	
	25.7	-	-	-	-	-	-	-	-	2	2	2	3	3	3	3	3	3	2	3	3	2	3	3	7	12	11	8	4	3	3	3	3	2	2	2	2	-	-
	26.7	-	-	-	-	-	-	-	-	2	2	2	3	3	3	3	2	2	3	3	2	2	3	4	5	5	4	3	2	3	3	3	4	3	2	-	-	-	
	27.7	2	-	-	-	-	-	-	-	2	2	2	3	4	4	4	3	3	4	3	2	2	3	3	2	2	3	3	3	2	2	-	-	-	-	-			
	28.7	-	-	-	-	-	-	-	-	-	2	2	2	3	3	2	2	2	3	3	3	3	3	3	2	2	3	3	3	2	2	2	2	-	-	-	-		
	31.7	-	-	-	-	-	-	-	-	2	2	3	2	2	2	3	3	3	3	3	3	5	7	8	6	5	4	4	3	4	5	5	4	3	2	2	-	-	-

Table 68a

Coronal observations at Sacramento Peak, New Mexico (6374A), east limb

Date GCT	Degrees north of the solar equator															0°	Degrees south of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1953																																					
Oct	1.7	4	3	4	3	2	2	2	-	-	2	2	3	4	2	3	4	4	4	5	5	8	7	8	11	5	4	3	3	2	3	3	-	2	2	-	2
	2.7	4	2	3	3	4	3	2	2	2	2	2	2	5	4	2	3	10	4	4	5	4	5	6	5	3	3	2	3	2	2	3	2	-	-	2	
	3.7	2	3	3	2	3	2	2	-	-	2	2	3	3	4	3	3	8	2	2	3	2	3	4	3	3	2	3	2	3	2	2	3	-	2	2	
	4.7a	2	2	2	2	3	2	3	2	-	2	2	2	3	2	3	4	5	3	2	3	4	4	4	3	3	2	-	2	2	3	-	-	-	-		
	5.7a	2	2	2	2	-	3	2	-	2	2	3	3	3	3	2	3	5	4	3	3	4	4	4	3	3	2	-	-	-	-	-	-	-	-	-	
	6.7a	3	3	2	3	3	2	-	2	2	2	3	4	3	3	3	3	2	2	2	5	4	4	4	4	3	4	3	2	-	-	-	-	-	-	3	
	7.7	3	2	3	3	2	2	3	-	2	2	4	5	7	5	4	3	2	4	5	5	5	6	6	4	3	2	2	-	-	-	-	-	-	-	-	2
	8.7a	-	-	-	-	-	-	-	-	-	2	3	3	2	3	2	3	2	2	2	2	2	8	4	3	2	2	3	3	2	-	-	-	-	-		
	9.7a	2	2	-	2	-	-	2	2	3	4	3	4	3	3	2	-	2	2	3	6	8	5	2	2	2	-	2	-	-	-	-	-	-	-	-	
	10.7	2	-	2	-	-	2	-	2	2	5	5	4	4	4	5	3	2	2	5	4	4	10	6	4	2	3	2	2	2	-	-	-	-	-	2	
	11.7a	3	3	2	3	-	2	2	2	-	3	5	4	3	4	5	4	3	4	4	5	8	7	6	5	3	2	2	2	-	-	-	-	-	-	-	
	12.8	-	3	3	2	2	-	-	2	2	5	4	5	3	5	6	5	11	15	5	5	8	7	4	5	5	4	3	2	2	X	X	X	X	X	X	
	13.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	2	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	14.7	2	4	3	3	2	2	-	-	2	3	2	2	3	5	7	13	14	12	11	8	5	5	4	6	5	5	5	3	-	2	2	3	-	2	2	
	15.7	3	4	2	2	3	2	2	2	-	3	-	2	3	5	7	8	7	7	6	5	5	4	4	5	5	2	2	-	2	-	2	2	-	2	2	
	16.7a	2	3	3	2	-	-	2	-	-	3	2	3	-	3	3	5	3	4	8	7	5	3	4	4	3	2	3	2	2	3	-	-	-	-	-	
	23.9	4	3	3	4	4	3	2	2	2	2	3	4	5	5	4	3	4	5	11	8	7	6	6	8	8	9	7	4	3	3	2	3	3	2	2	
	25.7	3	3	2	5	3	3	2	2	2	4	4	4	4	4	3	3	2	3	4	4	4	5	5	3	2	2	3	4	-	-	-	-	2			
	26.7	3	3	3	2	3	2	-	-	2	2	2	4	4	2	3	2	2	4	5	5	8	6	4	2	2	3	3	2	2	2	2	2	3			
	27.7	3	4	4	4	3	2	-	-	3	5	4	3	4	5	7	7	8	7	6	5	5	5	6	5	4	5	3	2	2	-	2	3	3	3		
	28.7	3	2	3	3	2	2	2	-	-	2	3	4	4	5	4	5	4	4	7	8	10	8	9	10	7	5	4	4	4	3	2	2	2	2	3	
	31.7	3	5	4	5	3	3	2	2	3	4	3	3	4	5	5	4	4	4	5	6	16	17	10	7	8	5	4	2	2	3	2	2	3	4	3	

Table 69a

Coronal observations at Sacramento Peak, New Mexico (6702A), east limb

Table 68b

### Coronal observations at Sacramento Peak, New Mexico (6374A), west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90			
1953																																					
Oct 1.7	2	2	3	2	3	2	3	2	2	2	3	3	6	6	5	5	5	7	8	5	6	8	7	4	5	5	4	3	3	2	2	3	2	3	4	4	4
2.7	2	3	2	2	3	3	2	2	3	4	5	4	3	4	5	5	12	11	8	7	6	5	5	4	4	4	3	2	2	2	3	4	4	4	4	4	
3.7a	2	3	2	2	-	-	2	-	2	2	3	4	3	3	4	4	5	4	4	5	X	4	4	3	5	2	2	-	2	3	-	2	2	2	2	2	
4.7a	-	2	-	-	-	-	-	-	-	3	3	4	3	3	4	4	5	4	5	6	8	7	5	2	2	2	3	-	2	2	-	2	2	2	2	2	
5.7a	-	2	3	3	2	2	2	2	-	2	2	3	3	3	4	4	5	6	5	6	8	7	5	2	2	2	3	-	2	-	2	2	2	2	2		
6.7	3	2	2	2	2	3	-	-	2	3	2	3	4	5	7	8	6	4	4	3	14	10	-	-	2	2	3	2	-	-	-	-	-	-	-	-	
7.7	2	3	2	3	-	3	2	-	-	3	2	2	4	5	10	9	7	4	4	3	14	16	-	2	-	2	-	-	-	3	2	-	2	2	3		
8.7	-	2	-	2	-	2	2	2	-	2	-	2	2	3	4	3	3	4	2	2	3	2	6	2	3	-	2	2	3	2	2	3	2	3	2		
9.7a	-	2	2	2	-	-	-	-	-	2	2	2	3	3	3	2	3	3	4	3	3	2	2	3	2	-	-	-	-	-	-	-	-	-	-	-	
10.7a	2	2	3	2	2	2	-	-	2	2	2	2	3	3	2	3	2	3	3	5	3	2	-	-	2	2	3	3	2	2	2	2	2	2	2		
11.7a	2	2	2	2	3	2	-	-	3	2	3	4	3	4	2	4	4	4	2	2	2	3	2	2	2	3	2	2	-	-	-	-	-	-	-		
12.8a	X	X	X	X	X	3	3	2	-	2	3	3	2	2	2	2	3	3	3	3	3	2	2	2	3	2	2	-	-	-	-	-	2	3	2		
13.7	-	-	-	-	-	2	2	2	3	3	4	2	3	2	-	2	3	4	4	5	4	4	3	-	4	3	4	-	-	-	-	-	-	-	-	-	
14.7	2	2	3	2	3	3	2	-	2	3	3	3	3	3	4	8	20	20	15	5	7	5	3	2	3	5	8	4	2	-	2	-	2	3	2		
15.7	2	3	3	3	3	2	-	-	2	-	3	2	3	3	4	5	8	20	16	5	4	3	2	3	2	3	3	2	2	3	2	2	3	3	2		
16.7	2	2	2	2	-	-	2	-	-	2	-	2	3	5	8	11	12	13	4	3	3	-	-	2	2	-	-	-	-	-	-	-	-	-	-		
23.9	3	4	3	2	3	3	2	3	2	2	4	8	9	8	8	7	7	5	6	5	6	5	4	4	4	7	8	6	6	4	3	2	2	4	3	4	4
25.7	2	2	2	-	2	3	2	-	-	3	3	4	5	5	5	7	7	6	5	6	5	4	4	4	3	6	4	4	4	3	2	2	3	2	3		
26.7	3	2	3	3	2	2	-	-	2	-	2	3	4	5	4	5	4	3	4	9	8	9	8	7	4	3	3	3	2	2	2	3	2	2	3		
27.7	3	3	4	3	3	3	2	3	5	2	5	5	6	6	11	7	7	8	7	10	11	12	10	8	5	4	5	4	3	3	2	2	3	3	3		
28.7	3	-	-	2	3	2	2	-	-	4	8	3	5	6	5	5	6	7	8	8	7	7	6	4	2	3	6	4	2	2	3	2	2	3	4	3	
31.7	3	4	3	2	2	3	2	2	3	8	9	8	7	10	12	11	13	12	11	10	7	4	3	4	4	5	3	3	2	2	2	2	2	2	3		

Table 69b

### Coronal observations at Sacramento Peak, New Mexico (6702A), west limb

Table 70Sudden Ionosphere Disturbances Observed at Washington, D. C.October 1953

1953 Day	GCT	Location of transmitters	Relative intensity at minimum*	Other phenomena
	Beginning End			
October 14	1426 1500	Ohio, Mexico, North Dakota	0.2	Terr. mag. pulse* 1423-1430 Solar flare*** 1420

\*Ratio of received field intensity during SID to average field intensity before and after, for station KQZXAU, (formerly W8XAL), 6080 kilocycles, 600 kilometers distant.

\*\*As observed on Cheltenham magnetogram of the United States Coast and Geodetic Survey.

\*\*\*Time of observation at Sacramento Peak, New Mexico, and at McMath-Hulbert Observatory, Pontiac, Michigan.

Table 71Sudden Ionosphere Disturbances Reported by Engineer-in-Chief,Cable and Wireless, Ltd., as Observed in England

1953 Day	GCT		Receiving station	Location of transmitters	Other phenomena
	Beginning	End			
October 14	0955	1005	Brentwood	Austria, Bahrein I., Barbados, Belgian Congo, Brazil, Greece, India, Iraq, Kenya, New York, Palestine, Portugal, Southern Rhodesia, Spain, Switzerland, Syria, Turkey, U.S.S.R., Yugo- slavia, Zanzibar	Solar flare* 1012
14	0954	1010	Somerton	China, Cyprus, Egypt, India, Iran, Nigeria, Thailand, Union of South Africa	Solar flare* 1012

\*Time of observation at Wendelstein Observatory, Germany. Flare began before time of observation.

Table 72Sudden Ionosphere Disturbances Reported by the Netherlands Postal and Telecommunication Services, as Observed at Nederhorst den Berg, Netherlands

Day	GCT		Location of transmitters	Other phenomena
	Beginning	End		
1952 July 28 November 15 22	1040	1050	Surinam	
	1256	1305	Surinam	
	1046	1100	Surinam, Egypt	
	1155	1320	Egypt, Surinam	
1953 March 31 May 4 5 August 11	0931	1045	Washington, Egypt, Surinam	
	0445	0530	Egypt	
	1537	1555	Surinam	Solar flare* 1536
				Solar flare** 1538

\*Time of observation at McMath-Hulbert Observatory, Pontiac, Michigan.

\*\*Time of observation at Sacramento Peak, New Mexico.

Note: Observers are invited to send to the CRPL information on times of beginning and end of sudden ionosphere disturbances for publication as above. Address letters to the Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

Table 73  
Solar Flares, October 1953

Observe- tory	Date	Time Observed		Dura- tion	Position		Time of Maxi- mum	Int. ensity of radiation	Rela- tive area	Import- ance	SID Obser- ved
		Begin- ning	End- ing		Latit- ude (of Visible)	Long- itude Diff.					
		(GCT)	(GCT)	(Min.)	(Hemisph.)	(Deg.)	(Deg.)	(GCT)	(Tenths)		
Wendel.	Oct. 14	1012B	1024A	App. 12	120	S07	W35	1012	1	1	1425
Sac. Peak	14	1420	1433	13	178	S08	W38	1425	1	1	1425
McMath	14	1420	1530	70		S09	W39	---	1	1	
Sac. Peak	31	1600	1800	120	40	S08	E69	1640	10	5	1 -

Wendel. = Wendelstein.  
Sac. Peak ≈ Sacramento Peak.





B Flare began before given time.  
A Flare ended after given time.  
Q Time reported as questionable.

Table 74

## Indices of Geomagnetic Activity for September 1953

Preliminary values of international character-figures, C;  
Geomagnetic planetary three-hour-range indices, Kp;  
Magnetically selected quiet and disturbed days

## GRAPHS OF IONOSPHERIC DATA






Fig. 5. FAIRBANKS, ALASKA  
64.9°N, 147.8°W SEPTEMBER 1953










Fig. 7. ANCHORAGE, ALASKA  
61.2°N, 149.9°W SEPTEMBER 1953








Fig. 13. UPSALA, SWEDEN  
59.8°N, 17.6°E SEPTEMBER 1953




Fig. 14. UPSALA, SWEDEN SEPTEMBER 1953




Fig. 15. SAN FRANCISCO, CALIFORNIA  
37.4°N, 122.2°W SEPTEMBER 1953
















Fig. 16. SAN FRANCISCO, CALIFORNIA SEPTEMBER 1953










Fig. 29. RESOLUTE BAY, CANADA  
74.7°N, 94.9°W AUGUST 1953




Fig. 30. RESOLUTE BAY, CANADA AUGUST 1953




Fig. 31. BAKER LAKE, CANADA  
64.3°N, 96.0°W AUGUST 1953




Fig. 32. BAKER LAKE, CANADA AUGUST 1953




Fig. 33. CHURCHILL, CANADA

58.8°N, 94.2°W

AUGUST 1953




Fig. 34. CHURCHILL, CANADA

AUGUST 1953




Fig. 35. FORT CHIMO, CANADA

58.1°N, 68.3°W

AUGUST 1953




Fig. 36. FORT CHIMO, CANADA

AUGUST 1953




Fig. 37. PRINCE RUPERT, CANADA  
54.3°N, 130.3°W AUGUST 1953




Fig. 38. PRINCE RUPERT, CANADA AUGUST 1953




Fig. 39. DE BILT, HOLLAND  
52.1°N, 5.2°E AUGUST 1953




Fig. 40. DE BILT, HOLLAND AUGUST 1953




Fig. 41. WINNIPEG, CANADA  
49.9°N, 97.4°W AUGUST 1953





Fig. 43. ST. JOHN'S, NEWFOUNDLAND  
47.6°N, 52.7°W AUGUST 1953








Fig. 44. ST. JOHN'S, NEWFOUNDLAND






Fig. 49. BATON ROUGE, LOUISIANA  
30.5°N, 91.2°W

AUGUST 1953




Fig. 51. LEOPOLDVILLE, BELGIAN CONGO  
4.3°S, 153°E

AUGUST 1953




Fig. 50. BATON ROUGE, LOUISIANA

AUGUST 1953




Fig. 52. LEOPOLDVILLE, BELGIAN CONGO

AUGUST 1953




Fig. 53. BUENOS AIRES, ARGENTINA  
34.5° S, 58.5° W AUGUST 1953




Fig. 54. BUENOS AIRES, ARGENTINA




Fig. 55. DECEPTION I.  
63.0° S, 60.7° W AUGUST 1953
















Fig. 56. DECEPTION I. AUGUST 1953










Fig. 69. OTTAWA, CANADA  
45.4°N, 75.9°W

JULY 1953




Fig. 70. OTTAWA, CANADA

JULY 1953




Fig. 71. CALCUTTA, INDIA  
22.6°N, 88.4°E

JULY 1953








Fig. 72. CALCUTTA, INDIA

JULY 1953






Fig. 77. POINT BARROW, ALASKA  
71.3°N, 156.8°W JUNE 1953




Fig. 78. POINT BARROW, ALASKA JUNE 1953




Fig. 79. WAKKANAI, JAPAN  
45.4°N, 141.7°E JUNE 1953




Fig. 80. WAKKANAI, JAPAN JUNE 1953




Fig. 81. AKITA, JAPAN  
39.7°N, 140.1°E

JUNE 1953




Fig. 82. AKITA, JAPAN

JUNE 1953




Fig. 83. TOKYO, JAPAN  
35.7°N, 139.5°E

JUNE 1953




Fig. 84. TOKYO, JAPAN

JUNE 1953




Fig. 85. YAMAGAWA, JAPAN  
31.2°N, 130.6°E JUNE 1953




Fig. 86. YAMAGAWA, JAPAN JUNE 1953



Fig. 87. CALCUTTA, INDIA  
22.6°N, 88.4°E JUNE 1953




Fig. 88. CALCUTTA, INDIA JUNE 1953




Fig. 89 RAROTONGA I.

21.3°S, 159.8°W

JUNE 1953




Fig. 90 RAROTONGA I.

JUNE 1953




Fig. 91. CHRISTCHURCH, NEW ZEALAND

43.6°S, 172.7°E

JUNE 1953








Fig. 92. CHRISTCHURCH, NEW ZEALAND

JUNE 1953

NBS 490



Index of Tables and Graphs of Ionospheric Datain CRPL-F11

	<u>Table page</u>	<u>Figure page</u>
Akita, Japan		
June 1953 . . . . .	18	66
Anchorage, Alaska		
September 1953 . . . . .	12	47
Baguio, P. I.		
July 1953 . . . . .	18	64
Baker Lake, Canada		
August 1953 . . . . .	14	53
July 1953 . . . . .	16	60
Baton Rouge, Louisiana		
August 1953 . . . . .	16	58
Buenos Aires, Argentina		
August 1953 . . . . .	16	59
Calcutta, India		
July 1953 . . . . .	17	63
June 1953 . . . . .	19	67
Christchurch, New Zealand		
June 1953 . . . . .	19	68
May 1953 . . . . .	19	69
Churchill, Canada		
August 1953 . . . . .	14	54
De Bilt, Holland		
August 1953 . . . . .	15	55
Deception I.		
August 1953 . . . . .	16	59
Fairbanks, Alaska		
September 1953 . . . . .	12	47
Fort Chimo, Canada		
August 1953 . . . . .	14	54
July 1953 . . . . .	17	61
Guam I.		
September 1953 . . . . .	14	52
Leopoldville, Belgian Congo		
August 1953 . . . . .	16	58
Maui, Hawaii		
September 1953 . . . . .	13	51
Narsarssuak, Greenland		
September 1953 . . . . .	12	48
Okinawa I.		
September 1953 . . . . .	13	50
Oslo, Norway		
September 1953 . . . . .	12	48

Index (CRPL-III, concluded)

	<u>Table page</u>	<u>Figure page</u>
Ottawa, Canada		
August 1953 . . . . .	15	57
July 1953 . . . . .	17	63
Panama Canal Zone		
September 1953 . . . . .	14	52
Point Barrow, Alaska		
June 1953 . . . . .	18	65
Prince Rupert, Canada		
August 1953 . . . . .	15	55
July 1953 . . . . .	17	61
Puerto Rico, W. I.		
September 1953 . . . . .	13	51
Rarotonga I.		
June 1953 . . . . .	19	68
May 1953 . . . . .	19	69
Resolute Bay, Canada		
August 1953 . . . . .	14	53
July 1953 . . . . .	16	60
St. John's, Newfoundland		
August 1953 . . . . .	15	56
July 1953 . . . . .	17	62
San Francisco, California		
September 1953 . . . . .	13	49
Schwarzenburg, Switzerland		
August 1953 . . . . .	15	57
Tokyo, Japan		
June 1953 . . . . .	18	66
Tromso, Norway		
September 1953 . . . . .	12	46
Upsala, Sweden		
September 1953 . . . . .	13	49
Wakkanai, Japan		
June 1953 . . . . .	18	65
Washington, D. C.		
October 1953 . . . . .	12	46
Watheroo, W. Australia		
July 1953 . . . . .	18	64
White Sands, New Mexico		
September 1953 . . . . .	13	50
Winnipeg, Canada		
August 1953 . . . . .	15	56
July 1953 . . . . .	17	62
Yanagawa, Japan		
June 1953 . . . . .	19	67



## CRPL and IRPL Reports

[A list of CRPL Section Reports is available from the Central Radio Propagation Laboratory upon request]

### Daily:

Radio disturbance forecasts, every half hour from broadcast station WWV of the National Bureau of Standards. Telephoned and telegraphed reports of ionospheric, solar, geomagnetic, and radio propagation data.

### Semiweekly:

- CRPL—J. North Atlantic Radio Propagation Forecast (of days most likely to be disturbed during following month).  
CRPL—Jp. North Pacific Radio Propagation Forecast (of days most likely to be disturbed during following month).

### Semimonthly:

- CRPL—Ja. Semimonthly Frequency Revision Factors For CRPL Basic Radio Propagation Prediction Reports.

### Monthly:

- CRPL—D. Basic Radio Propagation Predictions—Three months in advance. (Dept. of the Army, TB 11-499-, monthly supplements to TM 11-499; Dept. of the Navy, DNC 13 ( ) series; Dept. of the Air Force, TO 16-1B-2 series.)  
CRPL—F. Ionospheric Data.  
\*IRPL—A. Recommended Frequency Bands for Ships and Aircraft in the Atlantic and Pacific.  
\*IRPL—H. Frequency Guide for Operating Personnel.

### Circulars of the National Bureau of Standards:

NBS Circular 462. Ionospheric Radio Propagation.

NBS Circular 465. Instructions for the Use of Basic Radio Propagation Predictions.

### Reports issued in past:

IRPL—C61. Report of the International Radio Propagation Conference, 17 April to 5 May 1944.

IRPL—G1 through G12. Correlation of D. F. Errors With Ionospheric Conditions.  
(G1, G3, available. Others out of print; see second footnote.)

IRPL—R. Nonscheduled reports:

- R4. Methods Used by IRPL for the Prediction of Ionosphere Characteristics and Maximum Usable Frequencies.  
R5. Criteria for Ionospheric Storminess.  
\*\*R6. Experimental Studies of Ionospheric Propagation as Applied to the Loran System.  
R7. Second Report on Experimental Studies of Ionospheric Propagation as Applied to the Loran System.  
R9. An Automatic Instantaneous Indicator of Skip Distance and MUF.  
R10. A Proposal for the Use of Rockets for the Study of the Ionosphere.  
\*\*R11. A Nomographic Method for both Prediction and Observation Correlation of Ionosphere Characteristics.  
\*\*R12. Short Time Variations in Ionosphere Characteristics.  
R14. A Graphical Method for Calculating Ground Reflection Coefficients.  
\*\*R15. Predicted Limits for F2-Layer Radio Transmission Throughout the Solar Cycle.  
\*\*R17. Japanese Ionospheric Data—1943.  
R18. Comparison of Geomagnetic Records and North Atlantic Radio Propagation Quality Figures—October 1943 Through May 1945.  
\*\*R21. Notes on the Preparation of Skip-Distance and MUF Charts for Use by Direction-Finder Stations.  
(For distances out to 4000 km.)  
\*\*R23. Solar-Cycle Data for Correlation with Radio Propagation Phenomena.  
\*\*R24. Relations Between Band Width, Pulse Shape and Usefulness of Pulses in the Loran System.  
\*\*R25. The Prediction of Solar Activity as a Basis for the Prediction of Radio Propagation Phenomena.  
\*\*R26. The Ionosphere as a Measure of Solar Activity.  
R27. Relationships Between Radio Propagation Disturbance and Central Meridian Passage of Sunspots Grouped by Distance From Center of Disc.  
\*\*R30. Disturbance Rating in Values of IRPL Quality-Figure Scale from A. T. & T. Co. Transmission Disturbance Reports to Replace T. D. Figures as Reported.  
\*\*R31. North Atlantic Radio Propagation Disturbances, October 1943 Through October 1945.  
\*\*R33. Ionospheric Data on File at IRPL.  
\*\*R34. The Interpretation of Recorded Values of fEs.  
\*\*R35. Comparison of Percentage of Total Time of Second-Multiple Es Reflections and That of fEs in Excess of 3 Mc.

IRPL—T. Reports on tropospheric propagation:

- T1. Radar operation and weather. (Superseded by JANP 101.)  
T2. Radar coverage and weather. (Superseded by JANP 102.)

CRPL—T3. Tropospheric Propagation and Radio-Meteorology. (Reissue of Columbia Wave Propagation Group WPG—5.)

\*Items bearing this symbol are distributed only by U. S. Navy. They are issued under one cover as the DNC 14 ( ) Series.

\*\*Out of print; information concerning cost of photostat or microfilm copies is available from CRPL upon request.

