

National Bureau of Standards
Library, N. W. Bldg.

8 MAR 1954

Reference is not to be
taken

IONOSPHERIC DATA

ISSUED
FEBRUARY 1954

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
CENTRAL RADIO PROPAGATION LABORATORY
WASHINGTON, D. C.

IONOSPHERIC DATA

CONTENTS

	<u>Page</u>
Symbols, Terminology, Conventions	2
World-Wide Sources of Ionospheric Data	5
Hourly Ionospheric Data at Washington, D. C.	7, 12, 20, 44
Ionospheric Storminess at Washington, D. C. .	7, 32
Sudden Ionosphere Disturbances	7, 33
Radio Propagation Quality Figures	8, 34
Observations of the Solar Corona	9, 36
Relative Sunspot Numbers	10, 40
Observations of Solar Flares	10, 42
Indices of Geomagnetic Activity	11, 43
Erratum	11
Tables of Ionospheric Data	12
Graphs of Ionospheric Data	44
Index of Tables and Graphs of Ionospheric Data in CRPL-F114	68

SYMBOLS, TERMINOLOGY, CONVENTIONS

Beginning with data reported for January 1952, the symbols, terminology, and conventions for the determination of median values used in this report (CRPL-F series) conform as far as practicable to those adopted at the Sixth Meeting of the International Radio Consultative Committee (C.C.I.R.) in Geneva, 1951. Excerpts concerning symbols and terminology from Document No. 626-E of this Meeting are given on pages 2-7 of the report CRPL-F89, "Ionospheric Data," issued January 1952. Reprints of these pages are available upon request.

Beginning with data for January 1945, median values are published wherever possible. Where averages are reported, they are, at any hour, the average for all the days during the month for which numerical data exist.

The following conventions are used in determining the medians for hours when no measured values are given because of equipment limitations and ionospheric irregularities. Symbols used are those given in Document No. 626-E referred to above.

a. For all ionospheric characteristics:

Values missing because of A, C, F, L, M, N, Q, S, or T are omitted from the median count.

b. For critical frequencies and virtual heights:

Values of f_{oF2} (and f_{oE} near sunrise and sunset) missing because of E are counted as equal to or less than the lower limit of the recorder. Values of $h'F2$ (and $h'E$ near sunrise and sunset) missing for this reason are counted usually as equal to or greater than the median. Other characteristics missing because of E are omitted from the median count.

Values missing because of D are counted as equal to or greater than the upper limit of the recorder.

Values missing because of G are counted:

1. For f_{oF2} , as equal to or less than f_{oF1} .
2. For $h'F2$, as equal to or greater than the median.

The symbol W is included in the median count only when it replaces a height characteristic. This practice represents a change from that listed in issues previous to CRPL-F78.

Values missing for any other reason are omitted from the median count.

c. For MUF factor (M-factors):

Values missing because of G or W are counted as equal to or less than the median.

Values missing for any other reason are omitted from the median count.

d. For sporadic E (E_s):

Values of fEs missing because of E or G (and B when applied to the daytime E region only) are counted as equal to or less than the median foE, or equal to or less than the lower frequency limit of the recorder.

Values of fEs missing for any other reason, and values of h'Es missing for any reason at all are omitted from the median count.

Beginning with data for November 1945, doubtful monthly median values for ionospheric observations at Washington, D. C., are indicated by parentheses, in accordance with the practice already in use for doubtful hourly values. The following are the conventions used to determine whether or not a median value is doubtful:

1. If only four values or less are available, the data are considered insufficient and no median value is computed.

2. For the F2 layer, if only five to nine values are available, the median is considered doubtful. The E and F1 layers are so regular in their characteristics that, as long as there are at least five values, the median is not considered doubtful.

3. For all layers, if more than half of the values used to compute the median are doubtful (either doubtful or interpolated), the median is considered doubtful.

The same conventions are used by the CRPL in computing the medians from tabulations of daily and hourly data for stations other than Washington, beginning with the tables in IRPL-F18.

The tables and graphs of ionospheric data are correct for the values reported to the CRPL, but, because of variations in practice in the interpretation of records and scaling and manner of reporting of values, may at times give an erroneous conception of typical ionospheric characteristics at the station. Some of the errors are due to:

- a. Differences in scaling records when spread echoes are present.
- b. Omission of values when f_{oF2} is less than or equal to f_{oF1} , leading to erroneously high values of monthly averages or median values.
- c. Omission of values when critical frequencies are less than the lower frequency limit of the recorder, also leading to erroneously high values of monthly average or median values.

These effects were discussed on pages 6 and 7 of the previous F-series report IRPL-F5.

Ordinarily, a blank space in the f_{Es} column of a table is the result of the fact that a majority of the readings for the month are below the lower limit of the recorder or less than the corresponding values of f_{oE} . Blank spaces at the beginning and end of columns of $h'F_1$, f_{oF1} , $h'E$, and f_{oE} are usually the result of diurnal variation in these characteristics. Complete absence of medians of $h'F_1$ and f_{oF1} is usually the result of seasonal effects.

The dashed-line prediction curves of the graphs of ionospheric data are obtained from the predicted zero-muf contour charts of the CRPL-D series publications. The following points are worthy of note:

- a. Predictions for individual stations used to construct the charts may be more accurate than the values read from the charts since some smoothing of the contours is necessary to allow for the longitude effect within a zone. Thus, inasmuch as the predicted contours are for the center of each zone, part of the discrepancy between the predicted and observed values as given in the F series may be caused by the fact that the station is not centrally located within the zone.
- b. The final presentation of the predictions is dependent upon the latest available ionospheric and radio propagation data, as well as upon predicted sunspot number.

c. There is no indication on the graphs of the relative reliability of the data; it is necessary to consult the tables for such information.

The following predicted smoothed 12-month running-average Zürich sunspot numbers were used in constructing the contour charts:

Month	Predicted Sunspot Number									
	1954	1953	1952	1951	1950	1949	1948	1947	1946	1945
December	15	33	53	86	108	114	126	85	38	
November	16	38	52	87	112	115	124	83	36	
October	17	43	52	90	114	116	119	81	23	
September	18	46	54	91	115	117	121	79	22	
August	18	49	57	96	111	123	122	77	20	
July	20	51	60	101	108	125	116	73		
June	21	52	63	103	108	129	112	67		
May	22	52	68	102	108	130	109	67		
April	24	52	74	101	109	133	107	62		
March	27	52	78	103	111	133	105	51		
February	29	51	82	103	113	133	90	46		
January	14	30	53	85	105	112	130	88	42	

WORLD - WIDE SOURCES OF IONOSPHERIC DATA

The ionospheric data given here in tables 1 to 48 and figures 1 to 96 were assembled by the Central Radio Propagation Laboratory for analysis and correlation, incidental to CRPL prediction of radio propagation conditions. The data are median values unless otherwise indicated. The following are the sources of the data in this issue:

República Argentina, Ministerio de Marina:
 Buenos Aires, Argentina
 Deception I.

University of Graz:
 Graz, Austria

Meteorological Service of the Belgian Congo and Ruanda-Urundi:
 Leopoldville, Belgian Congo

Defence Research Board, Canada:

Baker Lake, Canada
Churchill, Canada
Ottawa, Canada
Prince Rupert, Canada
Resolute Bay, Canada
St. John's, Newfoundland
Winnipeg, Canada

Radio Wave Research Laboratories, National Taiwan University, Taipeh,

Formosa, China:
Formosa, China

Institute for Ionospheric Research, Lindau Uber Northeim, Hannover, Germany:
Lindau/Harz, Germany

The Royal Netherlands Meteorological Institute:

De Bilt, Holland

Icelandic Post and Telegraph Administration:

Reykjavik, Iceland

Radio Research Laboratories, Tokyo, Japan:

Akita, Japan
Tokyo (Kokubunji), Japan
Wakkanai, Japan
Yamagawa, Japan

Norwegian Defence Research Establishment, Kjeller per Lillestrom, Norway:

Oslo, Norway
Tromso, Norway

South African Council for Scientific and Industrial Research:

Capetown, Union of South Africa
Johannesburg, Union of South Africa

Research Laboratory of Electronics, Chalmers University of Technology,

Gothenburg, Sweden:
Kiruna, Sweden

Research Institute of National Defence, Stockholm, Sweden:

Upsala, Sweden

Royal Board of Swedish Telegraphs, Radio Department, Stockholm, Sweden:

Lulea, Sweden

Post, Telephone and Telegraph Administration, Berne, Switzerland:

Schwarzenburg, Switzerland

United States Army Signal Corps:
Okinawa I.
White Sands, New Mexico

National Bureau of Standards (Central Radio Propagation Laboratory):
Anchorage, Alaska
Fairbanks, Alaska (Geophysical Institute of the University of Alaska)
Guam I.
Huancayo, Peru (Instituto Geofisico de Huancayo)
Maui, Hawaii
Narsarssauk, Greenland
Panama Canal Zone
Point Barrow, Alaska
Puerto Rico, W. I.
San Francisco, California (Stanford University)
Washington, D. C.

HOURLY IONOSPHERIC DATA AT WASHINGTON, D. C.

The data given in tables 49 through 60 follow the scaling practices given in the report IRPL-C61, "Report of International Radio Propagation Conference," pages 36 to 39, and the median values are determined by the conventions given above under "Symbols, Terminology, Conventions." Beginning with September 1949, the data are taken at Ft. Belvoir, Virginia.

IONOSPHERIC STORMINESS AT WASHINGTON, D.C.

Table 61 presents ionosphere character figures for Washington, D. C., during January 1954, as determined by the criteria given in the report IRPL-R5, "Criteria for Ionospheric Storminess," together with Cheltenham, Maryland, geomagnetic K-figures, which are usually covariant with them.

SUDDEN IONOSPHERE DISTURBANCES

Table 62 shows that no sudden ionosphere disturbances were observed at Ft. Belvoir, Virginia, during the month of January 1954. Table 63 lists the sudden ionosphere disturbances observed in Sweden for October 1953.

RADIO PROPAGATION QUALITY FIGURES

Tables 64a and 64b give for December 1953: the radio propagation quality figures for the North Atlantic area, CRPL advance and short-term forecasts, a summary geomagnetic activity index and sundry comparisons, specifically as follows:

- (a) radio propagation quality figures, separately for each 6-hour interval of the Greenwich day, viz., 00-06, 06-12, 12-18, 18-24 hours UT (Universal Time or GCT).
- (b) whole-day radio quality indices (beginning October 1952). Each index is a weighted average of the four quarter-day Q-figures, before rounding off, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which designate whenever possible the days when significant disturbance or unusually quiet conditions will occur.
- (c) short-term forecasts, issued by CRPL every six hours (nominally one hour before 00^h, 06^h, 12^h, 18^h UT) and applicable to the period 1 to 13 (especially 1 to 7) hours ahead. Note that new scoring rules have been adopted beginning with October 1952 data.
- (d) advance forecasts, issued semiweekly (CRPL-J reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole-day quality indices.
- (e) half-day averages of the geomagnetic X indices measured by the Cheltenham Magnetic Observatory of the U. S. Coast and Geodetic Survey.
- (f) illustration of the comparison of short-term forecasts with Q-figures and also with estimates of radio quality based on CRPL observations only.
- (g) illustration of the outcome of advance forecasts (1 to 3 or 4 days ahead) and, for comparison, the outcome of a type of "blind" forecast. For the latter the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

The radio propagation quality figures are prepared from radio traffic data reported to CRPL by American Telephone and Telegraph Company, Mackay Radio and Telegraph Company, RCA Communications, Inc., Marconi Company, British Admiralty Signal and Radar Establishment, and the following agencies of the U. S. Government:--Coast Guard, Navy, Army Signal Corps, and State Department. The method of calculation, summarized below, is similar to that described in a 1946 report, IRPL-R31, now out of print. Beginning with recalculated figures for January 1952, only reports of radio transmission on North Atlantic paths closely approximating New York-London are included in the estimation of quality. Observations of selected ionospheric characteristics, even though strongly correlated with radio transmission quality, and traffic reports for paths such as New York-Stockholm or New York-Tangier, previously included in the quality-figure determination with low weight, have been left out of the present calculations inasmuch as a sufficient number of homogeneous reports are now available.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the quality scale of the original reports. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year.

with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. The 6-hourly quality figures are (subjectively) weighted means of the reports received for that period. These 6-hourly quality figures replace, beginning January 1953, the half-daily quality figures which formerly appeared in this table.

These quality figures are, in effect, a consensus of reported radio propagation conditions in the North Atlantic area. The reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality because of multipath, interference, etc. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

Note. The North Pacific quality figures, which were published through October 1951, have been temporarily discontinued. Since the establishment of the North Pacific Radio Warning Service at Anchorage, Alaska, a larger number of reports are being received than were previously available in Washington. The preparation of the quality figures will be resumed when sufficient data have been accumulated for determination of conversion tables for these new reports.

OBSERVATIONS OF THE SOLAR CORONA

Tables 65 through 67 give the observations of the solar corona during January 1954, obtained at Climax, Colorado, by the High Altitude Observatory of Harvard University and the University of Colorado. Tables 68 through 70 list the coronal observations obtained at Sacramento Peak, New Mexico, during January 1954, derived by Harvard College Observatory as a part of its performance of a research contract with the Upper Air Research Observatory, Geophysical Research Directorate, Air Force Cambridge Research Center. The data are listed separately for east and west limbs at 5-degree intervals of position angle north and south of the Solar Equator at the limb. The time of observation is given to the nearest tenth of a day, GCT.

Table 65 gives the intensities of the green (5303A) line of the emission spectrum of the solar corona; table 66 gives similarly the intensities of the first red (6374A) coronal line; and table 67, the intensities of the second red (6702A) coronal line; all observed at Climax in January 1954.

Table 68 gives the intensities of the green (5303A) coronal line; table 69, the intensities of the first red (6374A) coronal line; and table 70, the intensities of the second red (6702A) coronal line; all observed at Sacramento Peak in January 1954.

The following symbols are used in tables 65 through 70: a, observation of low weight; -, corona not visible; and X, position angle not included in plate estimates.

Attention is called to the publication of "Particulars of Observations" at Climax, Colorado, and Sacramento Peak, New Mexico, for July through December 1953 in tables 71 and 72 of CRPL-F113. Mention of these tables was inadvertently omitted in last month's issue.

RELATIVE SUNSPOT NUMBERS

Table 71 lists the daily provisional Zürich relative sunspot number, R_Z , for January 1954, as communicated by the Swiss Federal Observatory. Table 72 contains the daily American relative sunspot number, R_A , for December 1953, as compiled by the Solar Division, American Association of Variable Star Observers.

OBSERVATIONS OF SOLAR FLARES

Table 73 gives the preliminary record of solar flares reported to the CRPL. These reports are communicated on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete records are published later in the Quarterly Bulletin of Solar Activity, I.A.U., in various observatory publications, and elsewhere. The present listing serves to identify and roughly describe the phenomena observed. Details should be sought from the reporting observatory.

Reporting directly to the CRPL are the following observatories: Mt. Wilson, McMath-Hulbert, U. S. Naval, Wendelstein, Kanzel and High Altitude at Sacramento Peak, New Mexico. The remainder report to Meudon (Paris) and the data are taken from the Paris-URSIgram broadcast, monitored fairly regularly by the CRPL. The data on solar flares reported from Sacramento Peak, New Mexico, communicated by the High Altitude Observatory at Boulder, Colorado, are provided by Harvard University as the result of work undertaken on an Air Materiel Command Research and Development Contract administered by the Air Force Cambridge Research Laboratories.

The table lists for each flare the reporting observatory, date, times of beginning and ending of observation, duration (when known), total area (corrected for foreshortening), and heliographic coordinates. For the maximum phase of the flare is given the time, intensity, area relative to the total area, and the importance. The column "SID observed" is to indicate when a sudden ionosphere disturbance, noted elsewhere in these reports, occurred at the time of a flare. Times are in Universal Time (GCT).

INDICES OF GEOMAGNETIC ACTIVITY

Table 74 lists various indices of geomagnetic activity based on data from magnetic observatories widely distributed throughout the world. The indices are: (1) preliminary international character-figures, C; (2) geomagnetic planetary three-hour-range indices, K_p; (3) magnetically selected quiet and disturbed days.

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm). The magnetically quiet and disturbed days are selected by the international scheme outlined on pages 219-227 in the December 1943 issue of Terrestrial Magnetism and Atmospheric Electricity. The details of the currently used method follow. For each day of a month, its geomagnetic activity is assigned by weighting equally the following three criteria: (1) the sum of the eight K_p's; (2) the greatest K_p; and (3) the sum of the squares of the eight K_p's.

K_p is the mean standardized K-index from 11 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g., 5- is $4 \frac{2}{3}$, 50 is $5 \frac{0}{3}$, and 5+ is $5 \frac{1}{3}$. This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of K_p has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948," published in Washington, D. C., 1949, by the Association of Terrestrial Magnetism and Electricity, International Union of Geodesy and Geophysics. K_p is available from 1937 to date as noted in F108.

The Committee on Characterization of Magnetic Disturbance, ATME, IUGG, has kindly supplied this table. The Meteorological Office, De Bilt, Holland, collects the data and compiles C and selected days. The Chairman of the Committee computes the planetary index. Current tables are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

ERRATUM

The De Bilt, Holland, height values published in the F series, numbers 104 through 113, for January 1953 through October 1953, should be increased by 5 percent.

TABLES OF IONOSPHERIC DATA

Table 1

Time	h'F2	foF2	h'F1	foF1	h'E	fOE	fEs	(M3000)F2
00	(270)	(2.4)						(3.1)
01	(260)	(2.3)						(3.1)
02	260	2.8						(3.1)
03	250	(2.8)						(3.1)
04	250	2.9						3.2
05	250	3.1						3.2
06	240	(3.0)						(3.0)
07	230	2.9						3.4
08	220	4.3	210	—	—	1.7		3.6
09	230	4.8	220	—	—	1.7	2.4	3.6
10	250	5.3	210	3.6	110	2.5		3.5
11	250	5.7	210	3.7	110	2.7		3.5
12	250	5.9	210	3.8	110	2.8		3.5
13	260	5.8	210	3.8	110	2.8		3.5
14	260	5.6	210	3.6	110	2.6		3.5
15	250	5.6	210	—	—	2.4		3.5
16	230	5.5	220	—	—	1.7	1.9	3.6
17	220	4.8						3.5
18	220	3.8						3.4
19	230	3.2						3.3
20	230	2.7						3.4
21	(250)	2.1						3.2
22	(270)	2.2						3.1
23	(270)	(2.3)						3.1

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3

Time	h'F2	foF2	h'F1	foF1	h'E	fOE	fEs	(M3000)F2
00	280	(2.6)						4.3 (3.0)
01	330	(2.2)						4.4 (3.0)
02	330	(2.2)						5.7 (2.9)
03	360	(2.1)						4.5 (2.8)
04	335	(2.2)						5.0 (2.9)
05	320	(2.4)						4.8 (2.8)
06	320	(2.0)						4.1 (2.8)
07	300	(2.0)						4.2 (3.0)
08	320	(1.9)						4.3 (2.9)
09	250	2.6						3.2 3.2
10	230	3.6						3.0 3.4
11	220	4.1						1.9 3.4
12	220	4.2						2.4 3.4
13	220	4.5						3.9 3.5
14	210	4.1						2.7 3.5
15	210	3.6						2.6 3.4
16	220	2.7						3.1 3.4
17	250	2.0						3.6 3.3
18	200	(1.6)						4.4 (3.1)
19	340	(1.8)						5.0 (2.6)
20	—	—						4.4 —
21	310	—						4.7 —
22	270	(2.8)						3.4 (3.2)
23	280	2.6						4.0 3.1

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 5

Time	h'F2	foF2	h'F1	foF1	h'E	fOE	fEs	(M3000)F2
00	—	—						5.0 —
01	—	—						4.9 —
02	—	—						6.0 —
03	—	—						6.0 —
04	—	—						5.1 —
05	(300)	(2.4)						6.0 (3.2)
06	(280)	(1.8)						4.4 (3.1)
07	(270)	(1.6)						4.4 (3.5)
08	250	(1.8)						2.4 3.3
09	220	3.0	—	—	110	—		2.0 3.6
10	230	3.8	220	—	140	—		3.6 3.6
11	230	4.3	210	—	130	1.7		3.6 3.6
12	230	4.7	220	—	140	1.8		3.6 3.6
13	230	4.7	220	—	130	1.6		3.6 3.6
14	230	4.7	—	—	—	—		3.6 3.6
15	230	(3.9)	—	—	—	2.0 (3.2)		2.0 3.6
16	240	(3.2)	—	—	—	3.6 (3.2)		3.6 3.4
17	280	(2.8)	—	—	—	3.7 (3.1)		3.7 3.6
18	(290)	(2.2)	—	—	—	4.2 (3.0)		4.2 3.3
19	—	—	—	—	—	4.8 —		4.8 3.4
20	—	—	—	—	—	6.8 —		6.8 3.4
21	—	—	—	—	—	6.1 —		6.1 3.4
22	—	—	—	—	—	6.6 —		6.6 3.4
23	—	—	—	—	—	6.2 —		6.2 3.4

Time: 45.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

TABLES OF IONOSPHERIC DATA

Table 2

Time	h'F2	foF2	h'F1	foF1	h'E	fOE	fEs	(M3000)F2
00	—	—	—	—	—	—		4.7
01	—	—	—	—	—	—		4.6
02	(320)	(2.4)						3.8 (3.0)
03	290	2.1						3.4 3.0
04	300	1.8						3.0 3.0
05	290	1.7						3.1 3.0
06	(280)	1.6						2.9 3.0
07	(275)	<1.6						3.0 (3.2)
08	—	<1.6						3.0 (3.2)
09	250	1.8						2.8 3.1
10	225	2.7						2.6 3.4
11	220	3.3						2.7 3.5
12	220	3.4						2.8 3.4
13	220	3.3						2.9 3.4
14	220	2.8						2.9 3.4
15	240	2.1						2.9 3.2
16	245	1.7						2.7 3.2
17	(280)	(1.6)						3.0 3.1
18	—	—						3.3 —
19	—	—						5.9 —
20	—	—						4.4 —
21	—	—						4.6 —
22	—	—						4.4 —
23	—	—						4.3 —

Time: 15.0°E.

Sweep: 0.6 Mc to 25.0 Mc in 5 minutes, automatic operation.

Table 4

Time	h'F2	foF2	h'F1	foF1	h'E	fOE	fEs	(M3000)F2
00	(300)	(1.5)						2.9 (3.1)
01	300	(1.4)						2.6 (3.0)
02	330	(1.6)						3.2 (2.8)
03	320	1.6						2.9 2.8
04	320	(2.0)						2.9 (3.0)
05	340	(1.6)						3.0 (2.9)
06	320	(1.8)						2.7 (2.9)
07	300	(1.7)						2.4 (3.0)
08	300	1.8						3.0
09	250	3.2						1.6 3.4
10	220	4.2						3.6 3.6
11	220	4.6	220	—	—	1.8	1.8	3.5 3.5
12	220	4.7	210	2.2	210	1.8	1.8	3.5 3.5
13	220	5.1	220	—	—	1.7	1.7	3.6 3.6
14	210	4.8	—	—	120	1.6	1.6	3.6 3.6
15	210	4.0	—	—	—	—	—	3.6 3.6
16	220	3.2	—	—	—	—	—	3.3 3.3
17	230	2.2	—	—	—	—	—	3.3 3.3
18	260	1.7	—	—	—	—	—	3.2 (3.2)
19	(280)	(1.6)	—	—	—	—	—	(2.7) —
20	—	—	—	—	—	—	—	2.9 —
21	—	—	—	—	—	—	—	3.6 —
22	—	—	—	—	—	—	—	2.7 —
23	—	—	—	—	—	—	—	—

Time: 160.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 16 seconds.

Table 6

Time	h'F2	foF2	h'F1	foF1	h'E	fOE	fEs	(M3000)F2
00	—	—	—	—	—	—		(3.1)
01	275	1.4						3.0
02	260	1.4						3.1
03	260	1.4						2.2 3.0
04	(255)	1.3						1.2 3.0
05	(260)	1.2						3.1
06	—	1.3						—
07	—	1.4						—
08	—	1.8						3.0
09	230	3.3						3.4 3.4
10	215	4.2						3.0 3.6
11	210	4.5	210	—	—	1.8	3.0	3.6 3.6
12	210	5.0	220	—	—	1.8	3.0	3.6 3.6
13	210	5.1	215	—	—	1.8	3.1	3.6 3.6
14	210	4.8	—	—	—	—		3.1 3.6
15	210	4.3	—	—	—	—		3.0 3.6
16	215	3.6	—	—	—	—		3.4 3.4
17	230	2.8	—	—	—	—		3.4 3.4
18	—	2.0	—	—	—	—		3.3 3.3
19	—	1.7	—	—	—	—		(3.1) —
20	—	1.6	—	—	—	—		(3.0) —
21	—	1.5	—	—	—	—		—
22	—	1.7	—	—	—	—		(3.1) —
23	—	1.6	—	—	—	—		(3.0) —

Time: 15.0°E.

Sweep: 0.6 Mc to 14.0 Mc in 8 minutes, automatic operation.

Uppsala, Sweden (59.8°N , 17.6°E)

Table 7

Time	December 1953					
	h'F2	foF2	h'F1	foF1	h'E	foE
00	305	1.8			2.3	2.9
01	305	1.8			2.3	2.9
02	290	1.7			2.5	2.9
03	(320)	1.6			2.6	2.9
04	325	1.4			3.0	(2.8)
05	—	1.4			2.9	—
06	—	1.4			3.2	—
07	—	1.4			2.6	—
08	255	2.2			2.8	3.0
09	220	3.9	—	—	E	3.1
10	220	4.5	220	2.4	120	1.6
11	220	5.0	215	2.4	115	1.8
12	215	5.1	220	2.5	115	1.9
13	215	5.0	<215	2.4	115	(1.8)
14	215	4.7	210	—	—	1.6
15	210	3.9			—	E
16	220	3.0			—	2.3
17	230	2.3			—	2.3
18	(260)	1.8			—	2.4
19	(280)	1.6			—	2.4
20	(300)	1.6			—	2.0
21	(280)	1.6			—	2.2
22	(295)	1.7			—	2.3
23	(300)	1.7			—	2.3

Time: 15.0°E .

Sweep: 1.4 Mc to 17.0 Mc in 6 minutes, automatic operation.

Table 9

San Francisco, California (37.4°N , 122.2°W)

Time	December 1953					
	h'F2	foF2	h'F1	foF1	h'E	foE
00	(250)	(2.9)			2.4	(3.2)
01	(240)	(2.9)			2.3	(3.3)
02	(230)	3.0			2.2	3.3
03	(240)	3.0			—	3.4
04	(230)	2.9			2.2	3.4
05	(240)	2.9			—	3.3
06	(250)	(2.9)			2.3	3.3
07	230	3.0			2.5	3.5
08	220	4.8	220	—	(120) (1.8)	3.1
09	230	5.2	220	—	(120) (2.3)	3.3
10	240	5.6	220	—	110 (2.6)	3.4
11	250	6.3	220	—	110 (2.8)	3.8
12	250	6.5	210	—	(120) (2.9)	3.7
13	240	6.0	220	—	110 (2.9)	3.7
14	240	5.7	230	—	110 (2.7)	3.8
15	230	5.5	220	—	120 (2.4)	3.6
16	220	5.0	—	—	—	3.6
17	210	4.2			—	3.6
18	(230)	2.9			—	3.5
19	(220)	2.6			—	3.5
20	(230)	2.6			—	3.5
21	(230)	2.6			—	3.3
22	(250)	2.6			—	3.2
23	(250)	2.9			—	3.2

Time: 120.0°W .

Sweep: 1.0 Mc to 25.0 Mc in 16 seconds.

Table 11

Okinawa I. (26.3°N , 127.8°E)

Time	December 1953					
	h'F2	foF2	h'F1	foF1	h'E	foE
00	300	(2.8)			—	(3.0)
01	300	2.8			—	(3.1)
02	270	4.0			—	3.2
03	250	2.8			—	3.2
04	240	2.8			—	3.6
05	250	2.7			—	—
06	240	(3.3)			—	—
07	220	4.4	—	—	—	3.6
08	240	5.2	220	—	120	3.0
09	250	6.0	220	3.8	110	3.2
10	250	6.4	220	4.0	110	2.8
11	260	6.7	200	4.2	110	3.0
12	260	8.0	200	4.2	110	3.0
13	260	8.8	(200)	4.1	110	3.0
14	250	8.2	220	4.0	110	2.9
15	240	7.1	220	—	—	4.0
16	230	6.6	—	—	—	3.6
17	210	5.4	—	—	—	3.4
18	220	4.0			—	3.5
19	240	3.6			—	3.3
20	250	3.5			—	3.3
21	250	3.4			—	3.4
22	240	3.2			—	(3.4)
23	(260)	(2.8)			—	(3.3)

Time: 127.5°E .

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 8

Graz, Austria (47.1°N , 16.5°E) December 1953

Time	December 1953					
	h'F2	foF2	h'F1	foF1	h'E	foE
00	280	—	—	—	3.0	—
01	270	—	—	—	3.0	—
02	280	—	—	—	2.9	—
03	280	—	—	—	2.9	—
04	270	—	—	—	2.7	—
05	240	—	—	—	2.3	—
06	240	—	—	—	2.3	—
07	250	—	—	—	2.4	—
08	200	—	—	—	4.3	—
09	200	—	—	—	5.1	—
10	200	—	—	—	6.0	—
11	200	—	—	—	6.0	—
12	200	—	—	—	5.0	—
13	200	—	—	—	6.2	—
14	200	—	—	—	5.1	—
15	200	—	—	—	5.0	—
16	200	—	—	—	4.3	—
17	220	—	—	—	3.4	—
18	260	—	—	—	2.8	—
19	260	—	—	—	3.0	—
20	240	—	—	—	2.9	—
21	260	—	—	—	2.9	—
22	280	—	—	—	3.0	—
23	290	—	—	—	2.9	—

Time: 16.0°E .

Sweep: 2.6 Mc to 12.0 Mc in 2 minutes.

Table 10

White Sands, New Mexico (32.3°N , 106.5°W) December 1953

Time	December 1953					
	h'F2	foF2	h'F1	foF1	h'E	foE
00	260	—	—	—	3.2	—
01	250	—	—	—	3.4	—
02	240	—	—	—	3.4	—
03	240	—	—	—	3.3	—
04	230	—	—	—	3.4	—
05	250	—	—	—	3.0	—
06	260	—	—	—	2.9	—
07	230	—	—	—	3.9	—
08	230	5.0	220	—	110	2.0
09	250	5.3	220	—	3.6	110
10	260	5.8	220	—	3.9	110
11	260	6.0	210	—	4.0	110
12	260	6.8	200	—	4.0	110
13	260	6.6	210	—	4.0	110
14	250	6.2	220	—	3.8	110
15	240	5.9	220	—	—	110
16	220	5.4	—	—	—	4.2
17	210	4.5	—	—	—	3.4
18	220	3.1	—	—	—	3.4
19	230	2.8	—	—	—	3.6
20	240	2.8	—	—	—	3.2
21	250	2.9	—	—	—	3.3
22	260	2.8	—	—	—	3.2
23	270	2.1	—	—	—	3.1

Time: 105.0°W .

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 12

Maui, Hawaii (20.8°N , 156.5°W) December 1953

Time	December 1953					
	h'F2	foF2	h'F1	foF1	h'E	foE
00	300	(2.5)	—	—	—	2.1 (3.0)
01	300	2.7	—	—	—	2.4 3.0
02	260	2.6	—	—	—	1.7 3.3
03	250	2.6	—	—	—	3.4
04	240	(2.5)	—	—	—	1.6 (3.4)
05	260	(2.0)	—	—	—	2.0 (3.3)
06	300	(2.0)	—	—	—	2.1 (2.9)
07	250	3.7	—	—	—	2.7 3.3
08	250	5.3	240	—	120	2.2 3.0
09	250	6.5	250	4.0	120	2.6 3.7
10	280	7.2	230	4.2	120	2.9 5.1
11	290	7.5	220	4.3	120	3.1 4.8
12	300	7.8	220	4.3	120	3.1 4.7
13	300	9.1	200	4.3	120	3.1 4.9
14	270	9.6	220	4.2	120	3.0 6.6
15	260	8.5	240	4.0	120	2.8 5.4
16	240	7.3	240	3.7	120	2.5 3.7
17	230	5.6	—	—	130	1.9 4.4
18	220	4.6	—	—	—	4.5 3.6
19	220	3.1	—	—	—	4.1 3.5
20	270	2.7	—	—	—	4.0 3.1
21	280	2.9	—	—	—	3.4 3.2
22	250	2.8	—	—	—	3.5 3.3
23	280	(2.8)	—	—	—	2.7 3.2

Time: 150.0°W .

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 13

Puerto Rico, W.I. (18.5°N, 67.2°W)							December 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	250	4.0			2.3	(3.1)		3.1
01	250	4.3			2.7	(3.3)		
02	240	(4.5)			3.0	(3.5)		
03	210	(4.5)			2.5	3.5		
04	210	3.5			2.7	3.1		
05	240	3.0			2.7	3.2		
06	240	3.0			2.5	3.2		
07	220	4.1				3.5		
08	240	4.9	230	—	110	2.0	2.8	3.5
09	250	5.7	230	—	110	2.5	2.9	3.5
10	260	6.3	220	4.1	110	2.8	3.0	3.5
11	260	6.2	210	4.2	110	3.0	3.2	3.5
12	260	6.0	210	4.2	110	3.1	3.5	
13	280	6.0	200	4.2	110	3.1	3.4	3.3
14	270	6.5	210	4.1	110	3.0	3.9	3.4
15	250	6.5	220	3.9	110	2.8	3.6	3.5
16	240	5.6	220	—	110	2.5	3.6	3.6
17	230	5.3	220	—	110	2.0	3.6	3.6
18	210	4.9				3.0	3.6	
19	220	3.9				2.9	3.5	
20	240	3.4				2.8	3.2	
21	240	3.6					3.1	
22	250	3.8				2.5	3.1	
23	260	4.0				2.5	3.2	

Time: 60.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 15

Panama Canal Zone (9.4°N, 79.9°W)							December 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	240	(3.0)			2.0	(3.3)		
01	230	(3.0)			2.2	(3.3)		
02	220	(2.8)			2.4	(3.3)		
03	220	(2.4)			2.4	(3.4)		
04	260	(2.2)			2.6	(3.1)		
05	260	(2.4)			4.2	(3.1)		
06	260	2.6			3.1	3.1		
07	240	4.6	—	—	120	1.8	4.1	3.5
08	280	5.7	240	3.8	120	2.4	4.0	3.4
09	280	6.8	230	4.1	110	2.8	4.2	3.3
10	270	7.4	210	4.2	110	3.1	4.3	3.4
11	280	7.6	210	4.3	100	3.2	4.3	3.3
12	290	7.6	210	4.3	110	3.3	4.8	3.3
13	290	7.2	220	4.3	100	3.3	5.0	3.2
14	290	7.6	220	4.2	110	3.2	5.6	3.2
15	280	7.3	220	4.2	110	3.0	5.1	3.3
16	250	6.8	220	3.9	110	2.7	4.4	3.6
17	230	5.7	230	—	120	2.2	4.2	3.6
18	220	4.4				4.0	3.6	
19	230	3.3				4.2	3.4	
20	240	3.0				3.7	3.4	
21	260	2.9				3.2	3.2	
22	260	2.9				2.6	3.1	
23	250	(3.0)				2.3	3.2	

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 17

Point Barrow, Alaska (71.3°N, 156.8°W)							November 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00		(2.4)				6.8		
01		(2.6)				6.5		
02		(2.5)				6.0		
03		2.7				5.0		
04		2.8				4.1		
05		(2.6)				4.2		
06		(3.0)				4.5		
07		3.3				4.4		
08		(2.8)				4.7		
09		3.0				4.5		
10		3.7				4.2		
11		4.1				3.8		
12		4.0				3.0		
13		4.4				2.5		
14		4.2				2.8		
15		4.0				2.3		
16		3.4				1.9		
17		2.8				3.2		
18		2.2				2.8		
19		1.9				4.0		
20		(2.0)				4.0		
21		(2.4)				4.4		
22		(2.7)				4.6		
23		(2.4)				6.9		

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 14

Guam I. (13.6°N, 144.9°E)							December 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	240	3.0						3.2
01	250	2.9						3.3
02	260	2.8						3.3
03	240	2.7						3.6
04	240	1.9						3.6
05	240	1.8						3.4
06	240	1.4					1.6	(3.3)
07	240	4.4	230	—	120	1.4		3.6
08	260	6.2	220	—	110	2.2	2.8	3.4
09	280	7.6	210	3.9	100	2.7	3.8	3.2
10	300	8.0	200	4.1	100	2.9	3.4	2.9
11	320	7.7	190	4.2	100	3.1	3.8	2.7
12	330	7.7	200	4.2	100	3.1	4.4	2.7
13	340	7.6	200	4.2	100	3.1	4.8	2.7
14	320	8.0	200	4.1	100	3.0	4.0	2.8
15	300	8.1	210	—	100	2.9	4.4	3.0
16	270	8.4	220	—	110	2.5	3.8	3.2
17	240	8.1	230	—	120	2.1	3.4	3.4
18	220	7.8						3.5
19	210	6.6						2.9
20	210	5.4						3.5
21	220	4.9						3.7
22	230	3.9						3.4
23	240	3.5						3.4

Time: 150.0°E.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 16

Huancayo, Peru (12.0°S, 75.3°W)							December 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	300	(3.3)					3.5	(3.0)
01	340	(1.4)						
02	240	< 1.0					4.0	
03	400	< 1.0					3.7	
04	360	< 1.0					4.1	
05	360	< 1.0					6.2	
06	240	4.5						
07	(280)	6.4	230	—	120	1.7	4.6	3.3
08	320	7.3	210	4.1	100	2.8	10.5	3.0
09	340	7.4	200	4.2	100	—	11.5	2.7
10	370	7.6	200	4.3	100	—	12.2	2.6
11	380	7.2	200	4.4	100	—	11.7	2.6
12	380	7.3	200	4.4	100	—	12.0	2.6
13	370	7.4	190	4.3	100	—	11.5	2.6
14	360	7.6	200	4.3	100	3.2	10.3	2.7
15	330	8.1	200	4.1	100	3.0	9.3	2.8
16	300	8.5	200	—	110	2.8	6.5	3.0
17	(280)	8.3	230	—	120	2.3	4.6	3.1
18	260	7.9						3.1
19	250	7.2						3.3
20	260	6.0						3.3
21	260	5.2						3.2
22	280	4.0						3.2
23	300	6.6						2.9

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 18

Kiruna, Sweden (67.8°N, 20.5°E)							November 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	340	3.5					4.2	3.1
01	(350)	3.4					4.1	3.0
02	340	3.3					2.2	3.0
03	330	2.2					2.7	3.0
04	310	2.2					1.6	(3.2)
05	(300)	(2.1)					1.6	(3.2)
06	(295)	(2.2)					2.0	(3.4)
07	275	2.8						3.2
08	235	3.2						3.4
09	235	4.0						3.5
10	230	4.2						3.5
11	230	4.3						3.7
12	225	4.7						3.6
13	225	4.4						3.5
14	225	4.4						3.5
15	260	3.7						3.4
16	240	3.3						3.4
17	1''	(250)	(3.5)					(3.4)
18	(270)	(3.1)		</td				

Table 19

Lulea, Sweden (65.6°N , 22.1°E) November 1953.						
Time	$\text{h}'\text{F}2$	$\text{foF}2$	$\text{h}'\text{F}1$	$\text{foF}1$	$\text{h}'\text{E}$	foE
00	320	(2.0)				3.0
01						
02	330	(1.9)				2.6
03						
04	(300)	---				3.2
05						
06	(290)	---				3.0
07						
08	250	2.6				2.2
09						
10	240	4.0	225	---	1.9	3.2
11						
12	225	4.7	215	---	130	2.0
13						2.8
14	225	4.0			---	3.3
15						
16	245	2.8				2.6
17						
18	(245)	---				3.2
19						
20	(310)	---				2.7
21						
22	(300)	---				3.4
23						

Time: 15.0°E .

Sweep: 1.5 Mc to 10.0 Mc in 6 minutes, automatic operation.

Table 21

Reykjavik, Iceland (64.1°N , 21.8°W) November 1953						
Time	$\text{h}'\text{F}2$	$\text{foF}2$	$\text{h}'\text{F}1$	$\text{foF}1$	$\text{h}'\text{E}$	foE
00	---	---			4.4	---
01	(370)	(2.6)			4.7	(2.8)
02	(350)	(2.9)			4.2	(3.0)
03	(320)	(2.4)			3.6	(2.9)
04	(320)	2.5			3.9	2.9
05	(300)	2.2			2.1	(3.2)
06	(330)	(1.9)			1.9	(3.2)
07	---	(1.6)			(2.0)	---
08	280	2.6			3.2	
09	250	3.6			3.3	
10	240	4.3			3.4	
11	250	4.7	240	---	---	3.4
12	250	4.7	240	---	---	3.4
13	250	4.4	250	---	---	3.4
14	260	4.2			3.3	
15	250	4.1			3.2	
16	250	(3.8)			(3.1)	
17	260	(4.0)			2.5	(3.2)
18	270	(3.2)			2.2	(3.2)
19	(280)	(2.6)			4.2	(3.2)
20	(320)	(2.3)			4.0	---
21	---	---			4.0	---
22	---	---			4.1	---
23	---	---			4.2	---

Time: 15.0°W .

Sweep: 1.0 Mc to 25.0 Mc in 18 seconds.

Table 23

Lindau/Zurz, Germany (51.6°N , 10.1°E) November 1953						
Time	$\text{h}'\text{F}2$	$\text{foF}2$	$\text{h}'\text{F}1$	$\text{foF}1$	$\text{h}'\text{E}$	foE
00	280	2.8			2.1	3.1
01	260	2.8			2.0	3.1
02	260	2.9			2.2	3.1
03	260	2.8			2.2	3.2
04	250	2.4			2.2	3.2
05	240	2.2			1.9	3.3
06	240	2.0				3.4
07	230	2.2			2.2	3.5
08	215	3.8			2.4	3.7
09	220	4.8	210	---	1.5	3.0
10	220	5.0	210	---	2.2	3.7
11	230	5.5	205	105	2.3	3.5
12	235	5.8	210	105	2.4	3.6
13	230	5.6	200	110	2.4	3.6
14	225	5.4	210	110	2.2	3.3
15	225	5.4	220	120	2.0	3.3
16	220	4.8			3.0	3.6
17	215	4.2			2.6	3.5
18	225	3.4			2.4	3.4
19	240	2.8			2.3	3.4
20	250	2.6			2.2	3.4
21	275	2.4			2.2	3.2
22	280	2.4			2.1	3.1
23	275	2.7			2.2	3.3

Time: 15.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 8 minutes.

Table 20

Fairbanks, Alaska (64.9°N , 147.8°W) November 1953						
Time	$\text{h}'\text{F}2$	$\text{foF}2$	$\text{h}'\text{F}1$	$\text{foF}1$	$\text{h}'\text{E}$	foE
00			(1.9)			5.0
01			—			5.2
02			(2.5)			5.8
03			(1.8)			(3.1)
04			—			5.4
05			(2.6)			(3.0)
06			(2.4)			5.0
07			2.4			4.5
08			3.0			(3.2)
09			3.6		1.5	3.1
10			4.4		1.7	2.2
11			5.0		1.9	2.3
12			4.8		1.8	2.2
13			5.0		1.8	(3.8)
14			4.8		1.9	(3.8)
15			4.0		1.9	(3.7)
16			3.5		1.8	(3.5)
17			3.0		4.0	(3.6)
18			2.0		4.6	(3.5)
19			1.8		4.4	(3.3)
20			(1.8)		4.4	(3.2)
21			(2.2)		4.5	(3.3)
22			(2.2)		4.0	(3.3)
23			(2.2)		4.8	(3.3)

Time: 150.0°W .

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 22

De Bilt, Holland (52.1°N , 5.2°E) November 1953						
Time	$\text{h}'\text{F}2$	$\text{foF}2$	$\text{h}'\text{F}1$	$\text{foF}1$	$\text{h}'\text{E}$	foE
00	250	2.8				3.0
01	255	(2.9)				3.0
02	250	(2.8)				3.1
03	< 250	(2.6)				3.1
04	230	(2.0)				3.1
05	230	(2.0)				3.3
06	(220)	(2.1)				3.3
07	< 220	3.0				3.5
08	210	4.4	210	2.4	1.9	3.6
09	230	4.8	200	3.2	2.2	2.4
10	230	5.2	200	3.3	105	2.5
11	230	5.8	200	3.4	100	2.4
12	230	5.6	210	3.5	105	2.5
13	230	5.3	205	3.3	100	2.4
14	230	5.2	215	2.8	110	2.2
15	220	4.9	210	2.3	140	1.9
16	210	4.4				3.6
17	210	3.5				3.4
18	220	2.8				3.3
19	220	2.5				3.3
20	(230)	2.4				3.2
21	< 240	2.5				3.0
22	< 260	2.6				3.0
23	260	2.6				3.0

Time: 0.0° .

Sweep: 1.4 Mc to 11.2 Mc in 6 minutes, automatic operation.

Table 24

Schwarzenburg, Switzerland (46.8°N , 7.3°E) November 1953						
Time	$\text{h}'\text{F}2$	$\text{foF}2$	$\text{h}'\text{F}1$	$\text{foF}1$	$\text{h}'\text{E}$	foE
00	260	3.0				3.3
01	250	3.0				3.3
02	250	3.0				3.4
03	250	3.0				3.4
04	220	3.0				3.5
05	210	2.4				3.6
06	200	2.5				3.8
07	200	2.8				3.8
08	200	4.0				4.0
09	200	4.8			2.2	4.0
10	200	5.2			2.4	4.0
11	200	5.6			2.6	4.0
12	200	5.8			2.6	4.0
13	200	5.8			2.5	4.0
14	200	5.4			2.4	3.9
15	200	5.5			2.2	4.0
16	200	5.5			2.2	4.0
17	200	4.5				4.0
18	200	3.6				3.8
19	210	3.0				3.6
20	210	2.8				3.6
21	210	2.7				3.6
22	250	2.8				3.4
23	270	2.8				3.4

Time: 15.0°E .

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

Table 25

Formosa, China (25.0°N, 121.5°E)							November 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	270	3.4					2.8	
01	240	3.7					3.1	
02	240	3.6					3.2	
03	230	3.5					2.0	3.4
04	230	2.8					1.9	3.2
05	260	2.3					2.1	3.0
06	280	2.8					2.0	2.9
07	240	5.6					2.2	3.5
08	240	6.6	240	(3.8)	130	1.9	2.2	3.4
09	270	7.4	240	4.0	120	2.4	3.2	3.6
10	280	8.5	220	4.3	120	3.0	4.1	3.2
11	280	9.2	220	4.3	120	3.2	4.6	3.6
12	280	10.3	220	4.4	120	3.2	4.8	3.2
13	280	11.9	240	4.3	—	—	4.6	3.3
14	260	11.9	230	(4.1)	—	—	4.4	3.6
15	240	10.2	220	4.0	—	—	4.4	3.6
16	230	8.0	—	—	—	—	3.9	3.5
17	220	6.7	—	—	—	—	3.7	3.6
18	210	5.7	—	—	—	—	3.0	3.5
19	230	4.6	—	—	—	—	2.6	3.3
20	240	4.7	—	—	—	—	2.4	3.1
21	240	4.0	—	—	—	—	2.0	3.4
22	240	3.4	—	—	—	—	1.7	3.1
23	280	3.3	—	—	—	—	3.0	3.0

Time: 120.0°E.

Sweep: 1.1 Mc to 19.5 Mc in 15 minutes, manual operation.

Table 27

Buenos Aires, Argentina (34.5°S, 58.5°W)							November 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	300	5.2					2.8	2.9
01	290	5.0					3.0	2.9
02	280	5.0					2.6	3.0
03	260	5.0					2.0	3.0
04	250	4.7					3.1	
05	240	4.6	—	—	130	1.8	2.1	3.4
06	240	5.2	220	—	110	2.3	3.6	3.4
07	280	5.8	220	—	110	2.7	3.7	3.3
08	300	5.9	210	—	100	3.0	4.0	3.0
09	340	6.4	210	4.5	100	(3.2)	4.2	2.9
10	400	7.0	200	4.5	—	—	4.5	2.7
11	390	8.4	200	4.5	100	3.3	5.2	2.7
12	350	9.8	200	4.5	—	—	5.0	2.9
13	310	10.5	200	4.4	—	—	4.4	3.0
14	290	11.0	200	4.4	—	—	4.4	3.1
15	280	11.0	200	4.2	100	3.0	4.4	3.3
16	260	10.2	220	—	—	—	3.9	3.3
17	260	10.0	220	—	—	—	3.8	3.4
18	240	8.6	230	—	—	—	2.9	3.4
19	240	7.6	—	—	—	—	3.2	
20	260	6.3	—	—	—	—	3.0	
21	290	5.8	—	—	—	—	2.9	
22	300	5.8	—	—	—	—	2.9	
23	300	5.7	—	—	—	—	2.9	

Time: 60.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

Table 29

Resolute Bay, Canada (74.7°N, 94.9°W)							October 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	260	2.8					4.0	3.1
01	260	2.7					3.5	3.0
02	270	2.5					3.6	3.0
03	270	2.4					3.6	(3.1)
04	280	2.5					3.9	3.0
05	280	2.1					3.5	3.0
06	260	2.7	—	—			3.8	3.0
07	270	3.0	—	—	1.3	3.7	3.1	
08	260	3.5	—	—	120	1.4	3.2	3.1
09	260	3.7	—	—	110	1.7	3.3	5.1
10	260	4.0	250	2.8	110	1.8	2.3	3.1
11	260	4.0	250	3.0	110	1.9	3.1	
12	280	4.0	240	3.0	110	1.9	2.6	3.1
13	270	4.0	240	3.0	110	1.9	2.2	3.1
14	270	4.1	240	—	100	1.8	2.1	3.1
15	260	3.9	250	—	100	1.7	2.9	3.2
16	260	4.0	—	—	110	1.4	2.8	3.1
17	260	4.0	—	—	—	—	3.0	
18	250	3.5	—	—	—	—	1.8	3.0
19	250	3.3	—	—	—	—	3.5	3.0
20	260	3.2	—	—	—	—	2.8	3.0
21	260	3.0	—	—	—	—	2.8	3.1
22	260	3.0	—	—	—	—	3.0	
23	250	2.9	—	—	—	—	2.5	3.1

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 26

Leopoldville, Belgian Congo (4.3°S, 15.3°E)							November 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M2000)F2
00	245	6.0						2.3
01	240	5.8						2.3
02	220	4.5						2.5
03	230	3.4						2.5
04	240	2.8						2.6
05	240	4.6	—	—	125	1.7		2.7
06	270	5.6	230	—	115	2.4	2.9	2.5
07	305	6.2	220	4.2	110	2.7	3.3	2.2
08	345	7.0	220	4.3	110	3.1	3.2	2.0
09	385	8.2	210	4.4	110	3.3	3.8	2.0
10	370	9.4	210	4.4	110	3.3	3.4	2.0
11	380	10.4	210	4.4	110	3.4	3.5	2.0
12	370	11.0	215	4.4	110	3.4	3.9	2.1
13	350	11.4	220	4.3	110	3.2	4.0	2.1
14	350	11.4	220	4.2	110	3.0	3.9	2.1
15	325	11.8	230	4.0	115	2.6	3.6	2.2
16	280	11.6	240	—	120	2.0	3.0	2.2
17	250	11.0	—	—	—	—	2.6	2.3
18	250	10.1	—	—	—	—	2.6	2.2
19	265	9.5	—	—	—	—	2.0	2.2
20	260	9.4	—	—	—	—	2.0	2.3
21	240	9.6	—	—	—	—	2.5	
22	225	9.4	—	—	—	—	2.6	
23	220	6.9	—	—	—	—	2.4	

Time: 0.0°.

Sweep: 1.0 Mc to 16.0 Mc in 7 seconds.

Table 28

Deception I. (63.0°S, 60.7°W)							November 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	270	5.8						(3.2)
01	280	5.4						(3.2)
02	280	5.2						(3.1)
03	280	5.5						(3.1)
04	260	5.5						(3.2)
05	250	5.8						(3.2)
06	250	5.4	—	—	—	—	2.0	(3.2)
07	260	5.4	—	—	—	—	3.3	(3.2)
08	250	5.6	—	—	—	—	4.0	(3.4)
09	260	5.6	—	—	—	—	4.5	(3.4)
10	260	5.6	—	—	—	—	4.7	(3.4)
11	240	5.7	—	—	—	—	3.8	(3.4)
12	240	5.6	—	—	—	—	4.0	(3.4)
13	240	5.8	—	—	—	—	4.0	(3.4)
14	230	5.4	—	—	—	—	3.8	(3.4)
15	230	5.3	—	—	—	—	3.8	(3.4)
16	300	3.4	—	—	—	—	4.6	3.2
09	290	3.6	240	—	—	—	4.4	3.2
10	280	4.0	240	—	100	—	3.6	3.3
11	270	4.1	230	—	100	—	2.6	3.3
12	260	4.4	230	—	100	—	2.6	3.3
13	260	4.6	230	—	100	1.9	2.0	3.3
14	260	4.6	220	—	—	—	—	3.4
15	250	4.4	220	—	110	1.6	—	3.4
16	250	4.3	—	—	—	—	2.4	3.4
17	240	3.8	—	—	—	—	1.9	3.3
18	250	3.3	—	—	—	—	3.4	3.4
19	280	3.0	—	—	—	—	3.6	3.2
20	300	2.4	—	—	—	—	3.8	3.1
21	320	2.4	—	—	—	—	4.6	3.0
22	(300)	(2.7)	—	—	—	—	4.4	(3.2)
23	(290)	(2.7)	—	—	—	—	5.9	(3.1)

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 31								
Baker Lake, Canada (64.3°N, 96.0°W)								October 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	260	2.0	—	—	E	6.0	2.9	
01	290	1.8	—	—	E	6.0	2.9	
02	300	2.0	—	—	E	8.0	3.0	
03	300	1.9	—	—	E	5.0	(2.9)	
04	290	2.1	—	—	E	4.5	(2.9)	
05	300	2.4	—	—	—	5.0	(3.0)	
06	270	2.7	—	—	—	4.3	(2.9)	
07	250	3.0	—	—	120	1.9	4.0	2.9
08	280	3.8	—	—	120	2.7	2.5	3.0
09	280	3.7	—	—	120	2.7	3.0	3.0
10	300	4.0	270	3.4	120	2.8	3.0	3.0
11	340	4.2	260	3.5	120	2.8	—	2.9
12	320	4.4	260	3.5	120	2.6	—	3.0
13	320	4.7	260	3.5	120	2.7	—	2.8
14	300	4.9	250	3.3	120	2.5	—	3.0
15	300	4.5	260	3.2	120	2.4	—	2.9
16	280	4.0	280	—	130	2.3	—	3.0
17	290	4.0	—	—	130	2.3	4.2	2.9
18	280	3.4	—	—	130	2.3	6.0	2.9
19	290	3.2	—	—	2.0	6.0	2.8	
20	290	3.0	—	—	—	4.7	2.9	
21	270	3.0	—	—	—	7.5	3.0	
22	270	2.8	—	—	—	7.2	2.9	
23	280	2.5	—	—	E	7.0	3.0	

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 33								
Churchill, Canada (58.8°N, 94.2°W)								October 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	2.8	—	—	—	5.0	(2.8)	
01	300	2.8	—	—	130	3.2	8.0	(2.8)
02	290	2.8	—	—	(1.8)	6.0	—	
03	300	2.8	—	—	110	(2.3)	6.0	(2.8)
04	300	3.0	—	—	120	2.8	4.5	—
05	320	3.2	—	—	110	3.0	4.0	(2.8)
06	340	3.2	—	—	120	3.3	4.8	(2.8)
07	300	3.6	—	—	110	2.8	4.8	3.0
08	300	4.0	230	—	110	2.8	5.0	3.0
09	290	4.3	240	3.7	110	2.9	5.0	3.0
10	320	4.5	220	3.8	110	2.6	5.0	3.0
11	300	4.8	220	3.8	110	2.7	—	2.9
12	310	5.0	240	3.8	110	2.8	—	2.9
13	310	5.0	240	3.8	110	2.7	—	3.0
14	300	5.0	240	3.8	110	2.6	—	3.0
15	300	5.0	240	3.5	110	2.5	—	3.0
18	270	4.8	260	—	110	2.3	—	3.0
17	270	4.6	—	—	110	2.0	—	3.0
18	270	3.8	—	—	120	2.2	—	3.0
19	300	3.4	—	—	120	2.7	2.8	2.9
20	330	3.0	—	—	120	2.8	4.4	2.8
21	320	3.0	—	—	120	2.6	6.0	(2.9)
22	290	2.8	—	—	130	2.2	8.0	(2.9)
23	300	2.7	—	(2.6)	—	9.0	(2.8)	

Time: 90.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 35								
De Bilt, Holland (52.1°N, 5.2°E)								October 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	270	2.9	—	—	—	—	3.1	
01	270	3.0	—	—	—	—	3.1	
02	265	3.0	—	—	—	1.5	3.1	
03	260	2.9	—	—	—	—	3.1	
04	240	2.3	—	—	—	2.1	3.2	
05	240	2.2	—	—	—	—	3.3	
06	210	2.9	—	—	E	—	3.4	
07	210	4.3	205	—	—	1.8	3.6	
08	215	4.9	200	3.2	100	2.3	2.3	3.6
09	230	5.5	200	3.6	100	2.5	2.9	3.8
10	225	5.8	200	3.9	100	2.6	3.1	3.8
11	230	6.2	200	3.9	100	2.7	3.1	3.8
12	230	6.0	200	3.9	100	2.6	3.1	3.6
13	230	6.1	200	3.7	105	2.6	2.7	3.8
14	220	6.1	205	3.6	105	2.5	2.6	3.6
15	215	5.8	210	3.0	105	2.2	2.2	3.6
18	210	5.6	—	—	E	2.1	3.6	
17	205	5.3	—	—	—	2.2	3.4	
18	210	5.0	—	—	—	—	3.5	
19	210	3.9	—	—	—	—	3.5	
20	210	3.5	—	—	—	—	3.4	
21	230	2.9	—	—	—	—	3.2	
22	230	2.7	—	—	—	—	3.2	
23	260	2.8	—	—	—	—	3.1	

Time: 0.0°W.

Sweep: 1.4 Mc to 11.2 Mc in 6 minutes, automatic operation.

Table 32								
Reykjavik, Iceland (64.1°N, 21.8°W)								October 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	—	—	—	—	—	—	—	—
01	—	—	(2.2)	—	—	—	—	4.4
02	—	—	(2.0)	—	—	—	—	4.0
03	—	—	(2.3)	—	—	—	—	4.8
04	—	—	(2.3)	—	—	—	—	4.4
05	—	—	(2.0)	—	—	—	—	2.2
06	—	—	2.2	—	—	—	—	—
07	—	—	2.9	—	—	—	—	—
08	—	—	3.6	—	—	—	—	1.6
09	—	—	4.0	—	—	—	—	1.7
10	—	—	4.5	—	—	—	—	2.0
11	—	—	4.7	—	—	—	—	2.2
12	—	—	4.9	—	—	—	—	2.2
13	—	—	5.0	—	—	—	—	2.3
14	—	—	5.0	—	—	—	—	2.2
15	—	—	4.8	—	—	—	—	2.0
16	—	—	4.5	—	—	—	—	—
17	—	—	(4.2)	—	—	—	—	3.0
18	—	—	(3.8)	—	—	—	—	3.5
19	—	—	(3.3)	—	—	—	—	3.9
20	—	—	—	—	—	—	—	3.9
21	—	—	(3.2)	—	—	—	—	4.0
22	—	—	—	—	—	—	—	4.2
23	—	—	—	—	—	—	—	4.8

Time: 15.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 18 seconds.

Table 34								
Prince Rupert, Canada (54.8°N, 130.3°W)								October 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	1.8	—	—	—	—	—	2.4
01	300	1.7	—	—	—	—	—	2.7
02	300	1.8	—	—	—	—	—	3.5
03	(300)	1.5	—	—	—	—	—	4.2
04	310	1.9	—	—	—	—	—	—
05	320	1.9	—	—	—	—	—	—
06	300	2.0	—	—	—	—	—	—
07	260	3.0	—	—	—	—	—	—
08	240	3.8	220	—	—	120	2.0	3.3
09	260	4.2	220	3.3	110	2.2	2.3	3.3
10	300	4.6	210	3.6	110	2.6	2.6	3.3
11	300	5.0	200	3.8	110	2.7	—	3.2
12	300	5.2	210	3.8	110	2.7	—	3.3
13	290	5.2	220	3.8	110	2.8	—	3.3
14	290	5.1	220	3.7	110	2.7	—	3.3
15	260	5.0	230	—	110	2.6	—	3.3
16	240	5.0	240	—	110	2.3	—	3.3
17	240	4.8	—	—	110	2.0	1.6	3.3
18	230	4.2	—	—	—	—	—	3.2
19	230	3.4	—	—	—	—	—	1.6
20	260	2.5	—	—	—	—	—	—
21	260	2.0	—	—	—	—	—	—
22	280	2.0	—	—	—	—	—	2.1
23	300	1.8	—	—	—	—	—	3.0

Time: 90.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 37

St. John's, Newfoundland (47.6°N, 52.7°W)								October 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	300	2.0					3.0	3.0	
01	320	2.0					3.5	3.0	
02	310	2.0					3.0	3.0	
03	300	1.8					3.1	3.0	
04	290	1.8					2.8	3.1	
05	300	1.8				E	3.0	3.0	
06	250	3.3	230	—	130	1.7		3.5	
07	240	4.3	240	—	120	2.2		3.6	
08	260	5.0	230	3.4	120	2.5		3.6	
09	280	5.2	210	3.7	120	2.7		3.5	
10	280	5.6	200	3.9	110	2.9		3.5	
11	290	5.8	200	4.0	110	2.9		3.5	
12	280	6.0	210	4.0	110	2.9		3.5	
13	280	5.8	220	3.8	110	2.8		3.5	
14	280	5.6	240	3.8	120	2.6		3.5	
15	260	5.7	240	3.3	120	2.4		3.5	
16	240	5.7	240	—	130	2.0		3.5	
17	230	5.2	—	—	—	E	2.0	3.5	
18	230	4.6	—	—	—		3.4		
19	240	3.9					3.3		
20	250	3.2					3.0		
21	280	2.7					2.9		
22	300	2.4					1.9	3.0	
23	310	2.1					2.9	2.9	

Time: 60.0°W.

Sweep: 0.8 Mc to 10.0 Mc in 18 seconds.

Table 39

Wakkanai, Japan (45.4°N, 141.7°E)								October 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	300	3.8					2.8	3.0	
01	280	3.8					3.0	3.0	
02	270	3.8					3.0	3.0	
03	260	3.7					2.6	3.0	
04	250	3.7					2.6	3.1	
05	240	3.6					2.4	3.2	
06	230	4.2					2.5	3.4	
07	230	5.3	—	—	120	2.0	3.2	3.5	
08	240	6.1	230	3.7	110	2.4	3.3	3.4	
09	250	6.5	220	4.0	110	2.7	4.4	3.4	
10	250	6.8	220	4.1	110	2.8	4.4	3.4	
11	250	7.0	220	4.2	110	2.9	4.3	3.3	
12	250	7.2	220	4.1	110	2.8	4.2	3.4	
13	250	6.6	230	4.0	110	2.7	3.6	3.4	
14	250	6.3	230	3.8	110	2.5	3.5	3.4	
15	240	6.3	240	3.6	110	2.3	3.6	3.4	
16	230	6.1	—	—	120	1.8	3.3	3.4	
17	230	5.7					3.0	3.4	
18	230	4.5					3.5	3.2	
19	250	4.6					3.3	3.2	
20	260	4.1					3.0	3.1	
21	270	3.9					3.0	3.0	
22	270	3.9					2.8	3.0	
23	280	4.2					2.8	3.1	

Time: 135.0°E.

Sweep: 1.0 Mc to 15.5 Mc in 2 minutes.

Table 41

Tokyo, Japan (35.7°N, 139.5°E)								October 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	280	3.5					2.6	3.0	
01	270	3.5					2.8	3.0	
02	270	3.4					2.8	3.0	
03	250	3.5					2.5	3.1	
04	220	3.2					2.6	3.8	
05	250	3.1					2.5	3.1	
06	220	4.4			150	1.7	2.5	3.4	
07	220	6.0	220	—	120	2.2	3.0	3.5	
08	230	6.9	220	4.0	110	2.6	4.0	3.5	
09	240	7.0	220	4.0	110	2.8	4.0	3.5	
10	260	7.0	210	4.3	110	3.0	4.5	3.3	
11	260	7.7	200	4.3	110	3.0	4.5	3.4	
12	250	8.4	210	4.3	110	3.0	4.0	3.4	
13	250	7.3	230	4.2	110	3.0	3.9	3.4	
14	260	7.1	230	4.0	110	2.9	3.9	3.3	
15	250	7.1	240	3.7	110	2.6	3.9	3.4	
16	230	6.8	230	—	120	2.2	4.0	3.5	
17	220	6.2	—	—	—	—	3.9	3.5	
18	220	4.5					3.5	3.4	
19	230	4.3					3.0	3.3	
20	260	3.8					3.0	3.1	
21	260	5.5					3.0	3.1	
22	270	3.3					3.0	3.0	
23	280	3.6					3.0	3.0	

Time: 135.0°E.

Sweep: 1.0 Mc to 17.2 Mc in 2 minutes.

Table 38

Ottawa, Canada (45.4°N, 75.8°W)								October 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	320	2.0							3.0
01	330	1.9							(2.9)
02	340	1.6							(2.8)
03	350	1.6							(2.9)
04	380	1.8							2.6
05	(320)	(1.8)							3.3
06	280	2.4							3.1
07	240	4.0	240	—	120	2.0			3.3
08	260	4.8	220	3.6	120	2.2			3.4
09	280	5.3	220	3.9	120	2.6			3.3
10	280	5.4	210	3.9	120	2.8			3.3
11	290	5.8	210	4.0	120	2.9			3.2
12	280	5.8	220	4.0	110	2.9			3.3
13	290	5.8	220	4.0	120	2.9			3.3
14	290	5.8	230	3.9	120	2.8			3.3
15	280	5.9	230	3.8	120	2.6			3.3
16	270	5.8	240	—	120	2.2			3.3
17	240	5.3	—	—	140	1.9			3.3
18	240	4.6							3.2
19	250	3.8							3.2
20	260	3.0							3.1
21	280	2.8							3.0
22	290	2.4							3.0
23	310	2.0							3.0

Time: 75.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 15 seconds.

Table 41

Akita, Japan (39.7°N, 140.1°E)								October 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	280	3.5							2.6
01	290	3.5							2.6
02	290	3.5							2.8
03	250	3.5							2.9
04	240	3.4							3.1
05	250	3.1							3.0
06	230	4.5							3.4
07	230	5.5	—	—	130	1.9	3.2		3.5
08	240	6.5	—	—	120	2.4	3.4		3.5
09	250	7.2	230	4.0	110	2.6	4.3		3.4
10	260	7.4	230	4.2	110	2.9	4.4		3.3
11	280	7.8	220	4.4	110	3.0	4.2		3.2
12	290	8.7	220	4.4	110	3.0	4.2		3.3
13	270	9.1	230	4.4	110	3.0	4.2		3.3
14	280	8.6	240	4.2	110	3.0	3.5		3.3
15	260	8.4	240	4.0	110	2.8	3.8		3.4
16	250	7.6	240	3.8	110	2.4	3.3		3.4
17	250	6.9	—	—	120	2.0	3.4		3.5
18	240	5.7							3.7
19	230	4.4							3.5
20	260	3.5							3.2
21	280	3.4							3.0
22	270	3.4							2.7
23	300	3.3							2.9

Time: 135.0°E.

Sweep: 0.8 Mc to 20.0 Mc in 15 minutes, manual operation.

Table 43

Johannesburg, Union of S. Africa (26.2° S, 28.1° E)								October 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	260	3.4					2.3	3.0	
01	270	3.4					2.1	3.0	
02	250	3.4					1.8	3.1	
03	250	3.1						3.0	
04	260	2.9						3.0	
05	260	2.8					1.9	3.0	
06	230	4.6	---	---	---	1.8		3.4	
07	250	5.6	220	3.6	110	2.4		3.3	
08	280	6.1	220	4.1	110	(2.8)		3.3	
09	290	6.6	210	4.3	110	3.0		3.2	
10	290	6.9	200	4.4	110	3.2		3.2	
11	310	7.1	200	4.5	110	3.4		3.1	
12	310	7.7	200	4.5	110	3.4		3.0	
13	300	8.0	210	4.5	110	3.4		3.0	
14	300	8.0	210	4.4	110	3.3		3.0	
15	290	8.0	210	4.2	110	3.1		3.1	
16	280	7.9	220	4.0	110	2.8		3.1	
17	260	8.0	230	3.4	120	2.3		3.2	
18	240	8.0	---	---	---	---	2.7	3.3	
19	220	6.8					2.2	3.3	
20	230	5.5						3.3	
21	240	4.2					1.7	3.2	
22	260	3.7					2.2	3.0	
23	260	3.6					1.6	3.0	

Time: 30.0° E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 45

Point Barrow, Alaska (71.3° N, 156.8° W)								September 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	280	3.0					6.6	3.1	
01	270	3.0					6.2	3.1	
02	250	(3.0)					5.8	3.2	
03	280	2.9					4.9	3.2	
04	310	3.0					4.0	3.2	
05	300	3.0					4.0	3.0	
06	330	3.2	---	---	---	---	3.9	3.1	
07	320	< 3.3	240	---	---	---	4.7	(3.2)	
08	(360)	3.8	250	---	---	---	4.0	(3.0)	
09	(340)	3.9	220	3.4	100	2.2	4.2	(3.0)	
10	440	3.8	220	3.5	100	2.3	4.1	2.8	
11	410	4.0	220	3.5	100	2.3	2.5	2.7	
12	420	4.0	220	3.5	100	2.4	2.4	2.7	
13	380	4.0	230	3.6	100	2.5		2.8	
14	360	4.0	230	3.6	100	2.4		3.0	
15	320	4.0	230	3.5	100	2.3		3.1	
16	320	4.2	240	3.4	100	2.2	2.2	3.1	
17	280	4.0	240	3.1	100	1.8	2.1	3.2	
18	250	3.8	220	---	110	1.5	2.6	3.3	
19	250	3.3	---	---	---	---	3.3	3.3	
20	270	3.0	---	---	---	---	4.2	3.2	
21	320	3.0	---	---	---	---	4.7	(3.1)	
22	(340)	(2.9)	---	---	---	---	5.8	---	
23	300	2.9	---	---	---	---	5.4	3.1	

Time: 150.0° W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 47

Point Barrow, Alaska (71.3° N, 156.8° W)								August 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	320	3.0					6.8	3.2	
01	280	3.2					5.0	3.3	
02	290	3.2	---	---	---	---	5.8	3.2	
03	270	3.3	---	---	---	---	4.9	3.2	
04	300	3.3	220	---	---	---	4.9	3.2	
05	310	3.5	220	---	---	---	4.6	3.1	
06	420	3.6	230	3.2	100	---	4.6	2.8	
07	390	3.8	230	3.5	100	---	4.6	2.9	
08	430	4.0	220	3.6	100	(2.2)	4.6	2.8	
09	400	4.0	210	3.6	100	2.4	4.8	2.8	
10	500	4.0	200	3.7	100	2.5	4.0	2.5	
11	470	4.1	220	3.8	100	2.5	3.1	2.7	
12	430	4.2	210	3.8	100	2.7		2.8	
13	440	4.0	220	3.8	100	2.7		2.8	
14	440	4.1	220	3.8	100	2.5		2.7	
15	430	4.1	220	3.7	100	2.5		2.8	
16	410	4.2	220	3.6	100	2.4		2.9	
17	370	4.1	230	3.5	110	2.2		3.0	
18	330	4.0	230	3.4	110	2.0		3.2	
19	310	3.9	240	3.3	100	1.7		3.2	
20	280	3.6	240	---	---	---	3.9	3.3	
21	300	3.4	---	---	---	---	4.4	3.2	
22	300	3.3	---	---	---	---	4.9	3.2	
23	320	3.1	---	---	---	---	6.2	3.1	

Time: 150.0° W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 44

Capetown, Union of S. Africa (34.2° S, 18.3° E)								October 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	260	3.2							3.0
01	270	3.2							3.0
02	270	3.2							3.0
03	260	3.1							3.0
04	260	3.1							3.0
05	260	3.1							3.0
06	250	3.6	---	---	1.8				3.2
07	240	5.1	240	---	---	---	3.3	120	2.0
08	260	5.7	230	3.8	120	2.5			3.3
09	280	6.1	230	4.1	120	2.9			3.2
10	300	6.4	210	4.3	110	3.1		3.3	3.2
11	320	7.0	210	4.4	110	3.2			3.0
12	320	7.4	200	4.5	110	3.3		3.3	3.0
13	320	8.1	200	4.5	110	3.3		3.5	3.0
14	300	8.1	220	4.4	110	3.2		3.3	3.0
15	300	8.2	220	4.3	110	3.1			3.1
16	280	8.0	220	4.1	110	2.9		3.2	3.1
17	270	7.4	230	3.8	120	2.6			3.2
18	250	7.1	240	3.1	120	2.0		2.7	3.3
19	230	7.1	---	---	---	---	1.6		3.1
20	220	5.6							3.2
21	230	4.4							3.2
22	250	3.7							3.1
23	250	3.4							3.1

Time: 30.0° E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 45

Reykjavik, Iceland (64.1° N, 21.8° W)								September 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00									4.9
01									4.9
02									5.0
03									4.2
04									4.6
05									3.6
06									
07									
08									
09									
10									
11									
12									
13									
14									
15									
16									
17									
18									
19									
20									
21									
22									
23									

Time: 15.0° W.

Sweep: 1.0 Mc to 25.0 Mc in 18 seconds.

Table 46

Reykjavik, Iceland (64.1° N, 21.8° W)								August 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00									4.3
01									4.9
02									

TABLE 49
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

$h^{\prime} F_2$, Km
(Characteristic)
Observed at Washington, D.C.

Lat 38°7' N, Long 77°10' W

Km
(Unit)
January, 1954
(Month)

National Bureau of Standards
(Institution)
McC. E.J.W., J.W.P.

Day	75° W												Mean Time													
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23		
1	A	S	S	(2.80) ⁵	(2.50) ⁵	1.40	2.20	2.30	2.40	2.40	2.50	(2.40) ^A	2.40	2.40	2.30	2.20	2.20	2.30	2.30	2.30	2.20	2.20	2.20	2.20		
2	(2.40) ⁵	(2.60) ⁵	2.70	(2.70) ⁵	2.50	2.60	2.50	2.40	(2.50) ^A	2.30	2.60	1.60	2.50	2.40	2.40	2.40	2.20	2.20	2.20	2.20	2.20	2.20	2.20	(2.70) ⁵		
3	2.70	(2.50) ⁵	2.50	2.50	2.40	2.20	2.20	2.40	2.20	2.30	2.40	2.40	2.40	2.40	2.40	2.40	2.20	2.20	2.20	2.20	2.20	2.20	2.20	(2.70) ⁵		
4	3.01 ⁵	(2.60) ⁵	(2.70) ⁵	2.40	2.30	2.20	2.20	2.20	2.20	2.40	2.40	2.50	2.50	2.60	2.60	2.60	2.60	2.20	2.20	2.20	2.20	2.20	2.20	2.20	(2.40) ⁵	
5	2.50	2.50 ⁵	2.20 ⁵	2.90 ^F	2.50	2.60	2.60	2.60	2.60	2.60	2.70	(2.30) ^A	2.70	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.50	
6	2.50	(2.70) ⁵	(2.40) ⁵	(2.80) ⁵	2.70	2.50	2.30	2.40	(2.40) ⁵	2.30	2.50	2.80	2.60	2.60	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	(2.70) ⁵	
7	(2.50) ⁵	(2.80) ⁵	2.70	2.70	(2.70) ⁵	2.50	2.30	2.20	2.30	2.30	(2.50) ^A	(2.50) ^L	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	(2.70) ⁵	
8	2.50	(2.50) ⁵	(2.70) ⁵	2.70	3.60	(A)	2.70	2.40	2.20	2.20	2.40	2.40	2.40	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	(3.00) ⁵	
9	2.50 ⁵	(3.00) ⁵	2.70	2.50	(2.50) ⁵	(2.30) ⁵	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	S	
10	2.50 ⁵	(2.30) ⁵	(2.70) ⁵	2.50	(2.40) ⁵	2.30	2.20	2.20	2.20	2.30	(2.60) ^A	2.70	2.50	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	(2.60) ⁵	
11	(2.70) ⁵	(2.60) ⁵	(2.60) ⁵	2.50	(2.60) ⁵	2.50	2.30	2.20	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	S	
12	5	(2.60) ⁵	(2.60) ⁵	2.40	(2.60) ⁵	2.40	2.30	2.30	2.30	2.30	(2.30) ^A	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	(2.70) ⁵	
13	(2.70) ⁵	(2.50) ⁵	(2.60) ⁵	2.50	(2.50) ⁵	2.50	2.30	2.30	2.30	2.30	(2.50) ^A	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	(2.60) ⁵	
14	(2.70) ⁵	(2.60) ⁵	(2.60) ⁵	2.60	2.50	2.40	2.40	2.40	2.40	2.40	2.40	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	(2.60) ⁵	
15	(2.70) ⁵	(2.40) ⁵	(2.50) ⁵	2.50	(2.50) ⁵	2.50	2.30	2.20	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	S	
16	(2.70) ⁵	(2.40) ⁵	C	C	C	C	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	2.30	(2.70) ⁵	
17	5	(2.60) ⁵	(2.40) ⁵	2.30	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	2.20	(2.60) ⁵	
18	(2.70) ⁵	(2.60) ⁵	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	(2.60) ⁵	
19	4.0	(2.80) ⁵	(2.80) ⁵	2.20	2.50	(2.30) ⁵	(2.50) ⁵	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	S
20	(3.00) ⁵	(2.40) ⁵	(2.60) ⁵	(2.60) ⁵	(2.60) ⁵	(2.70) ⁵	(2.70) ⁵	(2.40) ⁵																		
21	(2.60) ⁵	(2.70) ⁵	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	(2.80) ⁵	
22	(2.60) ⁵	(2.60) ⁵	2.50	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	(2.80) ⁵	
23	5	(2.70) ⁵	2.50	2.50	(2.80) ⁵	(2.80) ⁵	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	(2.60) ⁵
24	5	(2.70) ⁵	(2.70) ⁵	(2.80) ⁵	(2.80) ⁵	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	S
25	5	(2.50) ⁵	(2.70) ⁵																							
26	(2.70) ⁵																									
27	1.70	2.40	2.40	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	(2.50) ⁵	
28	(2.70) ⁵	(2.20) ⁵	(2.80) ⁵																							
29	(2.90) ⁵	(2.80) ⁵	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	S	
30	5	(2.70) ⁵	(2.80) ⁵	(2.70) ⁵																						
31	(2.80) ⁵	(2.60) ⁵																								
Median	2.70	(2.60) ⁵	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	2.60	(2.70) ⁵	
Coupl	2.2	2.7	2.8	2.5	2.0	3.0	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	2.2	

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual □ Automatic ■

1.5 GIGAHERTZ PRINTING OFFICE 100-170319

TABLE 50
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

Form adopted June 1946

foF₂, Mc January, 1954

(Characteristic)

Mc (Unit)

January (Month)

Washington, D. C.

Lat. 38°7'N, Long. 77°1'W

75°W

Mean Time

National Bureau of Standards

(Institution)

E. J. W., J. W. P.

Scaled by: MC. E. J. W., J. W. P.

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	(1.7) ^s	[1.8] ^s	(1.9) ^s	(2.0) ^s	(2.1) ^s	(2.2) ^s	(3.7) ^s	(3.3) ^s	4.8	4.7	(5.9) ^s	6.6	6.0	5.4	5.4	5.0	4.6F	(4.8) ^s	(5.1) ^s	4.2	3.2F	3.1F	(2.9) ^s		
2	(2.7) ^s	(3.1) ^s	3.3F	3.4F	3.6	3.2	3.1F	(3.0) ^s	4.3	5.2	6.0	6.3	7.0	7.0	6.2	6.3	6.5	5.8	4.9	4.4	3.3	2.6F	2.2F	2.3F	
3	2.7F	3.2F	(3.3) ^s	3.4F	3.6F	3.7F	3.7F	3.2	4.5	5.0	6.2	7.0	6.5	5.8	5.4	5.5	5.0	4.5	4.3	3.2	2.5	1.9	1.7	(1.9) ^s	
4	(1.9) ^s	2.0F	2.4F	(2.5) ^s	(2.6) ^s	(2.8) ^s	(3.0) ^s	(2.8) ^s	2.5	3.9	4.9	5.1H	5.4H	5.2	5.4	5.3F	5.6	4.3	4.6	4.2S	3.4F	2.8	2.6	(2.8)F	
5	3.2F	F	F	F	F	F	(2.9) ^s	(3.0) ^s	4.2	4.6	5.2	6.0	6.2	5.6	5.4	5.2	4.8	3.4	6.0	5.0	3.3F	3.3	3.2	3.0	
6	2.6F	2.5F	(2.8) ^s	(2.8) ^s	(2.8) ^s	(3.2) ^s	3.3	3.4F	2.5F	3.7	4.6	5.3	6.2	6.5	6.1	6.2	6.0	5.0	3.8	3.5	3.0	2.4	2.1	2.4	(2.6)F
7	2.3F	2.3F	2.4F	2.4F	2.9	2.9	3.1	2.7	4.2	4.4	4.7	5.2	5.2	5.8	6.4	5.4	4.8	3.9	2.4	2.1	2.0	2.0F	(2.3)F	2.6F	
8	2.6F	(2.6) ^s	2.8	(2.8) ^s	2.7	(2.7) ^s	3.2	(2.9) ^s	4.5	4.9	4.9	5.5	5.1	5.4	5.7	5.6	4.7	4.4	3.3	2.6	1.9F	(1.8) ^s	(1.8) ^s	(2.1)F	
9	2.2F	(2.1) ^s	2.2F	(2.8) ^s	3.4F	3.5F	(3.1) ^s	2.5F	4.1F	4.4	4.9H	5.2	6.0	6.1	5.0	5.4	4.8	3.8	2.7	2.4	(2.2)F	(1.9) ^s	1.8F	[1.8]F	
10	1.9F	[2.0] ^s	(2.0) ^s	(2.3) ^s	(2.3) ^s	(2.8) ^s	(2.8) ^s	3.1	3.0S	2.9F	3.7	4.4	5.1	6.2	5.9	5.5	5.7	5.8	5.1	4.3	3.5	(3.2)F	(2.4)F	2.5	3
11	(2.6) ^s	(2.5) ^s	(2.4) ^s	(2.4) ^s	3.1F	2.8	(2.8) ^s	3.2	2.8S	4.3	4.7	4.9	6.3	5.2F	5.6	5.6	5.9	5.0	4.4	3.1	2.6	(2.3)F	1.8F	1.7F	(1.7)F
12	1.8F	(1.8) ^s	(1.8) ^s	(1.8) ^s	3.6	(3.4) ^s	3.5	[3.9] ^s	4.3	4.7	5.5	5.6	6.2	5.8	6.4	6.2	5.4	5.2	4.4	2.4F	2.2	2.2	2.4F	2.5F	
13	(3.1) ^s	3.6	3.3S	(3.3) ^s	3.4F	(3.2) ^s	(3.2) ^s	(2.6) ^s	3.7	4.2	5.0S	5.7	[5.1] ^s	5.7	5.4	5.2	4.8	4.3	3.4F	2.8	2.4	2.1	2.0	2.2F	
14	2.0F	2.1F	(2.8) ^s	(2.8) ^s	3.0F	3.1S	2.6F	4.1	4.6	5.5	5.2	5.5	6.0	4.9	5.1	4.8	4.1	3.7	3.6	2.8F	(2.5)F	2.2F	2.2F	(2.4)F	
15	(2.7) ^s	(3.3) ^s	(3.5) ^s	(3.5) ^s	(3.0) ^s	3.6	(3.0) ^s	3.4	3.9	4.5	4.8S	5.4	5.5	5.4S	5.5	5.7	5.5	5.4	5.4	3.5	(3.3)F	3.4	2.8F	(2.8)F	
16	[2.4] ^s	(2.0) ^s	C	C	C	C	(3.6) ^s	(3.4) ^s	3.3	4.5	5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.3	5.6	4.4	(4.3)F	(3.1)F	(2.0)F	
17	(1.8) ^s	(2.2) ^s	3.1	(3.1) ^s	3.2S	3.1	2.5	2.9	(4.7)B	(5.3)B	(4.8)B	6.0	5.9	5.3	5.4	5.4	5.2	4.6	3.7	2.7F	2.6F	2.3F	2.4F	(2.6)P	
18	2.8F	3.1F	(2.8) ^s	(2.8) ^s	3.1F	3.2F	3.4F	3.3	3.2	4.4	4.9	5.4	5.4	6.2	6.0	5.5	5.5	5.6	5.0	4.7	5.4	3.2	(3.1)F	(3.2)F	(3.4)P
19	(3.6) ^s	(3.9) ^s	4.1	3.9	3.6F	2.8F	(2.5) ^s	(2.5) ^s	3.2F	5.0	(5.8) ^s	5.5F	6.2H	6.0	6.6	6.6	6.4	(7.2)S	7.2	(8.2)F	5.3	4.3F	2.8F	2.7	(2.8)F
20	(3.0)F	F	F	F	A	(2.9) ^s	2.7	(2.5) ^s	(2.8) ^s	4.5	5.9S	5.7	5.4	6.0	5.8	5.8	5.5	4.9	3.8	3.0	2.5	2.4	2.5	(2.8)F	
21	(2.8)S	3.1	3.2	3.1	2.7F	2.7F	2.0F	2.3	4.3	5.2	5.5	6.2	6.8	5.6S	5.8	6.0	5.7	4.7	4.4	3.4	2.7	2.1	2.0	2.2F	
22	(2.3) ^s	(2.7) ^s	2.4F	2.4F	2.4F	(2.2) ^s	(2.2) ^s	(2.6)A	4.8	4.9	5.6	5.6	6.0	6.3	6.6	5.8	4.9	4.9	3.6	2.7F	(2.5)S	(1.7)F	(2.0)S		
23	(1.9)F	2.3F	(3.0) ^s	[2.4] ^s	2.3F	(2.0) ^s	(2.0) ^s	(2.0) ^s	4.2	4.8	5.8	5.8	5.8	5.8	5.8	5.8	4.9	(2.7)F	2.8F	(2.7)F	2.1F	(1.8)S	(1.8)S		
24	2.0	2.0	2.0	2.1	2.4	2.6F	(3.2) ^s	(3.4) ^s	4.2	5.3	5.6	5.6	6.2	6.1	5.4	5.6	5.6	4.8	3.0F	2.4F	2.3	2.0F	1.7F	(1.8)F	
25	(1.8) ^s	2.1	(2.3) ^s	2.5F	(2.5) ^s	(2.9) ^s	(2.7) ^s	(2.7) ^s	4.4	5.0	5.6	6.5	5.7	5.4	5.8	5.6	4.5	4.5	3.3F	3.3	2.7	2.1	1.7	2.0	
26	2.0F	2.1F	2.3F	2.5	2.6F	2.8F	3.0	(3.2) ^s	(4.3) ^s	4.5H	5.4H	6.0	5.8	5.4	5.4	5.4	5.8	5.8	4.8	3.8	3.5	3.0	2.5	2.2	
27	2.4	2.7S	3.0	3.1F	3.2	3.3	3.3S	3.3S	4.7	4.9	5.1	5.7	5.6	5.7	6.0	5.8	6.0	4.9	4.3	3.9	3.0	2.4	2.3F		
28	(2.4) ^s	[2.1] ^s	(1.9) ^s	(1.9) ^s	(2.7) ^s	3.2F	3.2	2.8	3.2	5.0	5.8	6.3	6.0	5.8	5.6	5.6	5.6	5.5	3.8	2.3F	1.7F	2.6F	2.3F	2.3F	
29	2.1F	2.1F	(2.4) ^s	(2.9) ^s	3.2F	3.1F	3.2F	3.2F	4.9	5.4	5.4	5.7	5.6	5.5	6.2	6.8	6.8	4.9	3.6	2.4	2.2	1.8	S	S	
30	(1.7) ^s	[1.6] ^s	(1.6) ^s	(2.3) ^s	(2.6) ^s	2.8F	(2.3) ^s	2.8	4.7	4.9	4.9	5.7	6.0	(6.1)H	6.0H	5.4	5.6	4.6	4.2	3.7	3.1F	2.5F	2.2	2.2	
31	(2.5)F	(2.8)F	(2.8)F	(2.9)F	2.8	(2.8)F	(2.8)F	(2.8)F	4.2	5.4	5.0	5.1	(5.6)H	5.8	5.6	5.6	6.0	5.1	4.3	3.1F	2.4	2.1F	(2.0)F	1.7F	
Median	(2.4)	(2.3)	2.8	(2.5)	2.9	3.1	(3.0)	2.9	4.3	4.8	5.3	5.7	5.9	5.8	5.6	5.6	5.5	4.8	3.8	3.2	2.7	2.1	2.0	(2.3)	
Count	31	29	28	29	29	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	30	

Sweep 10 Mc to 25.0 Mc in 0.25 min

Manual □ Automatic □

N

TABLE 51

fo F₂, Mc (Unit)

January, 1954

(Month)

Washington, D.C.

Lat 38.7° N, Long 77.1° W

IONOSPHERIC DATA

National Bureau of Standards
(Institution)
E.J.W., J.W.P.

Scaled by:

M.C.

Calculated by:

M.C.

E.J.W.

J.W.P.

Day	75° W Moon Time												1830														
	0030	0130	0230	0330	0430	0530	0630	0730	0830	0930	1030	1130	1230	1330	1430	1530	1630	1730	1830	1930	2030	2130	2230	2330			
1	(1.4) S	(1.9) F	(1.9) P	(2.2) S	(3.2) F	(3.6) F	(3.2) F	(3.2) S	5.0	(5.4) S	6.4	6.6	5.8 F	6.0	5.2	4.1 F	(5.7) S	(5.4) S	4.4 S	3.2 S	3.0	(3.2) S	3.0	(2.8) F	(2.7) F		
2	(3.1) F	(3.3) P	(3.2) F	3.5	3.5	3.1	3.1 F	3.2 F	4.0	5.2	6.6	6.8	6.4	6.4	5.8	5.4	4.5	3.8	3.0 F	2.4 F	2.2 F	2.2 F	2.5 F				
3	2.8 S	3.2 F	3.4 F	3.3 F	3.2 F	3.4 F	3.4 F	3.8 F	5.0	6.3	6.6	6.8	6.1	5.4	5.5	5.4	4.6	4.2	3.7	3.0	3.0	3.0	3.0	1.3	(1.3) S		
4	1.7 S	2.3 F	2.7 F	2.6 F	2.9 F	3.0 F	(3.2) S	2.5 F	2.5	4.7	5.7	5.2 F	5.2 F	5.4	5.0	4.2	4.5	3.7	(3.9) F	2.7 F	(2.5) F	2.7 F	2.9 F	2.3 F			
5	3.3 F	F	F	F	F	F	(2.8) F	3.6 S	4.5	5.0	5.6	6.2	5.8	5.5	5.4	5.0	4.9	6.0	5.4	4.1	3.5 F	3.3 S	3.1	2.8 F			
6	(2.5) S	(4.6) F	(2.3) S	3.6 S	2	3.2 F	3.4	2.7 F	3.2	4.3	4.8 F	6.1	6.1	6.1	5.9	5.5	4.7	3.3	3.5	2.3	2.3	2.3	2.3	2.5	2.5		
7	2.3	2.2	2.4	2.8	2.9 S	3.1	3.1	3.3	4.1	4.7	5.0	5.4	5.4	6.4	6.6	5.4	4.3	3.1	(2.1) S	2.1	2.1	2.1	2.5 F	2.7			
8	(2.6) S	2.5	2.9 F	2.8	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	(1.8) F	(1.7) S	(1.7) F	(1.7) S	(1.7) F	(1.7) S			
9	(2.2) F	(2.0) F	(2.0) F	(2.5) F	3.1 F	3.4 S	3.4 S	3.3 S	2.5 S	3.7 S	4.2	4.4	4.9	5.7	6.2	5.5	5.2	5.3	4.3	3.1 S	2.4	2.3	2.3	(1.8) F			
10	(1.9) F	E	(2.1) F	2	3.5	(2.8) S	3.0	3.1 S	2.9	3.5	4.3	4.9	5.2	6.6	5.6	5.7	5.6	4.7	(3.2) S	3.3 S	2.8 F	(2.3) S	(2.4) F	(2.5) F			
11	2.7	2.4 S	2.8 F	(2.4) F	(2.9) F	(2.0) S	(2.0) F	(2.0) S	(2.8) F	3.5	4.6	4.6	5.6	6.5	5.5	5.5	5.5	4.9	3.8	3.1	2.3	1.8 S	1.7 S	(1.7) S			
12	(1.7) F	(2.3) F	3.3	3.7	3.5	3.4	3.3 S	3.4	3.7 S	4.4	4.9	4.9	5.7	6.4	6.5	6.5	5.9	5.1	5.0	3.4 F	2.3	2.3	2.4 F	2.8 F			
13	(3.2) S	3.7	3.3	(3.5) S	(2.8) F	(2.7) F	(2.7) F	3.1	4.2	(4.6) S	5.6	C	C	C	C	C	5.5	5.0	4.9	4.4	3.8	3.0	2.5	2.0 F	2.1 F		
14	2.1 S	2.2 F	2.3 S	2.3 F	2.9 F	3.0	3.0	3.0	3.0	3.5	4.5	4.7	5.2	5.4	5.8	5.6	5.0	5.2	4.4	3.9	3.7	3.5	2.3 S	2.2 F	(2.1) F		
15	(3.2) F	(3.4) F	3.5	(3.3) S	(3.1) F	(3.1) S	(3.1) F	(3.1) S	3.6	3.7	4.1	4.7	5.2	5.7	5.4	5.7	5.4	4.9	5.0	4.1	3.5 S	3.5 S	2.6 S	(2.9) S	(2.8) S		
16	F S	C	C	C	C	(3.5) S	(4.2) F	(4.2) S	(4.7) C	4.2	5.3	5.2	5.3	5.1	5.1	5.1	5.1	5.0	4.1	2.9 S	2.9 S	2.0 P	1.9 S	(1.8) F			
17	(1.7) F	(2.0) F	3.2	(3.4) S	3.2	2.8	2.5 P	3.2	3.8	5.3	5.2	5.6	6.0	5.2	5.2	5.5	4.9	4.3	2.9	2.5 S	2.5 S	2.3 F	2.6 S	2.6 S			
18	3.1	2.9 S	2.9 F	2.9 F	3.2 F	(3.3) S	3.2 F	3.2 F	3.9	5.1 S	5.3 H	5.0	5.8	6.1	5.4	5.8	5.6	5.7	4.7	5.1	4.5	3.1 S	(3.0) S	3.5 S	3.5 S		
19	(3.2) S	4.2	(3.7) S	(3.7) S	3.7 F	3.2 F	3.2 F	2.4 F	(2.4) S	(2.4) S	5.2	4.8	6.6	5.4 F	6.8	4.4	2.0	2.5	7.6 S	7.0	3.7	3.8	(2.6) S	(2.8) F	(2.7) F		
20	F	F	F	(1.2) F	(1.2) F	(1.2) F	(1.2) F	(1.2) F	(2.8) S	2.8	3.4 F	4.5	5.0	5.8	6.1	5.5	5.5	4.5	4.1	3.3	2.9	2.4	2.4 F	2.8 F			
21	2.9 F	3.2 F	3.3	2.8	2.5	3.0	2.0 S	2.0 S	2.6	3.2 F	5.2	5.0	5.8	6.4	6.4	5.8	5.9	5.2	5.0	4.5	4.1	3.3	2.9	2.4 F	2.8 F		
22	(2.5) F	(2.6) F	2.4 F	2.4 F	(2.3) F	2.2 F	2.3 F	2.3 F	2.3	3.2 F	5.0	5.1 S	5.8	7.0	6.4	5.8	5.9	5.9	5.7	5.5	5.5	5.5	5.5	5.5	5.5		
23	2.1 F	(2.2) S	(2.1) F	(2.1) F	(2.6) F	(2.6) F	(2.6) F	(2.6) F	(2.0) S	(2.0) S	5.2	5.2	5.2	5.2	5.2	5.2	5.4	5.4	5.4	5.4	5.4	5.4	5.4	5.4	5.4		
24	(2.0) S	2.0 F	2.0 F	2.0 F	2.2 F	2.2 F	2.2 F	2.2 F	2.1 F	4.1	4.8	5.4	(5.6) H	6.9	6.2	5.8	5.4	5.8	5.2	5.0	(2.4) S	2.4 F	2.0 F	1.8 F	1.8 F		
25	(2.0) F	2.2	2.2 F	2.2 F	2.4 F	2.6 F	2.7 F	2.7 F	2.7 F	2.8 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F	3.0 F			
26	2.1 F	2.3 F	2.3 F	2.7 F	3.0 F	3.7	4.7	5.5	6.0	6.0	5.8	5.4	5.2	5.6	4.4	3.4	3.3	2.8 F	2.5 S	2.3 S							
27	2.5	2.9	3.0	3.1 F	3.2 F	3.3	3.2 F	3.2 F	4.8	4.9	5.2 S	5.6	6.0	5.8 H	5.8	5.7	5.8	4.9	3.8	3.6	2.4	2.3	2.2	2.2	2.2		
28	2.2 F	2.0 F	(1.7) F	P	(2.5) F	3.0	2.8	4.0	4.9	6.0	(6.2) H	6.2	6.2	5.7	5.8	5.5 F	5.8	4.8	3.2	2.8 F	2.4 F	2.3 F	2.1 F	2.1 F			
29	(2.0) S	2.2 F	(2.6) F	3.1 F	3.1 F	3.2 F	2.9 F	4.0	5.3	5.9	5.5 H	5.7	5.4	6.0	6.0	6.0	6.0	4.4	3.0	2.8	(1.9) F	S	S	S	S		
30	(1.7) S	(1.9) S	(2.0) S	(2.4) F	P	(2.7) S	(2.8) S	(2.2) F	(3.8) S	4.9 H	4.5	6.0	6.0	5.9	5.2	5.4	5.8 S	4.9	4.4	3.4	2.8	2.3	2.3	2.3 F	2.3 F		
31	2.5	2.7 F	(2.9) F	(2.9) F	(2.0) F	2.3 F	2.3 F	3.7	4.6	(1.5) S	5.4	5.4	5.4	5.8 H	5.8	5.5	5.8 H	5.8	5.2	4.7	2.7	2.4 F	(1.3) S	1.7 S			
Median	2.4	2.8	2.9	3.0	3.0	2.8	3.7	4.7	5.0	5.6	6.0	5.8	5.6	5.7	5.5	5.0	4.3	3.5	2.9	2.4	2.2	2.2	2.2	2.2	2.2		
Count	2.9	2.8	2.8	2.9	2.9	2.9	2.9	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.0	3.0	3.0	3.0	3.0	3.0	

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual □ Automatic ■

Form adopted June 1946

1946 O - 1010

U.S. GOVERNMENT PRINTING OFFICE 1946 O - 1010

TABLE 53
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

IONOSPHERIC DATA

fo F1 Mc January, 1954

(Characteristic) (Unit) (Month)

Observed at Washington, D. C.

Lat 38°N, Long 77°W

75°W

Mean Time

Day 00 01 02 03 04 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23

		National Bureau of Standards																						
		McC. E. J.W., J.W.P.																						
		Calculated by: McC. E. J.W., J.W.P.																						
Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1																								
2																								
3																								
4																								
5																								
6																								
7																								
8																								
9																								
10																								
11																								
12																								
13																								
14																								
15																								
16																								
17																								
18																								
19																								
20																								
21																								
22																								
23																								
24																								
25																								
26																								
27																								
28																								
29																								
30																								
31																								
Median	-	-	3.6	3.7	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	
Count	1	2	7	19	21	20	14	4	4	1	-	-	-	-	-	-	-	-	-	-	-	-	-	

Sweep I.O. Mc 10 25 O. Mc in. 0.25 min
Manual Automatic

TABLE 54
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.
IONOSPHERIC DATA

<u>$h'E$</u> , <u>Km</u> (Characteristic)		<u>January</u> , 1954 (Month)		<u>Washington</u> , D. C. Observed at		Lat <u>38.7°N</u> , Long <u>77.1°W</u>												National Bureau of Standards						
						7 5°W Mean Time												Scaled by:						
						7 5°W Mean Time												MC.						
						7 5°W Mean Time												E.J.W., J.W.P. (Institution)						
Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1																								
2																								
3																								
4																								
5																								
6																								
7																								
8																								
9																								
10																								
11																								
12																								
13																								
14																								
15																								
16																								
17																								
18																								
19																								
20																								
21																								
22																								
23																								
24																								
25																								
26																								
27																								
28																								
29																								
30																								
31																								
Median																								
Count																								

Manual Automatic

Sweep 1.0 Mc to 25.0 Mc in 0.25-min

U. S. GOVERNMENT PRINTING OFFICE: 1946 - 2014

TABLE 56
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

Es, **Mc Km** **January**, 1954

(Characteristic) (Month)

Observed at **Washington, D.C.**

Lat **38.7°N**, Long **77.1°W**

National Bureau of Standards
(Institution)

Scaled by **MCC.**

E.J.W.

J.W.P.

7.5°W												Mean Time													
Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	24/120	24/120	E	20/120	24/110	G	22/110	E	22/110	G	28/120	3/1	4/2	4/2	4/2	4/2	3/8	1/20	4/1	1/10	2/4	1/10	3/4	1/10	
2	E	E	E	E	3/0/140	3/4/140	H	2/8/100	3/1/100	3/2/100	4/9/100	3/7/100	3/3/100	4/3/100	2/9/100	3/6/100	3/9/100	4/7/100	2/9/100	3/4/100	2/9/100	2/9/100	E	E	
3	E	E	E	E	E	E	E	E	E	E	G	3/7/100	G	G	G	G	G	G	G	G	G	G	G	E	
4	E	E	E	E	9/0/10	6/8/10	E	2/6/120	3/0/120	3/0/110	G	1/3/100	G	G	G	G	3/6/110	2/4/120	1/9	1/10	2/0/120	1/2/100	3/3/100	2/4/110	E
5	E	E	E	E	2/4/100	2/4/100	E	2/4/100	2/4/100	2/3/100	0/3/05	2/9/110	G	G	G	G	1/4/100	G	G	G	E	E	E	E	E
6	E	E	E	E	2/4/110	E	E	2/3/110	E	G	2/4/110	2/6/110	2/4/3/10	1/0	2/7/110	G	G	2/3/110	G	G	2/6/110	2/4/110	2/7/110	E	E
7	24/100	27/100	22/100	E	22/100	E	E	24/110	G	G	G	G	G	G	G	G	2/1/20	E	E	E	2/4/110	E	E	E	E
8	E	E	24/120	3/9/110	2/7/110	3/2/100	2/4/110	3/8/100	3/9/100	G	G	G	2/4/100	G	G	G	1/9/100	2/4/110	1/9/100	2/4/110	2/3/100	E	E	E	E
9	E	E	E	E	E	E	E	2/0/120	E	5/5/110	G	G	G	G	G	G	2/4/120	E	E	E	E	E	E	3/2/100	
10	E	E	E	E	E	E	E	E	E	E	2/4/130	3/2/110	G	G	G	G	3/8/120	G	G	G	E	E	E	E	E
11	E	E	E	E	E	E	E	E	E	E	9/0/110	3/4/120	G	G	G	G	G	G	G	G	E	E	E	E	E
12	E	E	E	E	E	E	E	E	E	E	C	G	G	G	G	3/6/100	G	G	G	2/6/100	2/4/100	2/3/100	E	E	
13	E	E	E	E	E	E	E	E	E	E	3/2/110	3/6/110	3/5/110	C	G	3/1/110	G	G	G	E	E	E	E	3/4/110	
14	2/2/120	2/4/110	E	E	3/7/120	E	E	2/3/110	E	3/6/110	3/7/110	3/6/110	G	G	G	3/7/110	G	G	E	E	E	E	E	E	
15	E	E	22/110	E	E	E	E	E	E	E	3/2/110	3/7/110	3/3/110	3/6/110	3/7/120	G	G	3/5/110	3/1/110	2/9/110	2/3/110	2/3/110	2/5/100		
16	2/9/100	2/8/100	C	C	C	C	C	2/4/110	G	2/8/110	3/0/110	3/0/110	3/0/110	3/5/110	G	G	3/1/110	G	2/4/110	3/5/110	3/1/110	E	E	E	
17	E	E	E	E	E	E	E	E	E	E	E	3/6/120	3/3/110	3/5/110	1/0	17/2/120	4/9/110	3/7/120	3/3/110	2/8/110	2/4/110	3/1/100	2/3/100	E	
18	E	E	E	E	E	E	E	E	E	E	E	2/4/110	2/4/110	2/3/110	G	G	G	G	G	E	E	E	E	E	E
19	E	E	2/2/20	3/0/110	E	2/4/100	E	2/4/100	E	G	G	G	G	G	G	3/7/100	G	G	G	2/8/110	2/4/110	3/0/110	E	E	
20	E	E	3/1/30	4/2/20	7/5/110	E	E	4/2/110	G	G	G	G	G	G	G	2/1/100	2/8/110	3/8/100	3/0/100	E	E	E	2/6/110	2/9/100	
21	2/6/100	E	E	2/3/100	2/7/120	2/3/120	2/4/110	E	2/0/100	G	3/7/110	1/0/2/100	5/4/100	4/7/110	4/2/110	4/2/110	4/2/110	3/2/110	2/5/120	3/0/110	2/8/120	2/7/120	2/4/120	E	
22	2/3/10	3/0/110	2/7/110	H	2/2/20	E	5/0/120	4/4/120	4/1/110	3/7/110	2/9/110	G	G	G	G	G	2/3/20	E	E	E	E	E	E	2/2/100	
23	E	E	E	E	E	E	E	E	E	E	E	3/6/110	3/6/110	3/9/100	3/1/100	3/8/110	3/3/110	3/8/120	E	E	E	E	E	E	2/7/100
24	E	E	E	E	2/4/100	E	E	2/3/110	2/3/100	E	G	G	G	G	G	G	G	G	E	E	E	E	E	2/8/100	
25	3/4/100	2/6/100	E	E	E	E	E	E	E	E	E	2/1/100	E	G	G	3/4/110	G	2/6/110	2/4/110	3/0/110	2/4/110	4/3/100	2/9/100	E	
26	E	2/3/100	E	4/1/100	3/8/110	E	E	2/4/100	2/0/120	4/0/20	4/2/20	3/5/110	G	3/3/110	3/4/110	4/0/200	3/3/100	2/2/100	E	E	E	E	E	E	2/2/100
27	2/6/100	E	E	E	E	E	E	E	E	E	E	2/0/120	2/9/120	3/9/100	0/0	2/8/110	2/7/110	2/7/120	E	E	E	E	E	E	2/1/100
28	2/2/100	E	E	E	E	E	E	E	E	E	E	3/2/120	E	G	G	G	G	G	G	G	E	E	E	2/9/100	
29	3/4/100	2/4/100	2/3/100	E	2/1/100	4/4/100	2/1/100	2/3/100	2/1/100	G	4/1/110	G	G	G	G	G	3/1/110	E	2/4/110	3/0/110	3/7/100	2/9/100	E	E	
30	E	E	E	E	E	E	E	E	E	E	E	2/8/100	6/2/100	3/7/100	G	G	G	2/0/110	G	E	E	3/1/100	3/6/100	E	E
31	E	2/3/100	E	E	E	E	E	E	E	E	E	2/2/110	G	G	G	G	G	G	G	2/3/20	E	2/2/100	E	E	
Median	**	**	**	**	**	**	**	**	**	**	2/1	**	**	2/4	**	**	**	**	**	**	**	**	**	**	
Count	31	31	30	30	31	30	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	

XX MEDIAN F LESS THAN MEDIAN F_{oe}
OR LESS THAN LOWER FREQUENCY LIMIT OF RECORDER

Sweep 10 Mc 1025.0 Mc in 0.5 min
Manual □ Automatic ■

TABLE 57
IONOSPHERIC DATA
 Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
 (Month) **January**, 1954
 Lat **38.7°N**, Long **77.0°W**

(M1500) F2
 (Characteristic)
 (Unit)
 Observed at **Washington, D.C.**

Mean Time

Day	75°W												Mean Time												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	A	(2.1)F	(2.0)F	(2.3)F	F	(2.3)F	F	2.6	2.6	2.5	2.4	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	
2	(2.5)F	(2.0)F	(2.0)F	(1.9)F	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
3	2.0	(2.2)F	(2.2)F	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
4	(2.0)S	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
5	2.2	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	
6	2.2	(2.1)F	(2.1)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F	(2.0)F											
7	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F	(2.2)F														
8	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
9	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
10	2.5	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	
11	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S														
12	5	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F													
13	(2.1)F	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
14	(2.2)F	2.1	2.1	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
15	5	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	
16	F	(2.0)F	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
17	F	(2.1)F	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
18	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
19	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S	(2.1)S														
20	(2.0)F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	
21	(2.1)S	2.1	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
22	(2.2)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F													
23	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
24	2.0	2.1	2.1	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
25	(2.1)F	2.0	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
26	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
27	2.1	2.1	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
28	(2.1)F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	
29	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
30	5	5	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	
31	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F	(2.1)F														
Median	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	(2.1)	
Count	24	24	27	28	29	31	27	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31

Calculated by: **Mc. E.J.W. J.W.P.**

Scaled by: **Automatic**

Manual Automatic

Sweep **10 Mc 102.50 Mc 10.25 min**

U. S. GOVERNMENT PRINTING OFFICE 16-1015-19

TABLE 58
IONOSPHERIC DATA

(M3000) F2, January, 1954
(Characteristic) (Month)
Observed at Washington, D.C.

Lat 38.7°N, Long 77.1°W

7.5°W Mean Time

Day	7.5°W												23												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
	M.C.	E.J.W.	J.W.P.																						
1	A	A	(3.2)F	(3.0)F	(3.3)F	F	(3.3)F	F	3.7	3.7 F	(3.4)F	3.5	3.5 F	3.4 F	3.4	3.6 F	3.6 F	3.4 F	3.3 F	3.4 F	3.3 F	3.4 F	3.3 F	3.3 F	
2	(3.5)F	(3.0)F	(3.0)F	(2.8)F	3.2	3.1	3.1 F	(3.4)F	3.5	3.4	3.6	3.3	3.4	3.5	3.5	3.5	3.5	3.4	3.3	3.3	3.3	3.2 F	3.2 F	3.3 F	
3	3.2 F	(3.1)F	(3.2)F	3.1 F	3.1 F	3.4 F	3.4 F	3.4	3.4	3.6	3.6	3.6	3.7	3.4	3.6	3.6	3.5	3.5	3.4	3.7	3.1	3.0	5	(3.1)F	
4	(3.0)F	3.1 F	3.1 F	(3.1)F	3.3 F	(3.3)F	(3.3)F	3.4	3.4	3.6	3.7	3.5 H	3.6 H	3.6	3.5 F	3.6	3.5	3.3	3.3 S	3.3	3.3	3.2	3.2	(3.0)F	
5	3.1 F	F	S	F	F	F	(3.2)F	F	3.7	3.6	3.5	3.5	3.5	3.5	3.4	3.4	3.4	3.4	3.2 F	3.3	3.2	3.2	3.2	3.3 F	
6	3.2 S	(3.1)F	(3.2)F	(2.8)F	(3.0)S	(3.0)S	(3.0)S	3.1	(3.5)F	(3.3)F	3.5	3.6	3.4	3.5	3.4	3.6	3.7	3.5	3.4	3.2	3.4	2.9	3.1	(3.2)F	
7	(3.2)F	(3.2)F	(3.1)F	(3.2)F	3.1	3.2	3.4	3.5	3.7	3.5	3.6	3.5	3.3	3.4	3.7	3.7	3.6	3.3	3.1	3.0	2.9 S	(2.9)F	3.1 F	3.1 F	
8	3.2 F	(3.1)S	3.0	(3.0)S	3.2	(3.0)F	(3.0)F	3.2	(3.4)F	3.5	3.6	3.6	3.4	3.5	3.4	3.5	3.6	3.7	3.5	3.4 F	3.5	3.4 F	F	(3.1)F	
9	3.0 F	(3.1)F	3.2 F	(3.1)F	3.1 F	3.1 F	3.3 F	(3.2)F	3.4 F	3.7	3.3	3.4 F	3.3	3.6	3.6	3.5	3.7	3.5	3.2 F	3.2	3.2	F	S	F	
10	F	S	(3.2)F	(3.2)F	(3.2)F	3.4	3.4	3.4	3.5	3.6	3.7	3.6	3.4	3.5	3.5	3.6	3.6	3.5	3.2	(3.4)F	3.2	3.2	(3.0)S	(3.0)F	3.1 S
11	(3.0)S	(3.2)F	(3.2)F	(3.2)F	(3.0)F	3.3 F	3.0	(3.2)F	3.5	3.7	3.7	3.4	3.5	3.6 F	2.9	3.5	3.5	3.5	3.2	3.4	5	5	5	5	5
12	S	(3.2)F	(3.2)F	(3.2)F	(3.2)F	3.3 F	3.0	(3.2)F	3.5	3.7	3.7	3.4	3.5	3.6 F	2.9	3.5	3.5	3.5	3.2	3.4	5	5	5	5	5
13	(3.1)F	3.1	3.2 S	3.2 F	3.1 F	3.1 F	3.1 F	(3.1)F	3.5	3.6	3.5	3.5	3.6	3.5	3.5	3.5	3.6	3.5	3.2 F	3.0	2.9	3.1 F	3.0 F	3.0 F	
14	(3.3)F	3.1 F	3.0 F	(3.2)F	(3.2)F	3.2 S	3.2 S	3.4 F	3.4 F	3.7	3.7	3.4	3.6	3.5	3.6	3.5	3.6	3.5	3.4 F	3.3	3.3	3.2 F	3.1 F	3.1 F	
15	S	F	F	(3.1)F	(3.0)F	3.1	(3.1)F	3.3	3.5	3.6	(3.4)F	3.5	3.5	3.4 S	3.6	3.6 S	3.4	3.6 S	3.4	(3.2)F	3.4	3.2 S	(3.1)F	(3.2)F	(3.2)F
16	F	S	(3.0)F	C	C	C	(3.4)F	3.6	3.7	(3.7)F	3.6	3.6	3.5	3.5	3.4	3.6	3.5	3.5	(3.5)F	(3.5)F	5	F	F	F	
17	F	(3.3)F	3.2	(3.4)F	3.3 S	3.4	3.2	(3.4)F	3.4	(3.7)F	3.7	3.7	3.7	3.4	3.3	3.6	3.5	3.3	3.2 F	3.5 F	3.3 F	3.2 F	3.2 F	(3.2)F	
18	3.1 S	3.3 F	(3.2)F	(3.2)F	3.1 F	3.2 F	3.2 F	3.3	3.7	3.7	3.7	3.6	3.5	3.4	3.5	3.4	3.5	3.4	3.0 F	3.5	3.4	(3.0)F	(3.1)F	(3.1)F	
19	(3.1)F	(3.1)F	3.2	3.2	3.2 F	3.4 F	3.4 F	F	3.7	(3.7)F	F	3.2	3.4 F	2.9	(3.1)F	3.1	(3.3)F	3.4	(3.5)F	3.3 S	3.1	3.2	F	F	
20	(3.0)F	F	F	A	(3.1)F	3.0	(3.0)F	(3.0)F	3.5	(3.4)F	3.5	3.6 S	3.6	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.3	3.0	3.0	(3.0)F	
21	(3.1)F	3.1	3.0	3.4	3.5 F	3.1	3.2 F	3.2 F	3.4	3.6	3.5	3.5	3.5	(3.6)F	3.5	3.4	3.6	3.5	3.5	3.2 F	3.3	3.2	3.1	3.1 F	3.1 F
22	(3.3)F	(3.3)F	3.2 F	3.4 F	3.2 F	3.2 F	(3.3)F	(3.1)F	3.4 F	3.5	3.7	3.6	3.3	3.4	3.3	3.5	3.6	3.5	(3.5)F	3.5	3.4 F	3.2 F	3.2 F	(3.2)F	
23	(3.1)F	F	3.1 F	(3.1)F	(3.1)F	3.2 F	(3.4)F	(3.2)F	(3.5)F	3.5	3.6	3.3	3.6	3.5	3.5	3.4	3.5	3.6	(3.3)F	3.3 F	3.3 F	(3.3)F	(3.3)F	(3.3)F	
24	3.0	3.1	3.1	3.0	3.2	3.1	3.1 F	(3.1)F	3.7	3.6	3.7	3.4	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	F	
25	(3.1)F	3.0	(3.1)F	P	3.1 F	(3.2)F	(3.4)F	(3.5)F	(3.4)F	3.7	3.5	3.4	3.5	3.6	3.5	3.4	3.6	3.5	3.5	3.5	3.5	3.5	3.5	3.1	
26	3.2 F	3.3 F	3.1 F	3.2 F	3.2 F	3.1 F	(3.3)F	(3.3)F	3.2 F	(3.6)F	3.4	3.4	3.4	3.4	3.5	3.5	3.6	3.5	3.5	3.4 F	3.4	3.3	3.3	3.1 F	
27	3	3.2	3.2	3.2	3.1 F	3.1	3.2	3.3 S	3.4 S	3.7	3.6	3.4	3.4	3.4	3.5	3.5	3.5	3.5	3.5	3.2	3.4 F	3.3	3.3	3.3 F	
28	(3.2)F	F	(3.1)F	(3.0)F	3.2 F	3.4 F	3.3	3.3	3.5	3.6	3.7	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.4 F	3.4	3.4	3.4 F	3.3 F	
29	3.1 F	3.2 F	(3.3)F	(3.3)F	3.1 F	3.1 F	3.3 F	3.3 F	3.5 F	3.6	3.6	3.5	3.5	3.5	3.4	3.4	3.4	3.4	3.4 F	3.4	3.4	3.4 F	3.3 F	3.3 F	
30	S	5	(3.3)F	(3.1)F	(3.1)F	F	(3.4)F	(3.5)F	3.6	3.5	3.7	3.3 H	3.5	3.4	(3.4)H	3.6	3.4	3.4	3.4	3.4	3.3	3.4	3.4 F	3.3 F	
31	(3.1)F	(3.0)F	(3.1)F	(3.1)F	(3.1)F	F	(3.0)F	(3.0)F	(3.3)F	3.3	3.3	3.3	3.3	3.3	(3.3)F	3.5	3.4	3.4	3.4	3.4	3.6	3.6	3.6	(3.0)S	
Median	(3.1)	(3.1)	(3.1)	3.2	3.2	(3.3)	3.4	3.4	3.6	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.6	3.5	3.4 F	3.3	3.4	3.2	3.1	3.1	
Count	24	24	27	28	28	27	31	31	31	30	31	31	31	31	31	31	31	31	31	30	30	30	25	24	

Sweep 10 Mc to 25.0 Mc in 0.25 min
Manual □ Automatic □

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

TABLE 59
IONOSPHERIC DATA

(M3000) Fe, (Unit)
(Characteristic) January, 1954

Observed at Washington, D.C.
(Month)

Lat 38°N, Long 77°W

Day	75°W												Mean Time												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1																									
2																									
3																									
4																									
5																									
6																									
7																									
8																									
9																									
10																									
11																									
12																									
13																									
14																									
15																									
16																									
17																									
18																									
19																									
20																									
21																									
22																									
23																									
24																									
25																									
26																									
27																									
28																									
29																									
30																									
31																									
	-	-	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	
	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

Sweep 10 Mc 1025.0 Mc in 025 min
Manual Automatic

U. S. GOVERNMENT PRINTING OFFICE: 1954 1-1719

(M1500) E., (Umt)
(Characteristic)
Observed at Washington, D.C.

January, 1954
(Month)
Lat. 38.7°N, Long 77.1°W

National Bureau of Standards
(Institution)
Scaled by: McC. E. J.W., J.W.P.

TABLE 60
IONOSPHERIC DATA

Day	75° W Mean Time												19	18	17	16	15	14	13	12	11	10	09	08	07	06	05	04	03	02	01	00	
	McC.	E.	J.W.	J.W.P.	McC.	E.	J.W.	J.W.P.	McC.	E.	J.W.	J.W.P.																					
1	S	(4.4)F	4.5H	4.4"	A	A	A	A	S	(4.5)A	4.5	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4				
2	S	4.2H	4.4H	4.4"	S	(4.2)H	4.4	4.4	S	4.2H	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4				
3	S	(4.2)H	4.4	4.4"	S	4.3	4.4	4.4	S	(4.2)H	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4				
4	S	3.9	4.1	4.2	S	4.1	4.2	4.2	S	3.9	4.1	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2				
5	S	4.1	4.2	4.3	S	4.1	4.2	4.3	S	4.1	4.2	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3				
6	S	4.1	4.2	4.3	S	4.1	4.2	4.3	S	4.1	4.2	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3				
7	S	4.1	4.2	4.3	S	4.1	4.2	4.3	S	4.1	4.2	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3				
8	S	4.4H	4.3	(4.2)P	S	4.4H	4.3	(4.2)P	S	4.4H	4.3	(4.2)P	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4			
9	S	4.2F	4.2	4.2	S	4.2F	4.2	4.2	S	4.2F	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2				
10	S	(4.3)P	4.3	4.3	S	(4.3)P	4.3	4.3	S	(4.3)P	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3				
11	S	4.3	4.2H	4.2	S	4.3	4.2H	4.2	S	4.3	4.2H	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2				
12	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H																					
13	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H																					
14	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H																					
15	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H	S	4.3	4.2H	4.3H																					
16	S	(4.4)P	(4.1)H	4.2H	S	(4.3)H	(4.2)H	(4.1)H	S	(4.3)H	(4.2)H	(4.1)H	(4.0)H	(4.1)H	(4.2)H	(4.3)H	(4.4)H	(4.5)H	(4.6)H	(4.7)H	(4.8)H	(4.9)H	(4.0)H	(4.1)H	(4.2)H	(4.3)H	(4.4)H	(4.5)H					
17	S	4.3	4.3	4.4H	S	4.3	4.3	4.4H	S	4.3	4.3	4.4H	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4			
18	S	4.3H	4.3	4.4H	S	4.3H	4.3	4.4H	S	4.3H	4.3	4.4H	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4			
19	S	4.3H	(4.1)P	4.1	S	4.3H	(4.1)P	4.1	S	4.3H	(4.1)P	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1			
20	S	4.3	(4.3)H	(4.4)H	S	4.3	(4.3)H	(4.4)H	S	4.3	(4.3)H	(4.4)H																					
21	S	4.3F	4.3	A	S	4.3F	4.3	A	S	4.3F	4.3	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A			
22	S	4.0	4.0	4.1	S	4.0	4.0	4.1	S	4.0	4.0	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1			
23	S	4.1	4.1	4.1	S	4.1	4.1	4.1	S	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1			
24	S	3.9	4.1	3.9	S	3.9	4.1	3.9	S	3.9	4.1	3.9	4.0	4.1	4.0	4.1	4.0	4.1	4.0	4.1	4.0	4.1	4.0	4.1	4.0	4.1	4.0	4.1	4.0	4.1	4.0		
25	S	4.4	(4.4)P	4.2	S	4.4	(4.4)P	4.2	S	4.4	(4.4)P	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2			
26	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A				
27	A	(4.2)A	(4.1)A	A	A	(4.2)A	(4.1)A	A	A	(4.2)A	(4.1)A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A			
28	(4.3)S	4.0	4.1	4.4	(4.3)S	4.0	4.1	4.4	(4.3)S	4.0	4.1	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4		
29	4.2H	4.1	4.0	4.2	4.2H	4.1	4.0	4.2	4.2H	4.1	4.0	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2			
30	(4.3)A	4.0H	4.0	4.3	(4.3)A	4.0H	4.0	4.3	(4.3)A	4.0H	4.0	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3			
31	F	A	4.2	4.2	F	A	4.2	4.2	F	A	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2		
Median Count	-	4.2	4.2	4.3	4.3	4.2	4.2	4.3	4.3	4.2	4.2	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.3		
Count	4	19	23	25	25	23	23	25	25	23	23	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25

Sweep 0 Mc 1025.0 Mc in 0.25 min
Manual Automatic

1 GOVERNMENT PRINTING OFFICE: 1946 O 705-19

Table 61

Ionospheric Storminess at Washington, D. C.January 1954

Day	Ionospheric character*		Principal storms		Geomagnetic character**	
	00-12 GCT	12-24 GCT	Beginning GCT	End GCT	00-12 GCT	12-24 GCT
1	3	1			2	1
2	1	1			3	4
3	1	1			2	2
4	2	3			0	0
5	2	2			0	2
6	2	1			2	2
7	2	2			1	1
8	1	3			3	2
9	3	3			2	2
10	2	2			2	2
11	1	2			2	1
12	2	1			2	2
13	1	2			3	2
14	2	2			1	2
15	1	3			2	2
16	3	2			2	2
17	1	2			1	2
18	1	2			2	3
19	2	2			4	3
20	2	2			3	3
21	1	1			3	3
22	1	1			3	2
23	1	2			4	2
24	3	1			2	1
25	3	1			1	2
26	2	2			1	1
27	1	2			1	2
28	3	1			2	0
29	2	2			2	1
30	3	1			1	1
31	2	2			2	2

*Ionosphere character figure (I-figure) for ionospheric storminess at Washington, D. C., during 12-hour period, on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

**Average for 12 hours of Cheltenham, Maryland, geomagnetic K-figures on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

Table 62Sudden Ionosphere Disturbances Observed at Washington, D. C.January 1954

No sudden ionosphere disturbances were observed during the month of January.

Table 63

Sudden Ionosphere Disturbances Reported by Direction Générale des Télécommunications de Suède, as Observed at Enköping, Sweden

1953 Day	GCT		Location of transmitters	Other phenomena
	Beginning	End		
October 10	1405	1425	Lebanon	
14	0955	1005	Argentina, Austria, Brazil, Bulgaria, Czechoslovakia, Netherlands, Peru, Switzerland, Tangier	Solar flare* 1012

*Time of observation at Wendelstein Observatory, Germany. Flare began before time of observation.

Note: Observers are invited to send to the CRPL information on times of beginning and end of sudden ionosphere disturbances for publication as above. Address letters to the Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

Table 64a

Radio Propagation Quality Figures
 (Including Comparisons with Short-Term and Advance Forecasts)

December 1953

Day	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day quality index	Advance forecasts (J-reports) for whole day; issued in advance by:			Geomag- netic K_{Ch}	
	00	06	12	18	00	06	12	18		1-4 days	4-7 days	8-25 days	Half day (1)	(2)
	ts	to	to	to	06	12	18	24						
1	6	6	7	6	6	5	7	7	6	7	7		1	1
2	7	6	7	6	6	6	7	7	6	6	6		2	1
3	6	6	7	7	6	5	7	6	7	6	6		2	2
4	6	6	7	6	6	6	7	7	7	7	7		2	2
5	6	6	7	7	6	6	7	7	6	7	7		2	1
6	5	6	7	7	6	5	7	6	7	7	7		1	2
7	6	6	7	6	6	6	7	7	7	7	7		2	1
8	6	6	7	7	6	7	7	6	6	7	7		2	2
9	6	6	6	6	6	5	6	6	6	5	5		1	2
10	6	6	7	6	(4)	5	6	5	6	(4)	(4)	X	1	2
11	6	6	7	6	5	(4)	6	5	6	(4)	(4)	X	3	(4)
12	5	5	6	6	(4)	(4)	5	5	6	(3)	(3)	X	(4)	3
13	5	5	6	6	(4)	(4)	5	5	6	(3)	(3)	X	2	3
14	5	5	6	5	5	(4)	6	5	5	(4)	(4)	X	2	2
15	5	6	7	6	(4)	(4)	6	6	6	5	(4)	X	1	2
16	5	6	7	6	5	5	7	6	6	(4)	(4)	X	2	2
17	6	6	7	6	5	5	7	6	6	(4)	(4)	X	1	2
18	5	6	7	6	6	5	6	6	6	6	5		2	1
19	6	6	6	6	6	6	7	7	6	6	5		1	2
20	6	6	7	6	6	6	7	6	6	5	5		1	2
21	5	6	7	7	5	5	7	6	6	5	5		1	1
22	6	5	7	6	6	6	6	6	6	6	6		2	2
23	5	6	7	6	6	5	7	6	6	6	6		0	2
24	5	6	7	6	5	5	6	6	6	6	5		2	1
25	6	6	7	6	6	6	7	6	6	6	6		1	2
26	6	5	7	7	6	6	7	7	6	6	6		1	2
27	6	6	7	6	6	6	7	7	6	7	7		1	2
28	6	6	7	6	6	6	7	7	6	7	7		2	2
29	6	6	7	6	7	6	7	7	6	7	7		1	3
30	6	5	7	7	6	6	7	7	6	7	7		1	1
31	6	6	6	6	7	6	7	7	6	7	7		2	1
Score:														
Quiet periods														
P	19	11	21	14						11	8			
S	11	18	10	17						14	16			
U	0	0	0	0						0	0			
F	1	2	0	0						6	7			
Disturbed periods														
P	0	0	0	0						0	0			
S	0	0	0	0						0	0			
U	0	0	0	0						0	0			
F	0	0	0	0						0	0			

Scales:

- Q-scale of Radio Propagation Quality
 (1) - useless
 (2) - very poor
 (3) - poor
 (4) - poor to fair
 5 - fair
 6 - fair to good
 7 - good
 8 - very good
 9 - excellent

K-scale of Geomagnetic Activity
 0 to 9, 9 representing the greatest disturbance; $K_{Ch} \geq 4$ indicates significant disturbance, enclosed in () for emphasis

Scoring: (beginning October 1952)


- P - Perfect: forecast quality equal to observed
 S - Satisfactory: (beginning October 1952)
 forecast quality one grade different
 from observed
 U - Unsatisfactory: forecast quality two or more
 grades different from observed when both
 forecast and observed were ≥ 5 , or both ≤ 5
 F - Failure: other times when forecast quality
 two or more grades different from observed

Symbols:

- X - probable disturbed date

Table 64b

Short-Term Forecasts---December 1953



Outcome of Advance Forecasts (1 to 4 days ahead) --- December 1953




Table 55a

Coronal observations at Climax, Colorado (5303A), east limb

Table 55.

Coronal observations at Climax, Colorado (6374A), east limb

Date GCT	Degrees north of the solar equator															0°	Degrees south of the solar equator																		
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90
1954																																			
Jan.	1.7	1	1	1	2	1	1	1	1	2	1	3	3	4	5	6	6	6	6	6	6	6	4	3	1	1	1	1	1	1	2	2	2		
	2.8	2	1	1	3	2	1	-	-	-	1	3	3	2	4	6	5	5	5	5	5	5	4	4	3	2	1	1	1	1	1	1	2	2	
	3.7a	1	1	2	3	2	1	1	1	1	2	3	3	1	2	4	7	6	6	6	6	6	6	4	3	1	1	1	1	2	2	3	3	2	
	5.7	2	2	1	1	1	1	1	1	1	2	1	2	4	5	6	7	6	6	6	5	3	4	4	5	3	2	1	1	1	2	2	4	3	
	6.7	2	1	1	1	1	1	1	1	1	2	2	2	3	4	5	6	6	7	8	8	5	5	5	5	3	2	1	1	1	2	2	2	3	
	7.7	2	2	1	1	1	1	1	1	1	2	2	2	3	5	7	6	6	9	8	8	9	7	6	6	6	5	4	2	2	3	4	4	4	
	8.9b	2	2	2	2	2	2	2	2	2	2	2	2	3	5	7	6	6	9	8	8	9	7	6	6	6	5	4	2	2	3	4	4	3	
	10.8	3	4	4	2	2	3	2	1	2	3	3	3	4	5	3	4	4	5	6	8	8	9	7	6	7	7	8	2	3	3	3	3		
	14.9	2	1	1	1	-	-	-	-	1	1	1	1	1	1	2	2	2	2	2	3	2	2	2	1	1	1	1	1	2	2	2	2		
	21.9	3	3	2	2	1	1	-	-	1	2	4	5	4	4	4	3	2	2	4	5	5	5	5	4	3	3	3	2	1	1	1	1	3	
	22.7	3	2	2	2	1	1	1	1	1	3	6	5	5	5	3	5	5	7	7	7	6	6	5	5	4	3	3	2	1	2	2	2	4	
	23.7	3	3	2	1	1	-	-	-	1	1	2	3	4	3	4	3	2	4	5	5	6	5	3	3	3	2	2	2	1	1	2	3	3	
	27.8	X	X	X	X	X	X	X	X	1	2	1	1	1	1	2	2	2	2	3	3	3	3	1	1	1	1	1	1	2	2	X	X	X	
	28.9	2	2	1	1	1	1	1	1	1	2	3	3	4	5	5	4	5	5	6	5	5	3	3	3	3	1	1	1	1	2	2	2		
	30.9	2	2	1	1	1	1	1	1	1	2	3	3	2	4	5	5	5	5	4	4	4	5	5	3	2	1	1	1	1	2	2	2		
	31.7	2	2	2	2	3	2	1	1	1	1	2	2	2	2	6	9	5	5	5	4	3	5	5	4	3	3	2	2	2	2	3	3	3	

Table 57a

Coronal observations at Climax, Colorado (6702A), east limb

a indicates low weight 190 through 597

t indicates low weight N9 through N27

Table 55b

Coronal observations at Climax, Colorado (5303A), west limb

Date GCT	Degrees south of the solar equator															00°	Degrees north of the solar equator																		
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Jan. 1.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	1	-	-	1	2	2	1	1	1	1	-	-	-	-	-	-	
2.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	5	3	2	-	-	1	2	2	1	1	1	1	-	-	-	-	-	
3.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	1	2	1	1	1	-			
6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
7.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	2	1	1	-	-	-	-	-	-	
8.9c	-	-	-	-	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-
10.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	2	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
14.9d	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	1	1	1	1	-
21.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	1	1	1	1	1	-			
22.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	3	2	1	1	1	1	1	-			
23.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	1	1	1	1	1	-			
27.8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	
28.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-
30.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-
31.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	1	1	1	1	2	1	-	-	-			

Table 66b

Coronal observations at Climax, Colorado (6374A), west limb

Date GCT	Degrees south of the solar equator															Degrees north of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1954																																				
Jan. 1.7	2	2	2	2	1	1	1	1	2	3	3	3	4	4	5	8	4	3	4	3	1	1	2	4	3	1	1	1	1	1	1	2	2	1		
2.8	2	2	2	1	1	1	1	1	3	2	3	2	4	4	4	7	6	6	5	3	2	3	3	5	4	3	1	1	1	1	1	1	2	2	2	
3.7a	2	2	2	2	2	1	1	1	2	3	3	3	4	4	3	5	5	5	4	4	3	3	2	6	5	3	1	1	1	1	1	2	2	1		
5.7	3	3	2	2	2	2	1	1	3	4	5	5	6	6	6	6	7	6	6	5	4	4	4	5	4	3	3	2	1	1	2	2	2	1		
6.7	3	3	2	2	2	2	1	2	1	3	4	5	5	5	5	4	4	5	4	4	3	4	5	4	2	2	2	2	2	3	3	2	1			
7.7	3	3	3	2	2	2	1	1	1	4	4	4	4	5	5	5	4	4	4	4	4	4	4	3	3	3	1	1	2	2	2	1	1			
8.9	-	-	-	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X			
10.8	3	4	4	4	3	2	1	2	1	2	3	3	3	5	5	5	5	4	2	3	3	3	4	6	5	4	3	2	2	1	2	3	3	3		
14.9c	2	1	-	-	-	-	-	-	1	2	3	2	3	2	1	3	3	4	4	3	3	2	2	1	1	1	1	1	1	1	1	2	2	2		
21.9	2	2	2	3	2	1	1	1	1	2	3	3	4	3	8	9	8	5	5	3	3	4	3	2	1	1	2	1	1	1	1	2	2	2		
22.7	4	2	2	2	2	2	1	1	1	3	5	4	4	5	6	6	6	5	4	4	3	2	2	1	1	1	1	1	1	1	1	2	2	2		
23.7	3	3	3	3	2	2	1	1	1	2	2	2	2	4	5	6	5	4	4	3	2	2	2	1	1	2	1	1	1	1	2	2	1			
27.8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X			
28.9	2	2	3	2	2	1	1	1	2	2	2	3	4	4	4	4	4	4	5	5	5	3	3	3	3	4	4	3	1	1	1	2	2	2		
30.9	2	2	2	2	2	2	1	1	1	1	4	4	4	4	4	4	4	5	5	5	4	4	3	3	2	2	1	1	1	1	2	2	2			
31.7	3	3	3	3	2	1	1	1	1	2	3	4	6	6	3	4	4	5	5	4	4	4	4	4	3	2	1	1	1	1	1	2	2	2		

a indicates low weight S90 through N90.

c indicates low weight N80 through N90.

d indicates low weight S90 through S70.

Table 67b

Coronal observations at Climax, Colorado (6702A), west limb

Table 6ba

Coronal observations at Sacramento Peak, New Mexico (5303A), east limb

Date GCT	Degrees north of the solar equator															Degrees south of the solar equator																					
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	00	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90
1954																																					
Jan. 1.7	-	-	-	-	-	2	2	2	3	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3	2	2	-	-	3	3	2	-	-	-			
2.7	-	-	-	-	-	-	-	-	-	-	2	2	2	2	2	2	-	2	2	2	2	3	3	2	2	-	2	3	3	2	2	-	-	-			
3.8	-	-	-	-	-	-	-	-	-	-	2	2	2	2	3	3	2	3	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-				
4.7	-	-	-	-	-	-	-	-	-	-	2	2	3	2	3	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5.7	-	-	-	-	-	-	-	-	-	-	2	3	4	4	4	3	4	3	3	2	2	3	3	3	2	2	2	3	-	-	-	-	-	-			
6.7	-	-	-	-	-	-	-	-	-	-	2	2	2	3	4	4	3	2	2	2	3	2	2	2	-	2	3	3	3	4	3	2	2	-	-		
7.9a	-	-	-	-	-	-	-	-	-	-	2	3	4	4	3	2	3	2	3	2	2	3	3	-	-	2	2	3	3	2	2	2	-	-	-		
10.7a	-	-	-	-	-	-	-	-	-	-	2	3	2	3	3	4	3	2	5	4	4	3	3	2	2	2	-	-	-	-	-	-	-	-	-	-	
11.7	-	-	-	-	-	-	-	-	-	-	2	3	5	7	6	5	5	5	4	5	4	3	2	3	-	-	3	3	2	3	2	2	-	-	-		
16.7	-	-	-	-	-	-	-	-	-	-	2	2	3	3	3	2	3	3	4	3	3	3	4	3	2	2	2	-	-	-	-	-	-	-	-		
21.8a	-	-	-	-	-	-	-	-	-	-	2	2	3	3	4	4	3	2	2	2	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-		
22.9	-	-	-	-	-	-	-	-	-	-	2	3	5	5	4	4	3	2	2	2	2	2	3	-	-	-	-	-	-	-	-	-	-	-	-		
23.7	-	-	-	-	-	-	-	-	-	-	2	3	4	5	4	5	5	4	2	2	2	2	2	2	3	2	2	-	2	3	3	3	3	2	-	-	
26.7	-	-	-	-	-	-	-	-	-	-	2	2	3	3	3	2	3	3	2	3	3	3	2	2	2	2	3	4	4	5	5	4	3	3	2	2	-
31.7	-	-	-	-	-	-	-	-	-	-	2	2	3	3	2	2	-	-	3	3	3	2	2	2	2	2	2	2	3	4	3	2	2	2	-	-	

Table 69a

Coronal observations at Sacramento Peak, New Mexico (6374A), east limb

Date GCT	Degrees north of the solar equator															0°	Degrees south of the solar equator																					
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90			
1954																																						
Jan.	1.7	3	2	2	2	2	-	2	2	-	3	3	4	5	4	5	10	8	9	8	7	8	8	7	7	6	5	5	5	3	2	-	-	-	2	-	2	
	2.7	2	2	3	2	3	2	-	2	3	2	3	4	5	3	4	13	11	10	11	11	14	11	8	7	5	6	5	3	3	2	2	-	2	-	2	3	3
	3.8	2	3	3	2	3	3	2	2	2	3	5	5	6	7	12	12	11	10	10	8	13	10	8	5	5	6	7	4	3	2	-	2	2	3	4	2	
	4.7	2	-	3	2	3	2	3	4	2	2	2	3	4	3	7	8	11	10	9	8	9	5	6	5	4	4	2	-	-	2	2	-	3	2	3		
	5.7	4	3	3	3	3	3	2	2	2	2	3	2	3	5	8	11	14	13	13	12	11	10	8	5	6	5	2	-	2	-	2	3	3	3	2		
	6.7	5	4	3	3	3	5	4	3	2	3	3	5	6	7	8	11	11	14	13	14	15	14	13	10	10	9	9	7	3	4	2	2	3	3	4	3	
	7.9a	5	4	3	3	3	4	3	3	2	3	5	5	4	4	6	8	11	12	11	11	12	14	15	11	12	12	8	7	5	5	4	2	2	3	3	2	
	10.7a	3	2	-	2	2	2	2	-	2	2	2	2	2	2	3	3	3	4	5	6	7	8	6	5	4	5	4	2	-	2	2	-	-	-	-		
	11.7	4	3	4	4	5	4	3	2	4	-	2	3	5	4	5	5	6	7	8	11	12	14	15	13	12	8	6	5	2	2	3	2	2	3	3	3	
	16.7	3	5	3	2	2	3	3	3	2	3	3	4	5	8	6	11	12	14	13	13	12	10	8	8	7	5	5	4	3	2	2	3	2	3	2		
	21.8a	-	-	-	2	3	-	-	-	-	-	-	4	3	5	6	7	3	3	2	3	3	2	3	2	2	2	3	-	-	-	-	-	-	-	-		
	22.9	3	5	4	4	4	3	2	2	2	3	3	4	5	7	8	8	11	8	7	6	10	12	13	13	9	7	5	5	4	4	3	3	2	2	3	2	
	23.7	5	4	4	3	3	3	2	2	3	3	2	4	8	16	15	13	12	11	8	9	16	14	14	13	11	8	5	4	4	4	3	3	2	2	3	3	2
	26.7	4	2	3	3	5	3	5	2	-	3	4	8	9	7	3	5	6	8	9	10	10	11	8	5	11	8	4	4	3	-	2	2	-	3	3	4	
	31.7	2	2	2	-	2	2	3	-	-	2	3	3	4	8	11	16	11	10	8	6	5	5	6	8	9	3	3	2	-	-	3	2	2	3	2	2	

Table 70a

Coronal observations at Sacramento Peak, New Mexico (6702A), east limb

Table 68b

Coronal observations at Sacramento Peak, New Mexico (5303A), west limb

Date GCT	Degrees south of the solar equator															Degrees north of the solar equator																						
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	3	2	2	2	2	3	3	2	3	3	3	2	3	3	2	-	-	-
Jan. 1.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	2	2	3	2	3	3	2	3	2	3	3	2	2	-	-	-	-	-
2.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	3	3	3	3	3	2	2	2	3	3	3	2	2	-	-	-	-	-
3.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	3	4	4	4	5	8	4	5	6	5	6	5	3	2	2	-	-	-
4.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	2	2	2	3	2	2	3	2	3	3	2	2	-	-	-	-	-	
5.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	4	4	4	5	8	4	5	6	5	6	5	3	2	2	-	-	-	-
6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	4	5	5	4	4	4	5	5	6	5	4	2	-	-	-	-
7.9a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	2	2	2	3	2	2	3	2	3	3	2	2	-	-	-	-	-	-
10.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	2	3	2	2	3	2	2	-	-	-	-	-	-	-	-	-
11.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	4	3	2	2	2	-	-	3	2	2	2	-	-	-	-	-	-
16.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	3	2	2	2	3	2	3	3	2	-	-	-	-	-	-	
21.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
22.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	3	4	4	5	4	4	4	5	5	4	3	2	-	-	-	-	-
23.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	5	5	6	7	5	6	6	7	7	5	3	-	-	-	-	-	-
26.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	3	2	3	3	4	5	6	5	4	3	2	3	2	-	-	-
31.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	2	3	3	4	3	2	2	2	3	3	4	3	2	-	-	-	-	-

Table 69b

Coronal observations at Sacramento Peak, New Mexico (6374A), west limb

Date GCT	Degrees south of the solar equator															Degrees north of the solar equator																						
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	3	4	3	2	2	3	5	4	2	3	2	-	2	2	2	2	2	3
Jan. 1.7	2	2	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	7	7	6	4	3	5	5	8	3	3	2	-	2	2	-	-	-	-
2.7a	3	3	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	8	7	7	6	4	3	5	5	8	3	3	2	-	2	2	-	-	-	-
3.8	2	-	-	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	5	5	3	4	6	8	11	10	5	3	2	2	2	2	2	3	2	2
4.7	3	-	-	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	7	5	3	4	5	4	5	4	5	4	3	2	2	3	2	2	2	2
5.7	2	3	3	2	3	3	3	4	2	2	3	3	8	8	9	11	10	11	13	11	7	5	5	8	7	5	4	3	2	2	2	2	3	3	2	3		
6.7	3	3	2	2	3	3	3	3	2	3	4	10	11	11	13	12	11	10	8	7	5	6	10	11	8	5	5	4	2	3	3	3	4	3	4	5		
7.9a	2	3	3	2	3	2	2	2	2	3	5	4	5	10	11	9	8	6	5	6	5	6	5	11	8	6	5	4	3	2	3	3	2	3	4	5		
10.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	2	2	3	3	5	4	3	2	2	-	2	2	2	2	3	2	3
11.7	3	2	3	2	3	2	-	-	-	-	-	-	-	-	-	-	-	-	-	4	3	3	4	3	5	8	8	5	3	2	2	3	3	2	3	3	4	5
16.7	2	2	3	3	2	3	-	-	-	-	-	-	-	-	-	-	-	-	-	6	5	6	7	7	6	5	5	4	3	2	3	3	2	-	-	-	-	-
21.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	4	3	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-
22.9	2	3	2	2	3	2	2	2	-	-	-	-	-	-	-	-	-	-	-	7	5	5	3	2	3	2	2	2	2	3	2	-	-	-	-	-	-	-
23.7	3	4	3	3	2	3	3	3	5	5	6	5	10	14	15	15	14	11	10	8	6	4	6	5	4	5	6	4	2	2	2	3	3	4	4	5		
26.7	4	2	3	4	-	-	2	3	3	3	5	8	9	7	6	8	8	9	8	8	5	4	3	4	3	6	5	3	2	-	-	2	3	3	3			
31.7	2	3	2	2	-	-	2	-	-	-	3	4	6	8	7	7	8	10	11	10	10	7	6	8	7	4	3	2	3	2	-	-	3	2	-	-		

Table 70b

Coronal observations at Sacramento Peak, New Mexico (6702A), west limb

Date GCT	Degrees south of the solar equator															Degrees north of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Jan. 1.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
3.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7.9a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10.7a	-	-	-	-</td																																

Table 71
Zurich Provisional Relative Sunspot Numbers
January 1954

Date	R _Z *	Date	R _Z *
1	0	17	0
2	0	18	0
3	0	19	0
4	0	20	0
5	0	21	0
6	0	22	0
7	0	23	0
8	0	24	0
9	0	25	0
10	0	26	0
11	0	27	0
12	0	28	0
13	0	29	0
14	0	30	0
15	0	31	0
16	0	Mean:	0.0

*Dependent on observations at Zürich Observatory and its stations at Locarno and Arosa.

Table 72
American Relative Sunspot Numbers
December 1953

Date	R _{A'}	Date	R _{A'}
1	0	17	0
2	0	18	0
3	0	19	0
4	0	20	0
5	0	21	1
6	4	22	0
7	6	23	0
8	1	24	3
9	0	25	7
10	0	26	3
11	0	27	1
12	0	28	15
13	0	29	9
14	0	30	0
15	0	31	0
16	0	Mean:	1.6

Table 73

Solar Flares, January 1954

No solar flares were reported for the month of January.

Table 74

Indices of Geomagnetic Activity for December 1953

Preliminary values of international character-figures, C;
Geomagnetic planetary three-hour-range indices, K_p;
Magnetically selected quiet and disturbed days

GRAPHS OF IONOSPHERIC DATA




Fig. 1. WASHINGTON, D. C.
38.7°N, 77.1°W JANUARY 1954




Fig. 2. WASHINGTON, D. C. JANUARY 1954




Fig. 3. TROMSO, NORWAY
69.7°N, 19.0°E DECEMBER 1953








Fig. 4. TROMSO, NORWAY DECEMBER 1953






Fig. 9 NARSARSSUAK, GREENLAND
61.2°N 45.4°W DECEMBER 1953



Fig. 10. NARSARSSUAK, GREENLAND

















Fig. 11. OSLO, NORWAY
60.0°N, 11.1°E DECEMBER 1953












Fig. 25. PUERTO RICO, W. I.
18.5°N, 67.2°W DECEMBER 1953




Fig. 26. PUERTO RICO, W. I. DECEMBER 1953




Fig. 27. GUAM I.
13.6°N, 144.9°E DECEMBER 1953




Fig. 28. GUAM I. DECEMBER 1953




Fig. 29. PANAMA CANAL ZONE
9.4°N, 79.9°W DECEMBER 1953




Fig. 30. PANAMA CANAL ZONE DECEMBER 1953




Fig. 31. HUANCAYO, PERU
12.0°S 75.3°W DECEMBER 1953




Fig. 32. HUANCAYO, PERU DECEMBER 1953




Fig 33. POINT BARROW, ALASKA
71.3°N, 156.8°W NOVEMBER 1953




Fig. 34. POINT BARROW, ALASKA




Fig 35. KIRUNA, SWEDEN
67.8°N, 20.5°E NOVEMBER 1953




Fig. 36. KIRUNA, SWEDEN NOVEMBER 1953




Fig. 37. LULEA, SWEDEN
65.6°N, 22.1°E NOVEMBER 1953




Fig. 38. LULEA, SWEDEN NOVEMBER 1953




Fig. 39. FAIRBANKS, ALASKA
64.9°N, 147.8°W NOVEMBER 1953




















Fig. 40. FAIRBANKS, ALASKA NOVEMBER 1953












Fig. 57. RESOLUTE BAY, CANADA
74.7°N, 94.9°W OCTOBER 1953




Fig. 58. RESOLUTE BAY, CANADA




Fig. 59. POINT BARROW, ALASKA
71.3°N, 156.8°W OCTOBER 1953




Fig. 60. POINT BARROW, ALASKA




Fig. 61. BAKER LAKE, CANADA
64.3°N, 96.0°W OCTOBER 1953




Fig. 62. BAKER LAKE, CANADA OCTOBER 1953




Fig. 63. REYKJAVIK, ICELAND
64.1°N, 21.8°W OCTOBER 1953




Fig. 64. REYKJAVIK, ICELAND OCTOBER 1953




Fig. 65. CHURCHILL, CANADA
58.8°N, 94.2°W OCTOBER 1953




Fig. 66. CHURCHILL, CANADA OCTOBER 1953




Fig. 67. PRINCE RUPERT, CANADA
54.3°N, 130.3°W OCTOBER 1953




Fig. 68. PRINCE RUPERT, CANADA OCTOBER 1953




Fig. 69. De BILT, HOLLAND
52.1°N, 5.2°E OCTOBER 1953




Fig. 70. De BILT, HOLLAND OCTOBER 1953




Fig. 71. WINNIPEG, CANADA
49.9°N, 97.4°W OCTOBER 1953




Fig. 72. WINNIPEG, CANADA OCTOBER 1953





Fig. 73. ST. JOHN'S, NEWFOUNDLAND
47.6°N, 52.7°W OCTOBER 1953



OCTOBER 1953
Fig. 74. ST. JOHN'S, NEWFOUNDLAND





Fig. 75. OTTAWA, CANADA
45.4°N, 75.9°W OCTOBER 1953



OCTOBER 1953
Fig. 76. OTTAWA, CANADA




Fig. 77. WAKKANAI, JAPAN
45.4°N, 141.7°E OCTOBER 1953




Fig. 78. WAKKANAI, JAPAN OCTOBER 1953




Fig. 79. AKITA, JAPAN
39.7°N, 140.1°E OCTOBER 1953




Fig. 80. AKITA, JAPAN OCTOBER 1953




Fig. 81. TOKYO, JAPAN
35.7°N 139.5°E OCTOBER 1953




Fig. 82. TOKYO, JAPAN OCTOBER 1953




Fig. 83. YAMAGAWA, JAPAN
31.2°N, 130.6°E OCTOBER 1953




Fig. 84. YAMAGAWA, JAPAN OCTOBER 1953




Fig. 85. JOHANNESBURG, UNION OF S. AFRICA
26.2°S, 28.1°E OCTOBER 1953




Fig. 86. JOHANNESBURG, UNION OF S. AFRICA




Fig. 87. CAPETOWN, UNION OF S. AFRICA
34.2°S, 18.3°E OCTOBER 1953




Fig. 88. CAPETOWN, UNION OF S. AFRICA




Fig. 89. POINT BARROW, ALASKA
71°30'N, 156°8'W SEPTEMBER 1953




Fig. 90. POINT BARROW, ALASKA SEPTEMBER 1953




Fig. 91. REYKJAVIK, ICELAND
64°10'N, 21°8'W SEPTEMBER 1953




Fig. 92. REYKJAVIK, ICELAND SEPTEMBER 1953




Fig. 93. POINT BARROW, ALASKA
71.3°N, 156.8°W AUGUST 1953




Fig. 94. POINT BARROW, ALASKA AUGUST 1953




Fig. 95. REYKJAVIK, ICELAND
64.1°N, 21.8°W AUGUST 1953





Fig. 96. REYKJAVIK, ICELAND AUGUST 1953

Index of Tables and Graphs of Ionospheric Datain CRPL-F114

	<u>Table page</u>	<u>Figure page</u>
Akita, Japan		
October 1953	18	63
Anchorage, Alaska		
December 1953	12	45
Baker Lake, Canada		
October 1953	17	59
Buenos Aires, Argentina		
November 1953	16	57
Capetown, Union of S. Africa		
October 1953	19	65
Churchill, Canada		
October 1953	17	60
De Bilt, Holland		
November 1953	15	54
October 1953	17	61
Deception I.		
November 1953	16	57
Fairbanks, Alaska		
December 1953	12	45
November 1953	15	53
Formosa, China		
November 1953	16	56
Graz, Austria		
December 1953	13	47
Guam I.		
December 1953	14	50
Huancayo, Peru		
December 1953	14	51
Johannesburg, Union of S. Africa		
October 1953	19	65
Kiruna, Sweden		
November 1953	14	52
Leopoldville, Belgian Congo		
November 1953	16	56
Lindau/Harz, Germany		
November 1953	15	55
Lulea, Sweden		
November 1953	15	53
Maui, Hawaii		
December 1953	13	49
Narsarsuaq, Greenland		
December 1953	12	46
Okinawa I.		
December 1953	13	49

Index (CRPL-F114, concluded)

	<u>Table page</u>	<u>Figure page</u>
Oslo, Norway		
December 1953	12	46
Ottawa, Canada		
October 1953	18	62
Panama Canal Zone		
December 1953	14	51
Point Barrow, Alaska		
November 1953	14	52
October 1953	16	58
September 1953	19	66
August 1953	19	67
Prince Rupert, Canada		
October 1953	17	60
Puerto Rico, W. I.		
December 1953	14	50
Resolute Bay, Canada		
October 1953	16	58
Reykjavik, Iceland		
November 1953	15	54
October 1953	17	59
September 1953	19	66
August 1953	19	67
St. John's, Newfoundland		
October 1953	18	62
San Francisco, California		
December 1953	13	48
Schwarzenburg, Switzerland		
November 1953	15	55
Tokyo, Japan		
October 1953	18	64
Tromso, Norway		
December 1953	12	44
Upsala, Sweden		
December 1953	13	47
Wakkanai, Japan		
October 1953	18	63
Washington, D. C.		
January 1954	12	44
White Sands, New Mexico		
December 1953	13	48
Winnipeg, Canada		
October 1953	17	61
Yamagawa, Japan		
October 1953	18	64



CRPL and IRPL Reports

[A list of CRPL Section Reports is available from the Central Radio Propagation Laboratory upon request]

Daily:

Radio disturbance forecasts, every half hour from broadcast station WWV of the National Bureau of Standards. Telephoned and telegraphed reports of ionospheric, solar, geomagnetic, and radio propagation data.

Semimonthly:

- CRPL—J. North Atlantic Radio Propagation Forecast (of days most likely to be disturbed during following month).
CRPL—Jp. North Pacific Radio Propagation Forecast (of days most likely to be disturbed during following month).

Semimonthly:

- CRPL—Ja. Semimonthly Frequency Revision Factors For CRPL Basic Radio Propagation Prediction Reports.

Monthly:

- CRPL—D. Basic Radio Propagation Predictions—Three months in advance. (Dept. of the Army, TB 11-499-, monthly supplements to TM 11-499; Dept. of the Navy, DNC 13 () series; Dept. of the Air Force, TO 16-1B-2 series.)
CRPL—F. Ionospheric Data.
*IRPL—A. Recommended Frequency Bands for Ships and Aircraft in the Atlantic and Pacific.
*IRPL—H. Frequency Guide for Operating Personnel.

Circulars of the National Bureau of Standards:

NBS Circular 462. Ionospheric Radio Propagation.

NBS Circular 465. Instructions for the Use of Basic Radio Propagation Predictions.

Reports issued in past:

- IRPL—C61. Report of the International Radio Propagation Conference, 17 April to 5 May 1944.
IRPL—G1 through G12. Correlation of D. F. Errors With Ionospheric Conditions.
(G1, G3, available. Others out of print; see second footnote.)
IRPL—R. Nonscheduled reports:
R4. Methods Used by IRPL for the Prediction of Ionosphere Characteristics and Maximum Usable Frequencies.
R5. Criteria for Ionospheric Storminess.
**R6. Experimental Studies of Ionospheric Propagation as Applied to the Loran System.
R7. Second Report on Experimental Studies of Ionospheric Propagation as Applied to the Loran System.
R9. An Automatic Instantaneous Indicator of Skip Distance and MUF.
R10. A Proposal for the Use of Rockets for the Study of the Ionosphere.
**R11. A Nomographic Method for both Prediction and Observation Correlation of Ionosphere Characteristics.
**R12. Short Time Variations in Ionosphere Characteristics.
R14. A Graphical Method for Calculating Ground Reflection Coefficients.
**R15. Predicted Limits for F2-Layer Radio Transmission Throughout the Solar Cycle.
**R17. Japanese Ionospheric Data—1943.
R18. Comparison of Geomagnetic Records and North Atlantic Radio Propagation Quality Figures—October 1943 Through May 1945.
**R21. Notes on the Preparation of Skip-Distance and MUF Charts for Use by Direction-Finder Stations.
(For distances out to 4000 km.)
**R23. Solar-Cycle Data for Correlation with Radio Propagation Phenomena.
**R24. Relations Between Band Width, Pulse Shape and Usefulness of Pulses in the Loran System.
**R25. The Prediction of Solar Activity as a Basis for the Prediction of Radio Propagation Phenomena.
**R26. The Ionosphere as a Measure of Solar Activity.
R27. Relationships Between Radio Propagation Disturbance and Central Meridian Passage of Sunspots Grouped by Distance From Center of Disc.
**R30. Disturbance Rating in Values of IRPL Quality-Figure Scale from A. T. & T. Co. Transmission Disturbance Reports to Replace T. D. Figures as Reported.
**R31. North Atlantic Radio Propagation Disturbances, October 1943 Through October 1945.
**R33. Ionospheric Data on File at IRPL.
**R34. The Interpretation of Recorded Values of fEs.
**R35. Comparison of Percentage of Total Time of Second-Multiple Es Reflections and That of fEs in Excess of 3 Mc.
IRPL—T. Reports on tropospheric propagation:
T1. Radar operation and weather. (Superseded by JANP 101.)
T2. Radar coverage and weather. (Superseded by JANP 102.)
CRPL—T3. Tropospheric Propagation and Radio-Meteorology. (Reissue of Columbia Wave Propagation Group WPG—5.)

*Items bearing this symbol are distributed only by U. S. Navy. They are issued under one cover as the DNC 14 () Series.
**Out of print; information concerning cost of photostat or microfilm copies is available from CRPL upon request.

