

Library, S. W. 360.

CRPL-F 117

4-1954

FOR OFFICIAL USE

Reference book not to be
taken from the Library.

IONOSPHERIC DATA

ISSUED
MAY 1954

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
CENTRAL RADIO PROPAGATION LABORATORY
BOULDER, COLORADO

IONOSPHERIC DATA

CONTENTS

	<u>Page</u>
Symbols, Terminology, Conventions	2
World-Wide Sources of Ionospheric Data	5
Hourly Ionospheric Data at Washington, D. C..	7, 12, 20, 45
Ionospheric Storminess at Washington, D. C. .	7, 32
Radio Propagation Quality Figures	8, 33
Observations of the Solar Corona	9, 36
Relative Sunspot Numbers	10, 40
Observations of Solar Flares	10, 42
Indices of Geomagnetic Activity	11, 43
Sudden Ionosphere Disturbances	11, 44
Tables of Ionospheric Data	12
Graphs of Ionospheric Data	45
Index of Tables and Graphs of Ionospheric Data in CRPL-F117	69

SYMBOLS, TERMINOLOGY, CONVENTIONS

Beginning with data reported for January 1952, the symbols, terminology, and conventions for the determination of median values used in this report (CRPL-F series) conform as far as practicable to those adopted at the Sixth Meeting of the International Radio Consultative Committee (C.C.I.R.) in Geneva, 1951. Excerpts concerning symbols and terminology from Document No. 626-E of this Meeting are given on pages 2-7 of the report CRPL-F89, "Ionospheric Data," issued January 1952. Reprints of these pages are available upon request.

Beginning with data for January 1945, median values are published wherever possible. Where averages are reported, they are, at any hour, the average for all the days during the month for which numerical data exist.

The following conventions are used in determining the medians for hours when no measured values are given because of equipment limitations and ionospheric irregularities. Symbols used are those given in Document No. 626-E referred to above.

a. For all ionospheric characteristics:

Values missing because of A, C, F, L, M, N, Q, S, or T are omitted from the median count.

b. For critical frequencies and virtual heights:

Values of f_{oF2} (and f_{oE} near sunrise and sunset) missing because of E are counted as equal to or less than the lower limit of the recorder. Values of $h'F2$ (and $h'E$ near sunrise and sunset) missing for this reason are counted usually as equal to or greater than the median. Other characteristics missing because of E are omitted from the median count.

Values missing because of D are counted as equal to or greater than the upper limit of the recorder.

Values missing because of G are counted:

1. For f_{oF2} , as equal to or less than f_{oFl} .
2. For $h'F2$, as equal to or greater than the median.

The symbol W is included in the median count only when it replaces a height characteristic. This practice represents a change from that listed in issues previous to CRPL-F78.

Values missing for any other reason are omitted from the median count.

c. For MUF factor (M-factors):

Values missing because of G or W are counted as equal to or less than the median.

Values missing for any other reason are omitted from the median count.

d. For sporadic E (Es):

Values of fEs missing because of E or G (and B when applied to the daytime E region only) are counted as equal to or less than the median foE, or equal to or less than the lower frequency limit of the recorder.

Values of fEs missing for any other reason, and values of h'Es missing for any reason at all are omitted from the median count.

Beginning with data for November 1945, doubtful monthly median values for ionospheric observations at Washington, D. C., are indicated by parentheses, in accordance with the practice already in use for doubtful hourly values. The following are the conventions used to determine whether or not a median value is doubtful:

1. If only four values or less are available, the data are considered insufficient and no median value is computed.

2. For the F2 layer, if only five to nine values are available, the median is considered doubtful. The E and F1 layers are so regular in their characteristics that, as long as there are at least five values, the median is not considered doubtful.

3. For all layers, if more than half of the values used to compute the median are doubtful (either doubtful or interpolated), the median is considered doubtful.

The same conventions are used by the CRPL in computing the medians from tabulations of daily and hourly data for stations other than Washington, beginning with the tables in IRPL-F18.

The tables and graphs of ionospheric data are correct for the values reported to the CRPL, but, because of variations in practice in the interpretation of records and scaling and manner of reporting of values, may at times give an erroneous conception of typical ionospheric characteristics at the station. Some of the errors are due to:

- a. Differences in scaling records when spread echoes are present.
- b. Omission of values when f_{oF2} is less than or equal to f_{oFl} , leading to erroneously high values of monthly averages or median values.
- c. Omission of values when critical frequencies are less than the lower frequency limit of the recorder, also leading to erroneously high values of monthly average or median values.

These effects were discussed on pages 6 and 7 of the previous F-series report IRPL-F5.

Ordinarily, a blank space in the fEs column of a table is the result of the fact that a majority of the readings for the month are below the lower limit of the recorder or less than the corresponding values of f_{oE} . Blank spaces at the beginning and end of columns of $h^{\circ}F1$, f_{oFl} , $h^{\circ}E$, and f_{oE} are usually the result of diurnal variation in these characteristics. Complete absence of medians of $h^{\circ}F1$ and f_{oFl} is usually the result of seasonal effects.

The dashed-line prediction curves of the graphs of ionospheric data are obtained from the predicted zero-muf contour charts of the CRPL-D series publications. The following points are worthy of note:

- a. Predictions for individual stations used to construct the charts may be more accurate than the values read from the charts since some smoothing of the contours is necessary to allow for the longitude effect within a zone. Thus, inasmuch as the predicted contours are for the center of each zone, part of the discrepancy between the predicted and observed values as given in the F series may be caused by the fact that the station is not centrally located within the zone.
- b. The final presentation of the predictions is dependent upon the latest available ionospheric and radio propagation data, as well as upon predicted sunspot number.

- c. There is no indication on the graphs of the relative reliability of the data; it is necessary to consult the tables for such information.

The following predicted smoothed 12-month running-average Zürich sunspot numbers were used in constructing the contour charts:

Month	Predicted Sunspot Number									
	1954	1953	1952	1951	1950	1949	1948	1947	1946	1945
December	15	33	53	86	108	114	126	85	38	
November	16	38	52	87	112	115	124	83	36	
October	17	43	52	90	114	116	119	81	23	
September	18	46	54	91	115	117	121	79	22	
August	18	49	57	96	111	123	122	77	20	
July	20	51	60	101	108	125	116	73		
June	21	52	63	103	108	129	112	67		
May	22	52	68	102	108	130	109	67		
April	10	24	52	74	101	109	133	107	62	
March	11	27	52	78	103	111	133	105	51	
February	12	29	51	82	103	113	133	90	46	
January	14	30	53	85	105	112	130	88	42	

WORLD - WIDE SOURCES OF IONOSPHERIC DATA

The ionospheric data given here in tables 1 to 48 and figures 1 to 96 were assembled by the Central Radio Propagation Laboratory for analysis and correlation, incidental to CRPL prediction of radio propagation conditions. The data are median values unless otherwise indicated. The following are the sources of the data in this issue:

Commonwealth of Australia, Ionospheric Prediction Service of the Commonwealth Observatory:

Brisbane, Australia
 Canberra, Australia
 Hobart, Tasmania
 Townsville, Australia

British Department of Scientific and Industrial Research, Radio Research Board:

Falkland Is.
Inverness, Scotland
Khartoum, Sudan (University College of Khartoum)
Singapore, British Malaya
Slough, England

Defence Research Board, Canada:

Churchill, Canada

National Laboratory of Radio-Electricity (French Ionospheric Bureau):

Casablanca, Morocco
Poitiers, France

Institute for Ionospheric Research, Lindau Über Northeim, Hannover, Germany:

Lindau/Harz, Germany

Christchurch Geophysical Observatory, New Zealand Department of Scientific and Industrial Research:

Christchurch, New Zealand
Rarotonga, Cook Is.

National Bureau of Standards (Central Radio Propagation Laboratory):

Washington, D. C.

HOURLY IONOSPHERIC DATA AT WASHINGTON, D. C.

The data given in tables 49 through 60 follow the scaling practices given in the report IRPL-C61, "Report of International Radio Propagation Conference," pages 36 to 39, and the median values are determined by the conventions given above under "Symbols, Terminology, Conventions." Beginning with September 1949, the data are taken at Ft. Belvoir, Virginia.

IONOSPHERIC STORMINESS AT WASHINGTON, D.C.

Table 61 presents ionosphere character figures for Washington, D. C., during April 1954, as determined by the criteria given in the report IRPL-R5, "Criteria for Ionospheric Storminess," together with Cheltenham, Maryland, geomagnetic K-figures, which are usually covariant with them.

RADIO PROPAGATION QUALITY FIGURES

Tables 63a and 63b give for March 1954 the radio propagation quality figures for the North Atlantic area, the relevant CRPL advance and short-term forecasts, a summary geomagnetic activity index and sundry comparisons, specifically as follows:

- (a) radio propagation quality figures, Q_a , separately for each 6-hour interval of the Greenwich day, viz., 00-06, 06-12, 12-18, 18-24 hours UT (Universal Time or GCT).
- (b) whole-day radio quality indices (beginning October 1952). Each index is a weighted average of the four quarter-day Q_a -figures, before rounding off, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which designate whenever possible the days when significant disturbance or unusually quiet conditions will occur.
- (c) short-term forecasts, issued by CRPL every six hours (nominally one hour before 00^h, 06^h, 12^h, 18^h UT) and applicable to the period 1 to 13 (especially 1 to 7) hours ahead. Note that new scoring rules have been adopted beginning with October 1952 data.
- (d) advance forecasts, issued semiweekly (CRPL-J reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole-day quality indices.
- (e) half-day averages of the geomagnetic K indices measured by the Cheltenham Magnetic Observatory of the U. S. Coast and Geodetic Survey.
- (f) illustration of the comparison of short-term forecasts with Q_a -figures and also with estimates of radio quality based on CRPL observations only.
- (g) illustration of the outcome of advance forecasts (1 to 3 or 4 days ahead) and, for comparison, the outcome of a type of "blind" forecast. For the latter the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

These radio propagation quality figures, Q_a , are prepared from radio traffic data reported to CRPL by American Telephone and Telegraph Company, Mackay Radio and Telegraph Company, RCA Communications, Inc., Marconi Company, British Admiralty Signal and Radar Establishment, and the following agencies of the U. S. Government:--Coast Guard, Navy, Army Signal Corps, and U. S. Information Agency. The method of calculation, summarized below, is similar to that described in a 1946 report, IRPL-R31, now out of print. Only reports of radio transmission on North Atlantic paths closely approximating New York-London are included in the estimation of quality.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the quality scale of the original reports. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year, with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. The 6-hourly quality figures are (subjectively) weighted means of the reports received for that period. These 6-hourly quality figures replace, beginning January 1953, the half-daily quality figures which formerly appeared in this table. (These forecasts and quality indices are prepared by the North Atlantic Radio Warning Service, the CRPL forecasting center at Ft. Belvoir, Virginia.)

Table 62 gives for March 1954, the radio propagation quality figures for the North Pacific area, the relevant CRPL advance and short-term forecasts, and sundry comparisons, specifically as follows:

- (a) radio propagation quality figures, Q_p , separately for each of three 9-hour intervals of the Greenwich day, viz., 03-12, 09-18 and 18-03 UT (Universal Time or GCT).
- (b) whole-day radio quality indices for each Greenwich day. These are derived from the same basic data as the 9-hour indices, separately reduced.
- (c) short-term forecasts, issued daily at 02, 09 and 18 hours UT.
- (d) advance forecasts, issued semiweekly (CRPL-Jp reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole day quality indices.

These radio quality indices, Q_p , refer to radio propagation on optimum frequencies over moderately long transmission paths in the North Pacific area. Typical paths are Anchorage (Alaska) to Seattle, or Anchorage to Tokyo. The indices are derived from reports submitted regularly by communications agencies of the U. S. Army and Air Force, and by Aeronautical Radio, Inc. The method of derivation of Q_p differs from that of Q_a . For Q_p , each reported index is converted into a deviation (usually) from the 3-monthly mean for that index, in units of the standard deviation. These deviations are averaged for all reports for a given 9-hour period. The average is then put on the 1 to 9 Q-scale with an assumed standard deviation of 1.25 and assumed means of 5.33, 5.33, and 6.00, respectively, for the 03-12, 09-18 and 18-03 periods, and 5.67 for the whole day period. (These forecasts and quality indices are prepared by the North Pacific Radio Warning Service, the CRPL forecasting center at Anchorage, Alaska.)

These quality figures are, in effect, a consensus of reported radio propagation conditions. The reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality because of multipath, interference, etc. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

OBSERVATIONS OF THE SOLAR CORONA

Tables 64 through 66 give the observations of the solar corona during April 1954, obtained at Climax, Colorado, by the High Altitude Observatory of Harvard University and the University of Colorado. Tables 67 through 69 list the coronal observations obtained at Sacramento Peak, New Mexico, during April 1954, derived by Harvard College Observatory as a part of its performance of a research contract with the Upper Air Research Observatory, Geophysical Research Directorate, Air Force Cambridge Research Center. The data are listed separately for east and west limbs at 5-degree intervals of position angle north and south of the Solar Equator at the limb. The time of observation is given to the nearest tenth of a day, GCT.

Table 64 gives the intensities of the green (5303A) line of the emission spectrum of the solar corona; table 65 gives similarly the intensities of the first red (6374A) coronal line; and table 66, the intensities of the second red (6702A) coronal line; all observed at Climax in April 1954.

Table 67 gives the intensities of the green (5303A) coronal line; table 68, the intensities of the first red (6374A) coronal line; and table 69, the intensities of the second red (6702A) coronal line; all observed at Sacramento Peak in April 1954.

The following symbols are used in tables 64 through 69: a, observation of low weight; -, corona not visible; and X, position angle not included in plate estimates.

RELATIVE SUNSPOT NUMBERS

Table 70 lists the daily provisional Zürich relative sunspot number, R_Z , for April 1954, as communicated by the Swiss Federal Observatory. Table 71 contains the daily American relative sunspot number, R_A' , for March 1954, as compiled by the Solar Division, American Association of Variable Star Observers.

OBSERVATIONS OF SOLAR FLARES

Table 72 gives the preliminary record of solar flares reported to the CRPL. These reports are communicated on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete records are published later in the Quarterly Bulletin of Solar Activity, I.A.U., in various observatory publications, and elsewhere. The present listing serves to identify and roughly describe the phenomena observed. Details should be sought from the reporting observatory.

Reporting directly to the CRPL are the following observatories: Mt. Wilson, McMath-Hulbert, U. S. Naval, Wendelstein, Kanzel and High Altitude at Sacramento Peak, New Mexico. The remainder report to Meudon (Paris) and the data are taken from the Paris-URSIgram broadcast, monitored fairly regularly by the CRPL. The data on solar flares reported from Sacramento Peak, New Mexico, communicated by the High Altitude Observatory at Boulder, Colorado, are provided by Harvard University as the result of work undertaken on an Air Materiel Command Research and Development Contract administered by the Air Force Cambridge Research Laboratories.

The table lists for each flare the reporting observatory, date, times of beginning and ending of observation, duration (when known), total area (corrected for foreshortening), and heliographic coordinates. For the maximum phase of the flare is given the time, intensity, area relative to the total area, and the importance. The column "SID observed" is to indicate when a sudden ionosphere disturbance, noted elsewhere in these reports, occurred at the time of a flare. Times are in Universal Time (GCT).

INDICES OF GEOMAGNETIC ACTIVITY

Table 73 lists various indices of geomagnetic activity based on data from magnetic observatories widely distributed throughout the world. The indices are: (1) preliminary international character-figures, C; (2) geomagnetic planetary three-hour-range indices, K_p; (3) magnetically selected quiet and disturbed days.

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm). The magnetically quiet and disturbed days are selected by the international scheme outlined on pages 219-227 in the December 1943 issue of Terrestrial Magnetism and Atmospheric Electricity. The details of the currently used method follow. For each day of a month, its geomagnetic activity is assigned by weighting equally the following three criteria: (1) the sum of the eight K_p's; (2) the greatest K_p; and (3) the sum of the squares of the eight K_p's.

K_p is the mean standardized K-index from 11 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g., 5 is $4 \frac{2}{3}$, 5_o is $5 \frac{0}{3}$, and 5+ is $5 \frac{1}{3}$. This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of K_p has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948," published in Washington, D. C., 1949, by the Association of Terrestrial Magnetism and Electricity, International Union of Geodesy and Geophysics. K_p is available from 1937 to date as noted in F108.

The Committee on Characterization of Magnetic Disturbance, ATME, IUGG, has kindly supplied this table. The Meteorological Office, De Bilt, Holland, collects the data and compiles C and selected days. The Chairman of the Committee computes the planetary index. Current tables are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

SUDDEN IONOSPHERE DISTURBANCES

It is hoped that the information scheduled for table 74 will be published in a future issue of the F series.

TABLES OF IONOSPHERIC DATA

Table 1

Washington, D. C., U.S.A. (57.1°W)							April 1954										
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	2.4						3.0	00	290	2.7						2.9
01	290	2.2						3.0	01	290	2.5						2.9
02	290	2.2						3.1	02	295	2.3						2.8
03	290	2.1						3.1	03	320	2.0						2.8
04	290	(1.9)						(3.1)	04	295	(2.2)						2.8
05	280	2.2						3.2	05	290	3.0	275	2.4	135	1.5	2.5	3.1
06	260	3.2	250	—	130	1.7		3.3	06	(300)	2.6	240	3.0	130	1.8	2.8	2.1
07	300	3.8	220	3.4	110	2.1	2.2	3.2	07	380	3.8	225	2.4	115	2.1	2.8	3.1
08	340	4.4	210	3.7	110	2.4	2.9	3.1	08	290	4.0	215	3.6	110	2.4	2.8	3.0
09	340	4.6	210	3.9	110	2.7	2.8	3.2	09	290	4.2	210	3.8	110	2.6	2.9	3.1
10	390	4.8	200	4.0	110	2.9	3.2	3.0	10	425	4.4	210	3.9	105	2.8	2.0	3.1
11	400	4.9	200	4.1	100	3.0	3.2	3.0	11	420	4.4	200	4.0	105	2.8	2.1	2.9
12	370	5.0	200	4.2	100	3.1	3.2	3.0	12	425	4.4	210	4.1	100	2.9	3.2	2.0
13	360	5.2	210	4.1	100	3.1		3.0	13	415	4.5	205	4.1	105	2.9	2.0	2.9
14	360	5.3	210	4.1	110	3.0	2.0	3.0	14	420	4.4	210	4.0	105	2.9	2.9	2.9
15	330	5.2	220	3.9	110	2.9		3.0	15	425	4.4	220	4.0	105	2.8	2.8	2.9
16	310	5.1	220	3.8	110	2.6		3.2	16	390	4.4	225	3.8	110	2.6	2.0	2.9
17	300	4.8	230	3.4	110	2.2		3.2	17	365	4.5	230	2.7	110	2.4	2.0	2.9
18	270	4.9	240	—	(120)	1.7		3.2	18	315	4.6	240	2.4	125	2.1	3.0	3.0
19	240	4.8							19	280	4.6	246	2.1	135	1.7	3.0	3.1
20	240	4.8							20	255	4.8					1.9	3.1
21	250	3.6							21	260	4.8						3.0
22	270	2.9							22	275	4.2						3.0
23	290	2.5							23	295	3.2						3.0

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3*

Slough, England (51.5°E, 0.5°W)							August 1953										
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	3.2						2.6	00	360	(2.7)					3.1	
01	280	2.9						3.1	01	380	(2.8)					3.1	
02	280	2.5						3.8	02	340	(2.8)					3.1	
03	290	2.5						4.1	03	300	(2.7)					3.1	
04	290	2.4						4.2	04	280	(2.5)					3.1	
05	300	2.2	(250)	(2.8)	(130)	(1.5)		3.9	05	260	(2.0)					4.0	
06	310	3.9	235	3.3	125	2.0	3.2	3.1	06	235	4.9					(2.3)	
07	355	4.2	235	3.6	120	2.2	4.2	3.2	07	345	6.0	215	2.8	160	1.9	4.0	
08	265	4.5	230	3.8	115	2.6	4.7	2.2	08	335	5.4	220	4.1			2.3	
09	410	4.7	220	4.0	115	2.8	4.7	3.2	09	370	5.9	220	4.4	125	2.4	5.9	
10	380	4.7	215	4.1	115	2.9	4.7	2.2	10	410	7.5	210	4.5	(120)	3.2	2.9	
11	365	4.8	215	4.2	115	2.0	4.5	3.2	11	400	7.7	210	4.4	(130)	3.6	2.5	
12	400	4.9	220	4.2	115	3.1	4.6	3.3	12	395	8.3	220	4.4	125	3.8	4.4	
13	420	4.7	220	4.2	115	3.1	4.7	3.1	13	380	8.8	220	4.3	125	3.4	2.5	
14	415	4.7	220	4.2	115	3.1	4.4	3.2	14	355	9.0	225	4.1	125	3.4	2.5	
15	380	4.7	220	4.1	115	2.9	4.4	3.0	15	340	9.7	225	4.0	2.8	6.7	2.8	
16	385	4.5	235	4.0	115	2.8	4.4	3.0	16	310	10.0					6.0	
17	340	4.8	240	3.7	115	2.4	4.4	3.0	17	225	9.0					4.0	
18	315	5.0	250	3.4	120	2.0	4.5	3.0	18	245	6.9					4.3	
19	280	5.5							19	225	9.0					3.0	
20	255	5.6							20	240	5.7					3.4	
21	255	5.1							21	240	5.7					2.9	
22	255	4.4							22	305	4.5					3.1	
23	270	3.7							23	345	3.8					2.7	

Time: 0.0°.

Sweep: 0.55 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 5*

Singapore, British Malaya (1.3°S, 103.6°E)							August 1953										
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	250	3.7						4.0	00	300	3.1					2.5	
01	250	3.7						4.2	01	290	3.2					2.9	
02	245	3.1						3.2	02	290	3.3					3.0	
03	255	2.6						3.7	03	250	3.3					3.2	
04	245	2.4						3.6	04	260	2.8					3.1	
05	(250)	2.0						3.8	05	270	2.4					2.2	
06	270	2.9						3.4	06	250	4.5					2.4	
07	250	5.0	235		125	2.2	3.8	3.1	07	260	5.5	210	2.9	120	2.3	3.4	
08	300	7.5	220	4.1	120	2.8	5.7	2.9	08	300	5.1	210	4.0	110	2.7	3.5	
09	330	8.5	215	4.4	110	3.1	5.7	2.7	09	270	6.8	210	4.2	110	3.0	3.3	
10	335	9.1	205	4.4	110	3.3	6.1	2.6	10	270	6.5	210	4.3	110	3.1	3.5	
11	345	9.3	200	4.5	110	3.4	6.0	2.5	11	270	6.3	200	4.3	110	3.2	3.4	
12	345	9.2	200	4.5	110	3.5	5.1	2.5	12	270	6.0	200	4.4	105	3.2	4.5	
13	340	9.2	200	4.5	110	3.4	5.7	2.8	13	270	5.8	200	4.3	105	3.1	4.5	
14	350	9.2	200	4.4	110	3.3	5.7	2.5	14	300	5.8	200	4.4	105	3.1	3.2	
15	325	9.2	205	4.3	110	3.2	5.5	2.8	15	280	5.9	200	4.1	110	3.0	4.3	
16	315	9.1	215	4.2	115	2.8	5.2	2.6	16	270	5.9	200	3.7	115	2.7	3.2	
17	280	9.0	230		125	2.3	4.4	2.6	17	250	5.4	240	2.6			4.0	
18	250	9.0							18	250	5.5					3.3	
19	245	8.7							19	250	5.1					3.4	
20	235	8.4							20	240	4.8					3.1	
21	225	7.0							21	280	3.8					2.9	
22	225	5.4							22	270	3.6					3.0	
23	230	4.8							23	270	3.2					2.5	

Time: 105.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 6

Rarotonga I. (21.3°S, 159.8°W)							August 1953										
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	3.1							00	300	3.1					2.5	
01	290	3.2							01	290	3.3					2.9	

Table 7

Time	Christchurch, New Zealand (43.6°S , 172.7°E)						August 1953	
	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	2.1				3.4	3.0	
01	280	2.3				3.0	3.0	
02	270	2.4				3.8	3.1	
03	270	2.1				3.2	3.0	
04	260	1.8				4.0	3.1	
05	250	1.7				3.4	3.2	
06	260	1.7				4.1	3.1	
07	250	3.0			1.6	3.7	3.4	
08	240	3.8	240	2.9		1.9	3.5	
09	260	4.2	230	3.5		2.3	4.2	3.4
10	280	4.5	220	3.7		2.6	4.3	3.3
11	320	4.7	220	3.9		2.7	4.3	3.3
12	310	5.0	220	3.9		2.8	4.3	3.3
13	300	5.0	230	3.9		2.7	4.3	3.3
14	300	5.0	230	3.8		2.6	4.3	3.3
15	280	4.9	220	3.6		2.4	4.2	3.4
16	260	4.7	240	3.2		2.0	3.8	3.4
17	240	4.3	230	2.2		1.6	2.0	3.4
18	240	3.4					3.1	
19	260	3.2					3.1	
20	270	2.7					3.1	
21	280	2.4					1.7	3.0
22	280	2.3					3.0	
23	280	2.2					2.2	3.0

Time: 172.5°E .

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 55 seconds.

Table 9

Time	Churchill, Canada (68.6°N , 94.2°W)						July 1953	
	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	270	3.5			—	—	8.0	
01	280	3.5			100	2.3	6.2	(3.0)
02	300	3.4			110	2.2	7.0	(3.0)
03	300	3.2			100	2.0	6.0	3.0
04	280	3.4	—	—	100	2.0	4.2	3.0
05	300	3.6	220	3.0	100	2.4	5.0	2.8
06	490	(3.6)	260	< 3.6	100	3.2	3.0	6
07	G	< 3.9	210	< 3.8	100	3.4	3.6	6
08	540	4.0	210	3.9	100	3.3	5.6	2.2
09	0	< 4.0	210	< 4.0	100	3.3	7.0	6
10	630	< 4.0	210	4.0	100	3.1	6.0	2.4
11	420	< 4.1	200	4.0	100	3.1	4.5	2.4
12	520	4.0	210	4.0	100	3.2	7.5	2.3
13	500	4.2	200	4.0	100	3.2	7.5	2.5
14	440	4.4	210	4.0	100	3.1	2.6	
15	400	4.6	220	4.0	100	3.0	2.8	
16	380	4.6	220	3.9	100	3.0	2.8	
17	390	4.5	220	3.8	100	2.8	2.8	
18	370	4.3	230	3.7	110	2.9	2.8	
19	340	4.2	240	3.5	110	2.8	2.9	
20	200	3.9	—	—	110	2.7	5.2	3.0
21	300	3.8	—	—	110	2.5	7.0	3.0
22	300	3.6	—	—	120	2.2	9.0	3.0
23	300	3.2	—	—	> 10.0		(2.9)	

Time: 90.0°W .

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 11

Time	Lindau/Hars, Germany (51.6°N , 10.1°E)						July 1953	
	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	260	3.8				2.3	3.1	
01	260	3.6				2.3	2.1	
02	250	3.2				2.8	3.1	
03	260	2.8				2.2	3.1	
04	270	2.8	250	—	E	2.6	3.2	
05	300	3.5	230	2.8	—	3.0	3.2	
06	320	3.8	225	3.3	115	2.0	3.5	
07	425	4.2	220	3.6	105	2.4	3.8	3.0
08	415	4.3	220	3.8	100	2.6	4.3	3.1
09	410	4.5	200	3.9	100	2.8	5.0	2.9
10	360	4.8	210	4.0	100	2.9	5.0	3.1
11	360	4.9	200	4.1	100	3.0	4.9	3.1
12	390	4.8	210	4.2	100	3.0	3.8	3.0
13	365	4.8	210	4.2	100	3.0	4.4	3.0
14	405	4.6	215	4.2	100	3.0	4.5	2.9
15	365	4.6	210	4.1	100	3.0	3.6	3.1
16	390	4.4	210	4.0	100	2.8	4.3	2.9
17	355	4.6	215	3.8	100	2.6	3.6	3.1
18	325	4.6	220	3.6	105	2.3	4.6	3.1
19	280	4.8	225	3.1	120	1.9	3.7	3.2
20	260	5.2	230	—	E	3.2	3.5	
21	240	5.2	—	—		3.8	3.2	
22	240	4.9	—	—		3.4	2.2	
23	260	4.2	—	—		2.4	3.1	

Time: 15.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 8 minutes.

Table 8*

Time	Falkland Is. (51.7°S , 57.8°W)						August 1953	
	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	310	2.6						
01	290	2.5						
02	290	2.5						
03	275	2.4						
04	255	2.4						
05	220	2.3						
06	230	1.9						
07	235	3.3						
08	225	4.3						
09	230	4.7	220	—				
10	235	5.0	220	(3.8)				
11	240	5.2	220	3.7				
12	250	5.7	220	3.8				
13	245	5.5	215	3.7	115	2.7		
14	240	5.1	210	3.4	120	2.5		
15	230	5.2	205	3.0	125	2.3		
16	225	4.9	(210)		155	2.0		
17	215	4.1						
18	230	3.0						
19	240	2.7						
20	255	2.4						
21	260	2.5						
22	285	2.6						
23	290	2.6						

Time: 60.0°W .

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 10*

Time	Inverness, Scotland (57.4°E , 4.2°W)						July 1953	
	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	270	(3.2)						
01	280	(2.6)						
02	275	(2.5)						
03	280	(2.5)						
04	(290)	(3.0)	255	2.5	(145)	(1.5)		
05	355	3.4	240	2.9	125	1.8		
06	285	3.5	225	3.3	115	2.1		
07	460	3.8	215	3.5	110	2.3		
08	410	4.2	220	3.7	105	2.6		
09	430	4.3	210	3.9	105	2.7		
10	455	4.4	220	4.1	100	3.0		
11	395	4.8	225	4.2	110	3.0		
12	410	4.7	225	4.2	115	3.1		
13	440	4.8	230	4.2	115	3.1		
14	430	4.7	230	4.2	115	3.1		
15	415	4.6	218	4.1	115	3.0		
16	400	4.7	230	4.0	115	2.8		
17	375	4.6	235	3.8	115	2.6		
18	340	4.8	245	3.6	120	2.2		
19	290	5.0	245	3.1	130	1.9		
20	260	5.1						
21	245	5.3						
22	255	4.8						
23	265	4.2						

Time: 0.0°E .

Sweep: 0.55 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 13*

Singapore, British Malaya (1.8°N, 102.8°E)							July 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	260	4.0					4.8	3.0
01	270	3.5					4.2	(3.3)
02	285	3.0					3.9	
03	245	2.5					3.7	—
04	240	2.3					4.0	—
05	(245)	2.3					4.4	—
06	260	2.9					4.8	3.2
07	250	5.6					4.8	3.2
08	290	6.9	225	4.1	120	2.7	5.9	3.1
09	330	7.7	210	4.3	115	3.0	6.0	2.8
10	340	8.8	205	4.3	110	3.2	6.7	2.7
11	350	9.0	200	4.4	110	3.3	6.7	2.6
12	360	9.1	200	4.4	110	3.4	6.7	2.5
13	365	8.9	200	4.4	110	3.4	6.3	2.5
14	350	9.1	200	4.3	110	3.3	6.0	2.5
15	335	8.6	205	4.2	110	3.1	5.2	2.6
16	315	8.5	215	4.1	115	2.8	4.3	2.6
17	375	8.3	225		120	2.3	3.8	2.8
18	345	8.0					3.9	2.9
19	235	8.0					3.5	3.1
20	226	7.2					3.7	3.3
21	215	5.4					3.7	2.5
22	225	4.3					4.1	3.3
23	345	3.5					4.7	3.0

Time: 105.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 15

Christchurch, New Zealand (43.6°S, 172.7°E)							July 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	2.2					3.0	3.0
01	280	2.1					4.0	3.1
02	270	2.1					2.8	3.1
03	260	2.0					3.6	3.2
04	260	1.5					3.9	3.2
05	240	1.5					3.4	3.3
06	—	1.5					4.0	3.1
07	270	2.0					4.0	3.2
08	240	3.4	240	—			1.6	3.5
09	250	4.0	230	3.0			2.0	3.5
10	250	4.3	220	3.4			2.3	3.4
11	280	4.5	220	3.7			2.5	3.4
12	280	4.9	230	3.7			2.6	4.3
13	280	4.9	220	3.7			2.5	3.4
14	270	4.9	240	3.6			2.4	3.4
15	260	4.7	230	3.6			2.1	3.5
16	240	4.4	240	2.7			1.7	3.5
17	240	3.5					3.5	3.3
18	260	2.8					2.9	3.1
19	260	2.4					2.4	3.1
20	270	2.3					3.5	3.1
21	270	2.2					2.1	3.1
22	280	2.1					3.5	3.0
23	280	2.1					2.7	3.0

Time: 172.5°E.

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 55 seconds.

Table 17*

Inverness, Scotland (57.4°N, 4.2°W)							June 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	255	3.9					1.6	2.9
01	260	3.5					2.4	2.9
02	285	3.2					2.5	2.9
03	280	3.0					2.4	2.9
04	300	3.5	265	2.7	145	1.5	2.8	2.9
05	365	3.9	235	3.1	120	1.8	3.0	3.0
06	370	4.2	215	2.5	110	2.1	3.1	2.9
07	380	4.3	215	3.8	105	2.5	3.7	3.0
08	410	4.5	210	3.9	100	2.7	3.2	3.0
09	415	4.8	205	4.0	100	2.8	3.8	2.9
10	420	4.9	210	4.1	100	2.9	3.8	2.9
11	400	4.8	210	4.2	100	2.0	3.6	2.9
12	375	4.9	215	4.2	100	3.0	4.0	2.9
13	410	4.7	210	4.2	100	3.0	4.0	2.9
14	435	4.7	210	4.2	100	2.9	3.4	2.9
15	405	4.8	210	4.1	100	2.9	3.3	2.8
16	375	4.8	220	4.0	105	2.9	3.3	2.9
17	380	4.8	220	2.9	100	2.6	3.8	3.0
18	340	5.0	220	3.7	110	2.4	3.2	3.0
19	310	5.1	235	2.4	120	2.0	2.8	3.1
20	275	5.2	245	2.9	145	1.7	2.7	3.1
21	260	5.0					1.7	3.1
22	250	4.9					3.0	3.0
23	260	4.6					3.0	3.0

Time: 0.0°.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 14

Barotonga I. (21.3°S, 159.8°W)							July 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	< 290	2.9						2.9
01	250	3.0						2.9
02	250	3.0						3.0
03	250	3.1						3.2
04	240	2.9						3.2
05	(240)	2.4						3.0
06	270	2.3						3.0
07	250	4.0					< 1.7	3.3
08	260	5.2	200	2.8	120		2.1	3.5
09	260	5.3	200	3.9	110		2.5	3.3
10	270	6.2	200	4.0	110		2.8	3.5
11	270	5.9	200	4.2	110		2.9	3.5
12	280	5.6	200	4.2	110		3.0	3.5
13	290	5.7	200	4.2	110		3.0	3.4
14	290	5.6	200	4.1	110		2.9	3.3
15	290	5.8	200	4.0	110		2.8	3.4
16	260	5.6	220	3.6	115		2.5	3.4
17	250	5.6					2.0	3.3
18	240	5.3						3.4
19	230	4.4						3.2
20	250	3.2						3.0
21	260	3.0						2.4
22	270	2.9						2.3
23	270	3.1						2.2

Time: 157.5°W.

Sweep: 2.0 Mc to 16.0 Mc, manual operation.

Table 16*

Falkland Is. (51.7°S, 57.8°W)							July 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	300	2.4						2.9
01	290	2.4						3.0
02	275	2.3						3.0
03	280	2.3						3.1
04	265	2.2						1.6 (3.2)
05	240	2.2						1.7 (3.7)
06	210	2.0						
07	235	1.8						
08	225	3.3						3.5
09	220	4.0						3.7
10	220	4.4						3.8
11	230	4.9						3.6
12	240	5.0	205	(3.4)	120	2.3	3.7	3.6
13	225	5.1	(220)	(3.2)	125	2.3	3.4	3.7
14	230	4.9	210	(125)	(2.1)	2.8	3.7	3.7
15	235	4.9	220	4.0	110	3.0	5.9	3.0
16	385	4.9	235	4.2	110	3.0	5.6	3.0
17	380	5.2	235	4.2	105	3.1	5.6	3.0
18	380	5.2	230	4.3	110	3.1	5.2	3.0
19	395	5.0	230	4.3	110	3.2	5.8	2.9
20	425	4.9	240	4.3	110	3.2	5.4	2.8
21	430	4.9	230	4.3	110	3.1	5.0	2.8
22	395	5.0	230	4.3	110	2.9	4.9	3.0
23	365	5.1	235	4.1	110	2.7	4.8	3.0
24	360	5.2	230	3.9	110	2.7	4.8	3.0
25	375	5.4	245	3.6	115	2.3	4.6	3.0
26	295	5.7	255	3.2	125	1.9	4.4	3.0
27	265	6.2	(255)	(2.5)				3.6
28	255	6.1						3.1
29	250	5.4						3.4
30	260	4.6						3.0

Time: 0.0°.

Sweep: 0.55 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 19*

Singapore, British Malaya (1.3°N, 103.8°E)								June 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	265	3.3				3.9	3.0		
01	245	3.9				4.8	(3.1)		
02	235	3.0				4.3	(3.3)		
03	245	2.4				3.5	(3.5)		
04	240	2.2				3.4	(3.3)		
05	260	1.7				3.8	—		
06	260	3.4			(160)	(1.2)	3.5	3.2	
07	260	6.0	235		120	2.2	5.2	3.1	
08	300	7.6	225	4.2	115	2.7	5.4	2.9	
09	310	8.4	215	4.3		3.0	5.4	2.8	
10	330	9.4	205	4.4		3.3	5.7	2.6	
11	345	9.5	200	4.4		3.4	5.7	2.5	
12	360	9.2	200	4.4		3.4	5.9	2.5	
13	345	9.1	200	4.4		3.4	5.8	2.5	
14	335	9.0	205	4.4		3.3	5.9	2.6	
15	325	8.8	210	4.3	115	3.1	5.5	2.7	
16	290	8.8	215	4.2	115	2.7	5.2	2.9	
17	255	8.6	230		120	2.2	5.4	3.0	
18	230	8.3			(140)	(1.5)	4.4	3.1	
19	225	7.6					3.8	3.2	
20	220	6.1					3.8	3.4	
21	220	5.0					4.1	3.4	
22	225	3.8					4.8	3.3	
23	255	3.6					4.4	3.0	

Time: 105.0°E.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 21

Canberra, Australia (35.3°S, 149.0°E)								June 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	—	3.6						3.2	
01	—	3.7						3.2	
02	—	3.8						3.2	
03	(240)	3.8						3.2	
04	(240)	3.7						3.2	
05	(200)	3.7						3.4	
06	(200)	3.1						3.4	
07	210	3.5						3.5	
08	210	4.8			—	(1.8)	3.2	3.7	
09	230	5.2	220	3.4	110	(2.2)	3.0	3.7	
10	240	5.3	200	(3.8)	100	2.6	3.4	3.6	
11	250	5.7	210	4.0	100	2.8	3.5	3.6	
12	240	5.5	210	4.0	100	2.9	3.5	3.8	
13	260	5.8	200	4.0	100	2.8	3.5	3.5	
14	250	5.8	200	3.8	100	2.7	3.5	3.5	
15	240	6.0	210	(3.6)	100	2.4	3.5	3.6	
16	220	5.6	220	—	—	(1.8)	3.3	3.8	
17	210	5.2					3.4	3.8	
18	200	3.9					3.2	3.4	
19	—	3.4					3.2	3.4	
20	—	3.2					3.0	3.4	
21	—	3.5					2.8	3.4	
22	—	3.2					2.8	3.3	
23	—	3.5					2.4	3.2	

Time: 150.0°E.

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 23*

Falkland Is. (51.7°S, 57.8°W)								June 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	300	2.4						2.8	
01	290	2.5						2.9	
02	290	2.4						2.9	
03	275	2.4						3.0	
04	260	2.4						3.1	
05	245	2.3				1.0		3.2	
06	230	2.2				2.5		3.8	
07	245	1.9			170	1.1	1.8	3.4	
08	220	3.4			160	1.6	2.8	3.6	
09	210	4.0			130	(1.8)	3.0	3.8	
10	220	4.6			120	(2.1)	3.8	3.7	
11	220	4.6			120	2.3	4.2	3.8	
12	225	5.0	(200)		120	2.4	3.0	3.7	
13	225	5.2	(215)	(3.3)	125	(2.4)	3.0	3.7	
14	220	5.1			130	2.2	3.0	3.7	
15	220	4.6			160	1.9	2.7	3.7	
16	215	3.9				2.7		3.7	
17	235	2.8				2.0		3.3	
18	240	2.4				2.5		3.2	
19	255	2.3				2.5		3.2	
20	250	2.4				2.8		3.3	
21	260	2.4				3.5		3.1	
22	270	2.4				3.8		3.0	
23	290	2.4				2.8		2.9	

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 20

Townsville, Australia (19.3°S, 146.6°E)								June 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	220	(3.0)						—	(3.0)
01	220	(3.0)						—	(3.1)
02	220	3.0						—	(3.2)
03	200	(3.0)						1.8	(3.4)
04	200	(3.0)						2.0	(3.2)
05	200	(2.8)						—	3.2
06	200	3.0						—	3.5
07	220	4.4						—	3.5
08	220	5.5						100	3.4
09	250	5.9	220	4.0				110	4.0
10	250	6.5	210	4.1				110	4.4
11	250	6.5	220	4.3				120	3.2
12	270	8.1	200	4.3				110	4.3
13	270	8.4	210	4.2				110	4.5
14	260	6.5	220	4.2				110	4.8
15	250	6.0	200	4.0				120	3.4
16	240	5.8	220	—				110	4.2
17	220	5.6	—	—				2.0	3.4
18	210	4.8						—	3.4
19	230	3.7						—	3.5
20	230	3.2						—	3.1
21	230	3.4						—	2.5
22	240	3.0						—	3.1
23	230	3.0						—	3.0

Time: 150.0°E.

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 22

Hobart, Tasmania (42.9°S, 147.3°E)								June 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	270	2.0						—	3.0
01	270	2.1						—	3.0
02	270	2.3						—	2.9
03	260	2.3						—	3.0
04	260	2.0						—	3.0
05	250	2.1						—	3.0
06	250	2.0						—	3.0
07	260	2.0						—	3.0
08	230	3.8	270	3.7				100	3.2
09	230	3.8	250	3.8				100	3.4
10	240	4.0	250	3.8				100	3.6
11	240	4.0	230	4.0				100	3.6
12	240	4.0	210	4.0				100	3.6
13	250	4.1	210	4.0				100	3.6
14	240	4.1	220	4.0				100	3.4
15	240	4.1	230	4.0				100	3.4
16	250	4.2	210	4.0				100	3.1
17	250	4.2	200	4.0				100	3.1
18	250	4.2	230	3.6				110	2.8
19	250	4.5	240	3.8				110	2.8
20	310	4.0	—	—				110	2.6
21	300	3.7	—	—				120	2.9
22	280	3.8	—	—				120	6.0
23	280	3.9	—	—				—	(3.0)

Time: 150.0°E.

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 24

Ottawa, Canada (45.8°N, 75.8°W)								May 1953	
Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2	
00	270	3.3						—	9.0
01	280	3.0						—	9.0
02	280	3.0						—	6.5
03	300	3.2						—	6.0
04	300	3.6						—	(2.9)
05	280	3.6	—	—				—	5.5
06	350	< 3.8	270	3.7				100	6.0
07	420	< 4.0	250	3.8				100	3.4
08	420	< 4.0	250	3.9					

Time	h'F2	foF2	h'F1	foF1	h'E	foE	May 1953	
							(M3000)F2	
00	275	3.4	—	—	—	—	2.9	
01	285	3.7	—	—	—	—	2.8	
02	290	2.3	—	—	—	—	2.8	
03	295	2.4	—	—	—	—	2.8	
04	290	2.7	—	—	—	—	2.9	
05	300	3.4	2.0	1.7	2.8	3.1		
06	365	3.7	220	1.5	2.1	2.9	2.9	
07	300	4.2	220	3.6	110	2.3	3.0	2.9
08	405	4.4	215	3.7	105	2.6	3.0	3.0
09	385	4.7	215	3.9	105	2.7	3.0	3.0
10	405	4.7	210	4.0	105	2.8	3.1	2.9
11	330	4.8	215	4.1	100	2.9	3.1	3.0
12	390	4.5	220	4.2	105	2.9	3.1	3.0
13	420	4.7	215	4.2	105	3.0	3.1	2.8
14	420	4.7	210	4.1	100	2.9	3.0	2.9
15	405	4.8	220	4.1	105	2.9	3.1	2.8
16	390	4.9	225	3.9	105	2.7	3.0	2.8
17	330	5.1	225	3.8	110	3.5	2.8	3.0
18	305	5.0	225	3.9	110	3.3	2.6	3.1
19	290	5.0	230	3.0	140	1.9	2.6	3.1
20	285	5.8	265	2.6	155	1.7	2.0	3.1
21	255	4.7	—	—	—	—	2.1	3.1
22	260	4.5	—	—	—	—	3.3	3.0
23	275	5.5	—	—	—	—	3.3	3.0

Time: 0.0°.

Sweep: 0.67 Mc to 20.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 27*

Time	Singapore, British Malaya (1.3°N, 103.8°E)						(M3000)F2
	h'F2	foF2	h'F1	foF1	h'E	foE	
00	255	3.6	—	—	3.5	3.2	
01	260	3.4	—	—	3.6	3.2	
02	245	3.8	—	—	3.0	3.3	
03	250	2.2	—	—	3.5	3.3	
04	260	3.1	—	—	3.0	(3.2)	
05	245	2.0	—	—	3.4	(3.2)	
06	265	3.6	—	—	3.0	3.2	
07	255	6.4	220	1.3	150	1.3	3.0
08	290	8.0	225	4.2	120	2.8	5.0
09	305	9.0	315	4.4	115	3.1	5.3
10	325	9.6	200	4.5	110	3.3	5.9
11	320	9.5	200	4.5	110	3.4	5.7
12	345	9.7	200	4.5	110	3.4	5.6
13	335	9.2	200	4.5	110	3.4	5.8
14	325	9.2	200	4.4	110	3.3	5.4
15	315	9.4	215	4.3	115	3.0	5.0
16	280	9.5	225	4.1	115	2.7	4.4
17	265	9.6	235	4.1	120	3.2	4.5
18	235	9.2	—	—	—	—	3.2
19	225	8.3	—	—	—	—	3.6
20	220	6.5	—	—	—	—	3.9
21	220	5.1	—	—	—	—	3.4
22	215	4.3	—	—	—	—	3.3
23	240	3.4	—	—	—	—	3.4

Time: 105.0°E.

Sweep: 0.67 Mc to 20.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 29

Time	Brisbane, Australia (27.5°S, 153.0°E)						(M3000)F2
	h'F2	foF2	h'F1	foF1	h'E	foE	
00	250	3.8	—	—	—	—	3.0
01	260	3.8	—	—	—	—	3.1
02	250	3.8	—	—	—	—	3.1
03	240	4.0	—	—	—	—	3.2
04	230	3.9	—	—	—	—	3.3
05	230	3.3	—	—	—	—	3.3
06	240	3.5	—	—	—	—	3.1
07	220	5.0	—	—	—	—	3.6
08	240	5.9	220	3.8	110	2.6	3.5
09	240	6.4	220	4.1	110	2.8	3.5
10	250	6.2	210	4.2	100	3.3	3.5
11	245	6.5	210	4.3	100	3.3	3.4
12	260	6.5	210	4.3	100	3.3	3.4
13	270	6.1	210	4.2	100	3.2	3.3
14	260	6.7	210	4.0	100	3.0	3.5
15	240	7.0	225	3.8	105	2.7	3.6
16	230	6.6	225	3.2	—	—	3.6
17	220	5.4	—	—	—	—	3.4
18	220	4.0	—	—	—	—	3.4
19	230	3.6	—	—	—	—	3.2
20	250	3.6	—	—	—	—	3.0
21	260	4.2	—	—	—	—	3.1
22	250	4.1	—	—	—	—	3.1
23	250	3.9	—	—	—	—	3.1

Time: 150.0°E.

Sweep: 1.6 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 26*

Time	Slough, England (51.5°N, 0.6°W)						(M3000)F2
	h'F2	foF2	h'F1	foF1	h'E	foE	
00	275	3.5	—	—	—	—	3.4
01	275	3.4	—	—	—	—	2.5
02	280	3.1	—	—	—	—	2.8
03	275	2.9	—	—	—	—	3.0
04	280	3.1	—	—	—	—	2.8
05	290	3.6	240	3.1	130	1.7	4.3
06	355	4.1	240	3.4	120	2.1	3.7
07	355	4.6	225	3.8	120	2.5	4.6
08	395	4.8	230	4.0	115	2.7	4.7
09	405	4.8	225	4.1	115	2.9	4.5
10	400	4.9	230	4.2	115	3.0	4.9
11	385	4.9	215	4.2	115	3.1	4.9
12	375	5.1	220	4.3	115	3.1	4.9
13	395	5.0	230	4.3	115	3.2	4.7
14	395	5.0	220	4.3	115	3.2	4.7
15	365	5.0	220	4.2	115	3.1	4.6
16	360	5.2	230	4.0	115	2.8	4.4
17	325	5.3	235	3.8	120	2.5	3.5
18	295	5.6	240	3.5	120	2.1	3.0
19	270	5.6	245	2.9	140	1.7	3.2
20	250	5.8	—	—	—	—	3.2
21	245	5.3	—	—	—	—	2.3
22	260	4.6	—	—	—	—	3.4
23	270	4.0	—	—	—	—	3.0

Time: 0.0°.

Sweep: 0.55 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 28

Time	Townsville, Australia (19.3°S, 146.8°E)						(M3000)F2
	h'F2	foF2	h'F1	foF1	h'E	foE	
00	220	3.0	—	—	—	—	3.2
01	235	3.0	—	—	—	—	3.1
02	230	3.2	—	—	—	—	3.1
03	220	3.2	—	—	—	—	3.2
04	220	2.8	—	—	—	—	3.1
05	230	2.7	—	—	—	—	3.1
06	220	3.0	—	—	—	—	3.3
07	220	4.9	—	—	—	—	3.5
08	230	5.9	210	3.7	110	2.4	3.7
09	250	6.7	210	4.0	110	2.8	3.7
10	250	7.0	210	4.2	110	3.0	4.0
11	260	7.4	200	4.3	115	3.2	4.4
12	250	7.1	200	4.4	110	3.2	4.4
13	260	7.2	220	4.3	110	3.2	4.4
14	250	7.1	200	4.2	120	3.0	4.5
15	250	6.7	200	4.0	120	2.8	4.5
16	240	6.0	—	—	—	—	3.4
17	230	6.0	—	—	—	—	3.4
18	210	4.8	—	—	—	—	3.3
19	220	3.8	—	—	—	—	3.2
20	225	3.2	—	—	—	—	2.7
21	240	3.2	—	—	—	—	3.1
22	240	3.0	—	—	—	—	3.0
23	230	3.1	—	—	—	—	3.3

Time: 150.0°E.

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 30

Time	Canberra, Australia (35.3°S, 149.0°E)						(M3000)F2
	h'F2	foF2	h'F1	foF1	h'E	foE	
00	—	3.4	—	—	—	—	2.9
01	(240)	3.2	—	—	—	—	3.1
02	(240)	3.5	—	—	—	—	3.1
03	(240)	3.4	—	—	—	—	3.2
04	230	3.4	—	—	—	—	2.4
05	200	3.2	—	—	—	—	3.4
06	(200)	2.8	—	—	—	—	3.6
07	220	3.6	—	—	—	—	3.6
08	220	5.0	210	—	—	1.6	2.9
09	230	5.6	220	3.5	100	3.5	3.6
10	240	5.7	210	(4.0)	100	2.8	3.5
11	250	5.8	210	4.0	100	2.9	3.5</td

Table 31

Hobart, Tasmania (42°S 147°E)							May 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	2.0						3.0
01	290	2.0						3.0
02	290	1.8						3.0
03	290	1.7						(3.0)
04	300	1.7						(3.0)
05	—	E						(2.9)
06	—	E						(3.0)
07	250	2.0						3.1
08	220	3.5	110	1.8				3.1
09	210	4.2	100	2.1				3.2
10	200	4.7	100	2.4				3.2
11	200	5.0	100	2.6				3.2
12	200	5.5	100	2.6				3.2
13	200	5.5	100	2.6				3.1
14	210	5.5	100	2.5				3.2
15	210	5.7	100	2.3				3.2
16	220	5.5	100	1.9				3.2
17	210	4.5						3.2
18	220	3.7						3.1
19	230	3.0						3.1
20	250	2.3						3.0
21	250	2.1						3.1
22	250	2.0						3.0
23	280	2.0						3.0

Time: 150.0°E.

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 55 seconds.

Table 32*

Inverness, Scotland (57°N 4°W)							April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	310	2.1						2.8
01	310	(1.6)						2.2
02	280	(1.7)						(2.7)
03	330	(1.6)						2.0
04	305	(1.7)						(2.7)
05	270	2.4						2.6
06	365	3.1	(230)	(2.7)	140	1.7	2.8	3.3
07	280	3.8	220	3.3	120	2.0	2.7	3.2
08	450	4.1	215	3.6	110	2.4	3.0	3.1
09	420	4.4	220	3.8	110	2.5	2.9	3.0
10	415	4.4	215	4.0	110	2.8	3.0	2.9
11	395	4.7	210	4.0	110	2.7	2.9	2.9
12	410	4.8	215	4.1	105	2.8	2.9	2.9
13	395	4.9	205	4.1	105	2.9	2.8	2.9
14	380	5.0	215	4.0	105	2.8	3.0	3.1
15	380	5.0	215	4.0	105	2.7	2.8	2.9
16	340	5.0	220	3.9	110	2.5	2.8	3.1
17	315	5.0	235	3.5	115	2.3	2.3	3.1
18	280	4.9	245	3.2	130	1.9	3.1	
19	255	4.7	(260)	(2.7)	(150)	(1.6)		
20	250	4.4						3.1
21	260	4.0						3.0
22	275	3.2						2.9
23	310	(2.6)						2.9

Time: 0.0°.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 33*

Singapore, British Malaya (1.3°S 103.8°E)							April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	240	7.1						2.8
01	230	8.4						3.3
02	225	5.0						2.4
03	230	3.8						2.3
04	245	2.9						2.2
05	250	2.3						2.2
06	260	3.2						3.1
07	240	6.5						3.2
08	290	8.2	225	125	2.2	4.2	3.2	
09	300	9.4	215	(4.3)	115	3.1	4.5	2.8
10	315	9.8	210	4.6	110	3.3	5.9	2.4
11	335	10.5	200	4.7	110	3.5	5.7	2.2
12	340	9.8	200	4.6	110	3.5	5.6	2.3
13	335	10.0	200	4.7	110	3.5	5.8	2.4
14	325	10.0	200	4.5	110	3.4	5.4	2.5
15	310	10.2	215	4.4	110	3.2	5.4	2.5
16	295	10.5	230	(115)	2.8	5.4	2.6	
17	275	10.6	235		2.3	5.0	2.7	
18	255	11.1						4.8
19	255	11.0						4.0
20	245	10.4						3.9
21	220	9.5						3.8
22	220	8.2						3.3
23	220	7.2						2.8

Time: 100.0°E.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 32*

Falkland Is. (51.7°S, 57.8°W)							May 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	2.6						1.6
01	310	2.5						2.8
02	295	2.6						2.3
03	285	2.6						2.9
04	270	2.6						2.1
05	240	2.5						1.0
06	205	2.4						1.4
07	235	2.6					(170)	1.4
08	210	4.1					(155)	1.3
09	210	4.6					(120)	2.2
10	225	5.0					(205)	3.7
11	220	5.3					(210)	3.7
12	225	6.0					(215)	3.7
13	220	5.8					(220)	3.7
14	220	5.6					(215)	3.7
15	220	5.4					(220)	3.7
16	220	5.4					(24)	3.1
17	210	4.6					(140)	3.1
18	215	3.4						3.8
19	235	2.6						3.4
20	235	2.7						3.3
21	255	2.3						3.0
22	280	2.5						2.9
23	290	2.6						2.9

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 33*

Slough, England (51.5°N, 0.6°W)							April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	3.1						2.4
01	290	2.9						2.4
02	265	2.8						2.8
03	265	2.6						2.8
04	275	2.5						2.9
05	265	2.6						3.0
06	270	3.5	235	3.0	130	1.8		3.0
07	315	4.2	230	3.5	120	2.2		3.2
08	365	4.4	225	3.8	120	2.6		3.0
09	365	4.8	220	4.0	115	2.8		3.0
10	365	5.0	220	4.2	115	3.0		3.1
11	385	5.2	220	4.2	115	3.1		3.1
12	365	5.3	220	4.3	115	3.1		3.0
13	365	5.5	225	4.3	115	3.1		3.0
14	335	5.6	230	4.2	115	3.0		3.1
15	330	5.4	220	4.1	120	2.9		3.1
16	315	5.5	230	3.9	120	2.6		3.1
17	280	5.4	235	3.6	120	2.3		3.1
18	265	5.4	245	3.2	130	1.9		2.6
19	260	5.4						2.4
20	250	5.0						2.5
21	250	4.3						2.3
22	270	3.4						2.2
23	285	3.2						2.4

Time: 0.0°.

Sweep: 0.67 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 34*

Townsville, Australia (19.3°S, 146.8°E)							April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	260	(3.4)						3.5
01	260	3.5						3.0
02	260	3.5						3.5
03	220	3.2						3.3
04	220	2.8						3.2
05	230	2.7						2.7
06	230	3.0						2.4
07	220	5.2						3.5
08	230	6.4	215	3.5	110	2.5		3.4
09	250	7.6	220	4.3	110	3.0		3.3
10	260	8.5	220	4.4	110	3.2		3.3
11	250	8.9	210	4.5	110	3.3		3.4
12	260	8.2	200	4.4	115	3.3		3.4
13	270	8.0	200	4.4	110	3.3		3.2
14	260	8.4	220	4.3	110	3.2		3.3
15</td								

Table 37

Brisbane, Australia (27.6°S, 153.0°E)								April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	270	3.9				3.5	3.0		
01	280	3.8				3.3	3.0		
02	270	3.0				3.2	3.0		
03	280	4.0				2.5	3.2		
04	220	3.7				2.6	3.2		
05	230	3.4				2.0	3.2		
06	230	3.9					3.3		
07	220	5.6					3.8		
08	240	6.1	220	4.0	120	2.2			
09	260	6.8	220	4.2	100	3.0			
10	260	7.5	210	4.4	100	3.2	3.4		
11	250	7.6	210	4.5	110	3.2	3.5		
12	285	7.5	200	4.5	110	3.2	3.5		
13	280	7.2	210	4.6	110	3.1	3.3		
14	260	7.8	220	4.3	105	3.1	3.4		
15	250	7.8	230	4.1	105	3.0	3.2		
16	240	7.4	230	3.5	110		3.4		
17	220	6.6					3.4		
18	220	5.4					3.2		
19	240	4.7					3.4		
20	250	4.4					3.1		
21	250	4.3					3.7		
22	250	4.2					3.4		
23	250	4.0					2.8		

Time: 150.0°E.

Sweep: 1.6 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 39

Hobart, Tasmania (42.8°S, 147.5°E)								April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	270	2.8					2.9		
01	280	2.6					2.9		
02	270	2.5					2.9		
03	280	2.0					2.9		
04	290	2.0					2.9		
05	270	1.8					3.0		
06	280	2.0					(3.0)		
07	225	3.6					3.1		
08	220	4.7					3.2		
09	210	5.5					3.1		
10	210	5.7					3.1		
11	200	6.4					3.0		
12	200	6.6					3.1		
13	200	7.0					3.1		
14	200	6.6					3.1		
15	215	6.6					3.1		
16	220	6.0					3.1		
17	230	6.0					3.1		
18	240	5.1					3.1		
19	250	4.6					3.0		
20	250	3.7					3.0		
21	250	3.4					3.0		
22	270	3.0					2.9		
23	280	2.8					2.9		

Time: 150.0°E.

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 55 seconds.

Table 41

Poitiers, France (46.6°N, 0.3°E)								March 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	< 280	3.0					3.0		
01	< 280	3.0					2.9		
02	< 280	2.9					2.9		
03	< 270	2.9					2.9		
04	< 270	2.6					3.0		
05	(245)	2.2					(3.2)		
06	245	2.8					3.2		
07	245	4.0	220	2.2	—	E	2.0		
08	260	4.6	225	3.4	115	2.3	2.0		
09	280	5.0	205	3.8	110	2.6	3.4		
10	280	5.3	210	4.0	110	2.7	2.8		
11	285	5.6	215	4.2	105	2.9	3.4		
12	300	5.6	210	4.1	110	3.0	3.3		
13	300	5.8	220	4.1	110	3.0	3.3		
14	295	5.8	230	4.0	110	2.9	3.4		
15	285	5.6	235	3.9	115	2.8	3.3		
16	270	5.5	235	3.6	115	2.4	3.4		
17	250	6.4	245	2.5	130	2.0	2.1		
18	240	5.1	—	—	—	—	2.1		
19	240	4.8					3.4		
20	245	4.4					3.2		
21	< 245	3.8					3.2		
22	< 260	3.4					3.0		
23	270	3.0					3.0		

Time: 0.0°.

Sweep: 1.6 Mc to 18.0 Mc in 1 minute.

Table 38

Canberra, Australia (35.5°S, 149.0°E)								April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	—	—	—	—	—	—	—	3.6	3.0
01	(240)	3.5						3.0	3.0
02	(240)	3.5						3.0	3.0
03	(240)	3.6						3.2	3.1
04	230	3.6						2.8	3.3
05	(230)	3.3						3.0	3.3
06	(215)	3.1						2.6	3.3
07	210	4.5						1.7	3.6
08	230	5.6	210	—	—	—	—	2.3	3.6
09	240	6.1	205	(4.0)	100	2.6	3.8		3.5
10	250	6.4	200	4.1	100	2.6	3.8		3.5
11	260	6.5	200	4.2	100	3.1	3.4		3.4
12	260	7.2	200	4.2	100	3.2	3.5		3.4
13	260	7.8	210	4.2	100	3.1	3.5		3.4
14	240	7.0	210	4.2	100	3.1	3.5		3.3
15	245	7.2	220	(4.0)	100	2.9	3.5		3.4
16	240	6.9	220	(3.7)	110	2.6	3.4		3.5
17	230	6.2	—	—	—	—	1.7		3.5
18	210	5.8							3.4
19	215	4.6							3.2
20	(230)	4.4							3.1
21	230	4.1							3.1
22	(240)	3.8							3.1
23	—	3.8							3.1

Time: 150.0°E.

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 40

Falkland Is. (51.7°S, 57.8°W)								April 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	300	3.2						2.0	2.8
01	300	3.2						2.2	2.8
02	290	3.1						1.8	2.8
03	285	3.1						1.8	2.8
04	280	3.1						1.6	2.9
05	255	3.2						2.4	3.1
06	225	3.2						(170) (1.2)	3.4
07	215	5.0						150	1.4
08	215	6.1	225	—	—	—	—	2.2	3.6
09	225	6.9	225	(3.8)	115	2.6	4.5		3.6
10	235	7.6	225	4.1	110	2.8	4.8		3.5
11	235	8.4	220	4.2	110	2.8	4.8		3.5
12	230	8.1	220	4.1	105	2.8	4.6		3.6
13	230	7.3	220	4.0	110	2.8	4.1		3.7
14	225	6.3	210	3.8	110	2.7	4.8		3.7
15	225	6.0	220	3.8	115	2.4	3.2		3.7
16	225	5.6	240	3.9	110	2.5	3.4		3.4
17	220	5.3	—	—	—	—	1.7		3.4
18	215	4.0	—	—	—	—	—		3.6
19	230	5.0	210	3.0	110	2.2	3.6		3.6
20	250	5.5	200	3.8	105	2.6	3.4		3.5
21	275	5.9	200	4.2	100	2.9	3.4		3.4
22	270	7.0	215	4.3	100	3.2	3.4		3.4
23	270	7.1	230	4.2	100	3.0	3.4		3.4

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 41

Casablanca, Morocco (33.6°N, 7.6°W)								March 1953	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	< 250	3.4						3.0	
01	< 275	3.2						3.0	
02	250	3.1						2.9	
03	< 250	3.1						3.0	
04	< 250	3.0							

Table 43*

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	7.2						(2.7)
01	270	6.0						(3.0)
02	250	5.7						(3.2)
03	230	5.1						(3.3)
04	(230)	3.7						(3.4)
05	(240)	2.2						(3.4)
06	260	3.9						3.4
07	(240)	6.6	(220)	(4.2)	120	2.2		3.2
08	(250)	(7.9)	(230)	(4.5)	110	2.8	5.0	(3.2)
09	(230)	(8.8)	220	(4.5)	120	3.2	(3.7)	(2.9)
10	(330)	(9.3)	220	(4.6)	110	3.3	(4.1)	(2.6)
11	(370)	(9.9)	220	4.6	110	3.4	(3.5)	(2.8)
12	(320)	9.8	220	4.6	110	3.5		2.8
13	320	10.2	200	4.6	110	3.4	4.5	(2.8)
14	310	10.7	210	4.5	110	3.3		2.9
15	300	11.5	220	4.4	110	3.1	4.1	3.1
16	310	11.4	230	(4.0)	110	2.8	5.2	3.2
17	260	10.0	230		120	2.2	5.2	(2.6)
18	260	10.0						3.0
19	250	9.6			110	(1.5)	4.7	(2.9)
20	260	9.2						(3.0)
21	250	8.8						(2.8)
22	280	(7.7)						(2.7)
23	290	(7.5)						(2.7)

Time: 30.0° E.

Sweep: 0.67 to 25.0 Mc in 5 minutes.

*Average values except for foF2 and fEs, which are median values.

Table 45

Time	Poitiers, France (46.6°N, 0.3°E)							February 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	265	3.3						3.0
01	265	3.2						3.0
02	265	3.3						3.0
03	265	3.3						3.0
04	< 270	3.1						3.0
05	< 250	2.6						(3.1)
06	< 230	2.5						(3.2)
07	225	3.6						3.4
08	220	4.7	220	2.3	150	2.0		3.8
09	230	5.2	215	3.2	115	2.4		3.8
10	245	5.5	210	3.6	115	2.6		3.8
11	250	5.8	210	3.9	115	2.8	2.0	3.5
12	250	6.0	220	4.0	115	2.9	1.8	3.6
13	260	5.8	230	3.9	115	2.8		3.4
14	250	5.6	225	3.8	115	2.7		3.5
15	250	5.8	230	3.4	120	2.5		3.5
16	240	5.6	235	—	125	2.1	2.3	3.5
17	225	5.2	—	1.9	—	—	2.0	3.5
18	220	4.2						3.3
19	240	4.0						3.2
20	< 240	3.9						3.2
21	245	3.4						3.2
22	< 240	3.2						(3.0)
23	260	3.1						3.1

Time: 0.0°.

Sweep: 1.8 Mc to 16.8 Mc in 1 minute.

Table 47

Time	Poitiers, France (46.6°N, 0.3°E)							January 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	< 260	3.2						3.0
01	260	3.4						2.9
02	270	3.2						3.0
03	270	3.3						3.0
04	250	3.0						3.2
05	230	2.6						3.2
06	< 230	2.3						3.1
07	< 240	2.6						3.1
08	220	4.9	< 185	2.0	—	—	2.0	3.6
09	230	5.8	210	2.7	125	2.1	2.3	3.6
10	235	5.9	230	3.5	120	2.5	2.0	3.6
11	240	6.8	230	3.8	120	2.6	2.0	3.7
12	235	6.2	220	3.8	120	2.7		3.6
13	245	6.2	220	3.7	120	2.7		3.6
14	245	6.0	235	3.6	120	2.5		3.5
15	235	5.7	230	3.0	125	2.2	2.0	3.6
16	220	5.4	230	2.0	—	—	2.1	3.6
17	215	4.5						3.4
18	225	4.0						3.4
19	230	3.3						3.0
20	250	3.0						(2.9)
21	< 255	3.1						3.0
22	< 250	3.2						3.0
23	250	3.3						3.0

Time: 0.0°.

Sweep: 1.6 Mc to 16.8 Mc in 1 minute.

Table 44*

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	310	3.9						2.6
01	310	3.8						2.7
02	310	3.7						2.7
03	300	3.5						2.6
04	275	3.8						3.0
05	265	3.6						3.1
06	240	4.0						3.1
07	235	4.8						3.5
08	240	5.0	235		115	2.4	3.2	3.4
09	290	5.6	225	(3.7)	105	2.7	4.8	3.3
10	300	6.1	225	4.1	105	2.8	4.9	3.3
11	275	6.7	225	4.2	105	2.9	5.3	3.4
12	290	7.0	225	4.2	(110)	(2.9)	5.3	3.3
13	265	6.8	225	4.2	(110)	(3.0)	5.0	3.4
14	280	6.1	225	4.1	(105)	(2.0)	5.0	3.5
15	250	5.9	220	3.9	(110)	(2.7)	4.9	3.5
16	250	5.7	240	(3.7)	(110)	2.5	4.7	3.5
17	240	5.6						3.6
18	240	5.5						(4.7)
19	260	5.3						3.1
20	265	5.3						3.0
21	260	4.8						3.1
22	280	4.3						2.9
23	295	4.1						2.8

Time: 60.0° W.

Sweep: 0.67 to 26.0 Mc in 5 minutes.

*Average values except for foF2 and fEs, which are median values.

Table 46

Time	Casablanca, Morocco (33.6°N, 7.6°W)							February 1953
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	< 260	3.3						3.0
01	< 250	3.3						3.0
02	—	3.2						3.0
03	< 260	3.2						3.0
04	< 250	3.1						3.1
05	< 220	2.6						3.5
06	< 250	2.2						3.2
07	< 260	2.4						3.2
08	225	4.8	—	—	—	—	—	3.8
09	230	5.7	210	—	120	2.3	—	3.8
10	250	7.2	210	4.1	100	2.7	—	3.5
11	250	8.1	210	4.3	110	2.9	—	3.6
12	250	6.7	200	4.3	105	3.1	—	3.6
13	250	6.6	200	4.3	105	3.0	—	3.5
14	255	8.5	200	4.1	110	3.0	—	3.5
15	250	6.5	220	4.0	110	2.8	—	3.6
16	240	8.1	220	3.6	110	2.5	—	3.6
17	225	5.5	—	—	120	2.0	—	3.6
18	220	4.8	—	—	—	—	2.3	3.6
19	< 220	4.2	—	—	—	—	2.5	3.3
20	< 230	3.4	—	—	—	—	2.4	3.8
21	< 230	3.1	—	—	—	—	2.2	3.0
22	250	3.1	—	—	—	—	2.3	2.9
23	< 280	3.2	—	—	—	—	2.0	2.9

Time: 0.0°.

Sweep: 1.6 Mc to 16.0 Mc in 1 minute 15 seconds.

Table 48

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	< 250	3.2						3.0
01	< 260	3.2						3.0
02	—	3.1						3.1
03	250	3.1						3.2
04	240	3.0						3.2
05	< 220	2.6						3.5
06	< 250	2.2						3.2
07	< 260	2.4						3.2
08	225	4.8	—	—	—	—	—	3.8
09	230	5.7	210	—	120	2.3	—	3.8
10	250	7.2	210	4.1	100	2.7	—	3.5
11	250	8.1	210	4.3	110	2.9	—	3.6
12	250	6.7	200	4.3	105	3.1	—	3.6
13	250	6.6	200	4.3	105	3.0	—	3.5
14	255	8.5	200	4.1	110	3.0	—	3.5
15	250	6.5	220	4.0	110	2.8	—	3.6
16	240	8.1	220	3.6	110	2.5	—	3.6
17	225	5.5	—	—	120	2.0	—	3.6
18	220	4.8	—	—	—	—	2.3	3.6
19	< 220	4.2	—	—	—	—		

TABLE 49
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

Observed at **Washington, D. C.**
(Characteristic) **h'F₂** . Km **490** . April, 1954
(Unit) (Month)

Lat. **38.7°N** Long. **77.0°W**

Day	00	75°W Mean Time																							
		01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	340 ^F	S	S	S	(280) ^S	(310) ^S	260	270	310	340	350	370	330	300	310	290	280	260	230	(260) ^A	260	270	270	270	
2	260	270	260	250	240 ^N	(300) ^S	240	260	280	350	450 ^K	500 ^K	430 ^K	420 ^K	6 ^K	H/200 ^K	380 ^K	380 ^K	300 ^K	270 ^K	240 ^K	220	(260) ^S		
3	S	A	S	S	S	S	S	S	S	390 ^K	310 ^K	360 ^K	330 ^K	330 ^K	320 ^K	320 ^K	320 ^K	320 ^K	310 ^K	310 ^K	310 ^K	310 ^K	310 ^K		
4	S ^K	S ^K	S ^K	S ^K	S ^K	S ^K	S ^K	S ^K	S ^K	250	G	G	470	500	480	400	380	380	350	280	250	250	250	250	
5	S	S	S	S	S	S	S	S	S	(260) ^S	420	480	370	460	400	340	370	330	330	310	270	240	250	270	
6	S	S	S	S	S	S	S	S	S	250	250	270	310	320	320 ^H	350	350	330	330	310	270	250	240	240	
7	(280) ^S	(280) ^S	(280) ^S	(280) ^S	(280) ^S	(280) ^S	(280) ^S	(280) ^S	(280) ^S	240	330	400	310	320	370	410	370	380	330	310	270	250	270	(280) ^S	
8	300	280	280 ^S	260	280	290	270	G	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
9	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
10	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
11	280 ^S	280	270	280	270	270	270	270	270	270	270	270	260	C	C	C	C	C	C	C	C	C	C	S ^K	
12	S ^K	S ^K	(310) ^S	320 ^K	S ^K	S ^K	(280) ^S	270 ^K	270 ^K	G ^K	6 ^K	6 ^K	6 ^K	6 ^K	6 ^K	6 ^K	6 ^K	6 ^K	460 ^K	320 ^K	250 ^K	300 ^K	320 ^K		
13	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	5260 ^K	6 ^K	6 ^K	5860 ^K	360	360	370	370	370	320	330	290	260	250	280 ^S	
14	290	290	280 ^S	280	310 ^S	(320) ^S	270	280	300	310	310	320	320	350	310	340	340	300	310	300	260	250	240	240	270
15	280	280	270	280	(300) ^S	(280) ^L	(280) ^L	(280) ^L	(280) ^L	340	310	330 ^H	340	320	320	310	310	350	350	320	330	270	240	230 ^A	(300) ^S
16	310	(300) ^S	(290) ^S	(290) ^S	(310) ^S	360	260	(210) ^L	360	300	290	320	320	320	310	290	270	230	230	230	270				
17	(100) ^S	(1340) ^S	(290)	270	290	(230) ^S	260	330	310	340	390	390	390	350	320	310	300	290	290	260	250	230	220	(280) ^S	
18	320	280	(330) ^S	[320] ^L	320	350	410	370	370	340	330	320	300	270	240	230	240	300							
19	(310) ^S	320	300	250	250 ^S	(280) ^S	(280) ^S	(280) ^S	(280) ^S	1 ^K	470 ^K	G ^K	G ^K	G ^K	G ^K	6 ^K	450 ^K	420 ^K	360 ^K	350 ^K	350 ^K	240 ^K	260 ^K	300 ^S	
20	300	300	310	290	270	280	250	[280] ^S	310	300	320	410	310	310	360	320	320	300	280	260	250	230	220	(280) ^S	
21	330	(300) ^S	300	310	290	270	[210] ^L	260	340	400	310	390 ^H	350	320	320	330	300	290	260	240	230	230	260	280	
22	(280) ^S	(280) ^S	(270) ^S	(270) ^S	(270) ^S	(270) ^S	(270) ^S	(270) ^S	(270) ^S	250 ^H	280	300	330	320	340	300	300	300	280	250	230	230	280		
23	280	270	(270) ^S	(310) ^S	(310) ^S	(300) ^S	(300) ^S	(300) ^S	(300) ^S	250	320	390	310	350	340	360	380	370	370	300	280	230	230	290	
24	A	A	A	A	A	A	A	A	A	250 ^K	G ^K	G ^K	440 ^K	G ^K	500 ^K	490 ^K	440 ^K	330 ^K	320 ^K	270 ^K	240 ^K	240 ^K	(280) ^S		
25	340	(320) ^S	320	S	A	270	(280) ^L	L ^K	G ^K	G ^K	500 ^K	460 ^K	490 ^K	400 ^K	380 ^K	370 ^K	360 ^K	360 ^K	310 ^K	260 ^K	(270) ^S	A	S		
26	S	(330) ^S	(310) ^S	(310) ^S	(260) ^S	S	G	350	G	G	G	420	450	440	410	460	360	330	300	300	[280] ^A	[280] ^A	[280] ^A		
27	280 ^S	300	300	300	300	S	S	(290) ^S	L	G	G	470	[420] ^A	380	320	310	320	300	300	[280] ^A	A	250	240	280	
28	290	280	300	300	300	(290) ^S	270	L	G	310	430	390	350	370	430	440 ^S	380	320	320	280	250	250	250	260	
29	260	280	260	270	270	250	(260) ^L	270 ^H	260	300	270	320	320	300	350	330	340	370	390	270	240	220	210	280	
30	(280) ^S	290	290	300	300	250	250	280 ^H	270	400	430 ^H	620	450	420	410	360	300	300	280	240	230	240	260	290	
31																									

Sweep 1.0 Mc to 250 Mc in 0.25 min
Manual Automatic

TABLE 50

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

f₀F₂, Mc (Unit)

— April, 1954

(Month)

D.C.

Observed at Washington, D.C.

Lat. 38.7°N, Long 77.1°W

75° W Mean Time

National Bureau of Standards
(Institution) E. J. W. J. W. P. F. J. M. J. S. —

Scaled by J. W. P. F. J. M. J. S. —

Calculated by J. W. P. F. J. M. J. S. —

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	(1.9) ⁵	(1.4) ⁶	(1.7) ³	(1.4) ⁵	(2.0) ⁵	(2.0) ⁵	3.0	4.3	4.7	4.9	5.1	5.0	5.0	5.3	5.2	5.0	5.0	5.0	5.0	4.2	3.9	3.3	3.1	
2	2.6	2.4	2.3	2.4	2.1	2.0	3.4	4.5	4.8	4.5	4.5	4.5	4.6	4.6	4.6	4.4	4.4	4.5	4.5	4.7	3.2	2.3	2.1	
3	(2.0) ⁵	(1.9) ⁴	(1.8) ⁵	1.6	1.6	1.6	1.6	1.5	2.7	3.3	<3.7	4.4	5.0	5.0	5.5	5.2	5.2	4.4	4.4	4.4	4.5	4.7	4.7	
4	(2.4) ⁵	(1.9) ³	(1.9) ²	1.9	1.7	1.7	1.6	1.5	2.6	3.4	<3.7	4.5	(4.5) ⁴	(4.4) ⁴	4.4	4.5	4.5	4.4	4.4	4.4	4.2	4.2	4.2	
5	1.8	(1.6) ³	(1.6) ²	1.6	1.5	1.5	1.4	1.3	2.0	3.5	3.8	4.2	4.2	4.6	5.0	5.0	5.2	4.9	4.9	4.9	4.2	4.2	4.2	
6	(2.3) ⁵	(2.2) ⁵	(2.2) ³	[2.2] ³	(2.0) ³	(2.2) ³	(2.2) ³	(2.2) ³	5.1	5.0	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	4.5	4.5	4.5	4.5	
7	(2.6) ⁵	2.6	2.9	(2.7) ⁴	2.5	2.5	3.2	(4.0) ³	4.0	4.9	4.6	5.0	(4.8) ⁹	(4.5) ⁹	4.7	4.9	4.7	4.8	4.8	4.8	4.2	4.2	4.2	4.2
8	2.5	2.6	2.4	(2.0) ⁵	(2.3) ⁵	3.2	<3.6	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
9	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
10	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
11	2.8	(2.5) ³	(2.4) ⁵	(2.4) ⁵	(2.4) ⁴	(2.4) ⁴	(2.4) ⁴	(2.4) ⁴	3.9	5.1	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	
12	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	
13	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	
14	3.1	2.7	2.5	2.5	2.5	2.5	(1.8) ²	(1.8) ²	3.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	
15	3.0	3.0	2.9	2.6	2.6	2.6	(2.3) ²	(2.3) ²	3.3	3.3	3.8	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	
16	2.5	2.5	2.4	1.8	1.9	1.9	1.9	1.9	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	
17	2.2	(2.0) ⁵	2.1	2.3	(2.7) ⁵	3.5	(2.7) ⁵	3.5	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	
18	2.3	(2.2) ⁵	1.9	1.5	2.0	2.2	3.2	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	
19	1.9	2.0	2.0	(1.9) ⁵	(1.8) ⁵	(1.8) ⁵	(1.8) ⁵	(1.8) ⁵	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	
20	2.4	2.2	2.0	2.0	2.0	2.0	2.0	2.0	2.6	3.6	4.1	4.7	5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2	
21	2.1	2.1	2.1	(2.1) ⁵	2.1	2.1	2.1	2.1	3.1	4.0	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	
22	2.6	2.6	2.3	2	1.9	1.9	1.9	1.9	3.8	4.3	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	
23	3.0	2.8	2.3	2.3	(1.8) ⁵	(1.8) ⁵	(1.8) ⁵	(1.8) ⁵	2.1	3.3	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	
24	A	A	A	A	A	A	A	A	3.2	<3.4	<3.6	<3.6	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	
25	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
26	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	1.97	
27	2.9	2.9	2.3	2.1	1.7	(1.5) ⁵	(1.5) ⁵	2.0	3.1	-3.2	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	
28	(2.2) ⁵	(2.2) ⁵	2.1	2.1	2.1	(2.0) ⁵	(2.0) ⁵	2.1	3.2	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	<3.4	
29	(2.6) ⁵	2.6	2.3	2.1	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	
30	2.7	2.4	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
31																								
Median	2.2	2.2	2.1	2.1	2.2	3.2	3.8	4.4	4.6	4.8	4.4	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	
Count	2.6	2.6	2.8	2.7	2.5	2.6	2.8	2.8	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	

Sweep 1.0 Mc to 2.0 Mc in 0.25 min
Manual □ Automatic ☒

TABLE 51
IONOSPHERIC DATA

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

f_{OF2} Mc April, 1954

(Characteristic) (Unit)

Observed at Washington, D.C.

Lat 38.7°N, Long 77.1°W

National Bureau of Standards
Institution E. J. W., J. W. P., F. J. M., J. J. S.
Scaled by J. W. P. Calculated by F. J. M., J. J. S.

Day	75°W Mean Time											
	0030	0130	0230	0330	0430	0530	0630	0730	0830	0930	1030	1130
1	1.9 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵
2	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
3	2.5	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵	1.7 ⁵
4	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵	1.4 ⁵
5	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵
6	(2.3) ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵	2.3 ⁵
7	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1
8	2.6	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
9	2	2	2	2	2	2	2	2	2	2	2	2
0	1	1	1	1	1	1	1	1	1	1	1	1
11	1.7 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵	1.2 ⁵
12	(2.7) ⁵	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
13	1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵	<1.0 ⁵
14	3.0 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵	2.7 ⁵
15	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵	2.9 ⁵
16	-	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3
17	-	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1
18	0.3	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
19	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵	1.9 ⁵
20	3.5	2.0 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵
21	2	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1
22	0.5 ⁵	-	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3
23	2.5 ⁵	-	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1
24	(2.9) ⁵	(1.7) ⁵	A	A	2.7	3.3 ⁵						
25	(2.1) ⁵	(1.97) ⁵	1.8	A	2.7	3.4 ⁵	3.38 ⁵	3.38 ⁵	3.38 ⁵	3.38 ⁵	3.38 ⁵	3.38 ⁵
26	1.9 ⁵	1.9 ⁵	1.6 ⁵	1.6 ⁵	2.7	<3.4 ⁵	<3.5 ⁵	<3.7 ⁵	<3.9 ⁵	4.2	4.3 ⁵	4.4 ⁵
27	2.6	1.9 ⁵	1.6 ⁵	1.6 ⁵	1.6 ⁵	<3.4 ⁵	<3.6 ⁵	<3.8 ⁵	4.8	5.2	5.6	5.7
28	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.4 ⁵	2.0 ⁵	2.0 ⁵	2.0 ⁵	4.6	4.5	4.6	4.6
29	2.5 ⁵	2.4	2.4	2.4	2.4	2.0 ⁵	2.0 ⁵	2.0 ⁵	4.7 ⁵	5.1	5.3	5.7
30	2.4	2.4	2.4	2.4	2.4	2.3	2.3	2.3	4.6	4.7	4.8	4.9
31	-	-	-	-	-	-	-	-	-	-	-	-
Median	2.3	2.1	2.0	2.0	2.1	2.1	2.1	2.1	4.6	4.8	5.0	5.2
Count	28	27	26	24	24	28	26	26	26	26	26	27

Sweep 1.0 Mc to 25.0 Mc in 0.55 min
Manual □ Automatic □

TABLE 52

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

National Bureau of Standards
(Institution) J.W.P., F.J.M., J.J.S.

Scaled by, E.J.W., J.W.P., F.J.M., J.J.S.

Calculated by, J.W.P., F.J.M., J.J.S.

$h'F_1$, KM April, 1954
 (Characteristic) (Month)
 Observed at Washington, D.C.
 Lat 38.7°N, Long 77.1°W

75°W Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23						
1								240	210 ^H	180 ^H	200 ^H	210 ^H	200 ^H	220 ^H	220 ^H	240 ^H	240 ^H	240 ^H	240 ^H	240 ^H	240 ^H	240 ^H								
2								240	240	240	230 ^K	230 ^K	240 ^K	250 ^K																
3								210 ^K	230 ^K	220 ^K	220 ^K	200 ^K	200 ^K	210 ^K	230 ^K	240 ^K	230 ^K													
4								Q	240	210	210	210	210	190 ^H	200 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H					
5								240	230	230	220	210 ^H	210 ^H	190 ^H	200 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H	210 ^H					
6								210	240	210 ^H	200 ^H	190 ^H	190 ^H	190 ^H	200 ^H	210 ^H	220 ^H	230 ^H												
7								240	210	210	210	210	210	190 ^H	200 ^H	200 ^H	210 ^H													
8								240	230	230	C	C	C	C	C	C	C	C	C	C	C	C	C	C						
9								C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C						
10								C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C						
11								Q	230	220 ^H	C	C	C	C	C	C	C	C	C	C	C	C	C	C						
12								Q	K	(240) ^S	230 ^H	230 ^K	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	200 ^H	
13								Q	K	230 ^K	220 ^K	210 ^K	190 ^K	190 ^K	240 ^H	240 ^H	210 ^H													
14								250	230 ^H	220	210	200 ^H	200 ^H	200 ^H	200 ^H	210	190 ^H	190 ^H	210 ^H											
15								230	230	210	210	200 ^H	200 ^H	210	210	200	200	210 ^H												
16								240	220	200	200 ^H	200 ^H	220	220	210 ^H	210 ^H	200	200	210 ^H											
17								240	220	210	200 ^H	200 ^H	220	220	200	200	210 ^H													
18								230	210	210	210	200 ^H	200 ^H	200	200	200	200	210 ^H												
19								240 ^K	220 ^K	210 ^K	200 ^K	190 ^K	190 ^K	200 ^K	200 ^K	210 ^K	200 ^K													
20								240	220 ^H	230 ^H	210 ^H	200 ^H	200 ^H	190 ^H	180 ^H	200	230 ^H	220 ^H												
21								230	220	210 ^H	190 ^H	180 ^H	180 ^H	180 ^H	(200) ^H	210 ^H														
22								230	220	210 ^H	210 ^H	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200				
23								230 ^H	220	210	210	200	200	200	200	210	210 ^H													
24								A	K	A	210 ^K	200 ^K	200 ^K	200 ^K	210 ^K	200 ^K	210 ^K	200 ^K	200 ^K	190 ^K	190 ^K	190 ^K	200 ^K							
25								210 ^K	200 ^H	200 ^K	200 ^K	200 ^K	230 ^K	200 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	210 ^K	A				
26								230	210 ^H	200 ^H	180 ^H	210 ^H	210 ^H	210 ^H	200 ^H	240 ^H	220 ^H													
27								230	210	200 ^H	190 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	A			
28								220	220	200	180 ^H	210	300 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H		
29								210 ^H	210	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H	180 ^H				
30								210 ^H	200	210	180 ^H	200	250 ^H	200	250 ^H	210 ^H	230 ^H													
31																														
Median	230	220	210	210	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200	200		
Count	17	26	28	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	

Sweep 10 Mc 1023.0 Mc 100.5 min

Manual □ Automatic ☒

TABLE 53
IONOSPHERIC DATA

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

f_{oF1} — Mc — April, 1954

(Characteristic) (Unit) (Month)

Observed at Washington, D.C.

Lat. 38.7°N, Long 77.1°W

National Bureau of Standards
Scaled by E. J. W., J. W. P. (Institution)
Calculated by J. W. P., F. J. M., J. J. S.

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	L	L	L	L	L	L	L	L	L	L	L	L
2	L	3.6	3.9	3.9	3.9	3.9	3.9	3.9	4.0	4.0	4.0	4.0
3	L	3.7	3.9	3.9	4.0	4.0	4.0	4.0	4.1	4.1	4.0	4.0
4	①	3.5	3.7	3.7	3.8	3.8	4.0	4.0	4.0	4.0	4.0	4.0
5	3.2	3.5	3.7	3.7	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
6	L	3.9	3.9	4.0	4.0	4.0	4.0	4.0	4.1	4.1	4.1	4.1
7	3.3	3.6	3.9	3.9	4.1	4.2	4.2	4.2	4.2	4.2	4.2	4.2
8	L	3.6	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7
9	C	C	C	C	C	C	C	C	C	C	C	C
10	C	C	C	C	C	C	C	C	C	C	C	C
11	C	C	C	C	C	C	C	C	C	C	C	C
12	②	3.8	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
13	Φ	2.8	3.4	3.7	3.7	3.7	3.9	3.9	3.9	3.9	3.9	3.9
14	Φ	3.3	3.5	3.8	3.8	3.8	3.9	3.9	3.9	3.9	3.9	3.9
15	L	3.4	3.7	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
16	L	3.7	4.0	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
17	L	3.6	3.8	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9
18	L	3.4	3.7	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9
19	L	3.4	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6
20	L	3.6	3.6	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
21	L	3.8	3.8	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7
22	L	3.6	3.6	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
23	L	3.5	3.6	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
24	A	3.4	3.6	3.7	3.7	3.9	4.0	4.0	4.0	4.0	4.0	4.0
25	L	3.3	3.6	3.6	3.8	4.0	4.0	4.0	4.0	4.0	4.0	4.0
26	3.4	3.6	3.6	3.8	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
27	L	3.2	3.5	3.7	3.7	3.9	3.9	3.9	3.9	3.9	3.9	3.9
28	L	3.4	3.5	3.8	4.0	4.1	4.1	4.1	4.1	4.1	4.1	4.1
29	L	3.5	3.8	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
30	L	3.2	3.4	3.9	3.9	4.0	4.0	4.0	4.0	4.0	4.0	4.0
31	—	3.4	3.7	3.9	4.0	4.1	4.2	4.2	4.2	4.2	4.2	4.2
Median	1	3.4	3.7	3.9	4.0	4.1	4.2	4.2	4.2	4.2	4.2	4.2
Count	1	17	26	26	26	26	26	26	26	26	26	26

Sweep 10 — Mc 10.25 Q. Mc in 0.25 min
Manual □ Automatic ☑

TABLE 54

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

(Characteristic)	Day	Washington, D.C.												38.7°N, Long 77.1°W													
		h'E	Km	April (Month)	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22
1																											
2																											
3																											
4																											
5																											
6																											
7																											
B																											
9																											
10																											
11																											
12																											
13																											
14																											
15																											
16																											
17																											
18																											
19																											
20																											
21																											
22																											
23																											
24																											
25																											
26																											
27																											
28																											
29																											
30																											
31																											

National Bureau of Standards
Institution J.W.P., F.J.M., J.J.S.
Calculated by E.J.W.

TABLE 55
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

Day	75°W Mean Time												75°W Mean Time																		
	Observed at Washington, D. C.				Lat 38.7°N, Long 77.1°W				Calculated by J.W.P.				Calculated by F.J.M., J.J.S.				Calculated by J.W.P.				Calculated by F.J.M., J.J.S.										
foE	Mc	Mc	Mc	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23				
1	1.9	2.3	2.6	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a	3.0 ^a						
2	(1.9) ^a	2.4 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a	2.5 ^a						
3	1.8	2.5	2.5	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8					
4	2.1	2.3	2.5 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a	2.6 ^a						
5	A	2.5	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a						
6	A	2.5	2.7 ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a						
7	S	2.0	2.4	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
8	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
9	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
10	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S				
11	(1.8) ^a	2.0 ^a	2.4 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a	2.8 ^a					
12	B ^a	2.0 ^a	2.4 ^a	2.5 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a	2.7 ^a					
13	1.7	2.0	2.4	2.4	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7				
14	1.7	2.0	2.4	2.4	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7				
15	(1.6) ^a	2.2 ^a	2.5 ^a	2.8	2.8	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9				
16	(1.6) ^a	(2.2) ^a	(2.6) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a	(2.8) ^a					
17	S	(2.0) ^a	2.3 ^a	2.8	2.9 ^a	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1				
18	1.7	2.1	2.5	2.6	2.6	2.7	3.1	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2				
19	S	2.1 ^a	2.4 ^a	2.8 ^a	2.8 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a	2.9 ^a				
20	S	2.1	2.5	2.7	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A				
21	1.6	1.9	2.3	(2.8) ^a	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A				
22	1.8	2.2	2.6	2.8	3.0	3.0	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2			
23	1.7	2.1	2.4	2.8	2.8	2.8	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0			
24	S ^a	2.1 ^a	2.4	2.7	(2.7) ^a	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A			
25	S ^a	2.0 ^a	2.5 ^a	(2.7) ^a	(2.7) ^a	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A			
26	1.8	(2.2) ^a	2.5 ^a	2.8	(2.7) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a				
27	1.7	2.1	2.4	2.7	(2.8) ^a	2.8	(2.7) ^a	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
28	1.7	2.2	2.5	2.7	2.9	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a	(3.0) ^a			
29	1.6	2.2	2.5	2.8	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
30	1.7	2.3	2.7	3.0	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
31																															
Median Count	1.7	2.1	2.4	2.7	3.1	3.0	3.1	3.1	3.0	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9		
	1.9	2.5	2.7	2.6	2.1	2.0	2.5	2.5	2.4	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	

Sweep 1.0—Mc to 25.0 Mc in 0.25 min
Manual □ Automatic ■

TABLE 56
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

Mc., Km (Unit) April, 1954 (Month)

Observed at Washington, D.C.

Lat. 38° 7' N., Long. 77° 18' W.

National Bureau of Standards
(Institution)
Scaled by: E.J.W. J.W.P. F.J.M. J.J.S.

Day	Calculated by: J.W.P.												Calculated by: J.W.P.											
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	E	25 110	E	35 110	E	20 130	35 120	G	G	38 130	34 120	35 120	32 130	30 130	26 130	28 120	35 120	24 120	28 120	33 130	33 130	E	E	
2	E	E	E	E	E	E	E	E	G	G	G	G	35 130	38 130	45 130	45 130	G	G	G	G	E	E	E	E
3	E	44N 120	42N 120	42N 120	24 130	E	E	E	G	G	G	G	G	G	G	G	47 130	56 120	47 110	32 110	47 110	E	E	E
4	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	18 130	E	E	E	
5	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	21 130	18 120	26 120	25 120	
6	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
7	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
8	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
9	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
10	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
11	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	B	B	E	E	
12	S	S	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	S	S	
13	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
14	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
15	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
16	E	27 110	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
17	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
18	E	24 110	30 110	34 110	46 110	E	E	E	G	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E
19	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
20	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
21	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
22	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
23	E	23 100	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
24	E	35 110	30 110	40 110	40 110	32 120	32 120	32 120	32 120	37 110	40 110	38 110	36 110	36 110	42 120	50 110	43 120	50 110	42 120	50 110	E	E	19 120	
25	E	22 120	39 120	31 110	44 110	17 110	20 110	35 110	37 110	33 110	34 110	31 110	G	34 110	36 110	35 110	36 110	37 110	38 110	38 110	38 110	38 110	E	E
26	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
27	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
28	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
29	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
30	E	E	E	E	E	E	E	E	G	G	G	G	G	G	G	G	G	G	G	E	E	E	E	
31																								
Median	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	
Count	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27	27

* * MEDIAN FEWER THAN MEDIAN FOR, OR LESS
THAN LOWER FREQUENCY LIMIT OF RECORDER

Manual □ Automatic ◻

Sweep 10 Mc/min 25 Mc/min

TABLE 57
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
ONOSPHERIC DATA

(M) 1500) F2, (Mn) 1954
April (Month)

Observed at Washington, D.C.

Lat. 38.7°N, Long. 77.1°W

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

National Bureau of Standards
Institution of Standards

Scaled by E.J.W. J.W.P. F.J.M. J.J.S.

Day	75°W Mean Time												75°W Mean Time														
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23			
1	(2.1) S (2.1) S	S	S	S	(2.1) F (2.1) S	2.3 F	2.4	2.3	2.2	2.1	2.0	2.1	2.2	2.3	2.1	2.1	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0			
2	2.1	2.2	2.2	2.2	2.1	2.1	2.4	2.3	2.1	1.9 K	1.7 K	1.7 K	1.9 K	G K	2.0 K	2.0 K	2.0 K	2.0 K	2.3	2.2	2.3	2.2	2.2	2.0			
3	2.1	A	S	J	J	J	J	J	K	2.3 K	2.3 K	2.0 K	2.0 K	2.1 K	2.1 K	2.0 K	2.0 K	2.2 K	A K	(2.2) S	(2.1) S	(2.2) S	(2.1) K	(1.9) F			
4	(2.0) S (2.0) S (1.9) K	S K	S K	S K	2.0 K	J	J	J	J	2.3	2.3	6	G	G	(1.7) H	(1.7) H	1.7	2.0	2.0	2.0	2.0	2.1	2.1	2.0	2.0		
5	2.0	S	S	S	S	J	S	J	S	(2.1) S	1.9	1.8	2.1	1.8	2.0	2.1	2.0	2.2	2.2	2.2	2.2	2.2	2.1	2.1	F (2.0) S		
6	(2.0) F (2.1) S	S	S	J	S	J	S	J	S	(2.3) F (2.1) H	2.5	2.2	2.2	2.2	2.1	2.1	2.2	2.3	2.3	2.2	2.2	2.1	2.1	F (2.0) S	2.0		
7	(2.1) F	2.1	2.0	(2.1) S	(2.0) F	(2.1) S	2.0 F	2.4	(2.2) S	1.9	2.3 H	2.3 S	(2.1) P	1.9	1.9	2.0	2.0	2.1	2.2	2.2	2.2	2.2	2.1	2.1	2.1	2.0	
8	2.1	2.1 F	2.2 F	(2.2) S	(2.1) F	2.4	G	2.1	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
9	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
10	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
11	2.0	(2.1) S	J	S	(2.1) S	(2.1) S	(2.1) S	(2.1) S	(2.1) S	(2.2) F	2.4	2.3	2.4	C	C	C	C	C	C	C	C	C	C	C	C	C	
12	S K	S K	J	S K	J	S K	J	S K	S K	(2.2) F	2.3 F	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	S K		
13	E K	E K	E K	E K	E K	E K	E K	E K	E K	2.2 K	G K	1.6 K	G K	G K	2.0	2.0	2.0	2.0	2.2	2.1	2.1	2.1	2.0	2.0	F E K	E K	
14	1.9 F	2.0 F	2.0 F	2.0 F	2.0 F	(2.0) F	(2.1) F	2.3	2.3	2.2	2.2	2.2	2.2	2.1	2.1	2.2	2.2	2.2	2.3	2.2	2.2	2.2	2.1	2.1	2.0	2.0	
15	2.1 F	2.0	2.1 F	(2.1) S	(2.1) F	(2.1) F	(2.1) F	2.3	2.3	2.1	2.3	2.1 H	2.0	2.1	2.2	2.2	2.2	2.1	2.1	2.2	2.2	2.2	2.2	2.2	(2.3) F	2.2	
16	1.9	2.0	2.0	2.0	1.9 F	2.2	2.2	2.2	2.2	2.1 H	(2.4) S	2.2 H	2.1 H	2.3	2.2	2.1	2.1	2.2	2.1	(2.3) S	2.2	2.2	2.2	2.1	2.1	2.1	
17	1.9	(1.8) S	2.0	2.1	2.1	(2.2) S	2.3	2.3	2.2	2.3	2.1	2.0 H	1.9	1.9	1.9	1.9	1.9	2.1	2.1	2.3	2.2	2.2	2.1	2.1	2.0	2.0	
18	1.9	(2.1) S	2.0	2.0	2.0	2.1	2.2 H	2.2	2.2	(2.0) S	2.2	2.0	1.9	2.0	2.0	2.0	2.0	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.0		
19	1.9	2.0 S	2.0	2.1 S	2.2 S	2.1	2.2 H	1.8 K	G K	G K	G K	G K	G K	1.9 K	1.9 K	2.0 K	2.0 K	2.2 K	2.1 H	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
20	2.1	2.1 F	(2.1) F	(2.1) F	2.2 F	2.1 F	2.2	2.4	2.2	2.2	2.2	2.1	2.2	2.1	2.0	2.1	2.2	2.2	2.2	2.2	2.2	2.3	2.3	2.3	2.3	2.3	
21	1.9	2.0	2.1	2.1	(2.1) S	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	S	(2.2) H	1.9	2.2	2.1	2.1	2.3	2.2	2.2	2.2	2.2	2.2	2.1	2.1
22	2.0	2.0	2.1	2.1	(2.1) S	(2.2) S	2.3 H	2.3	2.3 S	(2.0) S	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.3	2.2	2.2	2.2	2.2	2.2	2.2		
23	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1		
24	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
25	(1.9) S	2.1 S	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
26	S	(2.0) S	(2.1) S	(2.2) S	(2.0) S	(2.1) S	(2.2) S	2.3 H	2.3	2.3 S	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
27	2.0	2.0	2.0	2.0	2.0	J	2.1	2.4	G	G	G	G	G	1.8 S	2.0	2.0	2.1	2.1	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
28	(2.2) S	(2.2) S	(2.0) F	(2.0) F	(2.1) F	2.3 F	2.3	G	2.3	1.9 S	2.0	2.1	1.9	1.8	2.0 S	2.2	2.2	2.1	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
29	(2.2) F	2.1	2.2 F	(2.2) S	(2.3) S	2.3 F	2.3	2.4 H	2.4	2.3	2.3	2.3	2.3	2.3	2.3 H	2.0 K	2.0 K	2.1	2.1	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
30	2.0	2.0 S	(2.1) S	(2.1) S	(2.0) S	2.3 H	2.3	2.3	(2.2) S	2.3 H	2.2	1.9 H	1.9 H	1.9 H	1.8 S	(1.8) S	1.9	1.9	2.0	2.0	2.1	2.1	2.1	2.1	2.1	2.1	
31																											
Median	2.0	2.0	2.1	2.1	2.1	(2.1)	2.2	2.3	2.2	2.1	2.2	2.0	2.0	2.0	2.0	2.0	2.0	2.1	2.1	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
Count	24	23	19	22	19	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	

Sweep 10 Mc 10.255.0 Min 0.28 min

Manual □ Automatic ■

TABLE 58

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

(M3000)F2, April, 1954

(Characteristic Unit)

Observed at Washington, D.C.

IONOSPHERIC DATA

National Bureau of Standards

Scaled by E.J.W.

(Institution)

J.W.P., F.J.M., J.J.S.

Calculated by J.W.P.

F.J.M., J.J.S.

Lat 38°7'N. Long 77°1'W

Mean Time

75°W

75°W

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	(3.1) ^F	(3.1) ^F	J ^s	J ^s	J ^s	(3.2) ^F	3.3 ^F	3.5 ^F	3.4 ^F	3.3	3.1	3.0	3.3	3.3	3.4	3.3	3.3	3.1	3.2	3.4	3.0	3.0	3.0	
2	3.2	3.3	3.2	3.3	3.2	3.3	3.1	3.5 ^H	3.5 ^H	3.4	3.1	2.8 ^K	2.6 ^K	2.6 ^H	2.8 ^K	G ^K	3.0 ^K	2.9 ^K	3.0 ^K	3.2 ^K	3.3	3.2	3.0	3.0
3	(3.1) ^S	A	S	J ^s	G ^K	3.0 ^K	3.2 ^K	3.1 ^K	3.2 ^K	3.0 ^K	3.0 ^K	3.2 ^K	A ^K	(3.2) ^F	(3.0) ^P	(3.0) ^K	(3.0) ^F	(3.0) ^P						
4	(2.9) ^F	(2.7) ^S	J ^s	J ^s	J ^s	J ^s	J ^s	J ^s	J ^s	G	G	(2.6) ^H	(2.6) ^H	2.6	3.0	3.0	3.0	3.2	3.2	3.0	3.1	3.0	3.0	3.1
5	3.0	J ^s	3.1	2.7	3.0	3.2	3.0	3.2	3.1	3.2	3.2	3.3	3.2	3.1	3.2	3.0								
6	(3.0) ^F	(3.1) ^F	J ^s	(3.3) ^F	(3.1) ^H	3.6	3.3	3.3	3.2 ^H	3.1	3.0	3.1	3.2	3.3	3.3	3.1	(3.0) ^F							
7	(3.1) ^F	3.1	3.0	(3.0) ^F	(3.0) ^F	3.0 ^F	3.0 ^F	3.4	(3.2) ^S	2.9	3.3 ^H	3.4 ^S	(3.1) ^P	(2.9) ^P	2.9	3.0	3.0	3.1	3.3	3.2	3.3	3.2	C	3.0
8	3.2	3.1	3.1	3.1	3.2	3.2	3.2	(3.1) ^F	(3.1) ^F	J ^s	C	C	C	C	C	C	C	C	C	C	C	C	C	
9	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
10	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
11	3.0	(3.1) ^S	J ^s	J ^s	(3.2) ^S	(3.2) ^F	(3.2) ^F	(3.1) ^F	(3.1) ^F	S ^K	S ^K	(3.2) ^F	3.3 ^F	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	S ^K	
12	S ^K	S ^K	J ^s	J ^s	3.0 ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	3.2 ^K	2.4 ^K	2.4 ^K	3.0	3.0	3.0	3.2	3.1	3.2	3.1	3.1	3.0	E ^K
13	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K	E ^K															
14	2.9	F	3.0	F	3.0	F	3.0	F	3.0	F	3.0	F	3.0	F										
15	3.1	F	3.0	F	3.1	F	3.1	F	3.1	F	3.3	F	3.2	F	3.3	F	3.0	F	3.1	F	3.2	F	3.3	F
16	2.9	3.0	3.0	2.9	3.1	3.1	3.1	(3.3) ^S	(3.3) ^S	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	
17	2.9	(2.9) ^S	2.9	3.1	3.1	3.1	3.1	(3.3) ^S	(3.3) ^S	3.2	3.0 ^H	2.9	2.9	2.9	2.9	3.1	3.1	3.2	3.2	3.2	3.2	3.2	3.2	
18	2.9	(3.1) ^S	2.9	3.0	3.0	2.9	3.1	3.2	3.2	3.3	3.0	2.9	3.0	3.0	3.0	3.1	3.1	3.2	3.2	3.2	3.2	3.2	3.2	
19	2.9	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	
20	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	
21	2.9	3.0	3.0	3.1	(3.1) ^S	3.1	3.1	3.2	3.2	T ^S	(3.3) ^P	2.9	3.3	2.9 ^H	3.1	3.2	3.2	3.2	3.3	3.2	3.2	3.2	3.2	
22	3.0	3.0	3.0	3.2	3.1	3.1	(3.1) ^S	(3.3) ^P	3.3 ^H	3.4	3.4 ^H	3.2	3.2	3.2	3.0	3.3	3.2	3.3	3.2	3.4	3.2	3.2	3.0	
23	3.0	3.0	3.0	3.1	(3.2) ^S	3.4	3.1 ^H	3.0	3.3	3.1	3.0	2.9	2.9	3.1	3.1	3.2	3.4	3.3	3.2					
24	A	A	A	A	A	A	A	A	A	3.3 ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K	G ^K		
25	(2.9) ^S	3.1 ^S	3.1	A ^S	A ^S	A	3.3	3.3 ^K	3.2 ^K	G ^K	G ^K	2.7 ^K	2.8 ^K	2.7 ^K	3.0 ^K	3.0 ^K	3.0 ^K	3.2 ^K	3.2 ^K	3.2 ^K	3.1 ^K	3.0 ^K	(3.1) ^S	
26	S	(2.9) ^S	(3.1) ^S	(3.2) ^S	(3.0) ^S	(3.0) ^S	S	G	3.2	G	G	G	2.9	2.8	2.8	2.9	2.9	2.9	3.0 ^S	(3.0) ^S	(3.0) ^S	A	S	
27	3.0	3.0	3.0	3.0	3.0	3.0	3.0	J ^s	3.1	3.5	G	G	2.7 ^S	3.0	3.0	3.0	3.0	3.0	3.0	3.1	3.1	3.1	3.1	3.1 ^F
28	(3.2) ^F	(3.2) ^F	3.0 ^F	3.0 ^F	(3.1) ^F	(3.1) ^F	(3.1) ^F	3.3 ^F	3.3 ^F	3.4 ^F	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H		
29	(3.2) ^F	3.1 ^F	3.2 ^F	(3.3) ^F	(3.3) ^F	(3.3) ^F	3.3 ^F	3.3 ^F	3.4 ^H	3.5	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H	3.4 ^H		
30	3.0	3.0	3.0	(3.1) ^S	(3.2) ^S	(3.2) ^S	3.0 ^S	3.3 ^H	3.4 ^H	3.4 ^H	(3.2) ^S	3.2	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	2.8 ^H	
31																								
Median	3.0	3.1	3.1	(3.1)	3.2	3.3	3.2	3.1	3.2	3.2	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	
Count	24	23	19	22	19	20	21	22	23	24	26	26	26	26	26	26	26	26	26	26	26	26	26	25

Sweep LO Mc to 2500 Mc in 0.25 min

Manual □ Automatic ■

TABLE 59
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA
Lat. 38.7°N., Long. 77.1°W.

(Month) April 1954
(Year) 1954
Observed at Washington, D.C.

(Characteristic) F1, (Intra)

National Bureau of Standards
Scal'd by: E.J.W. J.W.P. [Institution] F.J.M. J.J.S.

Calculated by: J.W.P. E.J.W. F.J.M. J.J.S.

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	L	L	L	L	L	L	L	L	3.7 ⁴	3.8 ⁴	3.7 ⁴	3.7 ⁴
2	L	3.6	3.6	3.7 ⁴	3.6 ⁴	3.6 ⁴	3.7	3.7				
3	L	3.4 ⁴	3.6 ⁴	3.7 ⁴	3.7 ⁴	3.7 ⁴	3.7 ⁴	3.7 ⁴	3.5 ⁴	3.5 ⁴	3.4 ⁴	L
4	Q	3.5 ⁴	3.7	3.8 ⁴	3.7	3.7	3.6	3.4 ⁴				
5	3.4	3.8	3.7	3.8	3.8	3.9	3.8 ⁴	3.8 ⁴	3.7	3.7	3.6 ⁴	3.6
6	L	L	3.9 ⁴	3.9 ⁴	4.0 ⁴	4.0 ⁴	4.0 ⁴	4.0 ⁴	3.8 ⁴	3.8 ⁴	3.6 ⁴	L
7	3.5 ⁴	3.9	3.6 ⁴	3.7	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.7 ⁴	3.6	3.5 ⁴	B
8	L	3.4 ⁴	3.6	C	C	C	C	C	C	C	C	C
9	C	C	C	C	C	C	C	C	C	C	C	C
10	C	C	C	C	C	C	C	C	3.6 ⁴	(3.7) ³	3.6 ⁴	L
11	3.8	L	3.7 ⁴	C	C	C	C	C	3.4 ⁴	3.4 ⁴	3.6 ⁴	L
12	3.5 ⁴	4 ⁴	3.5 ⁴	3.8 ⁴	3.9 ⁴	3.9 ⁴	3.9 ⁴	3.9 ⁴	3.7 ⁴	3.8 ⁴	3.7 ⁴	3.5 ⁴
13	Q	3.7	3.8 ⁴	3.7 ⁴	4.0 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	(3.9) ³	3.6 ⁴	3.7 ⁴	3.4 ⁴
14	L	3.6 ⁴	3.7	3.7	3.7 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.7	3.7 ⁴	3.5 ⁴
15	L	L	3.7	3.6	3.9 ⁴	3.7	3.7	3.7	3.7	3.7	3.5 ⁴	L
16	L	L	3.6	3.8 ⁴	2 ⁷	2 ⁷	2 ⁷	2 ⁷	3.6 ⁴	3.8	3.7 ⁴	3.6
17	L	3.5 ⁴	3.6	3.8 ⁴	4.0 ⁴	3.9	3.9	3.9	3.8 ⁴	3.8 ⁴	3.7 ⁴	3.6
18	L	3.7	(3.8) ³	3.8 ⁴	3.8 ⁴	3.7 ⁴	3.8 ⁴	3.8 ⁴	3.6 ⁴	3.5 ⁴	3.7	3.6
19	L	3.8 ⁴	3.7 ⁴	3.7 ⁴	3.9 ⁴	3.9 ⁴	4.0 ⁴	4.0 ⁴	3.8 ⁴	3.7 ⁴	3.7 ⁴	3.6
20	L	3.6 ⁴	3.6 ⁴	3.6 ⁴	3.8 ⁴	3.6 ⁴	3.8 ⁴	3.8 ⁴	3.7 ⁴	3.6 ⁴	3.7	L
21	L	L	3.7 ⁴	4.2 ⁴	4.0 ⁴	4.0 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.7 ⁴	3.8 ⁴	L
22	L	L	3.8 ⁴	3.8	3.9	3.9	3.9	3.9	3.8	3.8	3.6 ⁴	L
23	L	3.5 ⁴	3.6	3.7	3.9	3.9	3.9	3.9	3.8	3.7	3.6	L
24	A	3.5 ⁴	3.7 ⁴	3.8 ⁴	3.9 ⁴	3.9 ⁴	3.9 ⁴	4 ²	3.9 ⁴	3.9 ⁴	3.7 ⁴	L
25	L	3.7 ⁴	3.8 ⁴	4.0 ⁴	4.0 ⁴	3.9 ⁴	3.9 ⁴	3.9 ⁴	3.9 ⁴	A	3.6 ⁴	A
26	3.5 ⁴	3.7 ⁴	3.9 ⁴	4.0 ⁴	3.7 ⁴	3.9 ⁴	3.8 ⁴	3.8 ⁴	3.7 ⁴	3.8	3.6	3.5 ⁴
27	L	3.5 ⁵	3.9 ⁴	4.0 ⁴	4 ⁰	A	3.8	3.8 ⁵	3.6	A	3.5 ⁴	A
28	L	3.9	4.0	3.8 ⁴	3.9	3.7 ⁴	4.0	3.8 ⁴	3.8	3.6 ⁴	3.6	L
29	L	3.7 ⁴	3.8	3.6 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.8 ⁴	3.6 ⁴	3.6 ⁴	3.5 ⁴	L
30	L	3.9 ⁴	3.6	3.7	3.8 ⁴	3.7	3.7	3.7	3.7	3.5 ⁴	3.5 ⁴	3.5 ⁴
31	—	3.6	3.7	3.8	3.8	3.8	3.8	3.8	3.7	3.6	3.6	—
Median	—	3.6	3.7	3.8	3.8	3.8	3.8	3.8	3.7	3.6	3.6	—
Count	1	17	24	26	26	25	26	26	27	26	27	2

Sweep 40 Mc to 25.0 Mc. In 0.25 min
Manual □ Automatic □

TABLE 60

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

(M1500) E, April, 1954
 (Characteristic) (Month)
 Observed at Washington, D.C.

IONOSPHERIC DATA

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1												
2												
3												
4												
5												
6												
7												
8												
9												
10												
11												
12												
13												
14												
15												
16												
17												
18												
19												
20												
21												
22												
23												
24												
25												
26												
27												
28												
29												
30												
31												

National Bureau of Standards
 Scaled by: E.J.W. J.W.P. E.J.M. J.J.S.
 Calculated by: J.W.P. E.J.M. J.J.S.

Manual Automatic

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Table 61Ionospheric Storminess at Washington, D. C.April 1954

Day	Ionospheric character*		Principal storms		Geomagnetic character†	
	00-12 GCT	12-24 GCT	Beginning GCT	End GCT	00-12 GCT	12-24 GCT
1	3	2				
2	1	5	1500	----		
3	2	1	----	0100		
			1000	----		
4	3	4	----	1000		
5	4	2				
6	1	2				
7	1	3				
8	1	-				
9	-	-				
10	-	1				
11	0	8	1800‡	----		
12	4	6	----	----		
13	8	3	----	1600		
14	1	1				
15	0	3				
16	2	3				
17	1	1				
18	2	1				
19	2	5	1100	----		
20	2	2	----	0100		
21	2	2				
22	1	3				
23	1	2				
24	2	5	1100	----		
25	2	4	----	0200		
			1100	2200		
26	3	3				
27	2	3				
28	2	2				
29	1	3				
30	2	3				

* Ionosphere character figure (I-figure) for ionospheric storminess at Washington, D. C., during 12-hour period, on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

† Time uncertain, insufficient data.

‡ K-figures unavailable at time of publication.

---- Dashes indicate continuing storm.

Table 62
 Radio Propagation Quality Figures
 (Including Comparisons with Short-Term and Advance Forecasts)

March 1954

Day	North Pacific 9-hourly quality figures			Short-term forecasts issued at:			Whole day quality index	Advance forecasts (J _p - reports) for whole day; issued in advance by:	
	03 to 12	09 to 18	18 to 03	02	09	18			
1	5	5	6	5	5	6	5	(3)	(4)
2	5	5	6	5	5	6	5	(4)	(4)
3	5	5	7	5	5	6	5	5	5
4	5	5	7	6	5	6	6	6	5
5	6	6	7	6	5	6	6	6	6
6	5	6	5	6	5	6	5	6	6
7	5	5	6	5	5	6	5	6	6
8	5	5	6	6	5	6	5	6	6
9	5	5	6	6	5	6	5	5	6
10	5	(4)	6	6	5	6	5	5	6
11	5	5	6	6	6	6	6	5	5
12	5	6	6	5	5	6	6	5	5
13	5	5	6	5	5	6	5	5	5
14	5	(4)	6	5	(4)	5	(4)	(4)	(4)
15	(4)	(3)	7	5	(3)	5	(4)	(4)	(4)
16	5	(4)	6	5	5	6	5	(4)	(4)
17	5	5	6	5	(4)	6	5	(4)	5
18	5	5	6	5	(4)	6	5	(4)	5
19	5	(4)	6	5	5	6	5	(4)	5
20	6	5	6	5	5	6	6	5	(4)
21	5	5	6	5	5	6	5	(4)	(4)
22	6	6	5	5	6	6	6	(4)	5
23	5	(4)	(4)	5	5	6	(4)	5	5
24	5	(4)	(4)	5	(4)	5	(4)	5	5
25	5	5	6	5	(4)	5	5	(4)	(4)
26	7	6	6	5	5	6	6	(4)	(4)
27	6	6	6	6	5	6	6	5	(4)
28	6	5	6	6	5	6	6	5	5
29	7	6	6	6	6	6	7	6	5
30	6	6	6	6	5	6	6	6	6
31	5	5	7	5	(4)	6	5	5	6

Score:

Quiet Periods	P	20	13	20		7	7
	S	9	11	8		17	16
	U	1	0	1		1	1
	F	0	0	0		2	3

Disturbed Periods	P	0	3	0		2	2
	S	1	4	1		2	2
	U	0	0	0		0	0
	F	0	0	1		0	0

Scales:

- Q-scale of Radio Propagation Quality
- (1) - useless
 - (2) - very poor
 - (3) - poor
 - (4) - poor to fair
 - 5 - fair
 - 6 - fair to good
 - 7 - good
 - 8 - very good
 - 9 - excellent

Scoring: (beginning October 1952)

- P - Perfect: forecast quality equal to observed
- S - Satisfactory: (beginning October 1952)
forecast quality one grade different
from observed
- U - Unsatisfactory: forecast quality two or more
grades different from observed when both
forecast and observed were ≥ 5 , or both ≤ 5
- F - Failure: other times when forecast quality
two or more grades different from observed

Symbols:

- X - probable disturbed date

Note: All times are UT (Universal Time or GCT)

Table 63a

Radio Propagation Quality Figures
(Including Comparisons with Short-Term and Advance Forecasts)

March 1954

Day	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day quality index	Advance forecasts (J-reports) for whole day; issued in advance by:			Geomag- netic K_{CH}
	00	06	12	18	00	06	12	18		1-4 days	4-7 days	8-25 days	
	to 06	to 12	to 18	to 24									
1	(4)	(3)	6	6	5	(4)	6	6	(4)	5	(4)	X	2 2
2	5	(4)	6	6	5	(4)	6	6	5	5	5	X	(4) 2
3	5	(4)	6	6	5	(4)	6	6	5	5	6		2 2
4	5	(3)	6	6	5	5	6	6	5	6	6		3 2
5	(4)	(4)	6	6	6	(4)	6	6	5	6	6		3 2
6	5	5	7	6	6	5	6	6	6	6	6		2 2
7	5	5	6	6	6	5	6	6	5	6	6		(4) 3
8	5	5	6	6	5	5	6	6	6	6	6		3 3
9	5	(4)	6	6	6	5	7	6	5	6	6		3 3
10	5	5	7	6	5	5	7	6	6	6	6		2 2
11	5	(4)	6	5	6	5	6	6	5	7	6		(4) 3
12	5	(4)	6	6	5	5	6	6	5	6	7		3 3
13	5	(4)	7	6	5	5	6	6	5	6	6		2 3
14	(4)	(3)	6	5	5	(4)	5	5	(4)	5	6		(4) (4)
15	(4)	(3)	6	6	5	(3)	5	5	(4)	(4)	(4)	X	(5) 3
16	(4)	(3)	6	6	5	(3)	6	6	(4)	(4)	(4)	X	3 3
17	5	5	6	6	5	(4)	7	6	5	(4)	(4)	X	3 (4)
18	(4)	(4)	6	6	5	(3)	6	6	(4)	5	5		(4) 2
19	(4)	(4)	6	6	5	(4)	6	6	(4)	5	5		2 3
20	5	(4)	6	6	6	(4)	6	6	5	5	5		(4) (4)
21	5	(4)	6	6	5	(4)	6	6	5	(3)	(3)	X	3 2
22	5	(4)	6	6	6	(4)	6	6	5	(3)	(3)	X	3 3
23	(4)	(3)	6	5	6	(4)	6	5	(4)	5	(4)	X	(4) (4)
24	(4)	(3)	6	5	(4)	(3)	6	5	(4)	5	5		(4) 3
25	(3)	(4)	6	5	5	(3)	5	5	(4)	5	5		3 2
26	(4)	(4)	6	6	(4)	(4)	6	6	5	(4)	(4)	X	(4) 3
27	(4)	5	6	6	5	(4)	6	7	5	(4)	(4)	X	3 2
28	5	5	7	6	6	5	7	7	6	6	6		3 1
29	6	5	6	6	6	6	7	6	6	6	6		2 2
30	7	5	6	5	6	5	7	6	6	6	6		3 3
31	6	(4)	6	6	5	5	6	6	5	6	6		3 2

Score:

Quiet periods	P	10	6	23	26		9	8
	S	9	3	8	5		10	11
	U	0	0	0	0		3	3
	F	0	0	0	0		0	0
Disturbed periods	P	2	11	0	0		2	4
	S	7	10	0	0		7	4
	U	1	1	0	0		0	0
	F	2	0	0	0		0	1

Scales:

- Q-scale of Radio Propagation Quality
- (1) - useless
 - (2) - very poor
 - (3) - poor
 - (4) - poor to fair
 - 5 - fair
 - 6 - fair to good
 - 7 - good
 - 8 - very good
 - 9 - excellent

K-scale of Geomagnetic Activity
0 to 9, representing the greatest disturbance; $K_{CH} \geq 4$ indicates significant disturbance, enclosed in () for emphasis

Scoring: (beginning October 1952)

- P - Perfect: forecast quality equal to observed
- S - Satisfactory: (beginning October 1952)
forecast quality one grade different from observed
- U - Unsatisfactory: forecast quality two or more grades different from observed when both forecast and observed were ≥ 5 , or both ≤ 5
- F - Failure: other times when forecast quality two or more grades different from observed


Symbols:

- X - probable disturbed date

Note: All times are UT (Universal Time or GCT)

Table 63b

Short-Term Forecasts---March 1954



Outcome of Advance Forecasts (1 to 4 days ahead) --- March 1954




Table 64a

Coronal observations at Climax, Colorado (5303A), east limb

Date GCT	Degrees north of the solar equator															Degrees south of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Apr 1.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
2.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	8	4	2	-	-	-	-	-	-	-	-	-	-	-	-	-	
6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	8	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
7.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	5	2	1	1	-	-	-	-	-	-	-	-	-	-	-	-	
8.9a	X	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
9.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
12.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
13.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
16.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
17.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
18.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
19.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
21.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
22.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
23.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
24.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
26.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
27.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
28.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
29.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

Table 65a

Coronal observations at Climax, Colorado (6374A), east limb

Date GCT	Degrees north of the solar equator															Degrees south of the solar equator																					
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0°	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	4	4	4	4	4	4	1	2	2	1	1	1	1	1	2	2	2	
Apr 1.7	2	2	1	1	1	1	1	1	-	1	1	2	2	2	2	2	3	4	4	5	5	4	4	3	3	3	2	2	2	2	1	1	2	2	2		
2.7	2	2	2	1	1	1	1	1	1	1	1	1	2	2	2	2	2	3	4	5	14	8	3	5	4	4	3	1	1	1	1	1	2	2	2	2	2
5.7a	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	2	2	3	6	12	5	3	3	4	3	2	2	2	2	2	2	2	2	2	2	2	
6.7	2	2	2	1	1	1	1	1	1	1	1	2	2	2	1	2	3	4	4	5	5	6	3	4	3	2	2	2	2	2	1	1	1	1			
7.8	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	2	3	5	5	6	3	4	3	2	2	2	2	2	2	1	1	1	1	1			
8.9a	X	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	2	2	2	2	1	1	1	1	1	1	1	1	1	2	2	2	
9.6a	2	2	2	2	2	1	1	1	1	1	1	2	2	2	2	2	2	3	2	2	2	3	2	2	2	2	2	2	2	1	1	1	1	2	2		
10.6a	2	2	2	2	2	2	2	1	1	1	1	1	2	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3		
12.7	2	2	2	1	1	1	1	1	1	1	1	2	2	2	3	3	3	3	3	3	4	4	3	3	3	3	3	3	3	3	3	3	3	3			
13.7	2	2	1	1	1	1	1	1	1	1	1	2	2	2	3	2	2	2	2	2	3	3	3	3	3	3	2	1	1	1	1	1	2	2			
15.6	2	2	2	1	1	1	1	1	1	1	1	1	5	3	3	1	2	2	3	4	3	3	2	2	2	2	1	1	1	1	1	1	2	2			
16.7	2	2	2	2	1	1	1	1	1	2	4	5	5	5	3	4	5	5	6	5	5	5	5	5	5	5	3	1	1	1	1	1	2	2	2		
17.6	2	2	1	1	1	1	-	-	-	-	1	5	3	2	3	3	4	3	3	2	3	4	5	4	4	2	1	1	1	1	1	1	1	1	1		
18.8	2	2	2	1	1	1	1	1	1	1	2	3	3	3	3	3	3	3	4	3	4	4	5	4	3	2	1	1	1	1	1	1	1	1	2		
19.7	2	2	2	2	1	1	1	1	1	2	2	2	3	3	3	3	3	3	3	4	4	4	3	3	2	1	1	1	1	1	1	1	1	2	2		
20.6a	1	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	4	4	3	3	2	2	2	2	1	1	1	1	1	1	2	2		
21.6	2	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	2	2	2	2	1	1	1	1	1	1	1	1	1	1	1		
22.7	3	2	2	1	1	1	1	2	2	3	3	3	4	4	5	5	5	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4			
23.6	2	2	2	1	1	2	2	2	3	3	3	4	4	5	6	7	7	4	5	5	5	5	5	5	3	2	2	1	1	1	1	1	1	1	2		
24.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
26.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
27.8	2	2	2	2	1	1	1	1	1	1	2	3	2	2	2	2	2	3	4	5	5	5	5	4	4	4	4	4	4	4	4	4	4	4	4		
28.6	2	2	2	1	1	1	1	1	1	1	2	2	2	2	3	3	3	5	5	4	3	2	2	2	3	2	1	1	1	1	1						

Table 64b

Coronal observations at Climax, Colorado (5303A), west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator															
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Apr 1.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.9a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
12.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
13.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
16.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
17.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
18.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
19.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
21.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
22.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
23.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
24.6	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
26.8a	2	1	1	1	1	1	1	1	2	2	4	5	5	4	5	5	5	5	5	5	4	4	7	10	3	2	2	2	2	2	2	2
27.8a	2	2	1	1	1	1	1	1	2	2	2	2	1	3	4	4	3	3	3	3	3	3	3	3	2	1	1	1	1	1	1	1
28.6	2	2	2	1	1	1	1	1	1	2	2	2	2	1	3	4	4	3	3	3	3	3	3	3	2	1	1	1	1	1	1	1
29.6	2	2	2	2	1	1	1	1	1	1	2	2	2	1	3	4	4	3	3	3	3	3	3	3	1	1	1	1	1	1	1	1
30.6	2	2	2	1	1	1	1	1	2	2	3	3	3	4	4	3	2	3	3	3	3	3	3	3	2	1	1	1	1	1	1	1

Table 65b

Coronal observations at Climax, Colorado (6374A) west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator																
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Apr 1.7	2	2	2	2	1	1	1	1	1	2	2	2	2	2	3	3	3	3	3	3	3	2	1	2	3	2	1	1	1	1	1	2	
2.7	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	3	3	3	2	2	2	2	3	2	1	1	1	1	2	2	2	
5.7	2	2	2	2	2	1	1	1	1	1	1	1	1	1	2	3	3	3	2	2	4	4	2	3	2	2	1	1	1	1	2	2	
6.7	2	2	2	2	2	1	1	1	1	1	1	1	1	1	2	3	4	4	4	4	4	4	4	4	2	2	2	1	1	1	1	2	
7.8a	2	2	3	3	1	1	1	1	1	1	2	2	3	3	3	3	4	4	4	3	3	3	4	4	3	3	2	1	1	1	1	2	
8.9a	2	1	1	1	-	-	-	-	-	-	2	2	2	2	2	2	2	2	2	2	2	2	-	-	-	-	-	-	-	-	-	-	
9.6a	3	3	2	2	2	1	1	1	1	2	2	2	3	3	3	3	4	4	3	3	3	2	2	2	2	2	2	1	1	1	1	2	
10.6a	2	2	2	2	1	1	1	1	1	1	2	2	2	2	2	2	2	3	4	3	3	3	3	2	1	1	1	1	1	1	1	1	
12.7	2	2	1	1	1	1	1	1	1	1	1	1	1	1	2	2	2	3	2	2	2	2	2	2	2	2	1	1	1	1	1	2	
13.7a	2	2	2	2	1	1	1	1	1	1	2	2	3	4	4	4	4	4	3	3	2	2	3	3	3	3	1	1	1	1	2	2	
15.6	2	2	1	1	1	1	1	1	1	1	2	2	3	3	3	3	4	3	3	4	3	3	2	1	1	1	1	1	1	1	1	2	
16.7	2	2	2	2	1	1	1	1	1	1	1	3	4	4	4	4	5	6	7	7	7	7	5	3	2	1	1	1	1	1	2	3	
17.6	2	2	1	1	1	1	1	1	1	2	2	1	2	3	3	3	3	4	4	7	7	7	5	3	2	1	1	1	1	1	2	2	
18.8	2	2	2	1	1	1	1	1	1	2	2	4	3	3	3	2	2	4	9	3	3	4	3	3	2	2	1	1	1	1	1	2	
19.7a	2	2	2	1	1	1	1	1	1	2	2	2	2	3	4	4	3	2	8	3	4	4	3	3	2	2	1	1	1	1	1	2	
20.6a	2	2	1	1	1	1	1	1	1	1	3	3	3	3	3	3	5	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	
21.6	2	-	-	-	-	-	-	-	-	1	2	3	3	3	3	3	3	3	3	3	3	3	3	2	1	1	1	1	1	1	1	2	
22.7	2	2	3	2	1	2	2	2	2	3	3	3	3	3	3	3	4	4	5	5	4	4	3	3	1	1	1	1	2	2	2	3	
23.6	2	2	2	1	1	1	1	2	2	2	3	3	3	3	3	3	4	7	6	9	5	5	5	5	3	2	1	1	1	1	1	1	
24.6	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
26.8a	2	1	1	1	1	1	1	1	2	2	4	5	5	4	5	5	5	5	5	5	4	4	4	7	10	3	2	2	2	2	2	2	
27.8a	2	2	1	1	1	1	1	1	2	2	2	2	1	3	4	4	3	4	4	4	4	3	3	3	3	3	3	3	1	1	1	1	
28.6	2	2	2	1	1	1	1	1	1	2	2	2	2	1	3	4	4	3	3	3	3	3	3	3	3	3	3	3	1	1	1	1	2
29.6	2	2	2	2	1	1	1	1	1	2	2	2	2	1	3	4	4	3	3	3	3	3	3	3	3	3	3	3	1	1	1	1	2
30.6	2	2	2	1	1	1	1	1	2	2	3	3	3	4	4	3	2	3	3	3	3	3	3	3	3	2	1	1	1	1	1	2	

Table 66b</div

Table 67a

Coronal observations at Sacramento Peak, New Mexico (5303A), east limb

Date GCT	Degrees north of the solar equator													0°	Degrees south of the solar equator																		
	90	85	80	75	70	65	60	55	50	45	40	35	30		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Apr 1.7	-	-	2	2	2	3	3	2	3	3	4	4	3	2	2	2	-	-	2	2	2	2	-	2	2	2	3	2	3	3	-	-	-
2.7	-	-	-	-	-	-	-	-	2	2	3	3	4	4	3	2	2	2	3	3	3	3	2	2	2	2	3	3	2	-	-	-	
3.7	-	-	-	-	-	-	-	-	2	2	3	2	2	-	-	2	2	-	-	2	3	2	2	2	2	3	2	2	-	-	-		
4.8	-	-	-	-	-	-	-	2	2	3	4	3	3	2	2	3	2	2	2	2	3	8	11	10	3	2	3	3	2	3	2	-	
6.7	-	-	-	-	-	-	-	2	3	3	2	3	2	2	3	3	2	2	2	14	28	36	16	4	3	2	2	3	4	3	2	-	
7.7	-	-	-	-	-	-	-	2	4	5	4	4	5	4	4	2	2	3	8	20	18	8	5	4	3	2	2	3	3	2	-	-	
15.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
18.7a	-	-	-	-	-	-	-	-	3	4	3	3	3	2	3	-	-	-	-	3	4	3	4	3	2	2	3	3	2	-	-		
19.8a	-	-	-	-	-	-	-	-	2	3	3	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
20.7a	-	-	-	-	-	-	-	2	2	3	3	3	2	2	-	-	-	-	-	-	-	-	-	-	2	3	3	2	2	-			
23.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
27.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	3	3	2	3	4	2	-
28.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	2	2	3	2	3	2	2	3	3	3	2	-	-	-		

Table 68a

Coronal observations at Sacramento Peak, New Mexico (6374A), east limb

Date	Degrees north of the solar equator															0°	Degrees south of the solar equator																					
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
GCT	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																						
Apr 1.7	3	2	2	2	2	2	-	2	-	2	2	-	-	3	4	5	5	4	8	7	8	6	5	4	4	5	3	4	3	2	2	2	3	2	3			
2.7	3	2	5	3	3	2	2	2	2	2	3	3	4	5	5	6	8	13	14	12	9	8	5	4	5	5	4	4	3	3	3	2	2	2	3			
3.7	3	2	3	2	3	-	2	2	2	2	3	3	4	5	4	5	5	8	11	10	8	6	6	5	5	4	4	5	3	2	-	-	2	3	-	3		
4.8	3	3	2	3	2	2	2	3	2	3	4	5	8	10	9	7	8	13	14	18	20	15	14	11	9	12	13	11	4	3	2	-	2	3	3	4		
6.7	4	2	3	2	3	2	2	2	2	3	4	8	7	5	5	4	6	13	20	39	20	16	5	10	11	8	9	8	4	3	2	2	3	2	3	4	4	
7.7	4	3	4	3	4	3	2	2	3	2	3	5	6	7	8	14	15	20	16	14	13	10	8	9	6	7	8	4	3	2	-	2	2	3	4			
15.8a	-	-	-	-	-	-	-	-	-	-	2	2	3	5	4	3	3	4	5	6	6	5	4	3	3	2	2	3	3	-	-	-	-	-	-			
18.7a	-	-	-	-	-	-	-	-	-	-	2	2	3	2	3	3	4	4	4	5	4	5	3	4	5	5	5	3	2	-	-	-	-	-	-			
19.8a	-	-	-	-	-	-	-	-	-	-	2	2	3	3	2	4	3	3	3	3	3	4	4	3	4	3	2	-	-	-	-	-	-	-	-			
20.7a	-	-	-	-	-	-	-	-	-	-	2	3	4	5	4	4	5	6	8	8	4	5	4	5	4	3	2	-	-	-	-	-	-	-	-	2	2	3
23.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
27.8a	3	2	3	2	-	2	-	2	3	3	2	2	3	3	3	5	4	5	5	4	5	4	4	4	4	4	5	-	-	-	-	-	-	-	-	-	-	
28.7	4	4	2	3	2	4	3	2	2	3	4	3	3	4	5	6	7	8	8	11	12	14	11	5	4	4	5	4	3	2	-	2	2	2	3	4		

Table 69a

Coronal observations at Sacramento Peak, New Mexico (6702A), east limb

Table 67b

Coronal observations at Sacramento Peak, New Mexico (5303A), west limb

Date	Degrees south of the solar equator															0°	Degrees north of the solar equator																				
	90	85	80	75	70	65	60	55	50	45	4	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	
1954	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
Apr 1.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	2	2	2	2	3	3	4	4	4	3	2	2	2	-	-		
2.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	2	2	3	3	2	2	2	2	3	2	-	-	-	-	-		
3.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	2	-	-	3	2	2	2	2	-	-	-	-	-	-	-	-	-	
4.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	2	-	-	-	-	-	2	3	3	2	-	-	-	-	-	-	
6.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	-	-	2	3	2	3	4	3	2	2	2	-	-	-	-	-	
7.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	2	2	3	2	2	-	-	2	4	3	2	3	2	-	-	-	
15.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	2	2	3	2	2	-	-	-	-	-	-	-	-	-	-	-	-	
18.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	3	3	2	2	2	3	3	2	4	4	4	3	2	-	-	-	-	
19.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	8	5	5	4	3	3	-	-	-	-	-	-	-	-	-	-	-	-
20.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	3	2	2	-	2	2	-	-	-	-	2	3	X	X	-	-	-	
23.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	3	4	5	4	3	2	2	3	2	3	-	-	-	-	-	-	
27.8a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	3	3	3	3	-	-	3	4	4	3	3	-	-	-	-	
28.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	-	-	-	-	-	-	-	2	5	8	4	3	2	-	-	-	-

Table 68b

Coronal observations at Sacramento Peak, New Mexico (6374A), west limb

Date GCT	Degrees south of the solar equator															0°	Degrees north of the solar equator																					
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																						
pr 1.7	3	3	3	4	3	2	2	3	2	2	3	4	5	4	3	4	5	8	7	8	8	7	5	4	4	5	4	2	3	3	2	2	3	3	3			
2.7	3	3	2	3	2	3	2	3	3	3	2	3	3	4	4	5	9	11	8	7	5	6	5	4	5	4	3	3	—	—	3	2	2	3	3			
3.7	3	2	3	2	2	2	—	—	3	3	2	3	3	6	7	5	6	7	8	7	8	7	6	5	5	5	4	3	2	2	2	—	2	—	3	3		
4.8	4	2	3	2	2	2	—	2	2	3	3	3	3	4	4	6	7	8	9	8	11	10	7	5	6	8	8	7	4	2	2	—	2	3	2	2	3	
6.7	4	2	2	2	2	2	3	2	3	2	3	5	7	8	11	8	7	7	11	14	18	15	12	16	14	15	12	8	6	3	2	2	2	3	3	4	4	4
7.7a	4	4	3	4	3	2	3	2	2	3	4	5	8	7	8	6	6	11	14	15	11	11	10	7	6	5	5	3	2	2	2	3	3	4	4	4		
15.8a	—	—	—	—	—	—	—	—	—	3	4	5	8	7	5	6	6	5	6	6	5	5	4	3	5	4	2	—	—	—	—	—	—	—	—	—		
18.7	—	2	2	3	2	2	—	—	2	3	2	4	5	4	3	5	4	8	14	11	7	6	5	4	3	2	2	2	2	—	—	—	—	—	—	—	—	
19.8a	—	—	—	—	—	—	—	—	—	—	—	3	3	4	4	4	4	3	5	4	4	4	3	2	3	3	2	2	—	—	—	—	—	—	—	—	—	
20.7a	3	2	2	—	2	2	2	—	2	2	3	3	2	3	4	4	5	5	8	5	4	4	3	3	2	2	—	—	—	—	X	X	—	—	—	—	—	
23.8a	—	—	—	—	—	—	—	—	—	2	3	3	2	3	3	4	3	4	5	4	5	4	3	4	4	4	4	3	—	—	—	—	—	—	—	—	—	—
27.8a	—	—	—	—	—	—	—	—	—	2	3	3	2	3	3	3	3	2	3	5	4	3	4	3	2	3	3	2	2	2	—	2	2	2	3	3	3	
28.7	4	2	3	4	2	3	3	2	3	3	5	4	5	4	5	6	11	12	11	9	8	6	4	4	5	5	5	3	2	—	2	3	3	3	3	4	4	

Table 69b

Coronal observations at Sacramento Peak, New Mexico (6702A), west limb

Table 70
Zürich Provisional Relative Sunspot Numbers
April 1954

Date	R _Z *	Date	R _Z *
1	0	17	0
2	0	18	0
3	0	19	0
4	0	20	8
5	0	21	0
6	0	22	0
7	8	23	0
8	8	24	0
9	15	25	0
10	0	26	0
11	0	27	0
12	0	28	0
13	0	29	0
14	0	30	0
15	7		
16	7	Mean:	1.8

*Dependent on observations at Zürich Observatory and its stations at Locarno and Arosa.

Table 71
American Relative Sunspot Numbers
March 1954

Date	R _A	Date	R _A
1	10	17	24
2	8	18	26
3	9	19	19
4	1	20	17
5	0	21	15
6	0	22	10
7	0	23	0
8	0	24	5
9	0	25	1
10	0	26	0
11	0	27	0
12	8	28	0
13	19	29	0
14	24	30	0
15	26	31	0
16	26	Mean:	8.0

Table 72Solar Flares, April 1954

It is hoped to publish the April 1954 solar flare data in a future issue of the F series.

Table 73

Indices of Geomagnetic Activity for March 1954





Preliminary values of international character-figures, C;
Geomagnetic planetary three-hour-range indices, Kp;
Magnetically selected quiet and disturbed days





Table 74
Sudden Ionosphere Disturbances

It is hoped to bring the SID data up to date in a future issue of the F series.

Note: Observers are invited to send to the CRPL information on times of beginning and end of sudden ionosphere disturbances for publication as above. Address letters to the Central Radio Propagation Laboratory, National Bureau of Standards, Boulder, Colorado.

GRAPHS OF IONOSPHERIC DATA








Fig. 9. SINGAPORE, BRITISH MALAYA
 1.3° N 103.8° E AUGUST 1953




Fig. 10. SINGAPORE, BRITISH MALAYA




Fig. II. RAROTONGA I.
21.3°S, 159.8°W AUGUST 1953




Fig. 12. RAROTONGA I. AUGUST 1953




Fig. 13. CHRISTCHURCH, NEW ZEALAND
43.6°S, 172.7°E AUGUST 1953




Fig. 14. CHRISTCHURCH, NEW ZEALAND




Fig. 15 FALKLAND IS.
51.7°S, 57.8°W AUGUST 1953




Fig. 16. FALKLAND IS.




Fig. 17. CHURCHILL, CANADA
58.8°N, 94.2°W JULY 1953




Fig. 18. CHURCHILL, CANADA JULY 1953




Fig. 19. INVERNESS, SCOTL AND
57.4°N, 4.2°W JULY 1953




Fig. 20. INVERNESS, SCOTL AND JULY 1953




Fig. 21. LINDAU/HARZ, GERMANY
51.6°N, 10.1°E JULY 1953




Fig. 22. LINDAU/HARZ, GERMANY JULY 1953




Fig. 23. SLOUGH, ENGLAND
51.5°N, 0.6°W JULY 1953
















Fig. 24. SLOUGH, ENGLAND JULY 1953










Fig. 37. SINGAPORE, BRITISH MALAYA
1.3°N, 103.8°E JUNE 1953




Fig. 38. SINGAPORE, BRITISH MALAYA JUNE 1953




Fig. 39. TOWNSVILLE, AUSTRALIA
19.3°S, 146.8°E JUNE 1953








Fig. 40. TOWNSVILLE, AUSTRALIA JUNE 1953






Fig. 45. FALKLAND IS.

51.7° S, 57.8° W

JUNE 1953




Fig. 46. FALKLAND IS.

JUNE 1953




Fig. 47. CHURCHILL, CANADA

58.8° N, 94.2° W

MAY 1953
















Fig. 48. CHURCHILL, CANADA

MAY 1953










Fig. 61. HOBART, TASMANIA
 42.9°S, 147.3°E MAY 1953




Fig. 62. HOBART, TASMANIA MAY 1953




Fig 63. FALKLAND IS.
51°7'S., 57.8°W MAY 1953




Fig. 64 FALKLAND IS. MAY 1953




Fig. 65. INVERNESS, SCOTLAND

57.4° N, 4.2° W

APRIL 1953




Fig. 66. INVERNESS, SCOTLAND

APRIL 1953




Fig. 67. SLOUGH, ENGLAND

51.5° N, 0.6° W

APRIL 1953




Fig. 68. SLOUGH, ENGLAND

APRIL 1953



Fig. 69. SINGAPORE, BRITISH MALAYA
1.3°N, 103.8°E APRIL 1953




Fig. 70. SINGAPORE, BRITISH MALAYA




Fig. 71. TOWNSVILLE, AUSTRALIA
19.3°S, 146.8°E APRIL 1953




Fig. 72. TOWNSVILLE, AUSTRALIA APRIL 1953







Fig 74 BRISBANE, AUSTRALIA APRIL 1953






Fig 77. HOBART, TASMANIA

42°9'S, 147.3°E

APRIL 1953




Fig 78. HOBART, TASMANIA

APRIL 1953



Fig 79. FALKLAND IS.

51.7°S, 57.8°W

APRIL 1953




Fig 80. FALKLAND IS.

APRIL 1953




Fig. 81. POITIERS, FRANCE
46.6°N, 0.3°E MARCH 1953




Fig. 82. POITIERS, FRANCE MARCH 1953




Fig. 83. CASABLANCA, MOROCCO
33.6°N, 7.6°W MARCH 1953




Fig. 84. CASABLANCA, MOROCCO MARCH 1953




Fig. 85 KHARTOUM, SUDAN
15.6°N, 32.6°E

MARCH 1953




Fig. 86. KHARTOUM, SUDAN

MARCH 1953




Fig. 87. FALKLAND IS
51°S, 57°8'W

MARCH 1953








Fig. 88 FALKLAND IS

MARCH 1953






Fig. 93. POITIERS, FRANCE
46.6° N, 0.3° E JANUARY 1953




Fig. 94. POITIERS, FRANCE JANUARY 1953





Fig. 95. CASABLANCA, MOROCCO
33.6° N, 76° W JANUARY 1953



JANUARY 1953

Fig. 96. CASABLANCA, MOROCCO

Index of Tables and Graphs of Ionospheric Datain CPPL-F117

	<u>Table page</u>	<u>Figure page</u>
Brisbane, Australia		
May 1953	16	59
April 1953	18	63
Canberra, Australia		
June 1953	15	55
May 1953	16	59
April 1953	18	63
Casablanca, Morocco		
March 1953	18	65
February 1953	19	67
January 1953	19	68
Christchurch, New Zealand		
August 1953	13	48
July 1953	14	52
Churchill, Canada		
July 1953	13	49
May 1953	15	56
Falkland Is.		
August 1953	13	48
July 1953	14	52
June 1953	15	56
May 1953	17	60
April 1953	18	64
March 1953	19	66
Hobart, Tasmania		
June 1953	15	55
May 1953	17	60
April 1953	18	64
Inverness, Scotland		
August 1953	12	45
July 1953	13	49
June 1953	14	53
May 1953	16	57
April 1953	17	61
Khartoum, Sudan		
August 1953	12	46
March 1953	19	66
Lindau/Harz, Germany		
July 1953	13	50

Index (CRPL-F117, concluded)

	<u>Table page</u>	<u>Figure page</u>
Poitiers, France		
March 1953	18	65
February 1953	19	67
January 1953	19	68
Rarotonga I.		
August 1953	12	47
July 1953	14	51
Singapore, British Malaya		
August 1953	12	47
July 1953	14	51
June 1953	15	54
May 1953	16	58
April 1953	17	62
Slough, England		
August 1953	12	46
July 1953	13	50
June 1953	14	53
May 1953	16	57
April 1953	17	61
Townsville, Australia		
June 1953	15	54
May 1953	16	58
April 1953	17	62
Washington, D. C.		
April 1954	12	45

CRPL Reports

[A detailed list of CRPL publications is available from the Central Radio Propagation Laboratory upon request]

Daily:

Radio disturbance forecasts, every half hour from broadcast stations WWV and WWVH of the National Bureau of Standards.

Telephoned and telegraphed reports of ionospheric, solar, geomagnetic, and radio propagation data.

Semiweekly:

CRPL—J. North Atlantic Radio Propagation Forecast (of days most likely to be disturbed during following month).

CRPL—Jp. North Pacific Radio Propagation Forecast (of days most likely to be disturbed during following month).

Semimonthly:

CRPL—Ja. Semimonthly Frequency Revision Factors For CRPL Basic Radio Propagation Prediction Reports.

Monthly:

CRPL—D. Basic Radio Propagation Predictions—Three months in advance. (Dept. of the Army, TB 11-499-, monthly supplements to TM 11-499; Dept. of the Navy, DNC 13 () series; Dept. of the Air Force, TO 16-1B-2 series.) On sale by Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C. Members of the Armed Forces should address cognizant military office.

CRPL—F. Ionospheric Data. Limited distribution. This publication is in general disseminated only to those individuals or scientific organizations which collaborate in the exchange of ionospheric, solar, geomagnetic or other radio propagation data or in exchange for copies of publications on radio, physics and geophysics for the CRPL library.

Circulars of the National Bureau of Standards pertaining to Radio Sky Wave Transmission:

NBS Circular 462. Ionospheric Radio Propagation.

NBS Circular 465. Instructions for the Use of Basic Radio Propagation Predictions.

These circulars are on sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C. Members of the Armed Forces should address the respective military office having cognizance of radio wave propagation.

The publications listed above may be obtained without charge from the Central Radio Propagation Laboratory, unless otherwise indicated.
