

CRPL-F49

**National Bureau of Standards
Library, N. W. Bldg.**

SEP 28 1948

**Reference book not to be
taken from the Library.**

IONOSPHERIC DATA

ISSUED

SEPTEMBER 1948

**PREPARED BY CENTRAL RADIO PROPAGATION LABORATORY
National Bureau of Standards
Washington, D.C.**

IONOSPHERIC DATA

CONTENTS

	Page
Terminology and Scaling Practices	2
Monthly Average and Median Values of World-Wide Ionospheric Data	4
Ionospheric Data for Every Day and Hour at Washington, D. C.	6
Ionosphere Disturbances	6
American and Zürich Provisional Relative Sunspot Numbers	7
Solar Coronal Intensities Observed at Climax, Colorado	8
Errata	8
Tables of Ionospheric Data	10
Graphs of Ionospheric Data	42
Index of Tables and Graphs of Ionospheric Data in CRPL-F49	64

TERMINOLOGY AND SCALING PRACTICES

The symbols and terminology used in this report are those adopted by the International Radio Propagation Conference, and given in detail on pages 24 to 26 of the report IRPL-C61, "Report of International Radio Propagation Conference," and in the section on "Terminology" in report IRPL-F5.

Beginning with IRPL-F14 the symbol L, defined as follows, is used in detailed tabulations of hourly values of ionosphere characteristics observed at Washington:

L or l = critical frequency, muf, or muf factor for F1 layer omitted because no definite and abrupt change in slope of the $h'f$ curve occurs either for the first reflection or for any of the multiples.

In the past, ionospheric conditions were summarized on a monthly basis by using average or mean values for each hour of the day for each month. However, following the recommendations of the International Radio Propagation Conference, held in Washington April 17 to May 5, 1944, beginning with data for January 1, 1945, median values are published wherever possible.

Where averages are reported, they are, at any hour, the average for all the days during the month for which numerical data exist.

The monthly median values used here are the values equaled or exceeded on half the days of the month at the given hour. The following conventions are used in determining the medians for hours when no measured values are given because of equipment limitations and ionospheric irregularities. Symbols used are those given in the report referred to above, IRPL-C61.

a. For all ionospheric characteristics:

Values missing because of A, B, C, or F (see terminology referred to above) are omitted from the median count.

b. For critical frequencies and virtual heights:

Values of f^oF2 (and f^oE near sunrise and sunset) missing because of E are counted as equal to or less than the lower limit of the recorder. Values of $h'F2$ (and $h'E$ near sunrise and sunset) missing for this reason are counted as equal to or greater than the median. Other characteristics missing because of E are omitted from the median count. See CRPL-F36, page 9.

Values missing because of D are counted as equal to or greater than the upper limit of the recorder.

Values missing because of G are counted:

1. For f^oF2 , as equal to or less than f^oF1 .
2. For $h'F2$, as equal to or greater than the median.

Values missing for any other reason are omitted from the median count.

c. For muf factors (N-factors):

Values missing because of G are counted as equal to or less than the median.

Values missing for any other reason are omitted from the median count.

d. For sporadic E (Es):

Values of f^oE missing because no Es reflections appeared, the equipment functioning normally otherwise, are counted as equal to or less than the median f^oE, or equal to or less than the lower frequency count of the recorder.

Values of f^oE missing for any other reason, and values of hEs missing for any reason at all are omitted from the median count.

Beginning with data for November 1945, doubtful monthly median values for ionospheric observations at Washington, D. C., are indicated by parentheses; in accordance with the practice already in use for doubtful hourly values. The following are the conventions used to determine whether or not a median value is doubtful:

1. If only four values or less are available, the data are considered insufficient and no median value is computed.

2. For the F2 layer, if only five to nine values are available, the median is considered doubtful. The N and F1 layers are so regular in their characteristics that, as long as there are at least five values, the median is not considered doubtful.

3. For all layers, if more than half of the values used to compute the median are doubtful (either doubtful or interpolated), the median is considered doubtful.

The same conventions are used by the CRPL in computing the medians from tabulations of daily and hourly data for stations other than Washington, beginning with the tables in CRPL-F18.

Beginning with CRPL-F33, an additional group of symbols is used in recording the Washington, D.C. data. The list of additional symbols and their meanings follows:

N - unable to make logical interpretation.

P - trace extrapolated to a critical frequency.

Q - the F1 layer not present as a distinct layer.

R - curve becomes incoherent near the F2 critical frequency.

S - no observation obtainable because of interference.

V - forked record (previously denoted by U. This change should also be made in CRPL-7-1).

Z - triple split near critical frequency.

For a more detailed explanation of the meaning and use of these symbols, see the report CRPL-7-1, "Preliminary Instructions for Obtaining and Reducing Manual Ionospheric Records."

MONTHLY AVERAGE AND MEDIAN VALUES OF WORLD-WIDE IONOSPHERIC DATA

The ionospheric data given here in tables 1 to 43 and figures 1 to 85 were assembled by the Central Radio Propagation Laboratory for analysis and correlation, incidental to CRPL predictions of radio propagation conditions. The data are median values unless otherwise indicated. The following are the sources of the data:

Australian Council for Scientific and Industrial Research,
Radio Research Board:

Brisbane, Australia
Canberra, Australia
Hobart, Tasmania

Australian Department of Supply and Shipping, Bureau of
Mineral Resources, Geophysical Section:
Watheroo, W. Australia

British Department of Scientific and Industrial Research,
Radio Research Board:

Falkland Is.
Fraserburgh, Scotland
Slough, England

New Zealand Radio Research Committee:

Christchurch, New Zealand (Canterbury University College Observatory)
Rarotonga I.

Japanese Physical Institute for Radio Waves (under supervision of
Supreme Commander, Allied Powers):

Fukaura, Japan
Shibata, Japan
Tokyo (Kokobunji), Japan
Wakkanai, Japan
Yamakawa, Japan

National Bureau of Standards (Central Radio Propagation Laboratory):

Baton Rouge, Louisiana (Louisiana State University)
Boston, Massachusetts (Harvard University)
Guam I.
Huancayo, Peru (Instituto Geofisico de Huancayo)
Palmyra I.
San Francisco, California (Stanford University)
San Juan, Puerto Rico (University of Puerto Rico)
Trinidad, British West Indies
Washington, D. C.
White Sands, New Mexico
Wuchang, China (National Wuhan University)

All India Radio (Government of India), New Delhi, India:
 Bombay, India
 Delhi, India
 Madras, India

Radio Wave Research Laboratory, Central Broadcasting Administration:
 Chungking, China
 Lanchow, China
 Nanking, China
 Peiping, China

The dashed-line prediction curves of the graphs of ionospheric data are obtained from the predicted zero-muf contour charts of the CRPL-D series publications. The following points are worthy of note:

- a. Predictions for individual stations used to construct the charts may be more accurate than the values read from the charts since some smoothing of the contours is necessary to allow for the longitude effect within a zone. Thus, inasmuch as the predicted contours are for the center of each zone, part of the discrepancy between the predicted and observed values as given in the F series may be caused by the fact that the station is not centrally located within the zone.
- b. The final presentation of the predictions is dependent upon the latest available ionospheric and radio propagation data, as well as upon predicted sunspot number.
- c. There is no indication on the graphs of the relative reliability of the data; it is necessary to consult the tables for such information.

The following predicted smoothed 12-month running-average Zurich sunspot numbers were used in constructing the contour charts:

Month	Predicted Sunspot No.			
	1948	1947	1946	1945
December	126	85	38	
November	124	83	36	
October	119	81	23	
September	121	79	22	
August	123	122	77	20
July	125	116	73	
June	129	112	67	
May	130	109	67	
April	133	107	62	
March	133	105	51	
February	133	90	46	
January	130	88	42	

IONOSPHERIC DATA FOR EVERY DAY AND HOUR AT WASHINGTON, D. C.

The data given in tables 44 to 55 follow the scaling practices given in the report IRPL-C61, "Report of International Radio Propagation Conference," pages 36 to 39, and the median values are determined by the conventions given above under "Terminology and Scaling Practices."

IONOSPHERE DISTURBANCES

Table 56 presents ionosphere character figures for Washington, D. C., during August 1948, as determined by the criteria presented in the report IRPL-R5, "Criteria for Ionospheric Storminess," together with Cheltenham, Maryland, geomagnetic K-figures, which are usually covariant with them.

Table 57 lists for the stations whose locations are given the sudden ionosphere disturbances observed on the continuous field intensity recordings made at the Sterling Radio Propagation Laboratory during August 1948.

Table 58 lists for the stations whose locations are given the sudden ionosphere disturbances observed at the Brentwood and Somerton, England, receiving stations of Cable and Wireless, Ltd., from August 5 to August 17, 1948.

Table 59 gives provisional radio propagation quality figures for the North Atlantic and North Pacific areas, for Cl to 12 and 13 to 24 GCT, July 1948, compared with the CRPL daily radio disturbance warnings, which are primarily for the North Atlantic paths, the CRPL weekly radio propagation forecasts of probable disturbed periods, and the half-day Cheltenham, Maryland, geomagnetic K-figures.

The radio propagation quality figures are prepared from radio traffic and ionospheric data reported to the CRPL, in a manner basically the same as that described in IRPL-R31, "North Atlantic Radio Propagation Disturbances, October 1943 through October 1945," issued February 1, 1946. The scale conversions for each report are revised for use with the data beginning January 1948, and statistical weighting replaces what was, in effect, subjective weighting. Separate master distribution curves of the type described in IRPL-R31 were derived for the part of 1946 covered by each report; data received only since 1946 are compared with the master curve for the period of the available data. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. Each report is given a statistical weight which is the reciprocal of the departure from linearity. The half-daily radio propagation quality figure, beginning January 1948, is the weighted mean of the reports received for that period.

These radio propagation quality figures give a consensus of opinion of actual radio propagation conditions as reported by the half day over the two general areas. It should be borne in mind, however, that though the quality may be disturbed according to the CRPL scale, the cause of the disturbance is not necessarily known. There are many variables that must be considered. In addition to ionospheric storminess itself as the cause, conditions may be reported as disturbed because of seasonal characteristics, such as are particularly evident in the pronounced day and night contrast over North Pacific paths during the winter months, or because of improper frequency usage for the path and time of day in question. Insofar as possible, frequency usage is included in rating the reports. Where the actual frequency is not shown in the report to the CRPL, it has been assumed that the report is made on the use of optimum working frequencies for the path and time of day in question. Since there is a possibility that all the disturbance shown by the quality figures is not due to ionospheric storminess alone, care should be taken in using the quality figures in research correlations with solar, auroral, geomagnetic, or other data. Nevertheless, these quality figures do reflect a consensus of opinion of actual radio propagation conditions as found on any one half day in either of the two general areas.

AMERICAN AND ZÜRICH PROVISIONAL RELATIVE SUNSPOT NUMBERS

Table 60 presents the daily American relative sunspot number, R_A , computed from observations communicated to CRPL by observers in America and abroad. Beginning with the observations for January 1948, a new method of reduction of observations is employed such that each observer is assigned a scale-determining "observatory coefficient," ultimately referred to Zürich observations in a standard period, December 1944 to September 1945, and a statistical weight, the reciprocal of the variance of the observatory coefficient. The daily numbers listed in the table are the weighted means of all observations received for each day. Details of the procedure will be published shortly. The American relative sunspot number computed in this way is designated R_A . It is noted that a number of observatories abroad, including the Zürich observatory, are included in R_A . The scale of R_A was referred specifically to that of the Zürich relative sunspot numbers in the standard comparison period; since that time, R_A is influenced by the Zürich observations only in that Zürich proves to be a consistent observer and receives a high statistical weight. In addition, this table lists the daily provisional Zürich sunspot numbers, R_Z .

SOLAR CORONAL INTENSITIES OBSERVED AT CLIMAX, COLORADO

In tables 61a and 61b are listed the intensities of the green (5303A) line of the emission spectrum of the solar corona as observed during August 1948 by the High Altitude Observatory of Harvard University and the University of Colorado at Climax, Colorado, for east and west limbs, respectively, at 5° intervals of position angle north and south of the solar equator at the limb computed to the nearest 5° . A correction, P, as listed, has been applied to the position angles of the actual observations which were on astronomical coordinates. The time of observation is given to the nearest tenth of a day, GCT. The tables of coronal observations in CRPL-F29 to F41 listed the data on astronomical coordinates; the present format on solar rotation coordinates is in conformity with the tables of CRPL-1-4, "Observations of the Solar Corona at Climax, 1944-46."

Tables 62a and 62b give similarly the intensities of the first red (6374A) coronal line; tables 63a and 63b list the intensities of the second red (6704A) coronal line. The following symbols are used in tables 61, 62, and 63: a, observation of low weight; -, corona not visible; and x, position angle not included in plate estimates.

ERRATA

1. In CRPL-F47, page 66, figures 76 and 77 should have been grouped with the figures under the heading, "Graphs Superseding Previously Published Graphs," on page 69.
2. In CRPL-F48, the values of fOF2 from 1800 to 2200 given in table 55, page 23, were inadvertently omitted from the graph in figure 106 on page 73 of that issue.

TABLES AND GRAPHS
OF
IONOSPHERIC DATA

TABLES OF IONOSPHERIC DATA

Table 1

Washington, D.C. (38.9°N, 77.5°W)

August 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$
00	270	5.7						2.7
01	270	5.4				2.3		2.8
02	270	5.1				1.9		2.6
03	260	4.9				2.2		2.8
04	270	4.4						2.8
05	270	4.1						2.9
06	240	5.4			100	2.1	2.5	3.1
07	270	6.0	230	4.1	100	2.7	3.5	3.0
08	310	6.7	210	4.5	100	3.1	3.7	3.0
09	355	6.7	200	4.9	100	3.3	3.7	2.8
10	390	6.6	195	5.1	100	3.6	3.9	2.8
11	450	6.6	200	5.4	100	(3.7)	3.7	2.6
12	435	6.7	200	5.4	100	(3.8)		2.7
13	430	6.8	200	5.4	100	3.9		2.6
14	420	6.9	200	5.3	100	3.8		2.7
15	400	7.0	210	5.3	100	3.7		2.7
16	360	7.0	210	4.9	100	3.5		2.8
17	335	7.1	220	4.7	100	3.0	3.2	2.9
18	260	7.3	230		100	2.4	3.3	2.9
19	250	7.1			100	1.9	1.8	3.0
20	240	7.0						2.9
21	250	6.8					1.9	2.9
22	250	6.2					1.8	2.8
23	260	5.7					2.0	2.8

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 2

Boston, Massachusetts (42.4°N, 71.2°W)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$
00	275	6.7						2.6
01	300	6.4						1.4
02	300	5.4						1.7
03	280	5.0						1.5
04	305	4.7						2.6
05	305	5.6	275	4.3	112	1.9		2.8
06	340	6.1	300					2.8
07	405	6.5	250					2.7
08	445	6.9	250					2.6
09	455	7.0	248					2.5
10	452	7.3	230					2.5
11	495	7.4	245					2.4
12	482	7.1	240					2.5
13	500	7.4	260					2.4
14	472	7.4	225					2.5
15	440	7.5	242					2.5
16	385	7.2	250					2.7
17	360	7.5	252					2.7
18	290	7.8						2.7
19	260	7.8						2.9
20	282	8.0						2.7
21	280	7.7						2.7
22	290	7.4						2.7
23	288	6.9						2.6

Time: 75.0°W.

Sweep: 0.8 Mc to 14.0 Mc in 1 minute.

Table 3

San Francisco, California (37.4°N, 122.2°W)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$
00	320	6.0				3.0	2.4	
01	320	5.9				2.5	2.4	
02	300	5.6				2.6	2.5	
03	310	5.4				2.4	2.5	
04	300	5.0				2.0	2.5	
05	300	5.0				2.5	2.6	
06	260	5.8	260	4.0	120	2.4		2.5
07	400	6.6	240	4.7	120	2.9	4.1	2.4
08	400	7.6	220	5.1	120	3.2	4.8	2.5
09	380	8.0	220	5.2	120	3.6	5.0	2.5
10	410	8.4	220	5.4	120	3.8	4.5	2.4
11	400	8.6	220	5.4	120	3.9	4.7	2.5
12	400	8.6	220	5.4	120	4.0		2.5
13	380	8.6	220	5.4	120	3.9		2.5
14	380	8.4	220	5.4	120	3.9	4.8	2.5
15	380	8.3	220	5.3	120	3.8		2.6
16	360	7.8	220	5.2	120	3.4	4.2	2.6
17	355	7.8	240	4.9	120	3.1	4.5	2.6
18	260	7.8	240		120	2.6	3.6	2.7
19	260	7.8				2.8	2.8	
20	260	7.6				2.6	2.7	
21	260	7.1				2.8	2.6	
22	280	6.4				3.8	2.6	
23	300	6.0				3.0	2.5	

Time: 120.0°W.

Sweep: 1.3 Mc to 18.5 Mc in 4 minutes 30 seconds.

Table 4

White Sands, New Mexico (32.3°N, 106.5°W)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$
00	320	6.4						2.8
01	300	6.3						2.3
02	300	6.3						2.3
03	290	6.0						2.6
04	290	5.6						2.6
05	300	5.3						3.3
06	255	6.3						2.6
07	320	6.8	230	4.9	110	2.9	5.1	2.7
08	370	7.8	220	5.2	110	3.4	5.2	2.6
09	410	8.2	220	5.4	110	3.6	5.2	2.5
10	405	8.6	210	5.5	110	3.9	5.6	2.5
11	400	9.3	230	5.6	115	3.9	5.5	2.4
12	400	9.6	220	5.5	120	3.8	5.3	2.5
13	400	9.8	220	5.5	120	3.9	4.9	2.5
14	400	9.6	220	5.4	120	3.9	4.7	2.5
15	380	9.2	220	5.4	110	3.8	4.7	2.6
16	370	8.8	220	5.2	110	3.6	4.5	2.6
17	350	8.1	240	5.0	110	3.2	4.4	2.6
18	280	7.9	245	(4.4)	110	2.6	4.0	2.7
19	280	8.0						3.0
20	255	7.9						2.4
21	260	7.1						3.1
22	280	6.8						2.6
23	300	6.5						2.8

Time: 105.0°W.

Sweep: 0.7 Mc to 14.0 Mc in 2 minutes.

Table 5

Wuchang, China (30.6°E , 114.4°E)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$F2-13000$
00	295	8.9			5.0	2.8	
01	290	6.6			4.6	2.8	
02	270	8.6			4.0	2.9	
03	270	6.0			3.4	2.9	
04	270	7.6			3.4	2.8	
05	270	7.1			2.8	2.8	
06	250	8.0			120	3.2	3.0
07	240	8.7			110	3.8	3.1
08	252	6.8	225		100	3.4	5.2
09	290	8.7	230	5.8	100	3.7	6.2
10	365	8.9	252	6.0	100	3.9	6.4
11	368	9.9	250	6.0	100	3.9	7.8
12	370	10.7	250	6.0	100	4.0	6.8
13	362	11.0	232	6.0	100	4.0	6.0
14	352	11.4	230	5.8	100	4.0	5.6
15	340	11.8	235	5.8	100	3.8	5.6
16	320	11.3	230	5.6	100	3.7	4.6
17	300	11.3	230	5.3	100	3.3	5.0
18	285	10.7	260		100	2.8	5.5
19	270	10.0			110	3.0	5.0
20	270	8.8				4.2	2.8
21	290	8.8				4.6	2.7
22	308	9.2				5.0	2.6
23	300	9.2				5.0	2.7

Time: 120.0°E .

Sweep: 1.2 Mc to 19.2 Mc, manual operation.

Table 6

Baton Rouge, Louisiana (30.5°N , 91.2°W)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$F2-13000$
00	300	6.6			300	6.6	
01	300	6.6			300	6.3	
02	300	6.3			300	6.0	
03	300	5.6			300	5.6	
04	300	5.6			300	5.6	
05	300	5.6			300	5.6	
06	290	6.2			120	2.3	3.5
07	300	7.4	240		120	3.1	3.7
08	320	8.4	270		120	3.5	4.3
09	330	8.6	220		120	3.7	
10	380	9.6	220		120	3.8	
11	400	9.2	230		120	(3.6)	
12	390	9.8	(240)		120	3.8	
13	390	9.7	(240)		120	(3.7)	
14	390	9.5	(240)		120	3.7	
15	380	9.0	240		120	3.7	
16	380	8.8	240		120	3.6	
17	340	8.6	240		120	3.4	
18	300	8.3			120	2.6	3.9
19	300	8.0					2.9
20	290	7.9					2.9
21	290	7.5					2.8
22	290	7.2					2.8
23	300	6.8					2.8

Time: 90.0°W .

Sweep: 2.12 Mc to 15.3 Mc in 5 minutes, automatic operation.

Table 7

San Juan, Puerto Rico (18.4°N , 66.1°W)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$F2-13000$
00		9.1				2.8	
01	8.8					2.9	
02	8.0					2.8	
03	7.5					2.7	
04	7.3					2.8	
05	7.0					2.8	
06	7.1					2.9	
07	280	8.0				2.9	
08	280	8.8			3.2	2.7	
09	330	9.2	5.3		3.5	2.6	
10	370	10.4	5.6		3.7	2.5	
11	380	11.0	5.8		3.9	2.5	
12	375	11.0	5.8		4.0	2.5	
13	385	11.2	5.8		4.0	2.5	
14	370	11.5	5.6		4.0	2.5	
15	370	11.5	5.4		3.8	2.6	
16	360	11.0	5.2		3.6	2.5	
17	335	10.3			3.2	4.3	2.6
18	300	10.0				2.7	
19	290	9.8				2.7	
20	290	9.4				2.7	
21	290	9.3				2.6	
22	280	9.1				(2.7)	
23	280	9.1				2.7	

Time: 60.0°W .

Sweep: 2.8 Mc to 13.0 Mc in 9 minutes, supplemented by manual operation.

Table 8

Guam I. (13.6°N , 144.9°E)

July 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$F2-13000$
00	335	9.4					3.8
01	320	8.7					3.0
02	310	(8.0)					2.3
03	280	(7.6)					(2.8)
04	240	7.6					1.8
05	225	8.0					3.0
06	250	6.8					3.6
07	240	8.4					3.1
08	230	9.3					4.8
09	220	9.9					5.0
10	220	10.5					6.2
11	220	10.6					6.2
12	220	11.5	310		6.3		5.8
13	210	12.2			6.1		5.5
14	415	12.6	310		6.0	110	5.0
15	220	13.0			(5.0)	110	4.0
16	220	13.7	220		5.8		5.0
17	240	13.7					2.4
18	260	13.4					5.2
19	300	12.3					2.5
20	370	11.0					4.8
21	390	10.4					4.6
22	370	(9.7)					2.5
23	350	(9.3)					(2.4)

Time: 150.0°E .

Sweep: 1.25 Mc to 19.0 Mc in 12 minutes, manual operation.

Table 9

Trinidad, Brit. West Indies (10.6°N, 61.2°W)

July 1948

Time	h°F2	f°F2	h°F1	f°F1	h°E	f°E	fEs	F2-M3000
00	260	10.8					2.9	
01	260	9.9					2.8	
02	250	9.2					2.8	
03	270	8.7					2.8	
04	260	8.2					2.9	
05	260	7.5					2.9	
06	260	7.5			120	(1.5)	2.4	2.9
07	250	8.2			120	2.8	3.4	3.0
08	230	8.6			120	3.4	4.3	2.8
09	300	9.7	220	5.2	120	3.7	4.4	2.7
10	340	10.7	220	5.8	120	3.9	4.5	2.6
11	340	11.3	220	5.8	120	4.1	4.6	2.5
12	380	11.9	220	5.8	120	4.2	4.6	2.5
13	370	12.2	220	5.9	120	4.1	4.8	2.6
14	370	12.2	220	5.8	120	4.1	5.0	2.6
15	360	12.0	230	5.7	120	3.9	4.9	2.6
16	330	11.6	230	5.2	120	3.6	4.8	2.6
17	300	11.2	240	4.9	120	3.1	4.7	2.6
18	270	10.7			120		4.3	2.6
19	290	10.5					2.8	2.6
20	320	11.1					2.8	2.5
21	300	11.4					2.4	2.6
22	300	11.5					2.0	2.7
23	280	11.3						2.8

Time: 60.0°W.

Sweep: 1.2 Mc to 18.0 Mc, manual operation.

Table 10

Palmyra I. (5.9°N, 162.1°W)

July 1948

Time	h°F2	f°F2	h°F1	f°F1	h°E	f°E	fEs	F2-M3000
00		270			(10.0)			(3.0)
01		270			(10.0)			(2.9)
02		270			(10.0)			(2.9)
03		260			(10.0)			(2.9)
04		240			(8.8)			1.6 (3.1)
05		240			7.3			1.7 (3.0)
06		250			5.6		130	2.1 (3.0)
07		255			7.5		110	2.5 2.0
08		240			8.4		110	3.3 2.7
09		250			9.4	230	120	3.8 2.5
10		270			10.0	230	120	4.0 (2.3)
11		270			10.7	220	120	4.3 (2.3)
12		280			11.2	220	120	4.4 (2.3)
13		280			11.3	230	120	4.3 2.2
14		270			11.4	220	120	4.3 (2.2)
15		260			11.5	220	120	4.0 2.3
16		250			11.8	230	115	3.5 4.2 (2.3)
17		240			(11.3)		120	3.1 4.4 (2.3)
18		270			(11.2)		120	3.3 3.8 (2.3)
19		330			(10.2)			2.6 (2.2)
20		420			(9.4)			1.7 (2.2)
21		400			(9.2)			(2.3)
22		340			(10.3)			1.8 (2.4)
23		300			(10.4)			1.8 (2.7)

Time: 157.5°W.

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 36 seconds, automatic operation; 13.0 Mc to 18.0 Mc, manual operation.

Table 11

Huancayo, Peru (12.0°S, 75.3°W)

July 1948

Time	h°F2	f°F2	h°F1	f°F1	h°E	f°E	fEs	F2-M3000
00	240	7.9					2.9	
01	240	7.7					3.0	
02	240	7.2					2.8	
03	250	6.0					2.4	
04	250	5.0					2.3	
05	270	4.3					2.7	
06	310	4.7					3.0	
07	270	7.8			1.4	2.8	2.7	
08	250	9.8			2.5	7.2	2.9	
09	240	10.2	235	5.4	3.1	11.8	2.7	
10	230	10.0	230	5.5	3.6	12.1	2.5	
11	230	10.1	220	5.5	3.9	12.8	2.3	
12	300	9.8	220	5.5	4.0	12.8	2.3	
13	290	9.8	220	5.5	4.0	12.8	2.2	
14	230	9.7	220	5.5	4.0	12.8	2.2	
15	230	9.8			3.9	12.6	2.2	
16	250	9.6			3.5	12.6	2.2	
17	280	9.4			3.0	11.9	2.2	
18	330	8.8			2.4	10.8	2.2	
19	370	8.0			1.3	1.8	2.2	
20	340	8.2					2.2	
21	285	8.4					2.3	
22	260	7.9					2.4	
23	245	8.2					2.6	
							2.8	

Time: 75.0°W.

Sweep: 1.6 Mc to 0.5 Mc in 15 minutes, automatic operation.

Table 12

Wakkanai, Japan (45.4°N, 141.7°E)

June 1948

Time	h°F2	f°F2	h°F1	f°F1	h°E	f°E	fEs	F2-M3000
00		270			8.1			3.0 2.7
01		280			7.9			2.8 2.7
02		260			7.6			2.4 2.7
03		270			7.5			2.3 2.7
04		280			7.5			1.7 2.5 2.7
05		280			8.3	225	100	2.5 3.4 2.8
06		300			8.7	215	100	3.0 4.1 2.8
07		300			8.7	210	100	3.4 4.8 2.8
08		345			8.4	205	100	3.6 6.8 2.8
09		375			7.9	200	100	3.8 6.7 2.7
10		390			7.9	210	5.2	3.9 6.6 2.7
11		390			8.0	200	5.7	5.6 2.7
12		380			8.0	200	5.6	5.6 2.7
13		395			7.8	200	5.6	100 6.0 2.7
14		380			7.8	200	5.6	100 5.0 2.7
15		375			7.9	200	5.2	100 5.6 2.7
16		345			8.0	220	4.8	100 3.6 6.2 2.8
17		310			8.0	215	100	3.2 6.4 2.8
18		290			8.1		100	2.5 7.1 2.9
19		270			8.0			5.6 2.9
20		260			8.3			6.0 2.7
21		270			8.2			4.1 2.7
22		290			8.2			3.8 2.7
23		280			8.2			3.6 2.7

Time: 135.0°E.

Sweep: 1.0 Mc to 17.0 Mc, manual operation.

Table 13

Fukaura, Japan (40.6°N , 139.9°E)

June 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}s$	F2-M3000
00	310	8.5				4.1	2.5	
01	310	8.6				3.6	2.6	
02	300	8.2				3.4	2.6	
03	300	7.8				3.2	2.6	
04	300	7.7			(1.5)	2.8	2.6	
05	300	8.3	260		110	2.2	3.2	2.7
06	300	9.1	260		115	2.8	4.3	2.6
07	330	9.4			110	3.4	5.0	2.8
08	350	9.3			120	3.6	6.4	2.7
09	390	9.0		(5.4)	120	4.0	6.2	2.6
10	400	9.0			120	4.0	6.4	2.6
11	400	9.0			5.6		6.2	2.6
12	400	9.0			5.6		6.8	2.5
13	400	9.1			5.6		6.8	2.6
14	400	9.0			110	3.9	6.8	2.6
15	390	9.1			110	3.4	6.5	2.5
16	390	9.0			5.5		6.8	2.6
17	350	8.6	260		110	3.5	6.6	2.6
18	320	8.5	240		110	2.5	6.4	2.6
19	300	8.6					6.8	2.7
20	300	8.3					6.0	2.6
21	310	8.4					5.0	2.5
22	320	8.5					5.0	2.5
23	320	8.6					3.8	2.5

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc, manual operation.

Table 14

Peiping, China (39.9°N , 116.4°E)

June 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}s$	F2-M3000
00								
01								
02								
03								
04								
05								
06								
07								
08								
09								
10								
11								
12								
13								
14								
15								
16								
17								
18								
19								
20								
21								
22								
23								

Time: 120.0°E .

Sweep: 1.7 Mc to 20.0 Mc in 15 minutes, manual operation.

Table 15

Shibata, Japan (37.9°N , 139.3°E)

June 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}s$	F2-M3000
00	(300)	9.0				4.2	2.6	
01	300	9.0				3.9	2.7	
02	280	8.8				3.9	2.8	
03	280	8.3				3.6	2.7	
04	285	8.1				3.0	2.7	
05	265	8.4	230		110	2.4	3.4	2.7
06	275	9.5	220		100	2.9	4.6	2.8
07	300	9.8	220		100	3.4	5.3	2.8
08	320	9.3	220		100	3.6	6.3	2.8
09	370	9.2	210	(5.2)	100	3.8	6.6	2.7
10	405	9.1	205	5.6	100	4.1	6.8	2.6
11	400	9.1	200	5.6	100	4.0	6.4	2.5
12	400	9.7	210	5.8	100	4.1	6.4	2.6
13	380	9.8	220	5.7	100	4.2	7.2	2.7
14	380	9.4	220	5.6	100	4.1	6.3	2.7
15	370	9.3	240	5.5	100	3.9	7.0	2.7
16	360	9.4	225		100	3.6	7.2	2.8
17	315	9.3			100	3.2	6.6	2.8
18	300	9.2	240		100	2.8	6.6	2.9
19	265	9.0				5.2	2.8	
20	280	9.0				4.8	2.8	
21	310	9.0				5.4	2.6	
22	315	8.9				5.0	2.6	
23	300	9.1				4.3	2.7	

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 16

Tokyo, Japan (35.7°N , 139.5°E)

June 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}s$	F2-M3000
00	325	9.1						5.1
01	310	9.1						4.6
02	300	8.8						4.0
03	305	8.3						3.2
04	300	8.1						3.0
05	280	8.3	250					2.6
06	290	9.2	250					2.7
07	300	9.8	230					2.7
08	335	9.6	230					2.6
09	375	9.4	220					2.5
10	420	8.2	215					2.5
11	420	9.7	200					2.5
12	410	10.0	240					2.5
13	400	10.1	250	5.8				2.5
14	380	10.0						2.5
15	380	10.0	250					2.6
16	370	9.4	250					2.6
17	350	9.5	250	5.0				2.7
18	320	9.4	245					2.7
19	300	8.8						2.7
20	320	8.7						2.6
21	320	8.7						2.5
22	340	9.0						2.5
23	330	9.1						2.5

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 17

Yamakawa, Japan (31.2°N , 130.6°E)

June 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{F1}$	$\text{f}^{\circ}\text{F1}$	h°E	f°E	$\text{f}^{\circ}\text{Es}$	F2-N3000
00	310	9.8				4.6	2.6	
01	300	10.0				5.0	2.7	
02	290	9.6				4.2	2.7	
03	300	8.6				4.2	2.6	
04	300	8.0				3.4	2.6	
05	300	7.9				3.2	2.6	
06	280	8.8	260		110	2.2	3.6	2.7
07	280	9.5	235		110	2.8	5.0	2.8
08	290	9.2			110	3.5	5.6	2.7
09	350	9.4	230		110	3.6	6.8	2.6
10	395	9.6	210		110	4.0	7.0	2.5
11	410	10.4	240	5.6	120	4.3	7.0	2.4
12	410	10.4	230	5.6			7.5	2.4
13	400	11.0	240	5.6		4.2	6.8	2.5
14	400	10.9	220	5.4		6.9	2.5	
15	390	10.9	230	5.3		4.2	6.2	2.6
16	380	11.0	230	5.0	110	3.6	5.8	2.6
17	350	10.6			110	3.5	5.4	2.6
18	300	10.6			120	3.0	5.5	2.7
19	290	9.9			110	2.2	4.9	2.8
20	300	9.4					5.0	2.7
21	320	9.2					4.0	2.5
22	320	9.5					3.6	2.5
23	330	9.6					3.8	2.5

Time: 135.0°E .

Sweep: 0.6 Mc to 18.5 Mc in 15 minutes, manual operation.

Table 18

Chungking, China (29.4°N , 106.8°E)

June 1949

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{F1}$	$\text{f}^{\circ}\text{F1}$	h°E	f°E	$\text{f}^{\circ}\text{Es}$	F2-N3000
00	320				10.3			4.5
01	300				9.7			4.9
02	300				8.8			4.2
03	300				8.2			4.0
04	290				8.0			2.8
05	265				8.2			2.5
06	250				9.0			3.2
07	270				9.6	240		2.6
08	300				10.0	240		2.5
09	300				10.1	240		2.5
10	360				10.5	235		2.4
11	420				11.2	220	6.4	2.4
12	400				11.9	240	6.3	2.5
13	390				12.4	215	6.2	2.5
14	370				13.0	220	6.2	2.5
15	360				13.2	220	5.8	2.5
16	340				12.9	240	5.6	2.5
17	320				12.4	240	9.0	2.6
18	310				12.2	280		(2.6)
19	290				11.7			4.5
20	290				11.0			2.6
21	305				10.2			4.5
22	300				10.1			2.5
23	310				10.2			4.3

Time: 105.0°E .

Sweep: 1.7 Mc to 20.0 Mc in 15 minutes, manual operation.

Table 19

Brisbane, Australia (27.5°S , 153.0°E)

June 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{F1}$	$\text{f}^{\circ}\text{F1}$	h°E	f°E	$\text{f}^{\circ}\text{Es}$	F2-N3000
00	250	4.9						2.8
01	270	4.7						2.8
02	260	4.8			2.0			2.8
03	270	4.9			2.0			2.8
04	260	4.9			2.0			2.8
05	250	4.7			2.0			2.8
06	240	4.6						3.0
07	220	7.6			145	2.2	3.3	
08	230	10.0			100	2.8	3.4	
09	230	11.0	220		100	3.4	3.3	
10	240	11.4	220		100	3.6	3.3	
11	240	10.9	210		100	3.7	3.1	
12	250	10.6	220	4.5	100	3.7	3.0	
13	250	10.3	210	5.2	100	3.6	3.0	
14	240	10.3	210	4.4	110	3.5	2.9	
15	250	10.4	220		110	3.2	2.1	2.9
16	240	10.0			110	2.7	2.0	3.0
17	230	9.5				2.5	3.0	
18	215	8.1				1.8	3.0	
19	230	7.0				1.7	3.0	
20	235	6.0						3.0
21	240	5.8						2.9
22	240	5.5						2.9
23	250	4.8						2.9

Time: 150.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 20

Watheroo, W. Australia (30.3°S , 115.9°E)

June 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{F1}$	$\text{f}^{\circ}\text{F1}$	h°E	f°E	$\text{f}^{\circ}\text{Es}$	F2-N3000
00	270				3.8			3.0
01	275				4.0			3.0
02	262				4.0			2.8
03	268				4.0			2.9
04	250				4.2			2.8
05	235				3.9			2.9
06	228				3.5			3.0
07	240				5.4			3.2
08	230				8.4			3.3
09	235				10.4			3.2
10	240				10.8	235	4.6	3.2
11	240				10.8	232	4.7	3.8
12	262				10.9	230	4.8	3.9
13	255				10.8	230	4.8	3.0
14	248				11.0	228	4.6	3.0
15	250				11.1	232	4.7	3.2
16	240				10.6			2.9
17	225				10.5			1.9
18	212				8.9			3.0
19	220				7.0			3.1
20	225				5.4			3.1
21	240				4.3			3.0
22	245				4.0			2.9
23	260				3.8			2.8

Time: 120.0°E .

Sweep: 16.0 Mc to 0.5 Mc in 15 minutes, automatic operation.

Table 21

Canberra, Australia (35.3°S , 149.0°E)

June 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{Fl}$	$\text{f}^{\circ}\text{Fl}$	h°E	f°E	fEs	F2-M3000
00	260	4.7				2.6	2.6	
01	275	4.7				2.7	2.7	
02	278	4.6				2.5	2.7	
03	280	4.6				2.5	2.7	
04	280	4.9				3.3	2.8	
05	240	4.7				2.8	2.9	
06	235	4.1				2.7	2.9	
07	240	5.6			1.7	2.6	3.0	
08	220	8.5			100	2.5	3.5	3.2
09	230	10.5			100	3.0	3.5	3.2
10	225	11.4			100	3.3	3.5	3.2
11	222	11.6			100	3.5	3.6	3.1
12	220	11.6			100	3.5	4.9	3.1
13	230	11.3	200	4.5	100	3.5	4.7	3.0
14	230	11.4		4.0	100	3.4	4.4	3.0
15	230	11.0			100	3.0	4.4	3.0
16	230	11.0			100	2.5	3.9	3.0
17	220	10.2				1.7	3.5	3.0
18	215	8.6				3.5	3.0	
19	220	7.4				3.4	3.0	
20	225	6.0				2.8	3.0	
21	240	5.4				2.9	2.9	
22	250	4.8				2.6	2.8	
23	250	4.6				2.6	2.7	

Time: 150.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 22

Christchurch, New Zealand (43.5°S , 172.7°E)

June 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{Fl}$	$\text{f}^{\circ}\text{Fl}$	h°E	f°E	fEs	F2-M3000
00	280	4.4				4.4		2.7
01	300	4.7				4.2		2.8
02	300	4.6				4.3		2.5
03	280	4.6				4.4		2.8
04	280	4.9				4.3		2.6
05	240	4.7				4.1		2.9
06	240	4.1				3.6		3.0
07	250	5.6				4.2	(1.2)	3.0
08	230	8.5	100	2.5	3.5	3.2	1.8	3.2
09	230	10.5	100	3.0	3.5	3.2	2.5	3.2
10	225	11.4	100	3.3	3.5	3.2	2.8	3.3
11	222	11.6	100	3.5	3.6	3.1	3.1	3.1
12	220	11.6	100	3.5	4.9	3.1	3.2	3.0
13	230	11.3	200	4.5	100	3.5	4.7	3.0
14	230	11.4		4.0	100	3.4	4.4	3.0
15	230	11.0			100	3.0	4.4	3.0
16	230	11.0			100	2.5	3.9	3.0
17	220	10.2				1.7	3.5	3.0
18	215	8.6				3.5	3.0	
19	220	7.4				3.4	3.0	
20	225	6.0				2.8	3.0	
21	240	5.4				2.9	2.8	
22	250	4.8				2.6	2.8	
23	250	4.6				2.6	2.7	

Time: 172.5°E .

Sweep: 1.0 Mc to 13.0 Mc.

Table 23*

Fraserburgh, Scotland (57.6°N , 2.1°W)

May 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{Fl}$	$\text{f}^{\circ}\text{Fl}$	h°E	f°E	fEs	F2-M3000
00	340	(6.5)					2.4	
01	350	(6.0)					2.3	
02	350	5.4		4.7*			2.3	
03	340	(5.4)					2.5	
04	320	5.4	320#	3.2#			2.5	
05	320	5.6	280	3.8	140	2.3	2.5	
06	330	6.0	250	4.4	110	2.8	3.6	2.6
07	380	6.3	230	4.8	110	3.2	4.0	2.6
08	6.7	230	5.1	110	3.4	4.1	2.5	
09	6.8	220	5.3	110	3.6	4.2	2.4	
10	6.8	220	5.5	100	3.8	4.8	2.5	
11	420	7.0	230	5.6	110	3.9	4.4	2.5
12	460	7.1	220	5.7	110	3.9	4.1	2.4
13	460	7.0	220	5.6	100	3.9	4.0	2.4
14	440	7.1	220	5.6	100	3.9	4.3	2.4
15	390	7.1	240	5.5	100	3.2	4.2	2.4
16	370	7.2	230	5.3	100	3.5	4.2	2.5
17	310	7.3	240	4.9	110	3.2	4.0	2.5
18	250	7.4			110	2.8	3.8	2.6
19	260	(7.3)			140	2.5	2.6	
20	280	(7.2)		120#	2.8#		2.6	
21	280	(7.4)					2.6	
22	310	(7.3)					2.4	
23	330	(7.0)					2.4	

Time: Local.

Sweep: 2.2 Mc to 16.0 Mc in 1 minute.

*Average values except for $\text{f}^{\circ}\text{F2}$ and fEs , which are median values.

#One or two observations only.

Table 24*

Slough, England (51.5°N , 0.6°W)

May 1948

Time	$\text{h}^{\circ}\text{F2}$	$\text{f}^{\circ}\text{F2}$	$\text{h}^{\circ}\text{Fl}$	$\text{f}^{\circ}\text{Fl}$	h°E	f°E	fEs	F2-M3000
00	317	6.9						3.2
01	316	6.6						2.6
02	318	6.3						2.3
03	324	6.0						2.4
04	327	5.8	316	3.3	121	1.6	3.4	2.5
05	328	6.2	270	3.9	116	2.1	4.6	2.5
06	366	6.4	248	4.5	113	2.8	4.8	2.6
07	394	6.7	233	5.0	111	3.2	5.8	2.6
08	424	7.1	234	5.5	109	3.5	5.1	2.5
09	438	7.7	233	5.6	110	3.7	6.9	2.5
10	434	8.2	236	5.8	109	3.9	7.0	2.5
11	423	8.0	232	5.9	108	4.0	7.2	2.5
12	422	8.1	238	6.0	109	4.0	6.8	2.5
13	426	8.2	234	5.9	109	4.0	6.0	2.5
14	401	8.1	237	5.9	109	3.8	5.3	2.5
15	390	8.1	233	5.8	109	3.7	5.4	2.5
16	370	8.2	239	5.6	110	3.5	4.9	2.5
17	319	8.2	243	5.1	111	3.2		2.6
18	283	8.1	259	4.6	115	2.7		2.6
19	271	8.0			127	2.1	3.5	2.6
20	272	7.8				1.7	3.9	2.6
21	284	8.0					2.8	2.5
22	296	7.7					2.6	2.4
23	304	7.4					1.9	2.4

Time: Local.

Sweep: 0.5 Mc to 16.5 Mc in 5 minutes.

*Average values except for $\text{f}^{\circ}\text{F2}$ and fEs , which are median values.

Table 25

Lanchow, China (36.1°N , 103.8°E)

May 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}S$	$F2-\text{M3000}$
00	420	9.8			5.1	2.2		
01	400	9.6			4.4	2.2		
02	400	9.0			4.2	2.2		
03	400	7.8			4.2	2.2		
04	420	7.4			3.5	2.2		
05	400	7.6			3.9	2.2		
06	380	9.2			3.8	2.3		
07	360	10.1	330		150	3.4	4.4	2.5
08	400	11.0	320		150	3.7	5.4	2.3
09	440	11.5	320		155	4.1	5.6	2.2
10	480	12.2	340	7.0		5.4	2.2	
11	480	12.5	320	6.6		5.0	2.2	
12	480	13.0	320	7.0		5.3	2.2	
13	480	13.0	350	7.0		6.0	2.2	
14	510	13.0	350	6.8		5.2	2.2	
15	480	12.5	340	8.4		4.9	2.2	
16	440	12.2	330	6.5	150	4.0	5.0	2.3
17	440	12.0	320	6.1	155	3.8	5.0	2.2
18	400	12.0	340		155	3.0	5.0	2.3
19	360	11.5				4.5	2.3	
20	(365)	(10.6)				(3.2)	(2.2)	
21	405	10.0				4.6	2.1	
22	420	9.8				4.6	2.2	
23	440	9.8				5.6	2.1	

Time: 105.0°E .

Sweep: 2.4 Mc to 16.0 Mc in 15 minutes, manual operation.

Table 26

Nanking, China (32.1°N , 119.0°E)

May 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}S$	$F2-\text{M3000}$
00								
01								
02								
03								
04		280						
05		280						
06		250						
07		280						
08		300						
09		320						
10		380						
11		400						
12		390						
13		370						
14		380						
15		380						
16		350						
17		340						
18		260						
19		280						
20		290						
21		280						
22								
23								

Time: 120.0°E .

Sweep: 1.7 Mc to 15.0 Mc in 15 minutes, manual operation.

Table 27

Delhi, India (28.6°N , 77.1°E)

May 1948

Time	*	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}S$	$F2-\text{M3000}$
00		480	10.6			2.4		
01		470	10.3					
02	(480)	10.2						
03								
04		480	8.4			2.4		
05		440	8.8					
06		400	9.8					
07		400	10.6					
08		440	11.3			2.4		
09		480	11.8					
10		520	12.1					
11		520	13.4					
12		520	13.4			2.1		
13		540	(13.5)					
14		530	(13.8)					
15		520	(13.5)					
16		520	13.0			2.2		
17		510	12.8					
18								
19								
20		490	11.4			2.3		
21		520	11.0					
22		520	10.5					
23		520	10.5					

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 28

Bombay, India (19.0°N , 73.0°E)

May 1948

Time	*	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}S$	$F2-\text{M3000}$
00								2.5
01								
02								
03								
04								2.6
05								
06								
07		330	11.0					
08		390	11.3					2.7
09		480	12.2					
10		510	13.0					
11		540	13.8					
12		555	14.0					
13		(570)						
14								
15		(570)	14.3					
16		540	14.5					2.3
17		510	14.5					
18		480	14.6					
19		540	14.3					
20		540	13.9					2.4
21		510	13.5					
22		480	13.4					
23		(480)	(13.4)					

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 29

Madras, India (13.0°N , 80.2°E)

May 1948

Time	•	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F_2-\text{N}3000$
00								
01								
02								
03								
04								
05								
06	420	(10.0)						
07	420	11.0						
08	540	12.2						
09	600	12.8						
10	600	12.6						
11	600	12.0						
12	630	11.8						
13	660	12.2						
14	660	12.4						
15	660	12.7						
16	660	12.8						
17	630	12.9						
18	600	12.9						
19	600	(12.3)						
20	(540)	(11.5)						
21		(11.0)						
22		(11.1)						
23								

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 30

Rarotonga I. (21.3°S , 159.8°W)

May 1948

Time	$h^{\circ}\text{F2}$	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F_2-\text{N}3000$
00								
01								
02								
03								
04								
05								
06								
07								
08								
09	240	14.3						
10	250	14.2						
11	250	14.0	250	7.5	110	3.3	3.8	3.0
12	250	14.2	235	(6.9)	110	3.9	4.2	2.8
13	305	14.0	250	7.2	105	3.8	5.1	2.7
14	320	14.2	250	7.0	110	3.7	4.4	2.6
15	340	14.0	250	6.6	110	3.5	5.2	2.6
16								
17								
18								
19								
20								
21								
22								
23								

Time: 157.5°E .

Sweep: 2.0 Mc to 16.0 Mc, manual operation.

Table 31

Brisbane, Australia (27.5°S , 153.0°E)

May 1948

Time	$h^{\circ}\text{F2}$	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F_2-\text{N}3000$
00	250	6.5			2.0	2.8		
01	250	8.4			2.4	2.8		
02	260	6.0			2.0	2.8		
03	260	5.6			2.8	2.8		
04	250	5.5			2.0	2.8		
05	250	5.2				2.8		
06	240	5.9				3.0		
07	230	9.4	180	3.4		3.2		
08	230	12.0	108	3.1	2.6	3.2		
09	235	13.0	220	100	3.5	3.2		
10	240	13.0	220	100	3.6	3.1		
11	240	12.8	220	100	3.8	2.9		
12	240	12.3	220	6.5	100	3.8	2.9	
13	270	12.4	220	6.6	110	3.8	3.1	2.9
14	250	12.4	230	100	3.8	2.9		
15	250	12.2	230	110	3.4	2.6	2.9	
16	240	11.9			110	2.9	2.7	2.9
17	240	11.2			130	2.1	3.0	2.9
18	230	10.0				2.1	2.9	
19	240	9.7				2.0	2.9	
20	250	8.3				1.6	2.9	
21	240	7.5				2.0	2.9	
22	245	7.0				2.8	2.9	
23	250	6.5				2.8		

Time: 150.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 32

Canberra, Australia (35.3°S , 149.0°E)

May 1948

Time	$h^{\circ}\text{F2}$	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F_2-\text{N}3000$
00								
01	260	6.0						
02	270	5.9						
03	272	5.7						
04	272	5.5						
05	250	5.5						
06	240	4.9						
07	240	7.2						
08	230	10.9						
09	230	12.2						
10	230	13.0						
11	225	13.1						
12	220	12.8						
13	230	13.0						
14	230	12.9						
15	240	12.3						
16	240	12.0						
17	228	11.4						
18	220	10.2						
19	240	8.6						
20	240	8.0						
21	240	7.0						
22	250	6.8						
23	260	6.4						

Time: 150.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 33

Hobart, Tasmania (42.8° S, 147.4° E)

May 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$
00	265	6.0			2.4	2.7		
01	275	5.5			2.5	2.7		
02	270	5.6			2.5	2.7		
03	260	5.1			3.0	2.7		
04	262	5.1			2.7	2.8		
05	252	4.8			3.5	2.8		
06	250	4.6			2.4	2.8		
07	245	5.5			2.7	3.0		
08	230	9.0			105	2.4	2.7	3.5
09	230	(10.5)			100	3.0	3.0	3.6
10	220	10.8			105	3.2	3.4	3.6
11	225	11.0	225		100	3.5	3.4	3.6
12	235	(11.0)	210		100	3.5	3.5	(3.6)
13	235	(11.0)	218		100	3.5	2.9	(3.5)
14	225	(11.0)			102	3.3	3.9	3.5
15	228	10.5			100	3.0	3.6	3.4
16	225	10.5			105	2.5	3.1	(3.5)
17	220	(10.5)					2.8	(3.4)
18	215	9.5					2.5	3.2
19	220	8.6					2.2	3.2
20	240	7.5						3.1
21	248	6.8						2.8
22	250	6.4						2.4
23	258	5.8						2.5

Time: 150.0° E.

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 55 seconds.

Table 34*

Fraserburgh, Scotland (57.6° N, 2.1° W)

April 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$	
00					340	(5.5)			2.4
01					350	(4.8)			2.4
02					365	4.4			2.4
03					365	4.6			2.3
04					325	4.4			2.4
05					295	5.0			2.3
06					265	5.9	100#	(2.3)†	2.3
07					270	6.7	260#	4.6#	2.7
08					275	7.2	235	5.2	2.7
09					260	7.8	225	5.4	2.6
10					275	8.6	210	5.6	2.7
11					285	8.8	210	5.5	2.7
12					270	8.8	225	5.7	2.6
13					280	8.9	220	5.6	2.6
14					265	9.3	225	5.7	2.6
15					240	9.2	230#	5.3#	2.6
16					235	8.6	225	(3.8)‡	2.6
17					240	9.0			2.7
18					250	9.0			2.7
19					260	(8.6)			2.7
20					265	(8.2)			2.6
21					280	(8.0)			2.5
22					300	(7.3)			2.6
23					320	(6.5)			2.4

Time: Local.

Sweep: 2.2 Mc to 16.0 Mc in 1 minute.

*Average values except for $f^{\circ}F2$ and $f^{\circ}Es$, which are median values.

†One or two values only.

Table 35*

Slough, England (51.5° N, 0.6° W)

April 1948

Time	$h^{\circ}F2$	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$
00	299	7.1				1.7	2.4	
01	303	6.7				2.5	2.4	
02	305	6.3				2.6	2.4	
03	306	5.7				2.5	2.4	
04	298	5.3				2.6	2.4	
05	288	5.5	290	3.1#	135	1.6	3.7	2.5
06	278	6.4	250	4.3	121	2.3	4.3	2.7
07	287	7.2	239	4.7	113	2.9	4.8	2.7
08		8.2	228	5.1	111	3.3	4.6	2.7
09	309	8.7	231	5.3	111	3.6	4.8	2.6
10	323	9.3	231	5.8	110	3.7	4.8	2.6
11	331	10.0	222	5.8	110	3.8		2.6
12	331	10.3	229	6.0	110	3.9	4.8	2.6
13	332	10.3	229	6.0	109	3.9	4.0	2.6
14	330	10.0	230	6.0	109	3.8	4.6	2.6
15	311	9.7	231	5.5	110	3.6	3.9	2.6
16	295	9.5	232	5.3	111	3.3	3.5	2.6
17	260	9.2	240	4.7	113	2.9		2.7
18	255	9.1			120	2.3		2.7
19	254	9.2			139	1.7	1.9	2.7
20	252	8.7				1.6	2.6	
21	261	8.2					2.5	
22	279	7.8					2.5	
23	290	7.5					2.5	2.4

Time: Local.

Sweep: 0.5 Mc to 16.5 Mc in 5 minutes.

*Average values except for $f^{\circ}F2$ and $f^{\circ}Es$, which are median values.

†One or two values only.

Table 36

Delhi, India (28.6° N, 77.1° E)

April 1948

Time	*	$f^{\circ}F2$	$h^{\circ}F1$	$f^{\circ}F1$	$h^{\circ}E$	$f^{\circ}E$	$f^{\circ}Es$	$F2-M3000$	
00		420	11.1						2.6
01		450	10.3						
02		(390)	(8.3)						
03		420	8.7						
04		390	8.2						
05		390	8.1						
06		360	9.6						
07		360	11.0						
08		390	11.6						
09		420	12.4						
10		450	13.1						
11		450	(13.5)						
12		450	(13.9)						
13		450	(13.7)						
14		450	(13.6)						
15		465	(13.6)						
16		420	(13.6)						
17		450	(13.2)						
18									
19									
20		420	(12.2)						
21		420	(11.7)						
22		420	11.5						
23		420	11.5						

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}F2$.

**Average values; other columns, median values.

Table 37

Bombay, India (19.0°N , 73.0°E)

April 1948

Time	*	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$	**
00									
01									
02									
03									
04									
05									
06									
07		330	10.7						
08		390	12.1						
09		435	13.2						
10		495	14.0						
11		(480)	(14.4)						
12		(14.6)							
13		(14.7)							
14		(14.8)							
15		(14.9)							
16		(15.1)							
17		(480)	(14.9)						
18		(480)	(14.7)						
19		510	(14.7)						
20		510	(14.5)						
21		(14.7)							
22									
23									

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 38

Madras, India (13.0°N , 80.2°E)

April 1948

Time	*	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$	**
00									
01									
02									
03									
04									
05									
06									
07		420	10.2						
08		480	12.6						2.5
09		540	13.1						
10		540	13.0						
11		600	12.7						
12		600	12.6						
13		600	12.6						
14		600	12.6						
15		600	13.2						
16		600	13.9						
17		600	13.4						
18		600	13.3						
19		540	11.9						
20			(10.5)						
21			(10.3)						
22			(10.0)						
23									

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 39*

Falkland Is. (51.7°S , 57.8°W)

April 1948

Time	$h^{\circ}\text{F2}$	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$	**
00	372	5.6				2.4			
01	370	5.5				2.4			
02	374	5.4				2.4			
03	366	5.2				2.4			
04	354	5.3				2.4			
05	341	5.0				2.4			
06	302	5.5				2.5			
07	245	(8.3)				2.5			
08	234	11.0			117	2.7			
09	238	(13.6)			119	2.9			
10	237	(14.4)			113#	3.1			
11	240	14.5			115#	3.3#			
12	241	(14.1)			117	3.3			
13	242	13.1			123#	3.1#			
14	247	12.4			124	3.1			
15	249	11.8			125#	2.9			
16	245	11.1				2.4#			
17	243	9.8				3.1			
18	246	8.2				3.1			
19	248	6.6				3.2			
20	263	5.6				2.8			
21	296	5.4				2.7			
22	332	5.3				2.6			
23	358	5.5				2.4			

Time: Local.

Sweep: 3.2 Mc to 16.0 Mc in 1 minute.

*Average values except for $f^{\circ}\text{F2}$ which are median values.

#One or two values only.

Table 40

Delhi, India (28.6°N , 77.1°E)

March 1948

Time	*	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$	**
00		390	7.2						
01		360	7.0						2.6
02		400	6.6						
03		(360)	(5.8)						
04		390	5.2						
05		390	5.0						
06		330	5.7						
07		300	8.7						
08		330	10.6						
09		330	11.4						
10		360	12.4						
11		360	13.0						
12		390	13.8						
13		390	(14.2)						
14		390	(14.2)						
15		420	(14.0)						
16		390	(14.0)						
17		390	(14.0)						
18									
19									
20		360	11.6						
21		390	10.6						
22		390	8.8						
23		390	7.8						

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 41

Bombay, India (19.0°N , 73.0°E)

March 1948

Time	**							
	*	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$
00								2.7
01								
02								
03								
04								2.9
05								
06	(330)	(6.4)						
07	330	9.2						
08	330	11.3						3.0
09	330	12.6						
10	420	14.2						
11	(450)	(14.7)						
12		(14.7)						
13		(14.9)						
14		(14.8)						
15	(480)	(14.9)						
16	(450)	(14.9)						2.5
17	(450)	(15.1)						
18	480	(15.0)						
19	(480)	(15.1)						
20		(15.1)						2.7
21		(14.8)						
22		(14.2)						
23		(12.2)						

Table 42

Madras, India (13.0°N , 80.2°E)

March 1948

Time	**							
	*	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$
00								
01								
02								
03								
04								
05								
06								
07		420						
08		480						2.5
09		540						
10		540						
11		540						
12		600						
13		600						2.2
14		600						
15		600						
16		600						2.1
17		600						
18		600						
19		600						
20		(570)	(11.2)					2.6
21			(11.0)					
22			(10.5)					
23								

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 $f^{\circ}\text{F2}$.

**Average values; other columns, median values.

Table 43*

Falkland Is. (51.7°S , 57.8°W)

March 1948

Time	$h^{\circ}\text{F2}$	$f^{\circ}\text{F2}$	$h^{\circ}\text{F1}$	$f^{\circ}\text{F1}$	$h^{\circ}\text{E}$	$f^{\circ}\text{E}$	$f^{\circ}\text{Es}$	$F2-\text{M3000}$
00	336	6.9						2.5
01	343	6.6						2.5
02	341	6.4						2.5
03	331	6.2						2.5
04	330	6.0						2.5
05	322	5.6						2.5
06	255	7.1						3.0
07	245	8.3						3.1
08	245	9.8						3.1
09	258	11.4	255#	5.1#	112	2.9	3.4	3.1
10	255	11.9	233#	5.5#	110	3.3	4.2	3.0
11	245	12.4						3.0
12	256	12.6	245#	5.2#	111	3.3	4.6	3.0
13	257	12.5	243#	5.7#	111	3.2	4.0	3.0
14	247	11.8						3.0
15	249	10.9						3.1
16	250	10.4						3.1
17	251	10.0						3.2
18	247	9.6						3.2
19	247	8.2						3.1
20	258	7.2						3.0
21	287	6.5						2.8
22	308	6.6						2.8
23	328	6.7						2.5

Time: Local.

Sweep: 2.2 Mc to 16.0 Mc in 1 minute.

*Average values except $f^{\circ}\text{F2}$ and $f^{\circ}\text{Es}$, which are median values.

#One or two values only.

National Bureau of Standards
(Institution)

J.M.C.

E.J.W.

TABLE 44

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

hF_2 , km
(Characteristic)
Observed at Washington, D.C.
(Unit)

August 1948
(Month)

Lat. 39°0' N., long 77°5' W

Mean Time

Day	75°W												75°W												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	2000 ^K	3000 ^A	290 ^K	240 ^K	320 ^K	300 ^K	240 ^K	550 ^K	650 ^K	650 ^K	700 ^K	G	K	570 ^K	520 ^K	420 ^K	450 ^K	420 ^K	350 ^K	250 ^K	250 ^K	250 ^K	270 ^K		
2	250 ^K	250 ^K	280 ^K	260 ^K	270 ^K	280 ^K	250 ^K	220 ^K	480 ^K	510 ^K	560 ^K	G	K	610 ^K	540 ^K	460 ^K	460 ^K	280 ^K	250	240	250	280	260		
3	(13.0) ^A	26.0	28.0	29.0	28.0	26.0	24.0	23.0	28.0	26.0	33.0	36.0	42.0	50.0	53.0	50.0	43.0	40.0	41.0	300	260	240	(32.0) ^A	26.0	
4	28.0	27.0	25.0	24.0	25.0	24.0	23.0	21.0	23.0	23.0	33.0	33.0	42.0	45.0	47.0	40.0	35.0	27.0	25.0	24.0	28.0	27.0	27.0	27.0	
5	28.0	27.0	25.0	24.0	23.0	24.0	22.0	20.0	30.0	35.0	39.0	42.0	45.0	50.0	50.0	40.0	37.0	36.0	300	250	250	(2.40) ^A	25.0	26.0	
6	27.0	24.0	26.0	25.0	25.0	24.0	28.0	28.0	28.0	35.0	35.0	38.0	37.0	33.0	34.0	34.0	34.0	34.0	(2.30) ^A	(2.50) ^A	230	240	(2.50) ^A		
7	28.0	3.00	2.90	3.00	2.80	2.70	3.80	5.00	5.00	4.80	5.00	5.90	6.30	A	K	6.30	5.00	4.50	4.50	4.00	3.30	3.30	2.70	2.70	2.70
8	26.0 ^K	4.00 ^K	3.70 ^K	N ^C	(4.30) ^K	N ^C	2.80 ^K	G	K	G	K	G	K	G	K	G	K	520 ^K	520 ^K	300	280	B	K	360 ^K	
9	38.0 ^K	3.60 ^K	3.20 ^K	3.00 ^K	3.10 ^K	3.00 ^K	2.80 ^K	G	K	G	K	G	K	G	K	6.00	4.80	4.70	380 ^K	300	250	210	230	270 ^K	
10	25.0 ^K	2.80 ^K	A ^K	(2.50) ^A	310 ^K	310 ^K	32.0 ^K	G	K	G	K	G	K	G	K	G	K	4.80	4.80	4.30	4.20	350	250	270 ^K	
11	2.50 ^K	(3.01) ^K	(3.80) ^K	F ^K	(4.00) ^K	F ^K	(3.00) ^K	2.10 ^M	G	K	4.50	K	4.00	K	3.70	K	3.70	3.70	3.70	3.70	3.70	3.70	3.70	(2.50) ^A	
12	2.50 ^K	(3.00) ^A	2.80 ^K	(3.00) ^K	(3.00) ^K	(3.00) ^K	3.80 ^K	4.30 ^K	G	K	G	K	G	K	G	K	C	B	K	4.80	4.60	3.80	2.70	2.50 ^K	
13	3.10 ^K	3.00 ^K	A ^K	(2.80) ^K	2.90 ^K	A ^K	2.30 ^K	3.30 ^K	4.30 ^K	3.60 ^K	5.30 ^K	G	K	4.30	K	C	K	3.70	K	3.70	3.70	2.80	(2.30) ^A	A	
14	A	A	A	2.60	2.50	2.50	2.50	2.20	2.10	3.00	4.70	4.70	N	S	5.00	5.00	4.70	4.70	3.70	3.70	3.70	3.70	2.30	2.40	2.50
15	2.60	2.50	2.70	A	2.60	2.50	2.30	2.80	2.80	3.20	3.20	3.70	C	3.60	3.60	3.60	3.60	3.60	3.30	3.30	2.50	2.30	2.20	2.00	(2.50) ^A
16	2.60	2.60	2.50	(2.50) ^A	2.40	2.50	(2.40) ^A	3.10	3.00	3.40	4.50	4.30	4.00	3.60	3.80	3.80	3.50	3.50	3.30	2.40	2.40	2.30	2.30	2.50	2.50
17	(2.50) ^A	(2.80) ^A	2.60	2.50	2.50	2.50	(2.30) ^A	(2.30) ^A	3.20	3.40	4.10	(4.44) ^C	3.60	4.00	3.90	3.90	3.90	(2.44) ^A	2.50	2.50	2.30	2.30	(2.50) ^A	2.40	
18	2.60	2.50	2.50	2.50	3.00	2.30	2.20	3.00	C	C	4.60	4.70	4.50	4.60	4.20	4.20	3.60	2.80	2.80	2.80	2.80	2.80	2.80		
19	2.70	2.70	2.60	2.60	2.40	2.70	2.70	2.70	2.70	2.70	3.50	3.50	3.50	3.50	3.50	3.50	3.50	3.50	3.50	3.50	2.60	2.60	2.70		
20	2.80	2.80	2.70	2.50	2.40	2.40	2.30	2.30	3.50	3.20	4.50	3.90	(4.00) ^C	4.20	(4.60) ^C	4.00	(3.40) ^C	2.50	2.50	2.40	2.40	2.80	2.80	2.80	
21	3.30	3.00	2.80	2.70	2.70	2.70	2.70	2.70	2.70	2.70	3.30	5.60	4.00	3.70	3.90	4.00	3.50	3.00	2.60	2.50	2.20	2.30	2.60		
22	2.50	3.00	3.00	2.70	2.70	2.70	2.70	2.40	C	C	3.30	3.70	4.00	3.70	3.70	3.50	3.50	3.30	2.50	2.40	2.50	2.50	2.70		
23	3.00	2.50	2.70	2.60	2.60	2.70	2.70	3.10	3.00	4.40	C	3.80	3.70	3.50	3.50	3.50	3.40	3.00	2.30	2.50	2.30	2.60	2.70		
24	2.70	2.70	2.60	2.50	2.70	2.80	2.50	2.30	3.10	3.00	4.30	4.70	C	4.30	3.80	3.90	3.60	3.10	2.40	2.40	2.40	2.50	2.70		
25	2.70	2.50	2.30	2.40	2.50	2.50	2.40	2.50	3.00	3.00	2.80	(3.00) ^C	3.10	3.50	3.00	3.20	3.00	2.80	2.30	2.30	2.30	2.30	2.50		
26	2.90	2.60	2.60	2.50	2.40	2.50	2.90	C	C	C	C	3.20	3.20	3.40	C	3.30	2.70	(2.70) ^A	2.30	A	(2.30) ^A	(2.40) ^A	A		
27	A	A	(3.10) ^A	2.80	(2.70) ^A	2.30	F	2.50	3.00	H	C	2.90	H	A	C	(3.50) ^C	3.10	3.30	2.60	2.20	2.20	2.20	2.20	2.50	
28	2.50	2.70	2.70	2.60	2.70	2.50	2.20	2.60	2.60	2.60	3.40	3.40	3.20	3.00	2.70	H	2.80	2.30	2.30	2.40	2.40	2.40	2.40		
29	2.60	2.60	2.50	(2.60) ^A	2.20	K	2.50	2.60	2.70	3.30	K	4.50	K	5.90	K	5.70	K	4.50	K	4.00	A	(2.60) ^A	2.20	2.60	
30	3.10	2.90	3.00	3.00	3.00	K	2.60	K	2.30	3.00	K	4.30	K	5.60	K	4.90	K	5.00	K	4.70	K	4.30	3.60	2.40	(2.50) ^A
31	2.80	K	(2.60) ^A	2.70	3.00	K	2.70	3.50	3.90	K	4.20	K	6.20	K	4.10	K	3.80	K	4.00	K	3.50	3.00	2.30	2.60	2.90
Median	2.70	2.70	2.60	2.70	2.70	2.70	2.40	2.70	3.10	3.55	3.90	4.50	4.35	4.30	4.20	4.00	3.60	3.35	2.60	2.50	2.40	2.50	2.60		
Count	29	29	28	29	30	29	29	31	29	28	27	28	29	29	29	29	29	30	31	30	31	30	29	29	

Sweep 10 Mc sec 0.25 mm

Automatic

Manual

TABLE 45
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

f_{eff}^2 , Mc (Month) August, 1948
Mc (Unit) Observed at Washington, D.C.

Lat 39°0'N Long 77°5'W

National Bureau of Standards
(Institution)
J.M.S., J.M.C.
Scaled by: E.J.W.
Calculated by: M.C.E.
G.G.H.

Day	75° W Mean Time														21	22	23				
	00	01	02	03	04	05	06	07	08	09	10	11	12	13							
1	F	K	K(4.6)F	(3.9)F	3.7	3.1	4.1	K	4.9	K	5.0	K	4.9	K	5.2	K	G	K			
2	6.0	F	5.3	K	5.1	4.9	K	4.5	K	4.3	K	4.7	K	4.9	K	5.5	K	5.8	K		
3	5.2	4.9	4.6	4.4	4.0	4.1	5.5	6.9	7.2	7.5	7.3	6.5	(5.9)F	6.0	5.7	K	6.3	K			
4	6.1	5.5	F	5.6	F	4.9	F	4.7	F	4.5	(5.0)S	7.0	(6.6)F	7.2	6.7	K	5.9	K			
5	(5.7)F	5.5	F	5.3	F	(5.0)F	4.3	4.2	6.5	7.4	7.6	7.5	6.9	(6.1)F	6.4	6.6	K	5.5	K		
6	5.5	F	5.4	F	5.1	F	4.9	F	4.3	F	4.1	F	5.7	7.0	6.7	K	6.7	(6.2)F			
7	5.8	5.6	5.7	5.5	6.1	F	4.3	F	4.5	K	4.6	K	5.1	K	5.3	K	(5.4)F	(5.9)F			
8	5.9	F	K	F	K	N	C	2.9	K	[3]C	N	3.3	F	G	K	G	K	G	K		
9	(2.7)F	(2.7)F	2.8	F	2.3	F	2.3	F	3.7	F	G	K	G	K	G	K	(3.6)F	2.3	F		
10	(6.5)F	F	K	F	K	(2.8)F	3.1	F	3.1	F	3.8	F	G	K	C	K	5.3	K	7.9	K	
11	(3.6)F	(2.3)F	E	(1.9)F	K	(1.7)F	F	K	(2.5)F	3.9	K	G	K	5.3	K	5.8	K	5.6	F		
12	5.5	F	4.9	F	(4.2)F	2.8	F	(1.7)F	2.2	F	(4.0)K	4.9	F	G	K	G	K	5.6	K		
13	3.9	F	3.8	F	(3.7)F	2.9	F	1.1	K	(2.7)F	(4.7)F	5.7	K	5.5	K	5.7	K	5.7	K		
14	5.3	F	5.5	F	4.8	F	4.5	F	4.3	F	5.6	F	(5.6)F	5.9	[5.9]F	5.8	K	(6.3)F	(5.0)F		
15	(4.9)F	4.7	F	4.5	F	4.5	F	4.1	F	4.1	F	5.0	6.0	(6.7)F	(6.7)F	7.2	K	(6.3)F	(6.7)F		
16	5.5	5.5	5.4	F	5.2	F	4.9	F	4.5	S	5.0	F	5.5	F	5.5	K	5.6	F	5.5	F	
17	5.8	6.6	5.5	F	4.9	4.7	F	4.5	F	5.6	F	6.2	F	6.7	F	6.9	F	6.7	F	5.5	F
18	5.7	F	5.7	F	4.7	F	4.0	F	3.8	F	5.7	F	6.3	F	(6.5)F	(6.3)F	7.0	[6.8]F	6.8	F	
19	6.0	(5.8)F	J	5.7	F	5.4	F	5.1	4.9	F	6.1	7.2	8.8	(9.6)F	(9.1)F	8.8	8.7	8.7	(8.1)F	8.2	F
20	7.3	7.1	6.9	6.8	6.7	(6.2)F	5.6	6.8	7.3	8.0	(7.7)F	7.6	8.0	7.6	7.5	7.5	7.5	7.5	7.5	6.2	
21	(5.4)F	C	4.8	(4.7)F	3.9	4.8	5.6	6.3	6.7	(5.8)F	7.1	6.9	7.1	7.4	7.4	7.4	7.4	7.4	7.4	5.3	
22	4.6	F	4.3	F	4.5	4.2	4.1	3.7	5.3	C	6.8	7.0	7.3	7.9	8.2	8.5	8.6	8.7	8.7	5.7	
23	(6.2)F	(5.9)F	=	6.2	6.0	5.3	4.5	5.6	6.8	7.6	7.3	7.0	7.5	7.5	7.8	7.8	7.8	7.8	(6.4)F		
24	(6.1)F	J	5.7	5.7	4.9	4.7	F	4.3	5.4	F	6.3	6.7	(6.7)F	(6.5)F	6.6	(6.8)F	(6.6)F	(6.5)F	(6.5)F		
25	(5.9)F	(6.0)F	J	(5.9)F	(4.6)F	4.5	F	4.0	F	6.0	F	6.9	7.6	7.9	7.7	7.5	7.5	7.5	7.5		
26	5.4	F	5.3	4.9	F	4.9	F	4.5	4.1	(5.9)F	C	C	(8.5)F	8.6	[8.4]F	8.4	8.5	8.9	(8.8)F		
27	A	(5.4)F	J	5.1	F	5.0	F	4.6	F	5.7	6.6	(6.7)F	(6.7)F	7.7	7.9	8.0	8.1	7.9	7.7	(6.4)F	
28	(6.1)F	J	5.8	5.8	5.3	5.1	5.0	5.5	(6.5)F	7.8	7.7	7.9	8.1	8.7	8.7	9.3	9.0	8.9	8.8		
29	(6.3)F	6.3	K	6.5	K	6.2	K	5.2	K	3.1	K	4.9	K	5.0	K	5.9	K	6.4	K		
30	3.3	F	3.9	K	3.6	K	3.3	K	3.1	K	4.5	K	5.3	K	5.5	K	5.6	K	6.7	K	
31	4.0	F	3.9	F	3.6	F	(2.9)F	(2.7)F	4.1	F	5.2	F	5.5	K	5.7	K	6.0	F	6.3	F	
Median	5.7	5.4	5.1	4.9	4.4	4.1	3.4	6.0	6.7	6.4	6.6	6.7	6.8	6.9	7.0	7.1	7.0	6.8	6.7		
Count	29	29	29	30	30	30	31	31	29	29	28	30	31	31	31	31	31	30	30		

Sweep 1.0 Mc to 25.0 Mc in 0.25 min
Manual □ Automatic □

TABLE 46
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

Form adopted June 1946

IONOSPHERIC DATA

National Bureau of Standards
(Institution)
J.J.S., J.M.C.

f^oF2 Mc August, 1948
(Characteristic) (Unit) (Month)

Observed at **Washington, D.C.**
Lat. **39.0°N**, Long **77.5°W**

Scaled by: **E.J.W.** Calculated by: **G.G.H.** **K.L.W.**

75°W

Mean Time

1130 1230 0930 0630 0330 0030 0730 0830 0930 1030 1130 1230 1330 1430 1530 1630 1730 1830 1930 2030 2130 2230 2330

Day	0030	0130	0230	0330	0430	0530	0630	0730	0830	0930	1030	1130	1230	1330	1430	1530	1630	1730	1830	1930	2030	2130	2230	2330	
1	4.5 ^k (5.1) ^k	3.2 ^k (5.1) ^k	-3.4 ^k -3.1 ^k	3.1 ^k 4.7 ^k	3.7 ^k 4.7 ^k	4.9 ^k 4.5 ^k	5.2 ^k 4.8 ^k	5.3 ^k 4.8 ^k	5.3 ^k 5.3 ^k																
2	(5.9) ^k	5.1 ^k	5.7 ^k	4.7 ^k	4.5 ^k	4.5 ^k	4.8 ^k	5.1 ^k	5.3 ^k																
3	5.0	4.8	4.5 ^k	4.1	4.2	4.8	(5.9) ^j	2.1	2.6	2.5 ^k	6.7	6.5 ^k	6.5 ^k												
4	5.9 ^j (5.3) ^j	5.5 ^f	4.9 ^j	4.4 ^j	5.0 ^j	6.4 ^j	(6.7) ^j	6.7 ^j	7.2 ^j 7.6 ^j	8.1 ^j 8.6 ^j	(6.6) ^j	6.8 ^j	6.7 ^j												
5	(5.3) ^j	5.6 ^f	3.2 ^f	4.7 ^f	4.7 ^f	3.9 ^f	5.4 ^f	5.4 ^f	7.0	7.6	7.6 ^f	6.5 ^f	6.2 ^f												
6	5.3 ^f	5.1 ^f	4.9 ^f	4.7 ^f	4.1	4.9 ^f	6.5 ^f	7.1	2.2	2.2 ^f	(2.8) ^j	8.4 ^j	8.1 ^j	8.3 ^j											
7	5.6 ^j	5.7 ^f	5.7 ^f	5.1 ^j	5.1 ^j	4.5 ^k	4.5 ^k	4.5 ^k	5.0 ^k	5.7 ^k	5.5 ^k	5.7 ^k	5.5 ^k	5.7 ^k											
8	F ^k	2.9 ^f	N ^k	N ^k	N ^k	N ^k	N ^k	N ^k	G ^k																
9	(2.6) ^k	2.9 ^k	2.6 ^k	2.2 ^k	1.9 ^k																				
10	(5.6) ^j	F ^k	(2.9) ^k	2.9 ^k	3.1 ^k	3.3 ^k	3.9 ^k	G ^k	C ^k																
11	(3.1) ^j	F ^k	(1.8) ^k	F ^k	F ^k	(3.4) ^k	(4.5) ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k	G ^k		
12	4.9 ^f	4.5 ^f	(3.3) ^k	(2.1) ^k	F ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k	1.9 ^k		
13	3.9 ^k	3.5 ^f	(3.5) ^k	(3.1) ^k	2.7 ^k	3.2 ^k	4.2 ^k	5.2 ^k	5.2 ^k	5.5 ^k															
14	(5.5) ^f	(5.0) ^j	4.6 ^f	4.3 ^f	4.1 ^f	4.8 ^f	(5.7) ^c	5.7 ^c	5.7 ^c	5.9 ^j	(6.3) ^j	S	B	C	5.8 ^j	(6.2) ^j	(6.6) ^j	(6.3) ^c	(6.6) ^j						
15	(4.9) ^j	T ^f	(4.7) ^j	(4.7) ^j	(4.0) ^j	4.5 ^j	(5.5) ^j	(6.4) ^j	(6.5) ^j	(6.5) ^j	(6.5) ^j	20 ^f	(2.2) ^j	2/	2.2	2.8	2.7	2.9	2.2	2.8	2.2	2.8	2.2	2.8	2.2
16	5.1 ^j	(5.5) ^j	5.0 ^j	4.5 ^f	4.9 ^j	(5.7) ^j	(6.5) ^j	7.4 ^j	(6.5) ^j	(6.5) ^j	(6.5) ^j	6.8	20	2/	2.2	2.2	2.4	2.3	2.4	2.3	2.4	2.3	2.4	2.3	
17	5.7 ^f	5.5 ^f	5.3 ^f	4.9 ^f	4.4 ^f	4.8 ^f	6.2 ^f	6.6 ^f	6.7 ^f	6.7 ^f	2.4	2.6	2.8	2.5	2.6	2.7	2.5	2.7	2.5	2.7	2.5	2.7	2.5		
18	(5.5) ^f	5.1 ^f	4.5 ^f	4.3 ^f	3.8 ^f	4.5 ^f	6.0 ^f	6.0 ^f	20	7.2 ^j	(6.4) ^j	6.5 ^j	6.7	6.8	6.8	6.9	6.9	6.8	6.8	6.8	6.8	6.8	6.8		
19	6.0	5.9	5.6 ^f	5.3 ^f	5.0	5.0	5.5 ^f	6.1 ^f	8.0	(19) ^j	9.2 ^j	8.6	8.8	8.9	8.8	8.7	8.5	8.5	8.5	8.5	8.5	8.5	8.5		
20	7.4	6.9	(6.6) ^j	6.7	6.5 ^j	6.2	7.4	8.0	(2.5) ^c	(2.5) ^j	(2.5) ^j	8.2	7.6	7.3	7.0	7.4	7.5	7.4	7.4	7.4	7.4	7.4	7.4		
21	5.2 ^f	5.1 ^f	5.0 ^f	4.8 ^f	4.2 ^f	5.2	5.9	6.6	6.2	2.4	2.5	(6.8) ^j	7.5	7.5	7.5	7.5	7.5	7.5	7.5	7.5	7.5	7.5	7.5		
22	4.5 ^f	4.2 ^f	4.1 ^f	3.7	4.7	5.8 ^f	C	C	7.0	2.2	2.5	8.0	8.4	8.6	8.5	8.6	8.5	8.6	8.5	8.6	8.5	8.6	8.5		
23	6.5 ^j	(5.8) ^j	(6.0) ^j	5.4 ^j	4.9	4.9 ^j	(6.0) ^j	2.2	7.3	2.4	2.6 ^j	(2.2) ^j	8.0	7.8	8.2	7.8	8.2	7.7	8.2	7.7	8.2	7.7	8.2	7.7	
24	(5.8) ^j	5.7 ^f	5.3 ^f	4.9 ^f	(4.4) ^j	4.5 ^f	6.0 ^f	6.4	6.2	(6.5) ^j	6.8	6.9	20	2/	2.1	2.2	2.0	2.3	2.2	2.0	2.3	2.2	2.0	2.3	
25	(5.8) ^f	F ^k	(5.8) ^j	(4.6) ^j	4.5 ^f	5.3 ^f	6.7 ^f	2.3	8.1	8.2 ^j	8.9 ^j	(9.3) ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j	8.8 ^j		
26	5.4 ^f	5.2 ^f	4.9 ^f	4.7 ^f	4.2 ^f	3.8 ^f	6.2	C	C	C	C	(8.3) ^j													
27	(5.2) ^j	5.3 ^f	5.2 ^f	5.1 ^f	4.8	5.2	6.3	6.6	(6.4) ^j	2/	2.3	2.2	2.9	8.3	8.3	8.0	8.2	8.0	7.7	2.2	2.1	6.5	6.3		
28	(6.0) ^j	5.8 ^j	(5.7) ^j	5.3 ^j	4.9 ^j	4.9	6.3	20 ^j	(8.1) ^j	8.3	8.3	8.6	9.0	9.4	8.9	8.8	9.2	8.9	8.9	8.3	6.9	6.7	6.3		
29	6.5 ^k	6.3 ^k	6.3 ^k	5.3 ^f	4.9 ^k	4.9 ^k	5.6 ^f	6.8 ^k	G	C	5.6 ^k	5.6 ^k	5.2 ^k	G	6.0 ^k	6.0 ^k	5.8 ^k	6.0 ^k	6.0 ^k	6.1 ^k	5.6 ^k	5.6 ^k			
30	3.5 ^f	3.8 ^f	3.3 ^f	3.1 ^f	3.3 ^f	3.3 ^f	3.6 ^f	5.1 ^f	5.9 ^f	(6.0) ^j	5.3 ^f														
31	3.9 ^f	3.9 ^f	(3.4) ^f	3.0 ^f	3.0 ^f	3.2 ^f	4.5 ^f	5.3 ^f	5.6 ^f	5.6 ^f	(5.3) ^j	6.1 ^j	6.3 ^j	6.0 ^j	6.3 ^j	6.0 ^j	6.2 ^j	6.2 ^j	6.2 ^j	6.0	6.2 ^j	5.9 ^f	5.9 ^f		
Median	5.4	5.2	5.0	4.7	4.2	4.6	5.8	6.4	6.6	6.6	6.6	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	7.0	7.0	7.0	7.0	
Count	30	30	29	29	30	31	29	28	28	27	27	29	31	31	31	31	31	31	31	31	30	30	30	30	

Form adopted June 1946

5 COMPONENT PRINTING OFFICE 146 U-10219

2

3

Sweep 10. Mc 10.25.0 Mc in 0.25-min

Manual □ Automatic ■

TABLE 47
IONOSPHERIC DATA

$n^{\prime} F_1$. Km . August , 1948
 (Characteristic) (Unit) (Month)

Observed at **Washington, D.C.**

Lat **39°0'N.**, long **77°5'W**

Day	75° W. Mean Time												Calculated by:	N. N. M.	K. I. W.				
	00	01	02	03	04	05	06	07	08	09	10	11							
1	Q	K	(240) ^a	200	K	190	K	200	K	200	K	200	K	G	K	200			
2	Q	K	200	K	200	K	180	K	200	K	200	K	200	K	200	K			
3	Q	(200) ^a	(200) ^b	(200)	A	190	(220) ^b	200	180	H	200	K	220	K	220	K			
4	Q	200	200	200	200	200	200	200	200	200	200	200	210	210	210	230			
5	Q	210	200	180	180	180	180	180	180	180	180	180	200	200	200	230			
6	Q	(230) ^a	210	(200) ^a	A	(200) ^a	200	200	200	200	200	200	230	230	230	(230) ^a			
7	250	K	210	K	200	K	A	K	A	K	A	K	200	K	210	K			
8	Q	K	230	K	220	H	180	K	230	K	210	H	200	K	(210) ^a	(250) ^a			
9	Q	K	220	K	200	K	180	K	200	K	170	K	(180) ^a	200	K	A	K		
10	Q	K	200	K	220	K	200	K	C	K	C	K	230	K	210	K	240	K	
11			230	K	(250) ^a	190	K	A	K	200	K	180	200	210	S	200			
12			250	K	220	K	(200) ^c	(200) ^c	(180) ^c	C	K	200	K	200	K	(210) ^a	(200) ^c	230	K
13	Q	K	(210) ^a	(210) ^a	(210) ^a	240	K	A	K	180	K	170	K	200	K	C	K	A	K
14			Q	A	170	160	210	190	210	200	200	230	200	220	220	230	230		
15			Q	130	200	(200) ^a	250	210	200	200	A	210	210	210	210	210	(230) ^a		
16			Q	A	A	190	A	180	(200) ^a	200	200	200	210	210	210	210	210		
17			Q	200	A	210	(220) ^a	180	(170) ^a	200	200	210	(210) ^a	230	230				
18			Q	(210) ^a	C	C	C	200	200	200	210	210	220	220	220	A			
19			Q	230	210	180	180	190	220	200	190	190	(200) ^a	210	230	Q			
20			Q	Q	Q	210	210	210	(230) ^a	180	210	200	200	220	220	C	Q		
21			A	220	A	(200) ^a	200	200	190	220	220	220	210	230	230	A			
22			C	C	A	200	180	H	220	230	240	220	230	220	220	240			
23			250	230	220	200	C	180	200	(210) ^c	(200) ^c	220	220	210					
24			Q	230	210	200	180	200	200	200	200	200	200	210	220	Q			
25			220	200	210	200	230	C	200	200	200	210	210	210	230	Q			
26			C	C	C	C	C	200	200	210	C	(210) ^a	A	A	A	A			
27			200	A	A	200	A	(210) ^c	180	210	220	A	Q						
28			Q	200	H	200	180	220	200	200	200	240	220	200	220				
29			230	K	220	K	200	K	220	K	220	K	200	K	220	A	K	Q	
30			Q	K	210	K	200	K	180	K	210	K	230	K	220	K	230	Q	
31			(230) ^a	210	K	200	K	200	K	180	K	190	K	240	K	220	210		
Median				230	210	200	195	200	200	200	200	200	210	220	220	230			
Count				17	25	24	24	27	29	30	29	28	29	27	25	25	13		

Sweep 1.0 Mc to 25.0 Mc in 0.25 min
 Manual □ Automatic ■

¹⁰FI Mc August
(Characteristic) (Unit) (Month)

Observed at Washington, D.C.

Lat. 39.0°N., Long. 77.5°W.

75° W. Mean Time

National Bureau of Standards

(Institution)

Scaled by: E. J. W.

J. J. S., J. M. C.

Calculated by: M. C. E.

G. G. H.

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1								Q	K	42	K	4.5	K	4.7	K	4.9	K	5.0	K	5.2	K	5.1	K	(4.9)	K
2								Q	K	49	K	4.9	K	5.0	K	4.9	K	5.2	K	5.1	K	5.1	K	5.0	K
3								Q	L			5.2		5.0		5.3		5.4		5.1		5.1		(5.4)	
4								Q	Q			5.3		5.6		L		5.4		5.3		5.3		5.4	
5								Q	Q	48		5.3		5.0		5.4		5.3		5.3		5.5		4.9	
6								Q	L			5.7		5.7		5.6		5.7		5.3		5.3		L	
7								L	K	4.1	K	4.3	K	(4.7)	K	A	K	A	K	4.9	K	5.3	K	5.1	K
8								Q	K	3.7	K	3.9	K	4.3	K	4.5	K	4.6	K	(4.6)	K	4.6	K	4.7	K
9								Q	K	4.1	K	4.3	K	4.5	K	4.6	K	4.8	K	4.6	K	4.7	K	4.7	K
10								Q	K	4.1	K	4.3	K	4.5	K	4.6	K	4.7	K	4.7	K	4.7	K	4.7	K
11								Q	K	4.5	K	4.5	K	4.5	K	4.9	K	4.9	K	5.4	K	5.5	K	5.1	K
12								Q	K	4.1	K	4.4	K	4.4	K	4.7	K	4.7	K	(4.9)	K	5.0	K	[4.8]	K
13								Q	K	4.2	K	4.8	K	4.9	K	5.0	K	5.0	K	5.0	K	4.9	K	4.7	K
14								Q	Q	4.6		5.0		5.1		N	S	C	K	5.2		5.2		[5.2]	
15								Q	L			5.1		L		5.5		C	K	5.3		4.9		4.9	
16								Q	L			4.5		5.4		5.7		5.6		5.5		5.4		5.3	
17								Q	Q			5.5		5.5		(5.9)		5.8		5.5		5.7		5.9	
18								Q	Q			5.0		C		5.4		5.5		5.6		5.4		5.0	
19								Q	L			5.1		L		6.2		6.0		6.1		6.0		5.7	
20								Q	Q			(5.8)	P	L		6.1		5.8		5.8		5.5	H	5.5	L
21								A	9	5.1		5.4		5.7	H	5.5		5.6		5.6		5.6		5.9	
22								C	C	5.1		5.4		5.7		5.7		6.0		5.8		5.6		L	
23								L	L			(6.1)	F	[5.8]	C	5.5	H	5.9		5.6		5.7		5.6	
24								Q	L			5.5		5.5		5.7		5.5		5.4		5.5		5.3	
25								L	L			5.7		(5.7)	C	(5.9)	L	5.5		L		L		Q	
26								C	C	C	C	(5.7)	C	5.3	H	5.8	C	5.8		C	L	H	A	A	A
27								L	L			5.7		5.7		5.7		5.3		L		L		L	
28								Q	L			L		A		L		5.7		5.6		L		4.5	
29								L	K	L	K	4.7	F	(5.5)	C	5.9		5.5		L		4.5		4.5	
30								Q	K	L	K	4.7	K	4.9	K	4.8	K	4.9	K	4.9	K	4.7	F	S	K
31								L	K	4.5	K	4.9	K	4.8	K	4.9	K	4.9	K	4.9	K	4.9	K	4.9	L
Median		4.1	4.5	4.9	5.1	5.4	5.4	5.4	5.4	5.4	5.4	5.3	5.3	5.3	5.3	5.3	5.3	5.3	5.3	5.3	5.3	4.9	4.7		
Count		1	8	14	22	23	24	25	26	27	28	27	27	27	27	27	27	27	27	27	27	27	27	27	

Sweep 10 Mc to 25.0 Mc in 0.25 min
Manual Automatic

TABLE 49
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

$h'E$
(Characteristic)
km
(Unit)
August 1948
(Month)

Observed at Washington, D.C.

Lat 39.0°N

Long 77.5°W

National Bureau of Standards
(Institution)
Scaled by: E. J. W.
J. J. S.; J. M. C.

Calculated by: N. N. M., G. G. H.

75°W Mean Time												
Day	00	01	02	03	04	05	06	07	08	09	10	11
1												
2												
3												
4												
5												
6												
7												
8												
9												
10												
11												
12												
13												
14												
15												
16												
17												
18												
19												
20												
21												
22												
23												
24												
25												
26												
27												
28												
29												
30												
31												
Median	100	100	100	100	100	100	100	100	100	100	100	100
Count	26	28	29	28	27	29	30	29	29	31	31	18

Sweep LO Mc 1050 Mc in. 9.25 min
Manual Automatic

TABLE 50
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.Mc
(Characteristic)
f^oE
Mc
August
(month)
1948Washington, D. C.
Observed at

Lat 39.0°N, Long 77.5°W

IONOSPHERIC DATA

National Bureau of Standards
(Institution)

J.J.W.

Calculated by: E.J.W.

J.M.C.

K.L.W.

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23		
1								A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A		
2								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
3								2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	
4								2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
5								2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
6								2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
7								2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	
8								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
9								3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	
10								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
11								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
12								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
13								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
14								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
15								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
16								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
17								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
18								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
19								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
20								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
21								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
22								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
23								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
24								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
25								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
26								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
27								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
28								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
29								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
30								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
31								2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	
Median	2.7	3.1	3.3	3.6	3.7	3.8	3.9	3.8	3.7	3.5	3.0	2.4	1.9													
Count	1	23	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22	22

Sweep I.Q. Mc 1025.0 Mc in 3.25 min
Manual □ Automatic ☐

27

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

TABLE 51
IONOSPHERIC DATA
Es — **Mc/km** **August 1, 1948**
 (Characteristic) (Month)

Observed at **Washington, D.C.**
Lat. 39.0°N, Long. 77.5°W
National Bureau of Standards
 (Institution)

Scaled by **E. J. W.** **J. J. S.**, **J. M. C.**
Calculated by **J. T. D.** **F. H. L.**

75°W Mean Time																									
Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	2.9/100	2.3/100	4.3/100	3.9/100	3.4/100	6.5/100	3.7/100	4.0/100	4.2/100	5.6/100	4.0/100	4.0/100	5.6/100	4.0/100	7.0/100	3.2/100									
2		2.2/100			3.0/100	4.2/100	4.3/100	4.6/100	4.8/100	3.9/100	4.8/100	4.2/100	4.3/100	4.8/100	4.3/100	5.8/100	3.4/100								
3	3.4/100	2.2/100			2.0/100	3.6/100	5.4/100	5.2/100	4.9/100	4.6/100	5.0/100	4.0/100	4.3/100	5.1/100	4.6/100	3.7/100	3.2/100	5.8/100	3.4/100	3.7/100	3.8/100				
4	3.3/100	2.3/100			2.1/100	2.0/100	3.6/100	3.6/100	3.6/100	3.6/100	3.9/100	3.9/100	4.1/100												
5	3.2/100	1.9/100			1.8/100		3.4/100	4.2/100	3.6/100	4.0/100	5.2/100	4.6/100	4.8/100												
6	1.9/100	3.0/100			2.3/100	3.7/100	3.7/100	4.6/100	5.6/100	5.6/100	5.7/100	5.2/100	4.4/100	3.9/100	4.6/100	4.2/100	4.6/100	4.8/100	3.9/100	3.5/100	3.3/100				
7	1.9/100	2.3/100			1.9/100	2.5/100	3.8/100	3.8/100	5.4/100	4.8/100	7.0/100	7.0/100	7.4/100	4.5/100	5.6/100	5.9/100	3.8/100	3.0/100	3.4/100						
8	1.9/100	2.1/100			2.1/100	2.2/100			3.9/100	3.9/100	3.7/100	4.0/100													
9	3.9/100	3.9/100			2.3/100	2.3/100	3.0/100	3.0/100	3.7/100	4.2/100	5.2/100														
10	2.8/100	2.3/100	5.6/100	3.3/100	1.9/100	4.7/100	3.2/100	4.8/100	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
11	3.6/100	9.9/100							3.4/100	3.6/100	4.7/100	3.7/100													
12	2.0/100	3.6/100	7.5/100						5.8/100	2.9/100	3.1/100	3.4/100	3.6/100		B										
13	3.5/100	5.2/100	6.8/100	8.4/100	3.1/100	4.0/100	4.0/100	4.0/100	4.0/100	3.2/100	3.2/100	3.2/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100	3.7/100		
14	5.8/100	5.2/100	4.9/100	4.3/100						2.0/100	2.9/100	2.9/100	2.9/100	2.9/100											
15	1.9/100	3.1/100	4.9/100	4.7/100	2.3/100	2.3/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100	2.0/100		
16	2.4/100	3.4/100	3.6/100	3.1/100	3.5/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100	3.6/100		
17	3.2/100	3.4/100	3.3/100	3.1/100	3.5/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100	3.4/100		
18	2.4/100	2.0/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100	2.3/100		
19	3.1/100	3.7/100	3.1/100							2.0/100	2.0/100	2.0/100	2.0/100	2.0/100											
20										2.0/100	3.3/100	3.3/100	3.9/100	5.0/100											
21										3.5/100	5.9/100	4.2/100	5.3/100	5.3/100											
22		2.6/100	3.0/100							1.2/100	1.8/100	C	4.8/100	6.3/100	4.3/100										
23										2.0/100	3.0/100	3.4/100	3.5/100	3.5/100	C	C									
24										3.2/100															
25										2.0/100															
26		2.7/100	2.9/100	3.1/100	5.3/100	5.7/100	9.9/100	3.2/100	7.2/100	3.9/100	5.5/100	8.3/100	9.6/100	40/100	3.8/100	3.8/100	3.8/100	3.8/100	3.8/100	3.8/100	3.8/100	3.8/100	3.8/100		
27		6.9/100	5.7/100	5.3/100	5.7/100	5.7/100	9.9/100	3.2/100	7.2/100	3.9/100	5.5/100	8.3/100	9.6/100	40/100	3.1/90	4.3/90	3.9/100	3.9/100	3.9/100	3.9/100	3.9/100	3.9/100	3.9/100		
28										2.4/100															
29										3.1/100	3.2/100	3.2/100	3.8/100	3.8/100											
30										2.9/100	3.2/100	3.4/100	3.5/100	3.5/100											
31										2.9/100	3.1/100	3.0/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100	3.1/100		
Median	**	2.3	1.9	2.2	*	*	*	2.5	3.5	3.7	3.9	3.7	3.7	3.7	*	*	*	*	*	3.2	3.3	1.8	*	2.0	
Count	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	

 ** MEDIAN FEES LESS THAN MEDIAN FOR OR LESS THAN
LOWER FREQUENCY LIMIT OF RECORDER

 Sweep 1.0 Mc 1025.0 Mc in 0.25 min
Manual □ Automatic

TABLE 52
IONOSPHERIC DATA

National Bureau of Standards
 [Institution] J.J.S., J.M.C.
 Calculated by: E. J. W.

F2-M(500, (Unit)
 (Characteristic) Observed at Washington, D.C.

Lat 39.0°N Long 77.5°W

August 1, 1948
 (Month)

Day	75°W Mean Time																										
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23			
1	F K (1.0)F	(1.0)F	1.7K	1.8K	1.9K	G K	1.6K	1.5K	1.6K	G K	1.6K	G K	1.6K	G K	1.7K	1.7K	1.8K										
2	(1.0)K	1.9K	1.7K	1.7K	1.9K	1.9K	2.2K	1.7K	1.6K	G K	1.5K	1.7K	1.8K	1.9K	1.9K	2.0K	2.1	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9		
3	1.8	1.9	1.8	1.8	2.0	2.2	2.2	1.9	2.0	1.9	(1.6)J	1.8	1.6	1.7	1.7	1.8	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7		
4	1.8	1.9F	1.9F	1.9F	2.0	(2.2)S	2.3	(2.3)F	2.0	1.9	2.0	1.8	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7		
5	(1.7)F	1.8F	2.0F	(2.0)J	2.0	2.0	2.3	2.2	2.0	1.8	2.0	1.8	(1.8)J	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
6	1.8F	1.9F	1.9F	1.9F	1.9F	2.2K	2.1K	2.0K	2.1K	2.0K	(2.0)J	1.9K	(2.0)K	1.9K	1.9K	2.0K	2.0K	1.9K	2.0K	1.9K	2.0K	1.9K	2.0K	1.9K	2.0K	1.9K	
7	F(0.9)K	1.7K	1.8K	1.7K	1.8K	1.9K	1.9K	1.9K	1.7K	G K	1.6K	A K	1.5K	1.7K													
8	1.8K	F K	F K	N K	1.5K	N K	2.1K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K		
9	(1.6)K	1.7K	1.9K	1.9K	1.9K	2.1K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K		
10	(1.9)J	F K	E K	(1.7)F	1.7K	1.7K	2.0K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K		
11	(2.0)K	(2.0)F	1.7K	(1.7)F	F K	(1.9)F	2.2K	G K	1.7K	G K	1.6K	A K	1.5K	1.7K													
12	2.0F	1.8F	(1.8)K	1.8F	(1.6)K	(1.9)K	(1.9)K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K		
13	1.8K	1.7K	1.7K	1.8K	1.7K	1.9K	(2.0)F	(2.1)K	1.8K	G K	1.8K	2.1K	1.7K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K	G K		
14	1.8F	2.0F	1.9F	2.0F	1.9F	2.1F	2.2K	G K	1.9K	(2.1)F	(2.1)C	(2.1)K	1.8K	2.0	2.0	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
15	(1.8)J	1.9F	1.9F	1.9F	1.9F	2.0F	2.2	(2.3)J	(2.0)F	(2.2)J	(2.2)F	(2.2)C	2.1	2.0	1.8	1.9	1.9	1.9	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
16	1.8	1.9	1.9F	2.0	1.9F	2.0	2.3	(1.9)F	(2.0)F	(2.1)F	(2.0)F	(2.0)C	1.8	1.8	1.9	1.9	1.9	1.9	2.0	2.0	1.9	1.9	1.9	1.9	1.9	1.9	
17	1.9	1.9	1.9F	2.0	1.8F	2.0F	2.4F	(2.0)F	(2.2)F	(2.2)C	(2.2)J	1.9	2.1	1.8	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	
18	1.9F	1.9F	1.9F	1.9F	1.8F	2.0F	2.0F	2.1	2.0F	C	C	(1.7)J	1.7	N S	(1.7)C	1.7	C	1.7	(1.9)J	(1.9)J	2.0	N	(1.9)J	(1.9)J	1.9		
19	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	2.1	2.1	2.1	2.3	2.2	(1.8)J	(1.8)C	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9
20	1.6	1.7	1.7	1.6	1.8	1.8	1.9	2.0	1.9	1.9	2.0	(1.9)J	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	
21	1.7	1.7	1.7	1.8	1.8	1.8	1.9	2.0	1.9	1.9	2.1	(1.7)J	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
22	1.9F	1.7F	1.7F	1.7	1.9	1.9	2.0	C	C	C	C	C	C	C	C	C	C	C	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
23	(1.8)J	1.8	1.8	1.8	1.8	1.8	1.8	1.9	2.1	1.9	2.1	1.9	2.1	1.9	1.9	1.9	1.9	1.9	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
24	(1.8)J	1.8	1.8	2.0	1.8F	1.8	2.0F	2.0	(2.0)J																		
25	(1.8)J	(2.0)F	(1.9)F	(2.0)F	(1.9)F	(2.0)F																					
26	1.8	1.9	1.9F	1.9F	1.9F	2.0	2.0	(2.2)J	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
27	A	(1.9)A	1.9F																								
28	(1.8)J	1.8	1.8	1.8	1.8	1.9	2.0	(2.3)S	2.1	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2		
29	(2.0)J	1.7K	1.9K	1.9K	1.9K	2.0K	(1.8)F	2.0K	(2.0)J																		
30	1.7K	1.9K	1.7K	1.7K	1.8K	1.8K	2.2K	2.1K	2.2K																		
31	1.0K	1.9K																									
Median	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	2.1	2.1	2.0	1.9	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8		
Count	29	29	30	30	31	29	31	30	31	29	29	29	29	29	29	29	29	29	29	29	29	29	29	29	29	29	

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

TABLE 53
IONOSPHERIC DATA

F2-M3000, August, 1948
(Characteristic) (Month)
Observed at Washington, D.C.
Lat 39.0°N, Long 77.5°W

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

National Bureau of Standards
(Propagation)
Scaled by **E. J. W.** **J. M. C.**
Calculated by **J. L. K.** **N. N. M.**

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	F	K ^a (2.5) ^b (2.8) ^c	2.5K	2.6K	2.7K	2.7K	2.7K	2.7K	2.7K	2.7K	2.7K	2.7K
2	(2.9) ^d	2.8K	2.6K	2.6K	2.6K	2.7K	2.7K	2.7K	2.7K	2.7K	2.7K	2.7K
3	2.8	2.8	2.6	2.7	2.7	3.0	3.3	2.9	3.0	2.9	2.7	2.7
4	2.7	2.8	2.9	2.8	2.8	2.9	(3.2)	3.4	3.0	2.9	2.7	2.7
5	(2.7) ^e	2.8	2.9	3.0	2.9	3.4	3.2	3.0	2.8	2.7	2.7	(2.5) ^f
6	2.8	2.7	2.9	2.9	2.9	2.7	(2.8)	3.2	3.0	2.9	2.8	2.8
7	2.7	2.5	2.7	2.5	2.6	2.8	2.5K	2.6K	2.3K	2.3K	2.3K	2.3K
8	2.7	F	F	N	N	N	3.0K	3.0K	2.9K	2.9K	2.9K	2.9K
9	(2.3) ^g	(2.3) ^h	2.5	2.8	2.8	3.0	3.1	3.2	3.0	2.9	2.8	2.8
10	(2.9) ⁱ	F	F	(2.5) ^j	2.6	2.6	2.9K	2.9K	2.9K	2.9K	2.9K	2.9K
11	(3.0) ^k	(2.8) ^l	(2.5) ^m	F	(2.8) ⁿ	2.5K	2.6K	2.8K	(3.2) ^o	G	2.7	3.0
12	3.0	2.7	2.7	2.6	2.6	2.7	(2.9) ^p	(2.8) ^q	2.2K	8K	2.8K	3.0
13	2.7	2.6	2.7	2.6	2.6	2.7	3.0K	3.1K	2.5K	G	2.3K	2.9K
14	2.6	3.0	2.9	2.9	2.9	3.1	(3.1) ^r	(2.7) ^s	2.5	2.5	2.7	3.0
15	(2.8) ^t	2.8	2.9	2.9	2.8	2.9	(3.2) ^u	(3.0) ^v	3.1	3.0	2.9	3.0
16	2.7	2.9	2.9	2.9	2.9	3.3	(2.9) ^w	3.1	3.3	(2.9) ^x	2.7	3.1
17	2.8	2.8	2.9	2.9	2.9	3.4	3.2	3.1	2.7	2.8	2.8	3.0
18	2.9	2.9	2.8	2.8	2.8	3.0	C	C	(2.6) ^y	2.8K	2.8K	2.9
19	2.8	(2.9) ^z	2.8	2.8	2.9	3.1	3.1	3.1	(2.9) ^{aa}	(2.9) ^{ab}	3.0	N
20	2.5	2.6	2.5	2.5	2.7	(2.9) ^{cc}	2.9	3.0	(2.9) ^{dd}	2.6	2.6	2.7
21	2.4	(2.6) ^{ee}	2.7	2.7	(2.6) ^{ff}	2.7	2.9	3.1	(2.6) ^{gg}	2.7	2.7	2.7
22	2.7	2.6	2.5	2.6	2.7	2.8	3.2	C	C	2.6	2.6	2.9
23	(2.5) ^{hh}	(2.9) ⁱⁱ	2.7	2.8	2.8	3.0	2.8	3.2	(2.9) ^{jj}	2.8	2.8	2.9
24	(2.7) ^{kk}	2.7	2.8	2.7	2.7	3.0	3.0	(3.1) ^{ll}	(2.6) ^{mm}	2.7	2.7	2.7
25	(2.7) ⁿⁿ	(2.9) ^{oo}	(2.9) ^{pp}	2.8	3.0	3.3	3.2	3.0	3.1	2.7	2.7	2.7
26	2.7	2.8	2.9	2.9	2.9	(3.3) ^{qq}	C	C	(2.9) ^{rr}	2.9	2.9	(3.0) ^{tt}
27	A	(2.6) ^{uu}	2.8	2.9	2.9	3.2	(3.0) ^{vv}	(3.1) ^{ww}	3.2	A	3.0	3.0
28	(2.9) ^{xx}	2.8	2.8	2.8	2.9	3.2	(3.4) ^{yy}	3.4	3.2	3.1	2.9	2.9
29	(3.1) ^{zz}	2.7	2.8	2.9	2.9	2.9	(2.9) ^{aa}	2.7K	2.5K	2.4K	2.8K	2.8K
30	2.7	2.7	2.6	2.7	2.7	3.2	3.0	3.2	2.7	2.7	2.7	2.7
31	2.8	2.8	2.8	2.9	2.9	3.0	(3.0) ^{cc}	(4.4) ^{dd}	2.8K	2.8K	2.9K	2.6
Median	2.7	2.8	2.8	2.8	2.9	3.1	3.0	3.0	2.8	2.7	2.7	2.8
Count	29	27	29	30	30	31	29	29	27	28	30	30

Sweep 10—Mc 10 sec. 0 Mc 0.2 sec. min
Manual Automatic

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

TABLE 54
IONOSPHERIC DATAFI-M3000, August, 1948
(Characteristic) (Month)

Observed at Washington, D.C.

Lat. 39.0°N, Long. 77.5°W

Day	75°W											75°W											
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22
1	Q	K	36K	38K	40K	41K	38K	37K	37K	35K	35K	L	K										
2	Q	K	34K	36K	38K	43K	36K	37K	37K	34K	34K	L	K										
3	Q	L	L	35	38	38	38	38	38	38	38	38	38	38	38	38	38	37	37	35	L		
4	Q	Q	L	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	L		
5	Q	Q	Q	37	35	38	41	41	41	41	41	41	41	41	41	41	41	37	37	35	L		
6	Q	L	L	33	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	L		
7	L	K	38K	(37)K	A	K	A	K	A	K	A	K	A	K	A	K	3.4K	3.4K	3.4K	L	K		
8	Q	K	34K	36K	40K	39K	38K	38K	37K	37K	37K	A	K	3.8K									
9	Q	K	32K	37K	39K	40K	40K	37	37	37	3.6K	3.6K	3.4K										
10	Q	K	34K	37K	39K	C	K	C	K	C	K	C	K	C	K	C	3.8K	3.7K	3.9K	3.6K	3.6K	3.4K	L
11	Q	K	36K	36K	3.9K	3.8K	4.0K	4.0K	3.7	C	(3.5)K	(3.7)K	5	L									
12	2.9	K	36K	36K	38K	40K	B	K	3.2K	3.0K	2.9K	L	K										
13	Q	K	37K	35K	36K	A	K	C	K	C	K	C	K	C	K	C	(39)K	(38)K	(38)K	C	K	3.6K	
14	Q	Q	37	36	38	N	S	C	37	35	37	35	37	35	37	35	3.7	3.7	3.7	3.5	3.5	L	L
15	Q	L	L	39	L	L	L	L	37	C	L	3.5	L	L	L	L	3.5	3.5	3.5	L	L	L	
16	Q	L	L	42	35	34	36	36	36	36	36	36	36	36	36	36	36	36	36	36	3.4	L	
17	Q	Q	L	L	36	4.3	(3.4)K	3.6	3.6	3.6	3.5	3.5	L	F									
18	Q	Q	3.5	C	C	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.7	3.7	3.7	3.4	3.4	L
19	Q	L	L	L	L	34	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	Q
20	Q	Q	9	(36)P	L	3.3	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.3	L	Q
21	A	3.5	A	3.7	3.7	4	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.6	3.6	3.6	3.2	3.2	L	L
22	C	C	A	3.6	3.5	3.5	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	L	L
23	L	L	(31)F	C	3.8K	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	L	L	
24	Q	L	L	3.5	3.5	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	Q	
25	L	L	3.5	(3.8)K	(3.4)K	L	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	L	Q	
26	C	C	C	C	(3.6)K	3.8K	(3.3)K	C	C	C	C	C	C	C	C	C	C	C	C	A	A		
27	L	L	C	C	L	A	L	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5	3.5		
28	Q	L	L	3.9	(3.7)K	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	Q	
29	L	K	3.8K	K(3.6)K	K(3.8)K	3.7K	3.5K	3.7K	3.7K	3.7K	3.7K	3.7K	3.7K	A	K								
30	Q	K	3.8K	3.6K	3.9K	(3.8)K	3.6K	3.8K	3.8K	3.8K	3.8K	3.8K	3.8K	L	K								
31	L	K	3.5K	3.5K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	3.9K	L	
Median	3.5	3.6	3.8	3.8	3.8	3.8	3.7	3.7	3.6	3.7	3.7	3.6	3.7	3.7	3.6	3.7	3.6	3.5	3.4				
Count	8	14	20	21	24	27	27	27	26	26	26	26	26	26	26	26	26	26	26	26	26	26	

Sweep 1.0 Mc to 25.0 Mc in 0.25 min
Manual Automatic

**TABLE 55
IONOSPHERIC DATA**

E-MI500 August, 1948
(Characteristic) (Unit) (Month)
Observed at Washington, D.C.

Lat 39.0°N, Long 77.5°W

National Bureau of Standards
(Institution)

J.J.S., J.M.C.

Calculated by: J.L.K., N.N.M.

Day	75°W Mean Time													
	00	01	02	03	04	05	06	07	08	09	10	11		
1	A	K	4.4	A	K	A	K	4.5	K	A	K	4.5	K	
2	4.7	K	4.1	K	4.7	K	A	K	A	K	4.5	K	4.5	K
3	4.4	4.4	A	(4.5)	A	A	A	(4.6)	A	A	(4.6)	4.3	K	
4	4.4	4.3	(4.4)	A	4.4	4.7	A	4.4	(4.5)	A	4.7	4.2	4.2	
5	4.4	4.4	4.5	4.5	4.7	4.6	A	4.4	4.2	4.2	4.3	4.5	4.4	
6	4.4	4.5	4.7	(4.6)	A	4.6	A	4.5	4.5	4.4	4.3	(4.4)	A	
7	(4.5)	A	A	K	4.7	K	A	K	4.5	K	A	K	4.2	
8	A	K	4.2	K	4.5	K	(4.6)	K	(4.7)	K	(4.8)	(4.5)	K	
9	4.4	A	A	K	(4.5)	K	A	B	4.3	K	(4.5)	K	B	
10	4.2	K	(4.4)	K	4.3	K	C	K	(4.6)	K	4.6	K	4.2	
11	4.7	K	4.3	K	4.5	K	4.7	K	A	K	C	K	A	
12	A	H	A	K	(4.7)	C	A	K	C	K	B	K	(4.7)	
13	4.2	K	A	K	(4.8)	K	4.8	K	A	K	B	K	C	
14	4.0	A	4.7	4.5	C	A	(4.4)	C	4.6	4.2	4.1	4.3	4.3	
15	4.3	H	A	(4.9)	A	4.4	(4.7)	C	4.4	4.5	4.4	4.3	A	
16	A	A	A	(4.7)	A	C	A	C	(4.6)	C	(4.6)	C	A	
17	A	A	4.7	4.7	A	A	4.4	A	4.5	4.5	4.4	4.2	A	
18	4.2	(4.4)	A	C	C	(4.4)	C	C	C	4.5	4.2	4.2	4.3	
19	4.4	H	A	4.5	4.7	C	C	C	(4.3)	C	(4.5)	C	(4.4)	
20	4.0	(4.2)	A	(4.8)	A	4.7	C	C	(4.5)	C	(4.3)	C	C	
21	4.5	4.4	4.5	A	A	A	A	4.4	4.4	(4.3)	4.4	4.2	4.3	
22	4.4	C	C	4.6	A	A	(4.3)	C	C	4.2	4.2	4.2	4.2	
23	4.3	A	A	A	C	C	A	A	A	4.6	4.5	4.1	4.3	
24	4.3	4.5	F	4.3	4.3	(4.6)	C	C	A	(4.6)	C	4.5	4.3	
25	A	4.4	A	4.8	A	A	C	C	C	4.4	(4.5)	F	4.4	
26	4.3	H	C	C	C	C	C	C	C	(4.6)	C	(4.8)	A	
27	4.2	F	4.5	(4.5)	A	4.8	C	A	C	4.5	4.2	(4.3)	A	
28	4.2	H	4.1	H	4.7	H	A	4.5	C	4.2	4.2	(4.3)	4.5	
29	4.4	K	4.5	K	(4.1)	K	4.5	F	4.2	4.3	4.4	4.3	4.7	
30	A	K	4.3	K	4.5	K	4.5	K	C	K	4.5	K	(4.5)	
31	A	K	4.1	K	4.5	K	A	K	4.4	K	4.3	K	4.4	
Median	4.3	4.4	4.5	4.5	4.6	4.5	4.5	4.4	4.5	4.4	4.4	4.5	4.5	
Count	23	17	20	21	10	9	9	14	20	26	30	28	7	
Sweep 1.0 Mc to 25.0 Mc in 0.25 min														
Monaural □ Automatical ■														

Table 56

Ionospheric Storminess at Washington, D. C.August 1948

Day	Ionospheric character*		Principal storms		Geomagnetic character**	
	00-12 GCT	12-24 GCT	Beginning GCT	End GCT	00-12 GCT	12-24 GCT
1	4	5	---	---	3	3
2	4	4	---	2400	3	3
3	2	2			3	2
4	1	2			3	3
5	1	2			2	1
6	1	3			2	2
7	2	4	1100	---	3	3
8	4	6	----	---	6	6
9	5	5	----	---	5	4
10	4	5	----	---	5	5
11	5	2	----	1700	4	4
12	4	4	0700	---	5	3
13	4	4	---	2300	4	3
14	1	3			1	3
15	1	1			2	3
16	1	1			1	1
17	1	1			1	2
18	1	2			2	1
19	1	3			0	2
20	2	1			5	3
21	2	1			3	3
22	2	2			2	3
23	1	1			3	2
24	1	2			3	2
25	0	3			3	2
26	1	1			1	1
27	2	2			1	1
28	1	2			2	2
29	4	5	0300	---	4	4
30	4	5	----	---	5	2
31	4	5	---	2100	3	3

*Ionosphere character figure (I-figure) for ionospheric storminess at Washington, D. C., during 12-hour period, on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

**Average for 12 hours of Cheltenham, Maryland, geomagnetic K-figures on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

***No readable record. Refer to table 45 for detailed explanation.

/Dashes indicate continuing storm.

Table 57

Sudden Ionosphere Disturbances Observed at Washington, D. C.August 1948

Day	GCT		Location of transmitters	Relative intensity at minimum*	Other phenomena
	Beginning	End			
1	1612	1630	Ohio, D.C., England, New Brunswick	0.01	
1	1934	2020	Ohio, D.C., England, New Brunswick	0.02	
4	1716	1730	Ohio, D.C., England	0.5	
4	2011	2100	Ohio, D.C., England, New Brunswick	0.3	
5	1330	1420	Ohio, D.C., England, New Brunswick	0.2	
5	2011	2030	Ohio, D.C., England	0.0	Terr.mag.pulse** 2012-2025
6	1139	1205	Ohio, D.C.	0.3	
7	1348	1420	Ohio, D.C., England	0.03	Terr.mag.pulse** 1345-1352
16	1840	1905	Ohio, D.C.	0.2	
29	1452	1505	Ohio, D.C., England, New Brunswick	0.2	
31	1829	1845	Ohio, D.C., England, New Brunswick	0.0	

*Ratio of received field intensity during SID to average field intensity before and after, for station W8XAL, 6080 kilocycles, 600 kilometers distant.

**As observed on Cheltenham magnetogram of the United States Coast and Geodetic Survey.

Table 58

Sudden Ionosphere Disturbances Reported by Engineer-in-Chief.Cable and Wireless, Ltd., as Observed in England

1948 Day	GCT		Receiving station	Location of transmitters
	Beginning	End		
August	5	0825	0900	Brentwood
				Austria, Belgian Congo, Eritrea, Greece, India, Iran, Kenya, Madagascar, Palestine, Southern Rhodesia, Spain, Syria, U.S.S.R., Yugoslavia, Zanzibar
		0825	0842	Somerton
	5	1335	1355	Brentwood
				Afghanistan, Bahrein Is., Belgian Congo, Bulgaria, Canary Is., Chile, France, Greece, Iran, Kenya, Palestine, Portugal, Southern Rhodesia, Spain, Syria, Thailand, Turkey, U.S.S.R., Venezuela, Zanzibar
		1335	1415	Somerton
	6	1125	1220	Brentwood
				Austria, Bahrein Is., Belgian Congo, Chile, France, Greece, India, Iran, Kenya, Palestine, Panama, Portugal, Southern Rhodesia, Spain, Switzerland, Syria, Transjordan, Turkey, U.S.S.R., Yugoslavia, Zanzibar
		1140	1215	Somerton
9	9	1027	1135	Brentwood
				Austria, Belgian Congo, Chile, Greece, India, Iran, Kenya, Malta, Palestine, Panama, Portugal, Southern Rhodesia, Spain, Switzerland, Syria, Turkey, U.S.S.R., Yugoslavia, Zanzibar
		1035	1125	Somerton
	17			Australia, Brazil, Ceylon, India, New York, Union of S. Africa
		0550	0615	Brentwood
17	17	0550	0735	Somerton
				India
	17	1105	1200	Brentwood

Note: Observers are invited to send to the CRPL information on times of beginning and end of sudden ionosphere disturbances for publication as above. Address letters to the Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

Table 53

Provisional Radio Propagation Quality Figures
 (Including Comparisons with CRPL Warnings and CRPL Probable Disturbed Period Forecasts)
July 1948

Day	North Atlantic						North Pacific						Quality Figure Scale: 1 - Useless 2 - Very poor 3 - Poor 4 - Poor to fair 5 - Fair 6 - Fair to good 7 - Good 8 - Very good 9 - Excellent
	Quality figure	CRPL* Warning	Forecast of probable disturbed periods	Geo-magnetic K _{Ch}	Quality figure	CRPL* Warning	Forecast of probable disturbed periods	Geo-magnetic K _{Ch}					
	01-12 GCT	01-12 GCT	13-24 GCT		01-12 GCT	01-12 GCT	13-24 GCT		01-12 GCT	01-12 GCT	13-24 GCT		
1	6 7			2 2	8 7				2 2				
2	6 6			2 2	7 8				2 2				
3	6 6			2 3	7 8				2 3				
4	6 6			2 4	6 8				2 4				
5	6 6			3 3	7 7				3 3				
6	6 6			2 2	7 8				2 2				
7	7 6			2 2	8 7				2 2				
8	5 7			3 2	7 7				3 2				
9	7 6			2 2	7 8				2 2				
10	5 6			3 2	6 7				3 2				
11	6 7	X		2 2	7 8				X				
12	6 7	X		3 2	7 8				X				
13	7 7			2 3	7 7				2 2				
14	6 6			4 3	5 6				4 3				
15	7 6			2 3	6 6				2 3				
16	6 7	X		3 3	6 6				X				
17	7 6			3 3	7 7				3 3				
18	7 7			2 2	6 8				2 2				
19	6 6			1 1	6 7				1 1				
20	7 6			1 2	6 6				1 2				
21	7 6			2 3	6 7				2 3				
22	6 6			2 1	6 7				2 1				
23	7 6			2 2	6 7				2 2				
24	7 6			2 1	7 6				2 1				
25	6 6			2 2	6 7				2 2				
26	6 6	X		2 3	7 8				X				
27	6 6	X		2 2	6 5				X				
28	7 6			1 3	7 7				1 3				
29	5 5	X		4 2	6 7	X			4 2				
30	5 5	X X		3 3	7 7	X X			3 3				
31	(4) 5	X X	X	4 3	7 6	X X		X	4 3				
Score:													
H		1		1				0		0			
M		0		0				0		0			
G		28		25				28		25			
(S)		2		0				0		1			
S		0		5				3		5			

*Broadcast on WWV, Washington, D.C. Times of warnings recorded to nearest half day as broadcast.

**In addition to dates marked X, the following were designated as probable disturbed days on forecasts more than 8 days in advance of said dates: July 17 and 18.

Symbols:

X Warning given or probable disturbed date

H Quality 4 or worse on day or half day of warning

M Quality 4 or worse on day or half day of no warning

G Quality 5 or better on day of no warning

(S) Quality 5 on day of warning

S Quality 6 or better on day of warning

() Quality 4 or worse (disturbed)

Geomagnetic K_{Ch} on the standard scale of 0 to 9, 9 representing the greatest disturbance

Table 60American and Zürich Provisional Relative Sunspot NumbersAugust 1948

Date	R _A *	R _Z **	Date	R _A *	R _Z **
1	248	193	16	251	154
2	243	190	17	264	188
3	274	198	18	266	170
4	263	199	19	272	188
5	249	191	20	289	191
6	231	209	21	260	218
7	210.	175	22	223	198
8	175	150	23	178	163
9	150	130	24	179	159
10	135	117	25	197	136
11	146	117	26	175	135
12	145	88	27	174	121
13	146	115	28	167	132
14	162	107	29	142	100
15	210	134	30	160	116
			31	170	136
			Mean:	205.0	155.4

*Combination of 43 observers; see page 8.

**Dependent on observations at Zürich Observatory and its stations at Locarno and Arosa.

Table 61b

Coronal observations at Climax, Colorado (5303A), west limb

Table 62b

Coronal observations at Climax, Colorado (6374A), west limb

Table 63a

Coronal observations at Climax, Colorado (6704A), east limb

Table 63b

Coronal observations at Climax, Colorado (6704A), west limb

GRAPHS OF IONOSPHERIC DATA

42




Fig. 1. WASHINGTON, D. C.

39.0°N, 77.5°W

AUGUST 1948




Fig. 2. WASHINGTON, D. C.

AUGUST 1948




Fig. 3. BOSTON, MASSACHUSETTS

42.4°N, 71.2°W

JULY 1948




Fig. 4. BOSTON, MASSACHUSETTS

JULY 1948




Fig. 5. SAN FRANCISCO, CALIFORNIA
37.4°N, 122.2°W JULY 1948




Fig. 6. SAN FRANCISCO, CALIFORNIA JULY 1948




Fig. 7. WHITE SANDS, NEW MEXICO
32.3°N, 106.5°W JULY 1948








Fig. 8. WHITE SANDS, NEW MEXICO JULY 1948






Fig. 13. SAN JUAN, PUERTO RICO
18.4°N, 66.1°W JULY 1948




Fig. 14. SAN JUAN, PUERTO RICO JULY 1948




Fig. 15. GUAM I.
13.6°N, 144.9°E JULY 1948




Fig. 16. GUAM I. JULY 1948




Fig. 17. TRINIDAD, BRIT. WEST INDIES
10.6°N, 61.2°W JULY 1948




Fig. 18. TRINIDAD, BRIT. WEST INDIES JULY 1948




Fig. 19. PALMYRA I.
5.9°N, 162.1°W JULY 1948








Fig. 20. PALMYRA I. JULY 1948






Fig. 25. FUKAURA, JAPAN

40.6°N, 139.9°E

JUNE 1948




Fig. 26. FUKAURA, JAPAN

JUNE 1948
















Fig. 27. PEIPING, CHINA

39.9°N, 116.4°E

JUNE 1948










Fig. 40. CANBERRA, AUSTRALIA

35. 3°S, 149.0°E JUNE 1948




Fig. 41. CANBERRA, AUSTRALIA

JUNE 1948




Fig. 42. CHRISTCHURCH, N.Z.

43.5°S, 172.7°E JUNE 1948








Fig. 43. CHRISTCHURCH, N.Z.

JUNE 1948






Fig. 48. LANCHOW, CHINA
36.1°N, 103.8°E MAY 1948




Fig. 49. LANCHOW, CHINA MAY 1948




Fig. 50. NANKING, CHINA
32.1°N, 119.0°E MAY 1948









Fig. 51. NANKING, CHINA MAY 1948



Fig. 56. MADRAS, INDIA
13.0°N, 80.2°E

MAY 1948




Fig. 57. MADRAS, INDIA

MAY 1948




Fig. 58. RAROTONGA I.

21.3°S, 159.8°W

MAY 1948




Fig. 59. RAROTONGA I.

MAY 1948




Fig. 60. BRISBANE, AUSTRALIA
27.5°S, 153.0°E MAY 1948




Fig. 61. BRISBANE, AUSTRALIA MAY 1948




Fig. 62. CANBERRA, AUSTRALIA
35.3°S, 149.0°E MAY 1948




Fig. 63. CANBERRA, AUSTRALIA MAY 1948




Fig. 64. HOBART, TASMANIA

42.8°S, 147.4°E

MAY 1948




Fig. 65. HOBART, TASMANIA

MAY 1948




Fig. 66. FRASERBURGH, SCOTLAND

57.6°N, 2.1°W

APRIL 1948




Fig. 67. FRASERBURGH, SCOTLAND

APRIL 1948




Fig. 68. SLOUGH, ENGLAND
51.5°N, 0.6°W

APRIL 1948




Fig. 69. SLOUGH, ENGLAND

APRIL 1948




Fig. 70. DELHI, INDIA

28.6°N, 77.1°E

APRIL 1948




Fig. 71. DELHI, INDIA

APRIL 1948




Fig. 72. BOMBAY, INDIA

19. 0°N, 73. 0°E

APRIL 1948




Fig. 72. BOMBAY, INDIA

19. 0°N, 73. 0°E

Fig. 73. BOMBAY, INDIA




Fig. 74. MADRAS, INDIA

MADRAS, INDIA
13°0'N 80.2°F

APRIL 1948




Fig. 75. MADRAS, INDIA

APRIL 1948




Fig. 76. FALKLAND IS.

51.7°S, 57.8°W

APRIL 1948




Fig. 77. FALKLAND IS.

APRIL 1948




Fig. 78. DELHI, INDIA

28.6°N, 77.1°E

MARCH 1948




Fig. 79. DELHI, INDIA

MARCH 1948




Fig. 80. BOMBAY, INDIA
19.0°N, 73.0°E

MARCH 1948




Fig. 81. BOMBAY, INDIA

MARCH 1948




Fig. 82. MADRAS, INDIA
13.0°N, 80.2°E

MARCH 1948






Fig. 83. MADRAS, INDIA

MARCH 1948



Index of Tables and Graphs of Ionospheric Datain CRPL-F49

	<u>Table page</u>	<u>Figure page</u>
Baton Rouge, Louisiana		
July 1948	11	44
Bombay, India		
May 1948	16	55
April 1948	19	60
March 1948	20	62
Boston, Massachusetts		
July 1948	10	42
Brisbane, Australia		
June 1948	14	51
May 1948	17	57
Canberra, Australia		
June 1948	15	52
May 1948	17	57
Christchurch, New Zealand		
June 1948	15	52
Chungking, China		
June 1948	14	50
Delhi, India		
May 1948	16	55
April 1948	18	59
March 1948	19	61
Falkland Is.		
April 1948	19	61
March 1948	20	63
Fraserburgh, Scotland		
May 1948	15	53
April 1948	18	58
Fukaura, Japan		
June 1948	13	48
Guam I.		
July 1948	11	45
Hobart, Tasmania		
May 1948	18	58
Huancayo, Peru		
July 1948	12	47
Lanchow, China		
May 1948	16	54
Madras, India		
May 1948	17	56
April 1948	19	60
March 1948	20	62
Nanking, China		
May 1948	16	54

Index (CRPL-F49, continued)

	<u>Table page</u>	<u>Figure page</u>
Palmyra I.		
July 1948	12	46
Peiping, China		
June 1948	13	48
Rarotonga I.		
May 1948	17	56
San Francisco, California		
July 1948	10	43
San Juan, Puerto Rico		
July 1948	11	45
Shibata, Japan		
June 1948	13	49
Slough, England		
May 1948	15	53
April 1948	18	59
Tokyo, Japan		
June 1948	13	49
Trinidad, Brit. West Indies		
July 1948	12	46
Wakkanai, Japan		
June 1948	12	47
Washington, D. C.		
August 1948	10	42
Watheroo, W. Australia		
June 1948	14	51
White Sands, New Mexico		
July 1948	10	43
Wuchang, China		
July 1948	11	44
Yamakawa, Japan		
June 1948	14	40

CRPL and IRPL Reports

Daily:
Radio disturbance warnings, every half hour from broadcast station WWV of the National Bureau of Standards.
Telephoned and telegraphed reports of ionospheric, solar, geomagnetic, and radio propagation data.

Weekly:
CRPL-J. Radio Propagation Forecast (of days most likely to be disturbed during following month).

Semimonthly:
CRPL-Ja. Semimonthly Frequency Revision Factors for CRPL Basic Radio Propagation Prediction Reports.

Monthly:
CRPL-D. Basic Radio Propagation Predictions—Three months in advance. (Dept. of the Army, TB 11-499—
monthly supplements to TM 11-499; Dept. of the Navy, DNC-13-1 (), monthly supplements to
DNC-13-1.)
CRPL-F. Ionospheric Data.

Quarterly:
*IRPL-A. Recommended Frequency Bands for Ships and Aircraft in the Atlantic and Pacific.
*IRPL-H. Frequency Guide for Operating Personnel.

Nonscheduled reports:

- CRPL-1-1. Prediction of Annual Sunspot Numbers.
- CRPL-1-2, 3-1. High Frequency Radio Propagation Charts for Sunspot Minimum and Sunspot Maximum.
- CRPL-1-3. Some Methods for General Prediction of Sudden Ionospheric Disturbances.
- CRPL-1-4. Observations of the Solar Corona at Climax, 1944-46.
- CRPL-1-5. Comparison of Predictions of Radio Noise with Observed Noise Levels.
- CRPL-1-6. The Variability of Sky-Wave Field Intensities at Medium and High Frequencies.
- CRPL-7-1. Preliminary Instructions for Obtaining and Reducing Manual Ionospheric Records.
- NBS Circular 462. Ionospheric Radio Propagation.
- NBS Circular 465. Instructions for the Use of Basic Radio Propagation Predictions.

Reports issued in past:

- IRPL-C61. Report of the International Radio Propagation Conference, 17 April to 5 May 1944.
- IRPL-G1 through G12. Correlation of D. F. Errors With Ionospheric Conditions.

IRPL-R. Nonscheduled reports:

- R4. Methods Used by IRPL for the Prediction of Ionosphere Characteristics and Maximum Usable Frequencies.
 - R5. Criteria for Ionospheric Storminess.
 - R6. Experimental Studies of Ionospheric Propagation as Applied to the Loran System.
 - R7. Second Report on Experimental Studies of Ionospheric Propagation as Applied to the Loran System.
 - R9. An Automatic Instantaneous Indicator of Skip Distance and MUF.
 - R10. A Proposal for the Use of Rockets for the Study of the Ionosphere.
 - R11. A Nomographic Method for Both Prediction and Observation Correlation of Ionosphere Characteristics.
 - R12. Short Time Variations in Ionospheric Characteristics.
 - R14. A Graphical Method for Calculating Ground Reflection Coefficients.
 - R15. Predicted Limits for F2-layer Radio Transmission Throughout the Solar Cycle.
 - R16. Predicted F2-layer Frequencies Throughout the Solar Cycle, for Summer, Winter, and Equinox Season.
 - R17. Japanese Ionospheric Data—1943.
 - R18. Comparison of Geomagnetic Records and North Atlantic Radio Propagation Quality Figures—October 1943 Through May 1945.
 - R19. Nomographic Predictions of F2-layer Frequencies Throughout the Solar Cycle, for June.
 - R20. Nomographic Predictions of F2-layer Frequencies Throughout the Solar Cycle, for September.
 - R21. Notes on the Preparation of Skip-Distance and MUF Charts for Use by Direction-Finder Stations. (For distances out to 4000 km.)
 - R22. Nomographic Predictions of F2-layer Frequencies Throughout the Solar Cycle, for December.
 - R23. Solar-Cycle Data for Correlation with Radio Propagation Phenomena.
 - R24. Relations Between Band Width, Pulse Shape and Usefulness of Pulses in the Loran System.
 - R25. The Prediction of Solar Activity as a Basis for the Prediction of Radio Propagation Phenomena.
 - R26. The Ionosphere as a Measure of Solar Activity.
 - R27. Relationships Between Radio Propagation Disturbance and Central Meridian Passage of Sunspots Grouped by Distance From Center of Disc.
 - R28. Nomographic Predictions of F2-layer Frequencies Throughout the Solar Cycle, for January.
 - R30. Disturbance Rating in Values of IRPL Quality-Figure Scale from A. T. & T. Co. Transmission Disturbance Reports to Replace T. D. Figures as Reported.
 - R31. North Atlantic Radio Propagation Disturbances, October 1943 Through October 1945.
 - R32. Nomographic Predictions of F2-layer Frequencies Throughout the Solar Cycle, for February.
 - R33. Ionospheric Data on File at IRPL.
 - R34. The Interpretation of Recorded Values of fEs.
 - R35. Comparison of Percentage of Total Time of Second-Multiple Es Reflections and That of fEs in Excess of 3 Mc.
- IRPL-T. Reports on tropospheric propagation:
- T1. Radar operation and weather. (Superseded by JANP 101.)
 - T2. Radar coverage and weather. (Superseded by JANP 102.)
- CRPL-T3. Tropospheric Propagation and Radio-Meteorology. (Reissue of Columbia Wave Propagation Group WPG-5.)

*Items bearing this symbol are distributed only by U. S. Navy. They are issued under one cover as the DNC-14 series.

