

CRPL-F61

National Bureau of Standards
Library, N. W. 20234.

SEP 30 1949

IONOSPHERIC DATA

ISSUED
SEPTEMBER 1949

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
CENTRAL RADIO PROPAGATION LABORATORY
WASHINGTON, D. C.

IONOSPHERIC DATA

CONTENTS

	Page
Symbols and Terminology; Conventions for Determining Median Values	2
Monthly Average and Median Values of World-Wide Ionospheric Data	4
Ionospheric Data for Every Day and Hour at Washington, D. C.	6
Ionosphere Disturbances	7
American and Zürich Provisional Relative Sunspot Numbers	8
Solar Coronal Intensities Observed at Climax, Colorado. .	9
Errata	9
Tables of Ionospheric Data	10
Graphs of Ionospheric Data	44
Index of Tables and Graphs of Ionospheric Data in CRPL-F61	68

SYMBOLS AND TERMINOLOGY; CONVENTIONS FOR DETERMINING MEDIAN VALUES

Beginning with data reported for January 1949, the symbols, terminology, and conventions for the determination of median values used in this report (CRPL-F series) conform as far as practicable to those adopted at the Fifth Meeting of the International Radio Consultative Committee (C.C.I.R.) in Stockholm, 1948, and given in detail on pages 2 to 10 of the report CRPL-F53, "Ionospheric Data," issued January 1949.

For symbols and terminology used with data prior to January 1949, see report IRPL-061, "Report of International Radio Propagation Conference, Washington, 17 April to 5 May, 1944," previous issues of the F series, in particular, IRPL-F5, CRPL-F24, F33, F50, and report CRPL-7-1, "Preliminary Instructions for Obtaining and Reducing Manual Ionospheric Records."

Following the recommendations of the Washington (1944) and Stockholm (1948) conferences, beginning with data for January 1945, median values are published wherever possible. Where averages are reported, they are, at any hour, the average for all the days during the month for which numerical data exist.

In addition to the conventions for the determination of medians given in Appendix 5 of Document No. 293 E of the Stockholm conference, which are listed on pages 9 and 10 of CRPL-F53, the following conventions are used in determining the medians for hours when no measured values are given because of equipment limitations and ionospheric irregularities. Symbols used are those given on pages 2-9 of CRPL-F53 (Appendices 1-4 of Document No. 293 E referred to above).

a. For all ionospheric characteristics:

Values missing because of A, B, C, F, L, M, N, Q, R, S, or T (see terminology referred to above) are omitted from the median count.

b. For critical frequencies and virtual heights:

Values of foF2 (and foE near sunrise and sunset) missing because of E are counted as equal to or less than the lower limit of the recorder. Values of h'F2 (and h'E near sunrise and sunset) missing for this reason are counted as equal to or greater than the median. Other characteristics missing because of E are omitted from the median count. See CRPL-F38, page 9.

Values missing because of D are counted as equal to or greater than the upper limit of the recorder.

Values missing because of G are counted:

1. For foF2, as equal to or less than foF1.
2. For h'F2, as equal to or greater than the median.

Values missing because of W are counted:

1. For foF2, as equal to or less than the median when it is apparent that h'F2 is unusually high; otherwise, values missing because of W are omitted from the median count.
2. For h'F2, as equal to or greater than the median.

Values missing for any other reason are omitted from the median count.

c. For MUF factor (M-factors):

Values missing because of G or W are counted as equal to or less than the median.

Values missing for any other reason are omitted from the median count.

d. For sporadic E (Es):

Values of fEs missing because of G (no Es reflections observed, the equipment functioning normally otherwise) are counted as equal to or less than the median foE, or equal to or less than the lower frequency count of the recorder.

Values of fEs missing for any other reason, and values of h'Es missing for any reason at all are omitted from the median count.

Beginning with data for November 1945, doubtful monthly median values for ionospheric observations at Washington, D. C., are indicated by parentheses, in accordance with the practice already in use for doubtful hourly values. The following are the conventions used to determine whether or not a median value is doubtful:

1. If only four values or less are available, the data are considered insufficient and no median value is computed.

2. For the F2 layer, if only five to nine values are available, the median is considered doubtful. The E and F1 layers are so regular in their characteristics that, as long as there are at least five values, the median is not considered doubtful.

3. For all layers, if more than half of the values used to compute the median are doubtful (either doubtful or interpolated), the median is considered doubtful.

The same conventions are used by the CRPL in computing the medians from tabulations of daily and hourly data for stations other than Washington, beginning with the tables in IRPL-F18.

MONTHLY AVERAGE AND MEDIAN VALUES OF WORLD-WIDE IONOSPHERIC DATA

The ionospheric data given here in tables 1 to 48 and figures 1 to 96 were assembled by the Central Radio Propagation Laboratory for analysis and correlation, incidental to CRPL prediction of radio propagation conditions. The data are median values unless otherwise indicated. The following are the sources of the data in this issue:

Commonwealth of Australia, Ionospheric Prediction Service of the Commonwealth Observatory:

Brisbane, Australia
Canberra, Australia
Hobart, Tasmania

French Ministry of Naval Armaments (Section for Scientific Research):
Fribourg, Germany

National Laboratory of Radio-Electricity (French Ionospheric Bureau):
Poitiers, France

All India Radio (Government of India), New Delhi, India:
Bombay, India
Delhi, India
Madras, India
Tiruchirapalli, India

Electrical Communications Laboratory, Ministry of Communications:
Fukaura, Japan
Shibata, Japan
Tokyo (Kokubunji), Japan
Wakanai, Japan
Yamakawa, Japan

New Zealand Department of Scientific and Industrial Research:
Christchurch, New Zealand (Canterbury University College Observatory)

Norwegian Defense Research Establishment, Kjeller per Lillestrom, Norway:
Oslo, Norway

South African Council for Scientific and Industrial Research:
Johannesburg, Union of South Africa

United States Army Signal Corps:
Okinawa I.

National Bureau of Standards (Central Radio Propagation Laboratory):
Baton Rouge, Louisiana (Louisiana State University)
Boston, Massachusetts (Harvard University)
Guam I.
Huancayo, Peru (Instituto Geofisico de Huancayo)
Maui, Hawaii
Palmyra I.
San Francisco, California (Stanford University)
San Juan, Puerto Rico (University of Puerto Rico)
Trinidad, British West Indies
Washington, D. C.
White Sands, New Mexico

The tables and graphs of ionospheric data are correct for the values reported to the CRPL, but, because of variations in practice in the interpretation of records and scaling and manner of reporting of values, may at times give an erroneous conception of typical ionospheric characteristics at the station. Some of the errors are due to:

- a. Differences in scaling records when spread echoes are present.
- b. Omission of values when $foF2$ is less than or equal to $foF1$, leading to erroneously high values of monthly averages or median values.
- c. Omission of values when critical frequencies are less than the lower frequency limit of the recorder, also leading to erroneously high values of monthly average or median values.

These effects were discussed on pages 6 and 7 of the previous F-series report IRPL-F5.

Ordinarily a blank space in the fEs column of a table is the result of the fact that a majority of the readings for the month are below the lower limit of the recorder or less than the corresponding values of foE . Blank spaces at the beginning and end of columns of $h'F1$, $foF1$, $h'E$, and foE are usually the result of diurnal variation in these characteristics. Complete absence of medians of $h'F1$ and $foF1$ is usually the result of seasonal effects.

The dashed-line prediction curves of the graphs of ionospheric data are obtained from the predicted zero-muf contour charts of the CRPL-D series publications. The following points are worthy of note:

- a. Predictions for individual stations used to construct the charts may be more accurate than the values read from the charts since some smoothing of the contours is necessary to allow for the longitude effect within a zone. Thus, inasmuch as the predicted contours are for the center of each zone, part of the discrepancy between the predicted and observed values as given in the F series may be caused by the fact that the station is not centrally located within the zone.
- b. The final presentation of the predictions is dependent upon the latest available ionospheric and radio propagation data, as well as upon predicted sunspot number.
- c. There is no indication on the graphs of the relative reliability of the data; it is necessary to consult the tables for such information.

The following predicted smoothed 12-month running-average Zürich sunspot numbers were used in constructing the contour charts:

<u>Month</u>	<u>Predicted Sunspot No.</u>				
	1949	1948	1947	1946	1945
December		114	126	85	38
November		115	124	83	36
October		116	119	81	23
September		117	121	79	22
August	111	123	122	77	20
July	108	125	116	73	
June	108	129	112	67	
May	108	130	109	67	
April	109	133	107	62	
March	111	133	105	51	
February	113	133	90	46	
January	112	130	88	42	

IONOSPHERIC DATA FOR EVERY DAY AND HOUR AT WASHINGTON, D. C.

The data given in tables 49 to 60 follow the scaling practices given in the report IRPL-C61, "Report of International Radio Propagation Conference," pages 36 to 39, and the median values are determined by the conventions given above under "Symbols and Terminology; Conventions for Determining Median Values."

IONOSPHERE DISTURBANCES

Table 61 presents ionosphere character figures for Washington, D. C., during August 1949, as determined by the criteria presented in the report IRPL-R5, "Criteria for Ionospheric Storminess," together with Cheltenham, Maryland, geomagnetic K-figures, which are usually covariant with them.

Table 62 lists for the stations whose locations are given the sudden ionosphere disturbances observed on the continuous field intensity recordings made at the Sterling Radio Propagation Laboratory during August 1949.

Table 63 lists for the stations whose locations are given the sudden ionosphere disturbances observed at the Brentwood and Somerton, England, receiving stations of Cable and Wireless, Ltd., for various days in July and August, 1949.

Table 64 lists for the stations whose locations are given the sudden ionosphere disturbances observed at the Platanos, Argentina, receiving station of the International Telephone and Telegraph Corporation for two days in July 1949.

Table 65 lists for the stations whose locations are given the sudden ionosphere disturbances observed at the Point Reyes, California, receiving station of RCA Communications, Inc., for August 6, 19, and 30, and September 5 and 9, 1949.

Table 66 gives provisional radio propagation quality figures for the North Atlantic and North Pacific areas, for 01 to 12 and 13 to 24 GCT, July 1949, compared with the CRPL daily radio disturbance warnings, which are primarily for the North Atlantic paths, the CRPL weekly radio propagation forecasts of probable disturbed periods, and the half-day Cheltenham, Maryland, geomagnetic K-figures.

The radio propagation quality figures are prepared from radio traffic and ionospheric data reported to the CRPL, in a manner basically the same as that described in IRPL-R31, "North Atlantic Radio Propagation Disturbances, October 1943 through October 1945," issued February 1, 1946. The scale conversions for each report are revised for use with the data beginning January 1948, and statistical weighting replaces what was, in effect, subjective weighting. Separate master distribution curves of the type described in IRPL-R31 were derived for the part of 1946 covered by each report; data received only since 1946 are compared with the master curve for the period of the available data. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. Each report is given a statistical weight which is the reciprocal of the departure from linearity. The half-daily radio propagation quality figure, beginning January 1948, is the weighted mean of the reports received for that period.

These radio propagation quality figures give a consensus of opinion of actual radio propagation conditions as reported by the half day over the two general areas. It should be borne in mind, however, that though the quality may be disturbed according to the CRPL scale, the cause of the disturbance is not necessarily known. There are many variables that must be considered. In addition to ionospheric storminess itself as the cause, conditions may be reported as disturbed because of seasonal characteristics such as are particularly evident in the pronounced day and night contrast over North Pacific paths during the winter months, or because of improper frequency usage for the path and time of day in question. Insofar as possible, frequency usage is included in rating the reports. Where the actual frequency is not shown in the report to the CRPL, it has been assumed that the report is made on the use of optimum working frequencies for the path and time of day in question. Since there is a possibility that all the disturbance shown by the quality figures is not due to ionospheric storminess alone, care should be taken in using the quality figures in research correlations with solar, auroral, geomagnetic, or other data. Nevertheless, these quality figures do reflect a consensus of opinion of actual radio propagation conditions as found on any one half day in either of the two general areas.

AMERICAN AND ZÜRICH PROVISIONAL RELATIVE SUNSPOT NUMBERS

Table 67 presents the daily American relative sunspot number, R_A , computed from observations communicated to CRPL by observers in America and abroad. Beginning with the observations for January 1948, a new method of reduction of observations is employed such that each observer is assigned a scale-determining "observatory coefficient," ultimately referred to Zürich observations in a standard period, December 1944 to September 1945, and a statistical weight, the reciprocal of the variance of the observatory coefficient. The daily numbers listed in the table are the weighted means of all observations received for each day. Details of the procedure are given in the Publication of the Astronomical Society of the Pacific, issued February 1949, in an article entitled "Reduction of Sunspot-Number Observations." The American relative sunspot number computed in this way is designated R_A . It is noted that a number of observatories abroad, including the Zürich observatory, are included in R_A . The scale of R_A was referred specifically to that of the Zürich relative sunspot numbers in the standard comparison period; since that time, R_A is influenced by the Zürich observations only in that Zürich proves to be a consistent observer and receives a high statistical weight. In addition, this table lists the daily provisional Zürich sunspot numbers, R_Z .

SOLAR CORONAL INTENSITIES OBSERVED AT CLIMAX, COLORADO

In tables 68a and 68b are listed the intensities of the green (5303A) line of the emission spectrum of the solar corona as observed during August 1949 by the High Altitude Observatory of Harvard University and the University of Colorado at Climax, Colorado, for east and west limbs, respectively, at 5-degree intervals of position angle north and south of the solar equator at the limb. Beginning January 11, 1949, the actual measurements are on solar rotation coordinates rather than astronomical coordinates; thus values of the correction P given in previous coronal tables are omitted. The time of observation is given to the nearest tenth of a day, GCT. The tables of coronal observations in CRPL-F29 to F41 listed the data on astronomical coordinates; the present format on solar rotation coordinates is in conformity with the tables of CRPL-1-4, "Observations of the Solar Corona at Climax, 1944-46."

Tables 69a and 69b give similarly the intensities of the first red (6374A) coronal line; tables 70a and 70b list the intensities of the second red (6704A) coronal line. The following symbols are used in tables 68, 69 and 70: a, observation of low weight; -, corona not visible; and x, position angle not included in plate estimates.

ERRATA

1. CRPL-F60, p. 19, table 38: The latitude of Tiruchirapalli, India, should be 10.8°N instead of 12.0°N .
2. In the case of tables and graphs of data from Poitiers, France, first published in CRPL-F56 and continuing through this issue, the longitude should be changed from 2.0°W to 0.3°E . Points on the curves should be moved 0.13 of an hour to the right.

TABLES OF IONOSPHERIC DATA

Table 1

Weehington, D. C. (39.0°N, 77.5°W)

August 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	5.4				2.2	2.8	2.7
01	280	5.2				2.4	2.7	
02	280	5.0				2.7		
03	280	4.7				2.7		
04	270	4.3				2.8		
05	270	3.9			---	2.8		
06	260	4.9	250	---	120	2.1	1.9	3.1
07	280	5.8	240	4.1	110	2.7	3.4	3.1
08	330	6.5	220	4.7	110	3.1	3.8	3.0
09	350	7.6	220	4.7	110	3.4	3.8	2.9
10	330	7.3	205	4.9	100	3.7	3.5	2.9
11	350	7.5	210	(5.3)	100	3.8	3.4	2.8
12	350	7.8	200	5.3	100	4.0	3.0	2.8
13	350	7.9	220	5.4	100	3.9	3.0	2.8
14	350	7.8	220	(5.4)	110	3.8	3.0	2.8
15	340	7.8	230	4.9	110	3.6	2.1	2.8
16	320	7.8	230	4.8	110	3.3	2.3	2.8
17	300	7.7	235	4.4	110	2.9	3.6	2.8
18	280	7.9	250	---	120	2.3	3.3	2.9
19	250	8.4				2.9	2.9	
20	250	7.9				2.3	2.9	
21	250	7.0				2.3	2.8	
22	260	6.5					2.8	
23	270	5.9					2.8	

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3

Boston, Massachusetts (42.4°N, 71.2°W)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	276	6.7				2.7		
01	272	6.8				2.8		
02	272	5.4				2.8		
03	288	4.8				2.8		
04	280	4.4				2.8		
05	280	5.4				3.0		
06	300	6.0	---	---		3.0		
07	350	6.4	295	4.9		2.9		
08	400	6.8	240	4.8		2.7		
09	390	6.8	250	4.9		2.7		
10	400	6.9	250	4.9		2.6		
11	440	7.2	---	---		2.8		
12	475	7.2	---	---		2.6		
13	490	7.4	---	---		2.6		
14	430	7.4	250	6.0		2.7		
15	370	7.2	252	4.9		2.7		
16	365	7.3	270	4.7		2.7		
17	306	7.8	---	---		2.7		
18	290	8.0	---	---		2.8		
19	275	8.0	---	---		2.8		
20	270	8.3	---	---		2.8		
21	275	7.8				2.7		
22	280	7.4				2.8		
23	275	7.2				2.7		

Time: 75.0°W.

Sweep: 0.8 Mc to 14.0 Mc in 1 minute.

Table 2

Oslo, Norway (60.0°N, 11.0°E)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	275	6.9						
01	280	6.5						2.4
02	280	8.0						2.5
03	280	6.0						2.9
04	295	8.0	260		---	---	1.9	3.3
05	300	8.4	245		---	115	2.3	3.0
06	340	6.8	240		4.0	110	2.6	3.2
07	345	6.8	230		4.5	110	2.8	3.7
08	350	7.0	215		4.7	105	3.0	3.9
09	350	7.1	220		4.9	105	3.2	4.1
10	350	7.0	210		5.0	100	3.4	4.5
11	368	7.0	210		5.0	100	3.5	4.4
12	370	6.8	210		6.0	100	3.6	4.6
13	356	6.8	210		5.1	100	3.6	4.4
14	378	6.8	210		5.1	100	3.5	4.1
15	365	6.6	210		5.0	100	3.4	3.8
16	350	8.8	215		4.9	105	3.2	3.9
17	338	8.8	225		4.9	110	3.0	3.8
18	300	6.7	240		---	110	2.8	3.7
19	270	8.9	250		---	110	2.4	4.1
20	280	6.8	242		---	130	2.1	3.6
21	260	6.9	---	---	---	---	1.8	2.8
22	268	6.9						2.5
23	260	6.8						2.5

Time: 15.0°E.

Sweep: 1.8 Mc to 10.0 Mc in 6 minutes, automatic operation.

Table 4

San Francisco, California (37.4°N, 122.2°W)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	300	6.8						2.9
01	305	5.7						2.5
02	300	5.5						2.5
03	290	5.6						2.6
04	300	5.1						2.5
05	300	6.2						2.6
06	260	8.0	250		---	120	2.4	2.6
07	300	8.8	240		4.1	120	3.0	2.5
08	340	8.1	220		4.7	120	3.3	2.6
09	360	8.6	210		5.2	120	3.6	2.6
10	360	9.0	220		5.2	120	3.7	2.6
11	360	9.0	200		6.4	120	3.8	2.6
12	360	9.2	220		5.4	120	3.8	2.6
13	360	9.0	210		5.4	120	3.9	2.5
14	360	8.9	220		5.4	120	3.8	2.6
15	350	8.6	230		5.2	120	3.6	2.6
16	335	8.0	230		5.0	120	3.3	2.6
17	320	7.9	240		4.7	120	3.2	2.7
18	280	7.6	240		---	120	2.6	2.8
19	260	7.8			---	---	2.2	2.8
20	280	7.6					2.4	2.8
21	260	7.4					2.6	2.8
22	270	6.4					2.6	2.8
23	280	6.9					3.0	2.6

Time: 120.0°W.

Sweep: 1.3 Mc to 18.0 Mc in 4 minutes.

Table 5

White Sands, New Mexico (32.3°N , 106.5°W)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	310	6.4					5.2	2.5
01	300	6.4					4.2	2.6
02	280	5.9					3.2	2.6
03	280	5.5					3.0	2.6
04	280	5.3					2.9	2.6
05	300	5.0					3.9	2.7
06	250	6.3	250	---	110	(2.3)	4.9	2.8
07	300	7.4	(230)	4.4	110	2.8	5.0	2.7
08	340	8.3	230	5.0	110	3.2	5.1	2.6
09	360	8.6	220	5.3	110	3.5	5.4	2.5
10	380	9.2	210	5.3	110	3.7	5.9	2.5
11	380	9.4	220	5.3	110	3.8	5.4	2.5
12	380	10.0	220	5.3	110	3.9	6.0	2.5
13	370	9.8	220	5.3	110	3.9	5.4	2.6
14	360	9.8	220	6.2	110	3.9	5.4	2.6
15	360	9.4	230	5.1	110	3.7	5.2	2.6
16	360	9.1	230	5.0	110	3.4	5.0	2.6
17	340	8.5	230	4.6	110	3.0	4.7	2.7
18	(280)	8.2	---	—	110	2.4	5.1	2.7
19	280	8.1					5.0	2.8
20	260	8.0					4.8	2.8
21	270	7.3					3.3	2.7
22	280	6.9					3.8	2.7
23	300	6.3					4.9	2.6

Time: 105.0°W .

Sweep: 0.6 Mc to 14.0 Mc in 2 minutes.

Table 6

Baton Rouge, Louisiana (30.5°N , 91.2°W)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	6.3						3.1
01	280	6.2						3.0
02	270	5.8						2.7
03	270	5.5						3.0
04	280	4.8						3.0
05	280	5.1						3.1
06	280	6.2	240	---	120	2.4	3.7	3.2
07	290	7.0	230	4.4	120	2.9	4.2	3.1
08	300	7.4	220	4.6	120	3.3	4.2	3.0
09	340	7.6	215	5.0	120	3.5	2.9	
10	385	7.7	225	5.2	120	3.6	2.8	
11	380	8.7	220	5.3	120	(3.7)	4.0	2.8
12	375	8.8	200	(5.5)	(110)	(3.6)		2.8
13	370	9.1	230	5.5	---	(3.7)		2.8
14	350	9.2	235	5.3	115	3.6	4.0	2.8
15	350	8.8	230	5.0	110	3.5		2.8
16	330	8.3	225	4.8	110	3.4	3.9	2.9
17	310	8.3	230	4.4	120	(3.0)	4.0	2.9
18	290	7.9	260	---	120	2.5	4.0	3.0
19	270	7.6						3.6
20	240	7.5						3.5
21	265	6.9						3.8
22	280	8.6						3.5
23	290	6.4						4.0

Time: 90.0°W .

Sweep: 2.12 Mc to 15.3 Mc in 5 minutes, automatic operation.

Table 7

Okinawa I. (26.3°N , 127.7°E)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	9.6					4.1	2.8	
01	8.8					4.8	3.0	
02	(8.8)					3.7	2.9	
03	8.2					3.6	(3.0)	
04	7.0						2.9	
05	6.6						3.0	
06	7.1						3.2	
07	8.2					3.6	3.3	
08	7.8					4.8	3.1	
09	7.6					5.6	3.0	
10	8.0					6.0	2.6	
11	9.0	(5.4)				6.2	2.6	
12	10.2					5.8	2.7	
13	10.8	5.6				5.4	2.7	
14	11.2	(5.3)				5.5	2.7	
15	11.6	(5.3)				5.1	2.8	
16	12.0					5.9	2.8	
17	12.2					5.2	2.8	
18	12.0					5.4	3.0	
19	10.5					5.0	3.0	
20	9.4						2.8	
21	8.9						2.7	
22	8.6					3.4	2.7	
23	8.4						2.7	

Time: 135.0°E .

Sweep: 3.2 Mc to 18.0 Mc in 15 minutes, manual operation.

Table 8

Maui, Hawaii (20.6°N , 156.5°W)

July 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	9.3						2.0
01	280	(9.1)						2.3
02	280	8.5						2.8
03	260	8.1						1.7
04	270	7.2						1.5
05	270	6.5						1.6
06	260	6.5					130	2.9
07	230	7.0	---	---	110	2.5	3.9	2.6
08	225	7.4	215	5.1	100	3.1	4.0	2.6
09	290	8.4	200	5.3	100	3.4	4.2	2.4
10	440	9.0	210	5.6	110	3.8	4.4	2.3
11	470	9.7	210	5.5	110	3.8	4.3	2.3
12	420	10.4	210	5.4	100	3.9	4.2	2.4
13	410	11.0	215	5.4	105	3.9	4.6	(2.5)
14	410	11.2	210	5.4	100	3.9	4.9	2.5
15	380	11.6	210	5.3	100	3.7	4.4	2.7
16	350	12.2	230	5.0	108	3.4	4.6	(2.6)
17	310	12.0	230	4.6	110	3.0	4.6	(2.9)
18	280	11.5	240	---	110	2.4	4.1	3.0
19	280	10.4						4.0
20	270	9.5						3.9
21	290	9.3						3.6
22	290	9.5						2.8
23	290	9.4						3.0

Time: 150.0°W .

Sweep: 1.0 Mc to 26.0 Mc in 15 seconds.

Table 9

San Juan, Puerto Rico (18.4°N , 66.1°W)

July 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{F1}$	foF1	h°E	foE	fEs	(M3000) F2
00	295	9.0						2.7
01	280	9.2						2.8
02	270	8.8						2.8
03	275	8.1						2.8
04	260	8.1						2.8
05	255	7.3						2.9
06	265	7.2						2.8
07	270	7.8	---	---				2.9
08	295	8.5	4.5	3.2				2.9
09	330	9.0	5.0	3.5	4.3	2.7		
10	365	9.6	5.3	3.7	4.5	2.5		
11	390	10.2	5.5	3.8	5.0	2.5		
12	380	10.6	5.5	3.9	5.1	2.8		
13	380	11.1	5.5	3.9				2.8
14	360	11.4	5.4	3.9	5.1	2.5		
15	360	11.5	5.1	3.8	4.5	2.8		
16	350	11.2		4.9	3.6	4.7	2.6	
17	320	11.2	4.4	3.1	4.3	3.6		
18	300	10.2	---		4.3	2.7		
19	280	10.0			4.2	2.7		
20	290	9.7						2.7
21	290	9.5						2.6
22	300	9.5						2.6
23	300	9.3						2.7

Time: 60.0°W .

Sweep: 2.8 Mc to 13.0 Mc in 9 minutes, supplemented by manual operation.

Table 10

Guam 1. (13.6°E , 144.9°E)

July 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{F1}$	foF1	h°E	foE	fEs	(M3000) F2
00	300	(7.3)						2.6 (2.6)
01	300	(8.8)						(2.7)
02	290	7.2						2.7
03	275	(7.0)						(3.0)
04	250	7.0						2.0
05	220	6.6						3.3
06	240	6.3						3.3
07	245	7.4	220		120	---	---	2.3
08	260	8.2	205	---	100	3.2	3.8	2.8
09	270	9.1	200	---	100	3.6	4.0	2.6
10	400	9.6	210	5.3	100	3.8	4.4	2.4
11	400	10.0	200	5.4	---	---	4.5	2.3
12	405	10.0	200	5.4	---	---	4.4	2.3
13	430	10.7	200	5.5	110	(4.0)	4.8	2.3
14	430	(10.8)	200	5.4	100	4.0	4.8	(2.5)
15	445	11.0	210	5.4	100	4.0	5.2	2.4
16	400	11.0	215	---	100	3.4	5.3	2.4
17	400	(11.1)	230	---	---	---	5.8	2.5
18	410	(11.2)	260	---	---	---	5.3	(2.5)
19	345	(10.5)	285	---	---	---	4.2	2.5
20	370	10.0						3.8
21	360	(9.0)						---
22	340	---						2.2
23	340	(7.4)						(2.8)

Time: 150.0°E .

Sweep: 1.0 Mc to 28.0 Mc in 15 seconds.

Table 11

Trinidad, Brit. West Indies (10.6°N , 61.2°W)

July 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{F1}$	foF1	h°E	foE	fEs	(M3000) F2
00	260	10.8						2.9
01	260	9.7						3.0
02	260	9.0						3.0
03	260	8.6						3.0
04	250	8.4						3.3
05	260	7.6						3.1
06	260	7.3	---	---	3.4	3.2		
07	240	7.6		120	2.7	3.4	3.1	
08	260	7.6	220	4.6	120	3.2	3.8	3.0
09	300	8.6	220	5.2	120	3.7	4.6	2.8
10	340	9.4	220	5.6	120	3.8	4.8	2.6
11	375	10.2	200	6.5	120	4.0	4.9	2.6
12	380	11.1	220	6.5	110	4.1	5.0	2.6
13	370	11.8	210	5.5	120	4.1	5.0	2.7
14	350	12.2	220	6.5	120	5.1	2.7	2.7
15	340	11.9	220	6.2	120	3.8	5.0	2.8
16	330	11.8	220	5.2	120	3.4	4.8	2.8
17	285	11.2	220	4.5	120	3.1	4.4	2.7
18	260	10.8		120	2.8	4.1	3.7	
19	270	10.6			3.3	3.8		
20	290	10.8			3.0	2.6		
21	290	11.2			2.8	2.7		
22	280	11.0			2.0	2.8		
23	270	10.8			2.9			

Time: 60.0°W .

Sweep: 1.0 Mc to 18.0 Mc, manual operation.

Table 12

Palmyra I. (5.9°N , 162.1°W)

July 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{F1}$	foF1	h°E	foE	fEs	(M3000) F2
00	260	(10.1)						2.9 (3.0)
01	260	(9.2)						2.1
02	260	8.2						2.0 (2.8)
03	250	(8.4)						2.0
04	250	6.8						3.1
05	240	5.8						
06	270	4.7						2.8
07	260	6.9						2.8
08	240	8.3						2.6
09	240	8.7	220		120	3.6	3.8	2.4
10	250	9.1	215	---	120	3.8	3.2	2.2
11	320	9.4	220	---	130	4.1		
12	360	9.9	230	6.5	130	4.2		
13	360	10.0	230	5.3	130	---		
14	350	10.4	225	6.2	130	4.0		
15	300	10.8	220	6.2	120	2.8	4.2	2.2
16	250	11.0	220	---	120	2.5	4.3	2.3
17	240	10.9	---	---	120	3.0	4.0	2.3
18	280	10.7						3.7
19	340	10.0						3.8
20	380	9.3						2.8
21	370	9.4						2.1
22	330	(9.8)						2.3 (2.4)
23	290	(11.0)						3.1 (2.8)

Time: 157.5°W .

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 26 seconds, automatic operation; 13.0 Mc to 18.0 Mc, manual operation.

Table 13

Huancayo, Peru (12.0° S, 75.8° W)

July 1949

Time	$h^{\prime}F2$	$foF2$	$h^{\prime}Fl$	$foFl$	$h^{\prime}E$	foE	fEs	(M3000)F2
00	230	7.3				2.7	3.1	
01	220	6.8				2.7	3.1	
02	230	6.5				2.7	3.2	
03	230	5.6				2.7	3.2	
04	240	4.6				2.7	3.1	
05	250	3.9				2.8	3.1	
06	280	4.4				1.3	2.8	2.9
07	240	7.2				2.4	6.7	3.0
08	220	9.1				3.0	10.4	2.8
09	280	9.5	210	5.2		3.4	10.6	2.6
10	285	9.2	210	5.2		3.8	10.7	2.5
11	320	9.0	205	5.1		3.9	10.7	2.5
12	310	8.9	200	5.3		3.9	10.7	2.4
13	320	9.0	200	5.1		3.8	10.7	2.4
14	340	9.2	200	4.9		3.7	10.7	2.4
15	210	9.2				3.4	10.6	2.4
16	220	9.1				3.0	10.4	2.4
17	250	9.1				2.3	5.6	2.4
18	295	8.5				1.2	2.7	2.6
19	320	8.0						2.4
20	300	8.2						2.5
21	250	8.4						2.8
22	240	7.6						2.9
23	230	7.2						2.8

Time: 75.0° W.

Sweep: 16.0 Mc to 0.5 Mc in 15 minutes, automatic operation.

Table 14

Oslo, Norway (60.0° N, 11.0° E)

June 1949

Time	$h^{\prime}F2$	$foF2$	$h^{\prime}Fl$	$foFl$	$h^{\prime}E$	foE	fEs	(M3000)F2
00	290	6.8						
01	300	6.4						
02	300	6.2						
03	365	6.2	300					
04	340	6.4	250					
05	330	6.6	242					
06	355	6.5	240					
07	350	6.7	230					
08	370	6.8	225					
09	370	6.9	210					
10	400	6.9	210					
11	290	7.0	210					
12	400	6.7	210					
13	400	6.7	210					
14	400	6.4	202					
15	378	6.6	210					
16	350	6.6	210					
17	360	6.7	230					
18	320	6.7	235					
19	270	6.7	250					
20	260	6.8	—					
21	270	6.9	—					
22	270	7.0	—					
23	270	7.0	—					

Time: 15.0° E.

Sweep: 1.5 Mc to 10.0 Mc in 5 minutes, automatic operation.

Table 15

Okinawa I. (26.3° N, 127.7° E)

June 1949

Time	$h^{\prime}F2$	$foF2$	$h^{\prime}Fl$	$foFl$	$h^{\prime}E$	foE	fEs	(M3000)F2
00	9.4					2.7		
01	9.4					2.9		
02	9.0					3.0		
03	8.4					3.0		
04	7.8					2.9		
05	7.3					2.9		
06	7.4					2.0		
07	8.0				4.1	3.1		
08	8.1	—			4.4	3.0		
09	8.5				5.2	2.9		
10	9.1	—			6.0	2.7		
11	9.8		5.7		5.9	2.6		
12	10.5	(5.8)			6.6	2.6		
13	11.5	5.8			5.6	2.7		
14	11.6	(5.4)			5.0	2.7		
15	11.9	5.4			5.6	2.7		
16	12.5	—			5.6	2.8		
17	12.8				5.8	2.9		
18	12.6				5.4	2.9		
19	11.3				4.3	2.9		
20	10.2				4.4	2.8		
21	9.3				3.8	2.7		
22	10.0					2.6		
23	9.8					2.7		

Time: 135.0° E.

Sweep: 3.2 Mc to 18.0 Mc in 15 minutes, manual operation.

Table 16

Johannesburg, Union of S. Africa (26.2° S, 28.0° E)

June 1949

Time	$h^{\prime}F2$	$foF2$	$h^{\prime}Fl$	$foFl$	$h^{\prime}E$	foE	fEs	(M3000)F2
00	(260)	2.8						1.8 2.8
01	(280)	2.8						1.8 2.8
02	(260)	2.9						3.6 2.8
03	(285)	2.9						2.0 2.8
04	(250)	2.9						2.8 2.9
05	(280)	2.7						3.6 2.9
06	(250)	2.7						2.9
07	230	5.8					---	(1.8) 3.3
08	220	8.0	220		---	120	2.6	3.3
09	240	9.1	220		---	110	3.1	3.3
10	250	9.9	210		4.8	110	3.4	3.2
11	260	9.8	210		4.5	110	3.5	3.1
12	260	10.1	220		---	110	(3.6) 3.8	3.1
13	250	9.8	220		---	110	3.6	4.1 3.0
14	260	9.6	220		4.5	110	3.4	4.0 3.1
15	250	9.6	220		---	110	3.1	3.8 3.0
16	240	9.7	220		---	110	2.7	3.2 3.1
17	230	9.0					100	2.1 2.5
18	210	6.7						2.0 3.3
19	(220)	4.4						2.0 3.2
20	240	3.7						2.0 3.2
21	245	3.1						1.6 3.1
22	240	3.1						1.6 3.1
23	(250)	2.9						1.6 2.9

Time: 30.0° E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 17

Oslo, Norway (60.0°N , 11.0°E)

May 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	300	6.6						
01	300	6.0						
02	300	5.0						
03	300	5.9	---	---				
04	280	5.8	---	---	140	1.7		
05	260	6.0	260	---	130	2.2		
06	350	6.2	250	4.0	110	2.6		
07	350	7.0	240	4.5	110	2.8		
08	350	7.3	220	4.7	110	3.1		
09	350	7.4	220	5.0	110	3.3		
10	350	7.0	220	5.2	100	3.4		
11	360	8.1	210	5.3	100	3.5		
12	380	8.2	210	5.3	100	3.5		
13	350	7.8	210	5.3	100	3.5		
14	370	7.7	210	5.2	100	3.5		
15	340	7.5	210	5.1	100	3.4		
16	330	7.7	235	4.8	100	3.2		
17	250	7.6	230	---	110	3.0		
18	240	7.8	---	---	110	2.7		
19	250	7.8	---	---	120	2.3		
20	260	7.8	---	---	130	1.7		
21	260	7.6	---	---	---	---		
22	270	7.1						
23	280	6.9						

Time: 15.0°E .

Sweep: 1.6 Mc to 10.0 Mc in 5 minutes.

Table 18

Wakkanai, Japan (45.4°N , 141.7°E)

May 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	300	7.3						2.6
01	300	7.2						2.5
02	300	7.0						2.5
03	300	5.7						2.5
04	300	5.9						2.5
05	290	8.0	270	---	110	2.1	2.5	2.8
06	280	8.9	260	---	110	2.7	3.2	2.8
07	300	9.2	250	4.8	100	3.2	4.4	2.8
08	300	8.8	235	---	100	3.4	4.9	2.8
09	300	8.8	230	---	100	3.5	5.2	2.8
10	350	8.5	220	5.5	100	3.5	5.0	2.7
11	370	9.2	215	5.4	100	3.5	5.0	2.5
12	380	9.8	210	5.2	100	3.5	5.4	2.7
13	350	9.7	220	5.4	100	3.6	5.2	2.7
14	340	9.3	230	5.3	100	3.5	5.4	2.7
15	340	9.2	230	---	100	3.5	5.2	2.8
16	300	9.0	230	---	100	3.2	4.5	2.8
17	295	8.5	240	---	110	2.7	4.1	2.9
18	290	8.2	250	---	110	2.3	3.5	2.8
19	260	8.0	---	---	---	1.7	3.4	2.9
20	280	7.8						2.9
21	295	7.6						3.4
22	300	7.7						2.8
23	300	7.5						2.7

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 19

Fukaura, Japan (40.6°N , 139.9°E)

May 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	310	8.1				3.0	2.6	
01	300	8.0				3.0	2.7	
02	300	7.9				2.8	2.7	
03	300	7.5				2.7	2.7	
04	300	7.3				2.4	2.5	
05	270	8.5	255	---	120	2.1	2.8	
06	260	9.6	250	---	110	2.5	3.3	2.9
07	265	10.2	240	---	110	3.2	4.5	3.0
08	290	10.0	240	---	110	3.4	4.8	2.9
09	290	9.8	230	---	110	3.5	5.2	2.7
10	350	9.8	240	---	110	3.8	5.4	2.7
11	350	10.3	---	---	110	---	6.0	2.7
12	350	10.2	270	5.5	110	---	5.8	2.7
13	360	10.4	220	5.5	---	5.5	2.7	
14	350	10.0	230	5.0	110	---	5.8	
15	315	10.7	240	5.0	110	3.5	5.5	2.8
16	300	10.3	230	---	110	3.2	5.2	2.8
17	290	9.8	250	---	110	2.9	4.8	2.9
18	270	9.5	---	---	120	2.2	3.9	2.8
19	270	9.3	---	---	---	4.5	2.8	
20	280	8.5				5.0	2.8	
21	300	8.2				3.7	2.7	
22	300	8.3				4.0	2.7	
23	300	8.3				3.6	2.5	

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 20

Shibata, Japan (37.9°N , 139.3°E)

May 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	300	8.6						2.6
01	300	8.2						3.2
02	280	7.9						2.7
03	290	7.6						2.7
04	285	7.4						2.6
05	255	8.5						2.9
06	240	9.5	230	---	110	2.7	3.9	2.9
07	250	10.3	230	---	100	3.1	5.2	3.0
08	270	10.1	230	---	100	3.4	5.8	2.9
09	290	9.9	220	5.3	100	3.6	5.8	2.7
10	330	10.4	215	5.5	100	3.8	5.8	2.7
11	345	10.5	210	5.8	100	3.7	5.0	2.7
12	340	11.0	215	5.7	100	3.8	6.1	2.7
13	330	11.2	210	5.6	100	3.8	5.5	2.7
14	340	10.9	220	5.4	100	3.7	5.2	2.7
15	320	11.4	225	5.0	100	3.6	4.8	2.8
16	300	10.8	230	4.6	100	3.3	5.8	2.8
17	290	10.3	230	---	100	2.9	5.2	2.9
18	270	10.1	---	---	110	2.2	5.2	2.9
19	250	9.5	---	---	---	---	5.5	2.9
20	270	9.0						5.5
21	300	8.6						5.4
22	300	8.8						4.3
23	300	8.8						4.0

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 21

Tokyo, Japan (35.7°N , 139.5°E)

May 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	285	8.8					3.8	2.8
01	280	8.5					4.0	2.8
02	260	8.0					3.8	2.8
03	270	7.8					3.0	2.8
04	260	7.7	---	---			2.4	2.8
05	250	8.4	230	---	100	1.8	2.8	3.0
06	230	9.7	220	---	100	2.6	3.4	3.1
07	240	10.2	230	---	100	3.2	4.8	3.1
08	250	9.8	220	5.0	100	3.5	5.9	3.0
09	300	9.9	210	5.8	100	3.6	6.4	2.8
10	310	10.5	205	5.8	100	3.8	6.5	2.8
11	315	11.1	200	5.8	100	3.9	5.7	2.8
12	320	11.5	220	5.8	100	3.9	6.3	2.8
13	320	11.5	230	5.7	100	3.8	6.0	2.8
14	315	12.0	220	5.6	100	3.7	6.2	2.8
15	300	12.0	215	5.4	100	3.8	5.8	2.9
16	280	11.4	220	5.0	100	3.4	5.9	3.0
17	260	11.0	225	---	100	2.9	5.7	3.0
18	240	10.5	---	---	100	2.1	5.2	3.0
19	240	9.6	---	---	---	---	4.8	3.1
20	250	8.7	---	---	---	---	6.2	2.9
21	275	8.7	---	---	---	---	5.2	2.8
22	300	8.9	---	---	---	---	5.8	2.8
23	280	8.9	---	---	---	---	3.8	2.8

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 22

Yamakawa, Japan (31.2°N , 130.6°E)

May 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	300	10.0						3.6
01	300	10.0						3.6
02	295	9.1						3.4
03	280	8.5						3.6
04	280	8.2						2.7
05	280	8.1						2.7
06	280	9.2	250	---	125	2.0	3.0	2.7
07	250	9.4	245	---	110	2.7	3.7	3.0
08	250	9.5	250	---	110	3.1	5.0	2.9
09	300	9.7	230	---	110	3.4	5.9	2.7
10	330	10.5	245	---	110	3.6	6.5	2.6
11	360	10.9	235	---	110	(3.7)	6.4	2.6
12	350	11.7	240	---	110	---	6.3	2.6
13	350	12.2	250	---	110	4.1	5.6	2.6
14	350	13.0	245	5.2	110	3.8	5.2	2.7
15	350	13.0	240	---	110	3.6	5.8	2.7
16	330	12.6	245	---	110	3.6	5.2	2.8
17	300	12.0	240	---	110	3.2	5.0	2.8
18	290	11.3	250	---	110	2.5	4.6	2.8
19	270	10.9	---	---	110	2.1	4.6	2.8
20	280	10.1	---	---	---	---	4.8	2.7
21	290	9.8	---	---	---	---	4.6	2.6
22	305	9.8	---	---	---	---	4.4	2.6
23	305	9.8	---	---	---	---	4.2	2.6

Time: 135.0°E .

Sweep: 1.2 Mc to 18.5 Mc in 15 minutes, manual operation.

Table 23

Okinawa I. (26.3°N , 127.7°E)

May 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	13.9						2.8	
01	12.7						3.9	
02	10.2						3.0	
03	9.8						3.0	
04	9.2						3.0	
05	8.8						3.0	
06	8.6						3.0	
07	9.2				3.8	3.1		
08	9.3	---			4.6	2.9		
09	10.0	---			5.4	2.7		
10	11.2	---			5.8	2.8		
11	12.3	---			5.4	2.6		
12	12.9	---			5.7	2.7		
13	13.8	6.0	---		5.7	3.7		
14	14.7	(5.6)	---		5.6	2.8		
15	14.9	---			4.8	2.8		
16	14.6	---			4.6	2.8		
17	14.8	---			4.6	2.8		
18	14.8	---			4.4	2.8		
19	14.4	---			4.0	2.8		
20	13.0	---			4.4	2.8		
21	13.4	---			3.7	2.7		
22	13.8	---					2.8	
23	13.8	---			3.8	2.8		

Time: 135.0°E .

Sweep: 2.8 Mc to 18.0 Mc in 15 minutes, manual operation.

Table 24

Brisbane, Australia (27.5°S , 153.0°E)

May 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	270	5.0						2.0
01	260	5.0						2.0
02	270	5.2						2.0
03	260	5.1						2.9
04	240	4.6						2.1
05	240	4.3						2.9
06	240	4.8					---	1.5
07	230	7.9					120	2.4
08	240	10.1					110	3.0
09	240	11.5	230	5.0	100	3.3		3.2
10	250	12.0	225	5.0	110	3.5		3.2
11	250	11.6	215	5.0	100	3.6		3.0
12	250	12.0	220	5.0	100	3.6		3.0
13	250	12.0	220	5.0	110	3.5	4.0	3.0
14	250	11.9	220	4.5	100	3.5	4.0	3.0
15	240	11.6	220	4.0	100	3.2	3.1	3.0
16	230	10.9	---	---	100	2.7		3.0
17	230	10.0	---	---	150	1.8	2.5	3.0
18	220	9.0	---	---	---	<1.5	3.6	3.0
19	240	7.6	---	---	---	---	3.0	2.9
20	245	6.7	---	---	---	---	2.0	2.9
21	245	6.0	---	---	---	---	2.0	2.9
22	250	5.5	---	---	---	---	2.0	2.8
23	250	5.3	---	---	---	---	1.9	2.8

Time: 150.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 55 seconds.

Table 25

Canberra, Australia (36.3°S , 149.0°E)

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	260	4.5				2.6	2.7	
01	270	4.6				2.2	2.7	
02	260	4.6				2.3	2.7	
03	260	4.8				3.0	2.8	
04	240	4.8				2.6	2.9	
05	215	4.2				2.6	3.0	
06	235	3.8				<1.1	2.5	3.0
07	220	6.2			150	1.9	3.3	3.1
08	220	9.0	---	---	100	2.6	3.5	3.3
09	220	10.2	---	---	100	3.0	3.5	3.2
10	230	11.5	215	4.4	100	3.3	3.9	3.2
11	230	11.6	210	4.6	100	3.5	4.2	3.1
12	220	11.3	205	4.5	100	3.5	4.1	3.0
13	240	11.6	210	4.4	100	3.5	4.2	3.0
14	240	11.8	210	4.6	100	3.4	4.0	3.0
15	228	11.5	210	4.0	100	3.0	3.8	3.0
16	225	11.1	---	---	100	2.5	3.8	3.0
17	210	10.3			150	1.8	3.5	3.0
18	210	8.4				<1.5	3.5	3.0
19	220	7.5				3.4	3.0	
20	220	6.4				2.9	2.9	
21	240	5.2				2.6	2.9	
22	250	4.8				2.8	2.8	
23	250	4.6				2.6	2.7	

Time: 150.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 1 minute 65 seconds.

May 1949

Table 26

Hobart, Tasmania (42.8°S , 147.4°E)

May 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	4.0					2.6	2.7
01	290	4.2					2.0	2.6
02	280	4.0					2.5	2.7
03	270	3.9					3.0	2.8
04	255	3.8					2.2	2.8
05	240	3.8					2.1	2.9
06	240	3.4					2.5	3.0
07	250	4.4					2.0	3.1
08	240	7.4					2.2	3.4
09	240	9.6					2.8	3.4
10	240	10.2					3.0	3.7
11	240	(9.4)					2.2	(3.2)
12	240	---					2.3	---
13	240	(10.6)					2.3	3.1 (3.2)
14	240	(10.2)					2.0	(3.2)
15	240	(10.4)					2.8	3.0 (3.0)
16	230	(10.4)					2.3	(3.2)
17	230	10.0					2.8	3.3
18	240	(9.0)					2.1	(3.2)
19	240	7.0					2.1	3.2
20	230	6.0					2.0	3.1
21	240	6.3					2.7	3.0
22	250	(4.8)					2.8	(2.8)
23	265	4.0					2.4	2.6

Time: 150.0°E .

Sweep: 1.0 Mc to 13.0 Mc in 1 minute 55 seconds.

Table 27

Christchurch, New Zealand (43.5°S , 172.7°E)

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	4.9				3.0	2.7	
01	300	4.8				3.2	2.6	
02	300	4.8				3.5	2.6	
03	295	4.7				3.5	2.7	
04	280	4.5				3.5	2.8	
05	260	4.2				2.7	2.9	
06	250	3.6				3.5	2.9	
07	250	5.0			(1.6)	3.2	3.1	
08	230	8.3				1.8	4.3	3.2
09	230	9.9	---	---		2.6	4.4	3.2
10	230	10.7	---	---		2.9	4.4	3.2
11	240	11.3	230	4.5		3.2	5.1	3.1
12	230	11.8	---	---		3.2	5.2	3.0
13	240	12.0	---	---		3.2	6.0	3.0
14	240	11.8	---	---		3.0	4.5	3.0
15	240	11.7				2.6	4.4	3.0
16	230	11.3				2.3	4.4	3.0
17	230	9.6				1.6	3.8	3.0
18	230	8.0				4.2	2.9	
19	250	7.3				3.8	2.9	
20	250	6.2				3.5	2.8	
21	260	5.7				3.0	2.8	
22	270	5.2				3.3	2.7	
23	260	5.0				3.0	2.6	

Time: 172.5°E .

Sweep: 1.0 Mc to 12.0 Mc.

May 1949

Table 28

Poitiers, France (46.6°N , 2.0°W)

April 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	320	8.0						2.5
01	330	7.6						2.5
02	320	7.2						2.5
03	315	6.8						2.5
04	310	6.4						2.5
05	290	6.5						2.6
06	270	7.4	236					2.9
07	270	8.0	230					3.0
08	255	9.0	225					2.9
09	260	9.5	220	5.4	110	3.4	3.5	2.8
10	275	10.0	210	5.8	105	3.4	3.7	2.7
11	330	10.5	210	6.4	100	3.4	3.8	2.7
12	330	11.0	215	6.4	100	3.4	3.8	2.6
13	330	11.3	210	6.3	100	3.4	3.7	2.6
14	320	D	220	6.1	105	3.3	3.6	2.6
15	285	(11.0)	228	6.9	110	3.4		3.7
16	270	(10.8)	230					2.7
17	272	(10.6)	235					2.8
18	260	10.6	240					2.8
19	248	10.4						2.9
20	250	9.3						2.8
21	260	8.8						2.6
22	290	8.4						2.6
23	300	8.2						2.6

Time: 0.0° .

Sweep: 3.1 Mc to 11.8 Mc in 1 minute 15 seconds.

Table 29

Delhi, India (28.6°N , 77.1°E)

April 1949

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	340	10.5						
01	340	10.1						
02	---	---						
03	---	---						
04	---	---						
05	340	7.5						
06	335	9.1						
07	340	11.2						
08	340	11.8						
09	360	12.4						
10	400	13.0						
11	360	(13.5)						
12	---	(14.2)						
13	---	(14.1)						
14	---	(14.2)						
15	---	(14.2)						
16	---	(13.9)						
17	---	(13.9)						
18	---	(13.8)						
19	---	(12.9)						
20	360	(12.5)						
21	380	11.9						
22	400	11.7						
23	350	11.3						

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

Table 30

Bombay, India (19.0°N , 73.0°E)

April 1949

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06								
07	330	10.1						
08	---	---						
09	420	12.6						
10	480	(13.5)						
11	---	(14.1)						
12	---	(14.3)						
13	---	---						
14	---	(14.4)						
15	---	(14.7)						
16	---	(14.9)						
17	---	(15.1)						
18	---	(16.1)						
19	---	(14.9)						
20	480	(16.7)						
21	---	(14.8)						
22	---	---						
23	---	---						

Time: Local.

Sweep: 1.8 Mc to 16.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

Table 31

Madras, India (13.0°N , 80.3°E)

April 1949

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06								
07	420	10.4						
08	480	11.2						
09	540	12.3						
10	540	12.6						
11	600	12.9						
12	600	13.2						
13	600	13.4						
14	600	13.8						
15	600	13.9						
16	600	(13.9)						
17	600	(13.9)						
18	600	(13.6)						
19	800	(13.0)						
20	600	(13.0)						
21	---	(11.9)						
22	---	(11.6)						
23	---	---						

Time: Local.

Sweep: 1.8 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 32

Tiruchirapalli, India (10.6°N , 78.6°E)

April 1949

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	360	8.6						
07	360	10.0						
08	420	12.2						
09	540	12.7						
10	570	13.0						
11	600	11.7						
12	600	12.0						
13	600	12.0						
14	600	12.6						
15	600	13.9						
16	570	13.5						
17	555	13.4						
18	600	12.8						
19	660	12.0						
20	720	12.0						
21	650	12.0						
22	680	12.1						
23	---	---						

Time: Local.

Sweep: 1.8 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

Table 22

Fribourg, Germany (48.1°N , 7.9°E)

March 1949

Time	$\text{h}^1\text{F2}$	foF2	h^1Fl	foFl	h^1E	foE	fEs	(M3000)F2
00	290	(6.8)						(2.6)
01	295	6.6						(2.7)
02	290	(6.4)						(2.6)
03	300	6.0						(2.7)
04	295	5.6						2.6
05	280	5.2						2.7
06	270	5.8	---	---	E			(2.9)
07	240	(8.0)	---	---	120	2.1		(3.1)
08	230	10.1	230	3.3	111	2.7	2.6	3.0
09	230	11.3	222	4.0	110	3.1	3.4	3.0
10	240	12.0	216	4.4	106	3.4	4.1	3.0
11	240	(12.3)	230	4.2	110	(3.6)	4.0	(3.0)
12	230	(12.6)	222	(4.2)	110	3.7	4.0	(2.9)
13	240	(12.4)	220	4.5	110	3.6		(2.9)
14	235	(12.0)	230	--	110	3.5	1.7	(2.9)
15	235	11.6	230	--	108	3.2	3.0	(2.9)
16	240	11.7	---	---	110	2.9	1.7	(2.9)
17	240	(11.2)			110	2.4	3.1	(3.0)
18	240	10.8	---	E		3.0		(2.9)
19	230	(9.1)				2.4		(3.0)
20	240	(8.5)						(2.6)
21	250	(7.9)						(2.8)
22	260	(7.3)						(2.7)
23	290	(7.0)						(2.7)

Time: Local.

Sweep: 1.6 Mc to 17.6 Mc in 10 minutes, automatic operation.

Table 24

Poitiers, France (46.6°N , 2.0°W)

March 1949

Time	$\text{h}^1\text{F2}$	foF2	h^1Fl	foFl	h^1E	foE	fEs	(M3000)F2
00								
01								
02	305	6.6						2.6
03	300	6.5						2.6
04	290	6.0						2.6
05	290	5.4						2.6
06	280	5.6						2.8
07	250	8.0	230	--				3.0
08	240	9.8	230	--	105	3.3	3.4	3.1
09	240	(11.0)	220	--	120	3.3	3.4	2.9
10	240	D	220	--	110	3.4	3.4	(3.0)
11	240	D	220	--	110	3.4	3.6	--
12	250	D	220	--	110	3.4	3.6	--
13	250	D	230	--	110	3.4	3.6	--
14	250	D	225	--	120	3.4	3.5	(2.8)
15	250	D	230	--	120	3.4	3.4	(2.9)
16	255	D	230	--	--	--	3.4	(3.0)
17	260	(11.0)	235	--	--	--	3.5	(3.0)
18	250	10.0	230	--			3.4	3.0
19	240	9.5						(3.3)
20	255	8.8						(3.2)
21	262	8.1						2.7
22	288	7.6						2.6
23	290	7.5						2.6

Time: 0.0°.

Sweep: 3.1 Mc to 11.8 Mc in 1 minute 15 seconds.

Table 35

Fribourg, Germany (48.1°N , 7.6°E)

February 1949

Time	$\text{h}^1\text{F2}$	foF2	h^1Fl	foFl	h^1E	foE	fEs	(M3000)F2
00	288	(5.6)						
01	295	5.3			1.9	2.7		
02	295	5.1				2.6		
03	305	4.6				(2.7)		
04	300	4.6				2.7		
05	276	4.0				2.6		
06	272	3.9				2.6		
07	250	6.0				2.9		
08	230	9.4	---	---	120	2.0	3.2	
09	220	(11.6)	---	---	112	2.8	3.0	3.3
10	225	12.1	---	---	110	3.2	3.6	3.2
11	225	12.4	225	--	110	3.3	3.4	(3.2)
12	225	12.4	222	--	110	3.4	3.0	
13	220	12.4	220	--	110	3.3	3.0	
14	225	12.0	---	---	105	3.3	(3.1)	
15	225	(11.9)	---	---	108	3.0	3.0	(3.1)
16	230	11.7	---	---	112	2.6	3.1	(3.1)
17	220	11.1	---	---	1.7	2.6	(3.1)	
18	220	9.8				3.0		
19	225	8.4				2.2	(3.0)	
20	238	(7.3)				2.4	(3.0)	
21	248	6.6				2.2	(3.0)	
22	255	6.3					2.8	
23	270	5.8					2.9	

Time: Local.

Sweep: 1.6 Mc to 17.6 Mc in 10 minutes, automatic operation.

Table 36

Poitiers, France (46.6°N , 2.0°W)

February 1949

Time	$\text{h}^1\text{F2}$	foF2	h^1Fl	foFl	h^1E	foE	fEs	(M3000)F2
00								
01								
02					5.4			2.6
03					5.2			2.6
04					(4.7)			2.6
05					(4.2)			(2.7)
06					(4.0)			(2.6)
07	270	6.4						3.0
08	230	9.6	220	--				3.3
09	230	D	220	--				(3.1)
10	230	D	220	--				
11	230	D	220	--				
12	230	D	220	--	125	3.4	3.6	--
13	230	D	215	--	130	3.4	3.6	--
14	230	D	220	--	120	3.4	3.5	(3.0)
15	235	D	230	--				
16	230	D	230	--				
17	240	(10.2)	230	--				
18	230	9.5	--					
19	240	8.8						
20	250	7.6						
21	260	7.1						
22	280	6.4						
23	280	6.2						

Time: 0.0°.

Sweep: 3.1 Mc to 11.8 Mc in 1 minute 16 seconds.

Table 37

Wakkanai, Japan (45.4°N , 141.7°E)

February 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	5.1					2.8	
01	280	5.2					2.8	
02	290	5.0					2.7	
03	270	4.9					2.8	
04	260	4.7					2.9	
05	260	4.8					2.8	
06	230	5.0					3.0	
07	(230)	(8.7)	---	---	---	E	2.0	(3.1)
08	215	11.6	---	---	100	2.8	3.2	
09	210	12.3	---	---	100	3.2	3.2	
10	220	12.8	---	---	100	3.4	(3.2)	
11	220	12.4	210	---	100	3.7	3.2	
12	230	12.4	---	---	100	3.5	(3.6)	3.1
13	(240)	(12.0)	---	---	(100)	(3.6)	(3.1)	
14	220	11.7	---	---	100	3.4	3.0	
15	220	11.2	---	---	100	3.2	3.0	
16	220	10.3	---	---	100	2.5	3.1	
17	220	9.6	---	---	100	1.8	1.5	3.1
18	210	8.3	---	---	100	E	1.6	3.1
19	210	7.2					3.2	
20	210	6.4					3.0	
21	230	6.2					3.0	
22	240	5.6					2.9	
23	260	5.3					2.8	

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 38

Fukaura, Japan (40.6°N , 139.9°E)

February 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	300	5.4					1.6	2.6
01	300	5.2					1.9	2.7
02	300	5.1					2.2	2.6
03	290	5.1					2.6	
04	280	5.0					2.8	
05	270	4.9					2.7	
06	275	5.0					2.9	
07	235	8.2	---	---	115	2.0	3.2	
08	225	10.0	---	---	110	2.6	3.2	
09	230	11.7	---	---	110	3.1	3.1	
10	230	11.9	---	---	110	3.3	(2.6)	3.2
11	230	12.0	---	---	110	3.5	(4.3)	3.0
12	240	12.0	---	---	(110)	---	3.0	
13	240	11.6	---	---	110	3.4	3.8	2.9
14	245	11.2	230	---	110	2.4	2.9	
15	250	11.2	230	---	110	3.0	3.1	3.0
16	240	10.6	---	---	110	2.6	2.8	3.0
17	230	10.0	---	---	120	2.0	2.2	3.0
18	230	8.7	---	---	E	2.5	2.9	
19	240	8.0					3.0	
20	240	7.4					2.2	2.9
21	250	6.6					2.0	2.8
22	270	6.1					2.8	
23	290	5.6					2.7	

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 39

Shibata, Japan (37.9°N , 139.3°E)

February 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	280	5.6				2.4	2.7	
01	270	5.6				2.1	2.8	
02	285	6.2				2.5	2.7	
03	270	5.1				2.4	2.8	
04	260	4.9				2.3	2.7	
05	270	4.6				1.5	2.8	
06	250	4.8	---	---	E	2.2	2.9	
07	220	8.4	---	---	130	2.1	2.3	3.3
08	210	10.9	---	---	100	2.8	3.4	
09	220	12.2	---	---	100	3.3	4.0	3.2
10	230	12.8	220	---	100	3.6	3.8	3.1
11	230	13.4	215	---	100	3.7	4.0	3.0
12	230	13.2	220	---	100	3.8	3.9	3.0
13	230	12.5	210	---	100	3.7	2.9	
14	230	12.2	210	---	100	3.6	2.9	
15	220	11.9	220	---	100	3.3	3.9	2.9
16	230	11.2	---	---	100	2.8	3.4	3.0
17	220	10.4	---	---	105	2.2	2.6	3.1
18	220	9.0	---	---	---	1.4	2.8	3.0
19	220	8.4	---	---	---	2.4	3.0	
20	230	7.8	---	---	---	2.4	3.1	
21	230	6.7	---	---	---	2.3	3.0	
22	250	6.1	---	---	---	2.9		
23	270	5.7	---	---	---	2.8		

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 40

Tokyo, Japan (35.7°N , 139.5°E)

February 1949

Time	h'F2	foF2	h'Fl	foFl	h'E	foE	fEs	(M3000)F2
00	260	5.8					1.8	2.9
01	250	5.6					1.7	2.9
02	250	5.5					2.0	2.9
03	240	5.0					1.8	2.9
04	230	4.8					1.6	2.9
05	245	4.4					1.8	2.8
06	250	4.7	---	---	120	2.2	3.4	
07	220	8.4	210	---	100	2.8	2.8	3.4
08	220	11.1	210	---	100	3.2	3.6	3.3
09	210	12.5	210	---	100	3.5		
10	220	13.0	205	---	100	3.7	4.0	3.0
11	225	13.5	210	---	100	4.0	4.2	3.1
12	230	13.5	205	---	100	4.0	4.2	
13	230	12.9	200	---	100	3.8	4.2	3.0
14	240	12.7	210	---	100	3.7	3.9	3.0
15	220	12.2	220	---	100	3.4	3.6	3.0
16	220	11.6	210	---	100	2.9	3.2	3.0
17	215	11.0	200	---	100	2.2		
18	210	9.5	---	---	100	1.4	2.4	3.2
19	215	8.8	---	---	100		1.6	3.1
20	220	8.0	---	---	100			3.2
21	220	7.2	---	---	100			3.1
22	230	6.5	---	---	100			3.0
23	240	6.1	---	---	100			3.0

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 41

Yamakawa, Japan (31.2°N , 130.6°E)

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	280	6.7						2.8
01	280	6.5						2.8
02	280	6.2						2.8
03	285	5.6						2.9
04	250	4.8						3.1
05	280	4.2						2.7
06	300	4.1						2.7
07	270	6.4	240	---	175	8		2.7
08	240	9.9	---	---	110	2.6		3.3
09	240	11.8	230	---	110	3.1	3.4	3.2
10	250	12.8	230	---	110	3.4	3.9	3.0
11	250	13.3	230	---	110	3.6	4.2	3.0
12	290	13.6	220	---	110	3.7	4.6	2.9
13	300	14.3	220	---	110	3.8	4.4	2.8
14	300	13.8	230	---	110	3.8	4.2	2.9
15	300	13.9	220	---	110	3.5	3.8	2.8
16	290	13.6	230	---	110	3.3	3.6	2.8
17	270	13.2	240	---	110	2.7	2.4	2.8
18	250	12.4	---	---	100	2.0		2.9
19	230	11.5					2.2	2.9
20	240	10.4						3.0
21	240	9.7						3.0
22	230	8.3						3.0
23	250	7.6						2.9

Time: 135.0°E .

Sweep: 1.2 Mc to 18.5 Mc in 15 minutes, manual operation.

Table 42

Fribourg, Germany (48.1°N , 7.8°E)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	330	3.3						2.0
01	320	3.3						2.0
02	320	3.2						(2.6)
03	315	3.3						2.7
04	300	3.0						2.8
05	285	(2.8)						2.8
06	290	2.7	---	---	---	---	---	2.8
07	250	3.4	---	---	---	---	---	2.9
08	225	(7.0)	---	---	---	---	1.8	2.5
09	225	(9.4)	---	---	120	2.3	3.2	(3.3)
10	230	10.6	---	---	115	2.7	3.3	3.3
11	230	(10.8)	---	---	116	3.0	3.3	3.2
12	230	10.6	---	---	119	3.0	3.3	3.2
13	230	10.4	230	---	124	3.0	3.2	3.2
14	240	10.8	---	---	120	2.9	3.2	3.1
15	230	10.0	---	---	120	2.6	3.2	3.2
16	220	9.0	---	---	125	1.9	3.2	3.3
17	215	7.8	---	---				3.1
18	220	(6.4)						(3.3)
19	238	(4.9)						2.5
20	255	3.8						3.2
21	282	3.4						2.6
22	320	3.3						2.8
23	355	3.2						2.6

Time: Local.

Sweep: 1.6 Mc to 17.6 Mc in 10 minutes, automatic operation.

Table 43

Poitiers, France (46.6°N , 2.0°W)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00								
01								
02	---	3.4			(2.7)			
03	---	3.2			(2.8)			
04	---	3.0						
05	---	3.0						
06	---	3.0						
07	---	4.0						
08	230	7.5			3.0			
09	220	9.4			3.4			
10	228	10.4			(3.6)	3.4		
11	230	10.5			3.6	3.3		
12	220	10.2			3.4	3.2		
13	230	10.4			3.5	3.2		
14	230	10.2			(3.5)	3.2		
15	230	10.0			(4.9)	3.3		
16	220	9.1				3.2		
17	230	8.6			(4.7)	3.2		
18	240	6.7			(4.7)	3.2		
19	240	5.4			(4.7)	3.2		
20	260	4.1			(4.7)	3.0		
21	---	3.7			(4.8)	2.8		
22	---	3.6				2.6		
23	---	3.6				2.6		

Time: 0.0° .

Sweep: 3.1 Mc to 11.8 Mc in 1 minute 15 seconds.

Table 44

Wakkanai, Japan (45.4°N , 141.7°E)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	300	3.4						2.4
01	300	3.2						2.7
02	300	3.2						2.7
03	290	3.2						2.8
04	290	3.0						2.7
05	275	3.4	---	---	---	---		2.9
06	260	3.2	---	---	---	---		2.9
07	220	5.2	---	---	110	1.6	1.8	(3.2)
08	210	(7.2)	---	---	---	2.4	(2.4)	(3.5)
09	200	(9.7)	---	---	100	2.6	(2.7)	(3.3)
10	205	10.6	---	---	---	---	---	(3.2)
11	220	11.0	---	---	---	---		3.3
12	220	10.1	---	---	---	---		3.4
13	230	10.0	---	---	---	---		3.2
14	220	9.9	---	---	---	---		3.3
15	220	9.7	---	---	---	---		3.2
16	210	8.0	---	---	---	2.0	2.4	3.4
17	210	6.8	---	---	---	1.4	2.4	3.2
18	210	6.0	---	---	---	2.0		3.3
19	210	4.5	---	---	---	1.7		3.3
20	225	5.6	---	---	---	1.9		3.1
21	280	3.2	---	---	---	1.6		2.9
22	295	3.3	---	---	---	1.8		2.8
23	290	3.4	---	---	---	1.7		2.8

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 45

Fukaura, Japan (40.6°N , 139.9°E)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	310	3.3				2.8	2.6	
01	320	3.4				2.8	2.6	
02	315	3.5				2.6	3.7	
03	300	3.2				3.4	3.7	
04	295	3.3				3.4	3.8	
05	300	3.3				3.2	3.8	
06	280	3.1				3.2	3.0	
07	340	6.0			110	1.5	2.3	3.1
08	225	8.9	---	---	110	3.3	3.6	3.3
09	230	10.2	---	---	110	3.8	3.0	3.3
10	240	11.5	320	---	110	3.3	3.2	3.3
11	240	11.5	230	---	110	3.1	3.3	3.3
12	330	11.0	220	---	110	3.2	3.6	3.3
13	245	10.6	220	---	110	3.2	3.4	3.1
14	240	10.6	230	---	110	3.0	3.2	3.0
15	230	10.0	---	---	110	2.7	3.3	3.1
16	330	9.1	---	---	110	2.3	2.9	3.3
17	330	8.0				2.8	3.0	
18	220	7.3				2.6	3.1	
19	310	5.5				2.4	3.1	
20	230	4.0				3.3	3.0	
21	270	3.3				2.3	3.7	
22	300	3.3				2.2	3.7	
23	310	3.3				3.4	2.6	

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 46

Shibata, Japan (37.9°N , 139.3°E)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	280	3.5						3.4
01	290	3.5						3.3
02	275	3.6						3.4
03	245	3.3						3.4
04	250	3.3						3.3
05	280	3.1						2.3
06	330	3.2						2.3
07	210	5.9	---	---	---	---	1.7	3.1
08	200	8.7	---	---	100	3.5	2.6	3.5
09	200	10.0	---	---	100	3.0	3.6	2.4
10	210	12.0	---	---	100	3.3	3.8	3.3
11	210	12.7	210	---	100	3.4	3.6	3.3
12	210	11.6	200	---	100	3.5	3.5	3.2
13	210	11.0	---	---	100	3.5	3.3	3.3
14	230	10.9	200	---	100	3.3	3.1	3.3
15	210	10.4	---	---	100	2.9	2.9	3.3
16	210	9.5	---	---	100	3.5	2.4	3.3
17	205	8.2	---	---	100	3.0	3.3	3.3
18	210	7.3	---	---	100	3.6	3.3	3.3
19	200	6.3	---	---	100	2.6	3.4	3.4
20	200	4.3	---	---	100	2.5	3.4	3.4
21	250	3.3	---	---	100	2.4	3.0	3.0
22	280	3.3	---	---	100	2.4	3.0	3.0
23	280	3.4	---	---	100	2.4	3.0	3.0

Time: 135.0°E .

Sweep: 1.0 Mc to 17.0 Mc in 15 minutes, manual operation.

Table 47

Tokyo, Japan (35.7°N , 139.5°E)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	300	3.5				1.8	2.8	
01	290	3.5				1.8	2.8	
02	290	2.5				1.8	3.9	
03	370	2.3				2.3	3.0	
04	250	2.3				1.6	2.8	
05	395	2.1				2.7		
06	345	3.2				3.0		
07	220	6.5	---	---	160	1.9	2.0	3.4
08	210	8.7	---	---	100	3.6	3.5	
09	220	10.7	310	---	100	3.0	3.3	3.4
10	220	12.3	220	---	100	3.5	4.0	3.3
11	230	12.3	210	---	100	3.6	4.0	3.3
12	230	11.6	220	---	100	3.6	3.6	3.3
13	230	11.4	210	---	100	3.6	4.0	3.1
14	230	11.4	215	---	100	3.4	3.6	3.1
15	230	10.9	210	---	100	3.0	3.4	3.1
16	320	9.6	210	---	100	2.6	3.2	
17	210	8.3	200	---	110	1.9	2.8	3.3
18	210	7.5	---	---		2.8	3.3	
19	210	6.8				2.6	2.6	
20	210	4.8				2.4	2.4	
21	330	3.8				3.3	3.1	
22	260	3.6				3.3	3.9	
23	380	3.6				1.8	3.8	

Time: 135.0°E .

Sweep: 1.0 Mc to 16.0 Mc in 15 minutes, manual operation.

Table 48

Yamakawa, Japan (31.3°N , 130.6°E)

January 1949

Time	$\text{h}^{\circ}\text{F2}$	foF2	$\text{h}^{\circ}\text{Fl}$	foFl	h°E	foE	fEs	(M3000)F2
00	300	4.0						3.7
01	300	3.7						2.7
02	300	3.7						2.8
03	290	3.8						2.9
04	280	3.4						3.0
05	320	2.8						2.6
06	315	3.0						3.7
07	270	4.6	---	---	110	3.3		2.9
08	330	8.3	220	---	110	2.8		2.3
09	330	10.0	225	---	110	3.3		3.3
10	230	10.9	230	---	110	3.3	3.6	3.3
11	250	11.8	325	---	110	3.6	4.5	3.3
12	270	13.0	220	---	110	4.0	4.4	3.0
13	265	12.4	330	---	110	4.2		3.0
14	280	12.5	220	---	100	4.3		2.9
15	290	12.3	225	---	100	3.3	4.0	2.9
16	260	13.0	230	---	110	3.0	3.2	2.0
17	230	10.7	220	---	110	3.4	2.8	3.1
18	315	9.0	---	---	110	1.6	2.4	3.3
19	320	8.7	---	---	110			3.3
20	210	8.0	---	---	100			3.3
21	310	6.4	---	---	100			3.1
22	330	4.7	---	---	100			3.0
23	300	4.4	---	---	100			3.0

Time: 135.0°E .

Sweep: 1.3 Mc to 18.5 Mc in 15 minutes, manual operation.

TABLE 49
IONOSPHERIC DATA
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

h'F2 , Km August, 1949
(Characteristic) (Month)

Observed at Lat 39°0'N, Long 77°50'W
Washington, D.C.

National Bureau of Standards
Scaled by J.J.S., B.E.B.
(Institution)
Calculated by B.E.B., N.C.H., J.M.W.

Day	75°W											Mean Time					
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	
1	280	280	270	270	250	240	250	250	320	330	330	330	330	340	320	320	
2	270	280	300	300	300	300	380	480	440 ^a	400 ^b	350	350	350	350	340	340	
3	330 ^c	340 ^c	320 ^c	480 ^c	380 ^c	350 ^c	300 ^c	320 ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	
4	350 ^c	340 ^c	450 ^c	450 ^c	450 ^c	450 ^c	400 ^c	400 ^c	390 ^c	330 ^c	330 ^c	330 ^c	330 ^c	330 ^c	330 ^c	330 ^c	
5	480	280	280	250	300	300	280	200	[280] ^c	350	[370] ^c	380	400	350	360	320	330
6	470	270	270	270	300	300	300	330	320	320	320	320	320	320	320	320	
7	260	260	250	260	260	280	450 ^c	350 ^c	360	370	360	360	360	360	360	360	
8	350 ^c	330 ^c	300 ^c	380 ^c	370 ^c	370 ^c	370 ^c	370 ^c	370 ^c	370 ^c	370 ^c						
9	260 ^c	310 ^c	300 ^c	280 ^c	280 ^c	270 ^c	270 ^c	270 ^c	260 ^c	330	330	360	360	360	360	360	360
10	280 ^c	280 ^c	270 ^c	270 ^c	280 ^c	300 ^c	350 ^c	350 ^c	450 ^c	450 ^c	450 ^c	450 ^c	450 ^c	450 ^c	450 ^c	450 ^c	
11	250 ^c	250 ^c	250 ^c	250 ^c	250 ^c	270 ^c	250 ^c	280	280	320	320	330	330	330	330	330	330
12	280	250	260	260	270	270	250	320	290	280	300	320	310	320	310	320	310
13	320	280	250	240	250	270	280	320	320	330	330	350	350	330	320	320	320
14	280	270	270	290	300	280	290	350	400	430	350	330	330	360	320	350	320
15	300 ^c	350 ^c	300 ^c	290 ^c	460 ^c	250 ^c	330 ^c	350 ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	
16	*300 ^c	[340 ^c]	[310 ^c]	[330 ^c]	[320 ^c]	[320 ^c]	[320 ^c]	[320 ^c]	490 ^c	500 ^c	480 ^c	450 ^c	450 ^c	450 ^c	430 ^c	370 ^c	360 ^c
17	330	330	300	280	260	250	250	260	290	300 ^c	340	350	340	350	350	350	350
18	310	[320] ^c	320	330	330	300	300	350 ^c	A ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	G ^c	
19	280	280	290	290	280	260	250	280	330	440	420	420	410	440	370	350	350
20	300	310	300	300	270	280	350	340	320	330	310	350	350	340	350	350	350
21	290	270	270	250	260	250	L	L	L	330	340	340	360	350	330	310	300
22	280	300	300	290	320	270	270	280	350	330	330	380	380	370	350	350	350
23	260	250	270	270	280	270	260	280	270	290	300	[310] ^c	320	320	330	320	320
24	270	270	270	270	260	260	250	[260] ^c	270	[320] ^c	320	340	320	330	350	370 ^c	370 ^c
25	260	250	250	250	260	250	L	L	380	270	300	360	360	340	340	320	320
26	270	270	290	270	250	250	[250] ^c	[320] ^c	350	350	350	350	350	350	350	350	350
27	270	270	250	230	250	260	L	L	280	290	330 ^c	310	320	320	320	320	320
28	[280] ^c	280	270	280	270	260	250	250	270	300	[320] ^c	360	360	370	[260] ^c	[260] ^c	[260] ^c
29	300	300	280	280	250	240	250	250	L	L	350	350	340	340	330	320	320
30	290	300	300	270	270	250	270	250	320	300	340	330	340	340	280 ^c	280 ^c	280 ^c
31	280	290	300	270	260	250	250	250	[280] ^c	300	[320] ^c	290	300	330	320	320	320
Median	280	280	280	270	270	260	280	330	330	350	350	340	320	320	280	280	270
Count	27	31	31	31	30	30	39	27	29	29	29	31	31	31	31	31	31

* SUPPLEMENTARY DATA FROM FT BELVOIR,
LAT. 38.7°N, LONG. 77.1°W.

Mc 10240 Mc m.0.25 min

Sweep 1.0 Manual □ Automatic ■

U. S. GOVERNMENT PRINTING OFFICE 1440-1200-2

TABLE 50
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.
IONOSPHERIC DATA

Form adopted June 1946
for J.J.S., B.E.B. (Institution)
Calculated by: B.E.B., N.C.H., J.M.W.

foF₂ Mc August 1, 1949
(Characteristic) (Unit)

Observed at Washington, D. C.

Lat. 39°N, Long. 77°W

75°W Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	6.7	6.3	5.9	5.6	5.5	5.2	6.1	7.5	7.3	(8.3) ³	8.6	8.6	8.4	8.5	8.4	(8.4) ³	8.4	8.4	8.4	(8.1) ³	(7.7) ³	6.7	6.8		
2	6.7	6.3	6.0	5.7	4.3	3.7	4.7 ^f	5.5 ^f	5.5 ^f	6.3 ^f	6.9	(7.2) ³	C	C	A	*49 ⁶ * A	*45 ⁶ * A	*45 ⁶ * A	*49 ⁶ * A	*5.4 ⁶ * A	*5.4 ⁶ * A	6.8	7.0		
3	4.9 ^K	4.2 ^K	2.3 ^K	1.9 ^K	1.9 ^K	2.4 ^K	3.1 ^K	4.7 ^K	5.1 ^K	5.9 ^K	<3.9 ^K	<4.3 ^K	*7.9	*7.2	*4.5 ^K	*4.5 ^K	*4.5 ^K	*4.5 ^K	*4.5 ^K	*4.5 ^K	6.2 ^K	6.2 ^K	5.9 ^K		
4	(3.8) ³	2.7 ^K	3.3 ^K	*2.6 ^K	1.9 ^K	(2.6) ³	4.1 ^F	5.7 ^F	(6.6) ³	7.9	(8.1) ³	(8.6) ³	8.5	8.5	8.4	8.4	8.4	8.4	8.4	8.4	7.3 ^F	(7.1) ³	6.6 ^F	6.5 ^F	
5	5.5 ^F	5.2 ^F	4.5 ^F	4.5 ^F	3.8 ^F	3.1 ^F	3.0 ^F	4.3	5.1	6.0	5.8	6.1	6.9	7.3	7.6	7.6	7.5	7.5	7.5	7.5	7.5	7.3	6.8 ^F	6.7	
6	6.0 ^F	5.6 ^F	5.0	4.4 ^F	4.4 ^F	3.7 ^F	3.7 ^F	4.9	5.3	5.2	5.7	6.5	6.4	6.9	7.3	7.5	7.4	7.8	7.9	7.7	7.3	7.5	6.5	6.1	5.7
7	5.3	5.0	4.5	3.9 ^F	3.9 ^F	3.2 ^F	3.4 ^F	4.9	5.6 ^K	5.4 ^K	5.8 ^K	6.7 ^K	6.1 ^K	6.1 ^K	6.5 ^K	6.2 ^K	5.9 ^K	5.9 ^K	5.9 ^K	5.9 ^K	6.7 ^K	7.0 ^K	6.8 ^K	6.5 ^K	
8	5.0 ^F	3.7 ^K	2.6 ^F	2.9 ^F	2.5 ^F	3.0 ^F	4.1 ^K	5.1 ^K	4.9 ^K	4.9 ^K	6.0 ^K	6.0 ^K	6.2 ^K	6.1 ^K	6.0 ^K	5.0 ^K	6.2 ^K	6.6 ^K	6.5 ^K	6.0 ^K	6.0 ^K	4.9 ^F	4.4 ^F		
9	4.1 ^F	3.9 ^K	3.4 ^F	3.3 ^K	2.8 ^K	(3.4) ³	4.8 ^K	5.5 ^F	6.2	6.5 ^F	6.9	6.8	7.1	(7.1) ³	7.4	7.3	7.4	7.2	7.1	7.1	6.8	6.2 ^K	5.3 ^K	4.8 ^F	
10	4.6 ^F	4.3 ^K	(4.4) ³	3.5 ^K	3.0 ^K	3.0 ^K	4.6 ^K	5.4 ^K	5.1 ^K	5.1 ^K	5.0 ^K	<47 ^K	(5.5) ³	5.5 ^K	5.3 ^K	5.7 ^K	5.6 ^K	5.7 ^K	6.5	6.5	6.7	6.4	5.3	4.7	
11	4.1	3.9	3.7	3.5	3.3	3.6	5.2	5.9	7.2	7.7 ^V	7.9	7.5	7.9	7.4	8.4	8.5	7.9	7.8	7.8	7.8	8.8	8.7	7.8	6.6	
12	5.4	5.2	4.8	4.7	4.1	4.2	4.9	5.9	7.3	7.8 ^V	8.6	8.2	7.8	7.8	7.8	7.6	7.8	8.0	8.6	8.4	7.9	7.0	6.0	5.5	
13	4.7 ^F	5.2 ^F	(5.4) ³	3.5 ^F	3.4 ^F	4.8	5.7	5.8	6.7	6.5	6.6	7.3	7.6	7.2	7.0	7.1	(7.0) ³	7.2	7.6	7.4	6.9	6.9	6.5	6.6	
14	6.6	6.4	5.7	5.7	4.7	4.3 ^K	4.9	5.7	5.5	6.4 ^F	6.5	7.2	6.7	7.1	7.2	6.7	(7.7) ³	8.1	8.8 ^K	(7.9) ³	(7.1) ³	5.7 ^K	5.3 ^K		
15	(4.0) ³	3.8 ^K	4.2 ^F	4.2 ^K	4.1 ^K	(4.3) ³	3.0 ^K	4.0 ^K	4.7 ^K	<4.2 ^K	<4.5 ^K	<4.7 ^K	<4.8 ^K	*5.5 ^K	*5.6 ^K	*5.4 ^K	*5.4 ^K	*5.4 ^K	*5.4 ^K	(6.0) ³	(6.0) ³	(6.0) ³	(5.5) ³		
16	(4.3) ³	(4.2) ³	4.0 ^K	*3.5 ^K	*3.2 ^K	*4.6 ^F	4.9 ^K	5.4 ^K	5.6 ^K	5.7 ^K	6.0 ^K	(6.4) ³	(6.3) ³	(6.0) ³	(6.0) ³	(5.8) ³	(5.4) ³								
17	5.2 ^F	5.0	4.9 ^F	4.7 ^F	(4.5) ³	(4.2) ³	(5.6) ³	(6.4) ³	7.0	7.6	7.9	7.6 ^H	8.6 ^H	8.6 ^H	8.0	*8.0	*8.1	8.6	8.4	8.5	7.4 ^F	(6.9) ³	5.7		
18	4.9 ^F	4.7 ^F	(4.4) ³	4.1 ^F	4.0 ^F	3.6 ^F	4.3	4.9	5.7	5.5	5.7	[5.7] ^A	<5.2 ^G	5.7 ^K	6.0 ^K	6.2 ^K	6.3 ^K	6.3 ^K	6.7 ^K	(6.0) ³	6.1 ^K	6.1 ^K	(5.6) ³		
19	(5.4) ³	4.8	(4.3) ³	4.1	3.6	3.5	4.7	5.4	5.4	(5.7) ^A	6.5	6.7	6.5	7.2	7.1	7.5	7.5	7.4	7.6	(7.1) ³	6.6	5.9	(5.6) ³		
20	(5.3) ³	4.9 ^F	4.9 ^F	4.7 ^F	4.2 ^F	4.8	5.8	6.5	7.3	7.8	8.2	8.3	8.4	8.4	8.2	8.4	8.4	8.6	8.7	8.8 ^V	8.8 ^V	8.9	(6.9) ³		
21	5.8 ^F	5.5 ^F	4.8 ^F	4.3 ^F	4.0	(6.0) ³	7.1	7.9	8.1	8.8	8.5	8.9	8.9	8.8	8.7	8.3 ^H	(8.6) ³	8.4	(7.0) ³	(7.0) ³	(7.0) ³	6.8			
22	(6.0) ³	6.3	5.9 ^F	(5.6) ³	5.0	4.7	5.3	6.1	6.5	7.1	7.3	7.5	7.7	7.4	7.4	(7.3) ³	(7.1) ³	6.9 ^F	6.6 ^F	(6.5) ³	6.3 ^F				
23	(5.9) ^F	(5.5) ³	5.2	4.9	4.7	4.5	5.8	7.4	8.5	8.8	9.0	9.0	9.5	9.5	9.4	9.1	8.8	8.6	8.7	(8.0) ³	(7.5) ³	6.9	6.5		
24	6	3	(6.0) ³	5.9	(5.5) ³	4.9	5.4	7.5	(8.2) ³	8.6	8.8	8.7	9.0	9.0	9.1	8.9	8.7	8.7	8.6	(8.7) ³	(8.6) ³	7.5	(7.2) ³		
25	(6.8) ³	6.5	(6.1) ³	5.5	5.0	4.9	6.3	8.1	9.0	9.5	8.9	9.0	9.1	9.4	9.4	9.2	9.1	9.0	9.0	(9.2) ³	(8.8) ³	8.1	7.6		
26	6	8	(6.4) ³	6.1	5.9	(5.6) ³	5.5	6.0	6.7	7.2	7.7	8.2	8.0	8.1	8.0	7.7	7.7	7.9	7.9	(8.0) ³	(8.0) ³	7.1	7.0		
27	6	8	6.9	6.8	(5.8) ³	5.3	5.3	6.0	7.7	(8.9) ³	9.8	9.3 ^H	9.2	10.1	10.2	9.9	9.4	9.2	9.2	(9.5) ³	(8.7) ³	7.3	6.9		
28	(6.2) ³	6.6	5.9	5.5	5.2	5.1	(6.0) ³	8.7	8.6	7.9	7.8	8.3	8.7	8.8	8.6	8.4	8.7	8.9	8.9	(9.2) ³	8.5	7.6	6.8		
29	5.8	5.7	5.6	5.2	(5.0) ³	4.8	(5.9) ³	7.1	8.2	8.5	8.9	8.7	8.8	9.2	9.5	9.7	9.7	9.7	(9.7) ³	(9.6) ³	7.9	6.6			
30	6	1	6.0	5.9	5.6	5.3	6.7	(8.4) ³	8.7	9.0	9.7	(8.9) ³	9.0	9.1	8.9	8.6	8.7	8.5	8.6	(7.9) ³	(6.9) ³	6.5	(5.8) ³		
31	5.4	(5.3) ³	5.1	5.2	4.9	(4.2) ³	(5.9) ³	7.1	8.3	9.6	9.1	9.3	9.2	9.1	9.3	9.2	9.2	9.3	(9.6) ³	9.5	8.6	6.9			
Median	6.4	5.2	5.0	4.7	4.3	3.9	4.9	6.5	7.1	7.3	7.5	7.8	7.9	7.7	7.8	7.8	7.7	7.9	8.4	7.9	7.0	6.5	5.9		
Count	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31		

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
LAT. 38°7'N, LONG. 77°10'W.

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual □ Automatic □

U. S. GOVERNMENT PRINTING OFFICE 1946 O - 17219

TABLE 51
IONOSPHERIC DATA
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

foF₂ Mc August 1, 1949
(Characteristic) (Unit) (Month)

Observed at Washington, D.C.
Lat. 39°0'N, Long. 75°W

Day	0030	0130	0230	0330	0430	0530	0630	0730	0830	0930	1030	1130	1230	1330	1430	1530	1630	1730	1830	1930	2030	2130	2230	2330	Mean Time		
																									National Bureau of Standards (Institution)		
1	6.5	6.0	(5.7) ³	(5.7) ³	5.3	5.8	7.2	7.6	(7.6) ³	(8.0) ³	8.7	8.6	(8.4) ³	8.7	8.4	8.6	8.5	8.1	8.7	(8.0) ³	(7.1) ³	6.9	6.9	6.9	6.9	6.9	6.9
2	6.7	6.0	6.0	5.2	4.0	4.1	4.9 ^f	4.5	6.0	6.8	5.9	7.2	C	6.6	6.6	6.8	6.9	(6.8) ³	8.1	8.0	6.8 ^k	5.8 ^k	5.6 ^k	5.6 ^k	5.6 ^k	5.6 ^k	
3	5.7 ^k	4.6 ^j	1.8 ^k	1.9 ^k	2.0 ^k	2.7 ^k	3.2 ^k	3.8 ^k	<3.9 ^k	B	A	A	A	<4.6 ^k	<4.6 ^k	<4.9 ^k	<4.9 ^k	*5.6 ^k	*5.5 ^k								
4	*[3.2] ^c	3.5 ^k	*3.5 ^k	N	K	*2.2 ^k	*3.1 ^k	*4.8 ^k	*4.8 ^k	*4.8 ^k	*4.8 ^k	*6.6 ^f	*6.6 ^f	(7.8) ³	*8.7	7.8	8.2	8.5	9.3	9.3	8.4	(7.5) ³	6.8	6.8 ^f	5.6 ^f	5.6 ^f	5.6 ^f
5	5.5 ^f	4.6 ^f	3.9 ^f	3.5 ^f	2.9 ^f	3.6 ^f	4.8 ^f	4.9	5.8	6.0	6.5	6.7	6.7	7.4	7.0	7.7	7.6	7.7	7.7	7.7	7.7	7.1	6.9	6.9	6.9	6.9	
6	5.7 ^f	5.3 ^f	4.7 ^f	4.2 ^f	3.7 ^f	4.3	5.3	5.6 ^k	5.6 ^k	5.7	6.2	6.7	6.9	7.6	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.2	7.2	6.4 ^f	6.4 ^f	6.4 ^f	
7	5.1	4.9	4.1	3.7 ^f	3.1 ^f	4.2	5.4 ^k	5.5 ^k	5.7	6.0 ^k	[6.0] ³	[6.0] ³	6.1 ^k	6.3 ^k	5.9 ^k	5.9 ^k	5.9 ^k	6.2 ^k	6.5 ^k	7.0 ^k	6.6 ^k	6.2 ^k	6.2 ^k	6.2 ^k	6.2 ^k		
8	3.7 ^k	4.3 ^k	2.8 ^k	2.6 ^k	2.6 ^k	4.5 ^k	4.5 ^k	5.5 ^k	5.5 ^k	5.7	6.4 ^k	6.4 ^k	6.4 ^k	6.5 ^k	6.5 ^k	6.5 ^k	6.6 ^k	6.6 ^k	6.6 ^k	6.6 ^k	6.5 ^k	6.5 ^k	6.5 ^k	6.5 ^k			
9	3.9 ^f	3.7 ^k	3.4 ^k	3.4 ^k	3.4 ^k	5.3 ^k	5.3 ^k	5.9 ^k	5.9 ^k	7.4	6.9	7.4	7.3	7.1	7.6	7.4	7.4	7.4	7.4	7.4	7.4	5.1 ^k	4.7 ^k	4.7 ^k	4.7 ^k		
10	4.6 ^k	4.3 ^k	4.0 ^k	3.1 ^k	2.8 ^k	3.7 ^k	5.1 ^k	5.7 ^k	5.7 ^k	5.3 ^k	<4.7 ^k	[5.5] ³	[5.5] ³	6.0 ^k	5.4 ^k	5.4 ^k	5.5 ^k	5.5 ^k	5.9 ^k	6.7	6.6	6.4	6.4	6.4	6.4		
11	3.9	3.7	3.5 ^f	3.3 ^f	3.3 ^f	4.3	5.7	6.5	7.4	7.8	7.3 ^H	7.7	8.1	8.7	8.0	7.8	7.7	7.5	8.4	8.9	9.2	7.4	5.9	5.4	5.4		
12	5.3	5.3 ^f	4.9	4.7	4.4 ^f	4.1	(4.3) ⁵	5.5	6.6	7.4	8.0	8.4	7.9	8.0	7.8	7.8	7.6	8.0	8.2	8.4	8.3	7.4	6.7	5.5	5.1		
13	5.3 ^f	5.3 ^f	4.8 ^f	4.0 ^f	3.4 ^f	4.1	(5.1) ⁵	(5.9) ⁵	(6.5) ⁵	6.7	6.7	6.9	7.6	7.6	7.2	7.0	7.0	7.3	7.9	7.9	7.3	6.6	6.7	(6.6) ³	(6.6) ³		
14	6.6	(5.9) ³	(5.0) ^f	4.4 ^f	4.2 ^f	3.7	(4.6) ³	4.9	5.7	6.5	6.6	6.7	(7.0) ³	(6.7) ⁵	7.2	(7.0) ³	8.0	7.8 ^k	(9.6) ³	7.6	7.6	(6.4) ³	(5.3) ³	4.9 ^k	4.9 ^k		
15	4.0 ^f	(4.0) ³	4.1 ^k	4.1 ^k	4.2 ^k	4.2 ^k	3.8 ^k	4.3 ^k	4.3 ^k	4.8 ^k	5.1 ^k	<4.6 ^k	<4.7 ^k	<4.8 ^k	<4.7 ^k	5.2 ^k	5.5 ^k	(5.6) ³	5.8 ^k	(5.9) ³	6.0 ^k	*5.9 ^k	5.3 ^k	*4.6 ^k	4.9 ^k		
16	*4.2 ^k	*4.0 ^k	*3.9 ^k	*3.2 ^k	*3.2 ^k	*3.2 ^k	*3.2 ^k	*3.7 ^k	*3.7 ^k	*4.2 ^k	*4.2 ^k	*4.7 ^k	*5.3 ^k	*5.6 ^k	*4.9 ^k	*4.9 ^k	6.0 ^k	(6.5) ³	6.5 ^k	6.2 ^k	5.2 ^k	4.9 ^k					
17	5.1 ^f	4.8	4.7 ^f	4.0 ^f	3.4 ^f	4.1	(5.1) ⁵	(5.9) ⁵	(6.5) ⁵	6.7	6.7	6.9	7.6	7.6	7.2	7.0	7.0	7.3	7.9	7.9	7.3	6.6	6.7	(6.6) ³	(6.6) ³		
18	4.9 ^f	4.6 ^f	4.4 ^f	4.1 ^f	3.8 ^f	4.1 ^f	(4.7) ³	5.3 ³	5.4 ^k	5.7 ^k	6.0 ^k	5.8 ^k	6.0 ^k	6.3 ^k	6.3 ^k	6.7 ^k	6.7 ^k	(6.6) ³	(6.6) ³	(6.6) ³	6.0	(5.7) ³	5.5 ^f	5.5 ^f	5.5 ^f		
19	(5.0) ³	(4.4) ³	4.0	3.9	3.5 ^f	4.1	5.3	5.7	6.2	6.3	6.9	6.9	6.9	7.3	7.6	(7.5) ³	7.4	(7.5) ³									
20	5.0	4.9 ^f	4.9 ^f	4.3 ^f	(4.5) ³	5.2	5.2	(6.0) ³	7.1	9.2	8.1	8.4	8.3	8.4	8.4	8.2	8.6	8.6	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	
21	5.7 ^f	5.6 ^f	5.0 ^f	4.7 ^f	3.9 ^f	4.7 ^f	6.6	7.3	7.4	7.9	8.5	8.5	8.5	8.5	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	
22	6.3	5.9	5.8	4.9	4.9	4.9	(6.4) ³	(6.4) ³	(6.4) ³	6.9	7.2	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	
23	5.8 ^f	5.2	(4.9) ³	4.9	4.4 ^f	4.8	6.5	(7.9) ⁵	8.9	8.9	(8.9) ⁵	9.1	9.5	9.6	9.1	9.1	(8.9) ³	8.8	(8.2) ³	(8.2) ³							
24	6.2	(6.0) ³	5.6	(5.4) ³	4.9	5.3	6.8	7.4	7.4	8.5	8.9	8.7	8.9	9.0	9.0	9.0	8.8	8.8	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	
25	(6.7) ⁵	6.4	(6.0) ³	5.4	5.0	(5.5) ³	7.2	8.8	9.2	9.0	8.9	9.3	9.3	9.2	9.3	9.0	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	
26	6.7	6.4	5.9	(5.6) ³	5.6	5.6	6.5	6.7	7.4	7.9	8.0	8.1	7.8	8.3	8.1	7.8	7.8	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	
27	6.8	6.9	6.3	5.7	(5.3) ³	5.3	6.7	v	8.7	9.4	9.7	10.2	9.7	9.9	10.0	(9.5) ⁵	9.3	9.2	9.1	9.0	9.0	8.9	8.9	8.9	8.9	8.9	
28	(6.4) ³	6.6	5.5	(5.3) ³	5.3	6.9	8.2	8.7	8.3	7.7	8.5	8.6	8.7	8.5	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	
29	5.7	5.6	5.2	4.9	5.3	6.5	7.6	8.1	(8.5) ³	8.9	9.1	9.5	9.7	(9.6) ³	9.3												
30	5.9	5.9	(6.0) ³	5.6	5.4 ^f	5.6	7.4	(8.8) ³	8.6	9.7	9.2	8.8	9.1	9.1	9.1	8.7	8.7	8.6	8.7	8.7	8.7	8.7	8.7	8.7	8.7	8.7	
31	(5.4) ³	(5.3) ³	5.2	5.2	(4.4) ³	4.7	6.8	7.3	9.0	9.7	9.0	9.3	(9.0) ³	(9.0) ³	(9.0) ³	9.4	9.4	9.5	(9.7) ³	8.9	8.9	8.9	8.9	8.9	8.9	8.9	
Median	5.5	5.2	4.8	4.5	4.0	4.3	5.4	6.5	7.1	7.4	7.8	8.0	7.8	7.8	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	
Count	31	31	31	30	31	31	31	31	30	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31

* SUPPLEMENTARY DATA FROM FT BELVOIR,
LAT. 38°7'N, LONG. 77.1°W.

Swept 0.0 Mc in 0.5 min

Manual □ Automatic ■

Form adopted June 1946

TABLE 52
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.
IONOSPHERIC DATA

h'F_I **Km** **August**, 1949
(Characteristic) (Unit) (Month)

Observed at **Washington, D. C.****Lat. 39.0°N, Long. 77.5°W**

Day	75°W												Mean Time												
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1																									
2																									
3																									
4																									
5																									
6																									
7																									
8																									
9																									
10																									
11																									
12																									
13																									
14																									
15																									
16																									
17																									
18																									
19																									
20																									
21																									
22																									
23																									
24																									
25																									
26																									
27																									
28																									
29																									
30																									
31																									
Median	250	240	220	220	205	210	205	210	200	190	180	170	160	150	140	130	120	110	100	90	80	70	60	50	
Count	9	27	28	49	48	26	27	29	28	27	26	25	24	23	22	21	20	21	22	23	24	25	26	27	28

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
LAT. 38.7°N, LONG. 77.1°W

Sweep I.Q. Mc 10-25.0 Mc in 9.25 min
Manual □ Automatic □

U. S. GOVERNMENT PRINTING OFFICE: 1946 O-700-18

National Bureau of Standards
(Institution)
Calculated by **B.E.B., N.G.H., J.M.W.**
Scaled by **J.J.S., B.E.B.**

Form adopted June 1946

TABLE 53
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

for f_{I}
(Characteristic) Mc
August (Month)
Observed at Washington, D.C.

Lat 39°0'N, Long 77°5'W

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	Q	Q	4.5	5.3	5.2	[5.2] ^P	5.4	5.3	5.4	5.3	5.3	5.3
2	L	4.5 ^F	4.7 ^F	5.0	[5.0] ^P	5.3	5.3 ^P	C	C	4.5	4.9	(4.2) ^A
3	Q	X	3.6 ^L	3.9 ^X	3.9 ^X	*[4.3] ^P	*[4.5] ^{K*}	*[4.6] ^{K*}	*[4.6] ^{K*}	*[4.5] ^{K*}	*[4.5] ^{K*}	(4.0) ^P
4	*Q	*Q	*4.8	*4.5	*4.8	*5.1	*5.0	*4.9	*5.0	*5.0	*5.0	4.6
5	Q	Q	L	4.8	A	B	4.9	(5.2) ^S	5.1	5.0	4.9	L
6	L	4.1	4.5	4.6	4.9	5.1	5.3	5.3	5.1	5.1	4.9	L
7	L	5.2 ^X	4.9 ^X	4.7 ^X	4.9 ^X	4.8 ^X	(4.7) ^S	4.8 ^X	4.7	4.8 ^X	4.7 ^K	4.4 ^K
8	Q	K	4.1 ^X	4.3 ^X	4.5 ^X	(4.7) ^B	4.9 ^K	5.0 ^X	A ^K	A ^K	4.8 ^X	3.9 ^K
9	Q	K	4.6 ^X	4.3	4.4	4.9	[5.0] ^A	5.0	4.8	5.0	4.8	4.6
10	Q	K	4.1 ^X	4.5 ^X	4.5 ^X	4.6 ^X	4.7 ^X	4.8 ^X	4.7 ^X	4.6 ^A	4.7 ^K	4.3 ^A
11	Q	*L	L	4.7	4.9	5.3 ^H	5.1	4.9 ^H	4.9	4.8	4.5	L
12	Q	L	4.7	4.9 ^H	4.9	5.4 ^H	5.0	4.7	4.9	4.8	L	Q
13	L	L	4.7	4.8	4.8	5.3	4.9	[5.0] ^N	(5.0) ^A	(4.3) ^S	4.3	4.0
14	Q	4.3	4.7	4.6	4.9	5.2	4.9	5.1	A ^K	L	L	K
15	L	K	3.9 ^X	(4.3) ^A	4.5 ^X	4.5 ^X	[4.7] ^{N*}	4.8 ^X	4.7 ^X	4.6 ^K	4.3 ^K	A ^K
16	*Q	*K	4.0 ^P	4.3 ^X	4.3 ^X	4.6 ^X	4.7 ^X	4.9 ^X	(5.1) ^K	4.6 ^K	5.0 ^K	L ^K
17	Q	L	L	(5.6) [?]	[5.6] ^A	(5.3) ^P	5.6	(5.5) ^P	(5.5) ^{P*}	4.9 ^L	(4.3) ^L	Q
18	L	L	X	(4.7) ^P	5.0 ^X	4.9 ^T	5.2 ^H	5.1 ^X	A ^X	A ^X	L	K
19	Q	L	4.5	4.9	5.6 ^H	5.5 ^H	5.2 ^H	5.7 ^H	5.4 ^H	5.3	4.7	L
20	L	A	(5.0) ^P	(5.4) ^P	L	L	(5.6) ^P	5.8 ^H	5.4	4.8	(4.9) ^P	L
21	Q	L	L	L	(5.3) ^P	(4.4) [?]	4.6	(5.9) ^P	5.8	(5.1) ^P	(5.6) ^P	L
22	Q	L	L	L	(5.5) ^P	(5.5) ^P	[5.6] ^T	5.5	(5.6) ^P	(5.8) ^P	(5.5) ^P	L
23	Q	L	L	L	(5.5) ^P	[5.6] ^L	(5.7) ^P	(5.9) ^P	(5.9) ^P	L	L	L
24	Q	L	L	L	(4.6) ^P	[5.1] ^L	5.6	(5.6) ^P	(5.8) ^P	(6.1) ^P	(4.5) ^P	L
25	L	L	L	L	(5.3) ^P	(5.9) ^P	(6.3) ^P	(5.8) ^P	(5.8) ^P	4.9	L	L
26	Q	L	L	L	(5.5) ^P	5.6	(6.2) ^P	(5.8) ^P	(5.4) ^P	(4.8) ^P	(5.0) ^P	L
27	L	L	L	(5.7) ^P	5.4	(6.0) ^P	(6.2) ^L	(5.9) ^P	(6.4) ^P	L	L	Q
28	Q	L	L	L	(5.3) ^P	(5.9) ^P	(5.5) ^P	(5.8) ^P	(5.7) ^P	(5.5) ^P	(5.2) ^P	L
29	Q	L	L	L	L	(6.0) ^P	(5.8) ^P	(5.2) ^P	(5.4) ^P	L	L	L
30	Q	L	L	L	L	(5.9) ^P	(5.6) ^P	(6.0) ^P	(5.5) ^P	4.9	L	L
31	Q	L	L	L	(5.3) ^P	L	(5.7) ^P	(5.1) ^P	5.7	(5.7) ^P	(5.6) ^P	L
Median Count	-	4.1	4.7	4.7	4.9	(5.3)	5.3	5.4	(5.4)	4.9	4.6	-
Count	-	10	19	23	25	29	31	28	27	26	23	14

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
LAT 38°7'N, LONG 77°10'W.

Swept I.Q. Mc 10-25.0 Mc in 0.25 min
Manual □ Automatic ☒

U.S. GOVERNMENT PRINTING OFFICE: 1946 O-17819

TABLE 54
 Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

Km **August[†], 1949**
 (Characteristic) (Month)

Observed at Washington, D.C.

Lat 39°N Long 77.5°W

National Bureau of Standards
 Scored by: J.J.S., B.E.B.
 (Institution)

Day	75°N Mean Time																								
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1								A									B								
2																	C								
3																									
4																									
5																									
6																									
7																									
8																									
9																									
10																									
11																									
12																									
13																									
14																									
15																									
16																									
17																									
18																									
19																									
20																									
21																									
22																									
23																									
24																									
25																									
26																									
27																									
28																									
29																									
30																									
31																									
Median	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Count	1	1	24	24	28	28	29	29	28	28	29	29	28	28	29	28	29	28	29	28	29	28	29	28	29

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
 LAT. 38°7'N, LONG. 77°10'W.

Sweep J.O. Mc 10-25.0 Mc in 0.25 min
 Manual □ Automatic ■

U.S. GOVERNMENT PRINTING OFFICE 1947 O-18219

Form adopted June 1946

Scored by: J.J.S., B.E.B.

Calculated by: B.E.B., N.C.H., J.M.W.

Farm adopted June 1946
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

fo_E, Mc (Characteristic)
August, 1949
(Month)

Washington, D.C.
Lat 39°0'N, Long 77°5'W

National Bureau of Standards
Scaled by J.J.S., B.E.B. (Institution)
Calculated by B.E.B., N.C.H., J.M.W.

IONOSPHERIC DATA

75°W Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1																								
2																								
3																								
4																								
5																								
6																								
7																								
8																								
9																								
10																								
11																								
12																								
13																								
14																								
15																								
16																								
17																								
18																								
19																								
20																								
21																								
22																								
23																								
24																								
25																								
26																								
27																								
28																								
29																								
30																								
31																								
Median	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Count	/	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
LAT 38°7'N, LONG 77°1'W.

Sweep 1.0 Mc to 25.0 Mc in 0.25 min
Manual □ Automatic ■

U. S. GOVERNMENT PRINTING OFFICE: 1946 O-170518

TABLE 56
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

National Bureau of Standards
(Institution)

Characteristic, Mc./Km. (Unit) August, 1949
Observed at Washington, D. C.
Lat 39°0'N., Long 77.5°W.

Day	75°W — Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	31/100	43/100	43/100	34/100	52/100	40/100	57/100	67/100	46y/20	39y/20	42/100	40/100
2	32/100	25/100	40/100	34/100	23/100	32/100	32/100	40/100	38/100	30/100	B	B
3	G	G	G	G	1/8/20	2/4/30	G	G	G	G	45/130	46/130
4	C	C	C	C	C	C	C	C	C	C	C	C
5	37/10	29y/20	30y/20	G	G	G	44y/20	34y/20	37/20	54y/100	34y/100	G
6	G	G	G	66y/20	30/100	26y/100	47/120	45y/120	47/120	38y/100	46y/100	G
7	G	G	G	G	G	21/130	27/130	35/130	34y/100	32/100	G	G
8	G	G	G	G	G	36y/20	44y/20	34y/20	36/110	36/100	46y/100	54/100
9	32/10	50y/20	33/30	33/110	31/110	60/20	35/120	46y/120	52/110	60/110	G	37/120
10	G	26/100	38/100	G	25/130	32/130	30/140	39/100	39/120	41/110	G	G
11	T	T	T	T	T	T	3/2/20	57/120	2/2/20	3/2/20	G	G
12	G	G	G	G	38/120	31/120	36/120	46/120	35/120	39/120	G	G
13	45/110	53y/110	G	G	62y/110	G	33y/20	34y/20	35/110	53/110	G	G
14	G	G	G	G	G	42/110	30/140	39/120	67/140	38/130	G	G
15	46/110	32/150	35y/110	26/120	G	G	20/130	31/130	39/120	G	G	G
16	C	C	C	C	C	C	37/110	40y/110	33/120	G	G	G
17	35/100	43/100	40y/110	26/110	G	21/120	35/130	46/120	55/120	45/120	G	G
18	36/110	37/110	31/110	G	G	21/150	G	41y/120	1/9/110	41/110	44/130	66/120
19	G	41/110	33/110	G	28/110	G	34y/30	32/120	32/110	47/110	G	G
20	32y/10	39/150	31/110	20/110	G	22/110	G	54/100	43/100	31/100	38/100	43/100
21	G	24/110	30/120	39/110	G	35/20	G	31/110	28/100	21/100	G	G
22	26y/110	26y/110	32/100	53/110	G	G	36/110	30/100	G	G	G	G
23	26/100	23/110	G	G	G	29/100	G	41/130	38/130	G	G	G
24	G	G	G	G	G	G	47/130	23/100	G	G	G	G
25	24/110	22/110	G	G	G	23/130	G	47/120	44/100	42/100	56/100	40/120
26	22/110	G	G	30/110	31/100	53/100	38/100	31/100	55/110	55/120	42/100	43/100
27	G	23/110	(23)y/100	G	G	G	26/110	42/100	32/100	26/100	21/100	26/100
28	55/100	38/100	32/100	23/100	G	63y/20	10/20	40/20	43/100	43/100	33/100	38/100
29	G	G	32/100	G	G	18/130	20/110	28/100	54y/100	29/100	30/100	27/100
30	G	G	G	G	G	34/120	39/120	30/100	32/100	24/100	27/100	23y/100
31	G	G	G	G	G	22/110	70y/100	26/100	22/100	36/100	30/100	28/100
Median	**	2.2	2.4	**	**	**	1.9	3.4	3.8	3.5	3.4	3.0
Count	28	28	28	28	28	28	30	30	30	30	28	29

** MEDIAN FEES LESS THAN MEDIAN FOR LESS THAN LOWER FREQUENCY LIMIT OF RECORDER

Sweep L.O. Mc 10.23.0 Mc in 0.25 min Manual □ Automatic ■

Manual □ Automatic ■

National Bureau of Standards

Scaled by: J.J.S., B.E.B. (Institution)

Calculated by: B.E.B., N.C.H., J.M.W.

TABLE 57
IONOSPHERIC DATA
Lat 39.0°N, Long 77.5°W
75W Mean Time

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

(M1500) F₂, h₄₉
(Characteristic) (Unit)
Washington, D.C.
Observed at

Lat 39.0°N, Long 77.5°W
Augus†, 1949
(Month)

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1	1.8	1.8	1.9	1.8	1.9	2.1	2.2	2.2	2.1	2.2	2.1	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	
2	1.8	1.7	1.7	1.8	1.6	1.8	1.8	1.8	1.7	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
3	1.6	1.7	1.7	1.5	1.5	1.6	1.6	1.6	1.6	1.7	1.7	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
4	1.6	1.7	1.7	1.5	1.5	1.6	1.6	1.6	1.6	1.7	1.7	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
5	1.8	1.7	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
6	1.8	1.8	1.9	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
7	1.8	1.8	1.8	1.9	1.9	1.9	2.0	2.0	2.1	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
8	1.6	1.6	1.6	1.8	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
9	1.9	1.9	1.8	1.8	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
10	1.8	1.8	1.8	1.9	1.9	1.9	2.0	2.0	2.0	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
11	1.9	1.9	1.8	1.9	1.9	2.0	2.0	2.3	2.1	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
12	1.9	1.9	1.9	1.7	1.7	1.9	2.0	2.3	2.3	2.0	2.0	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	2.1	
13	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
14	1.5	1.9	1.9	1.8	1.7	1.8	1.8	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	
15	1.6	1.7	1.7	1.7	1.7	1.7	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
16	K	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
17	1.7	F	1.8																						
18	1.7	F	1.7																						
19	1.8	S	1.9	S	1.7																				
20	1.7	S	1.6	F	1.7	F	1.8																		
21	1.7	F	1.6	F	1.7																				
22	1.7	S	1.7																						
23	1.8	F	1.8																						
24	1.6	F	1.6	F	1.7	F	1.7	F	1.8																
25	1.9	F	1.8																						
26	1.7	F	1.6	F	1.7																				
27	1.8	F	1.8																						
28	1.7	F	1.8	F	1.7	F	1.7	F	1.8																
29	1.7	I	1.7	I	1.8																				
30	1.7	I	1.7	I	1.7	I	1.8																		
31	1.7	I	1.7	I	1.8	I	1.8	I	1.9																
Median	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	
Count	51	31	31	30	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	31	

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
LAT. 38.7°N, LONG. 77.1°W.

†

Manual □ Automatic ■
Sweep I.O. Mc 1025.0 Mc in Q.25 min
U.S. GOVERNMENT PRINTING OFFICE 160-10-31811
U.S. GOVERNMENT PRINTING OFFICE 160-10-31811

TABLE 58
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

Form adopted June 1946
TABLE OF STANDARDS
August 1949
(Month)
Observed at Washington, D.C.
(Latitude)
Lot 39°0'N., Long 77°5'W.

(M3000) F2, (Unit)

(Characteristic) (Month)

August, 1949

National Bureau of Standards
(Institution)

Scaled by J.J.S., B.E.B.,

Calculated by B.E.B., N.G.H., J.M.W.

Day	75°W Mean Time											
	00	01	02	03	04	05	06	07	08	09	10	11
1	2.7	2.8	2.8	2.7	2.9	3.1	3.3	3.2	3.1	(3.3)J	2.7	(3.2)J
2	2.8	2.7	2.6	2.8	2.5	2.7	2.7	2.7	2.7	(2.8)P	C	C
3	2.4K	2.6K	2.2K	2.0K	2.2K	2.0K	2.2K	2.0K	2.2K	GK	GK	A
4	(2.4)S	2.6K	2.1K	N	N	2.6K	2.1K	2.7K	*3.2F	2.9	*3.1	(2.8)T
5	2.7F	2.7F	2.9F	2.6F	2.8F	3.0	3.4	3.0	2.8	2.8	2.7	(2.9)J
6	2.8F	2.8F	3.0	2.7F	2.8F	2.9F	3.0	2.5	2.9	2.9	2.9	2.9
7	2.8	2.8	3.0	2.8F	2.8F	2.9F	3.1	2.8K	2.9K	2.7K	2.9K	2.8K
8	2.4K	2.8F	2.7F	2.7F	2.8F	2.9K	2.9K	2.5K	2.7K	(3.0)P	2.8K	2.8K
9	2.7K	2.8F	2.6K	2.8K	2.9K	2.8K	2.9K	2.9	2.8	(2.9)J	3.0	2.9
10	2.8K	2.8F	(2.3)O	3.0F	2.7K	2.8K	2.9K	2.6K	2.7K	(2.7)S	2.7K	2.8K
11	2.8	2.8	2.7	2.9	2.9	3.0	3.4	3.2	3.0	3.0	3.0	3.0
12	2.8	2.8	2.9	2.7	2.9	3.3	3.3	3.1	3.3	3.2	3.2	3.0
13	2.8F	2.7F	(2.9)S	(3.0)P	2.9F	3.2F	3.1	3.0	3.1	2.9	3.0	3.1
14	2.7	2.8	2.8	2.7	2.6F	2.8F	2.9	2.8	2.6	3.0F	3.0	3.0
15	K(2.7)S	2.5K	2.5K	2.6K	2.6K	(3.0)F	2.9K	2.6K	3.0K	GK	GK	GK
16	K(2.7)S	K(2.7)S	2.8K	K(2.8)F	2.5K	K(2.8)F	2.6K	2.5K	2.6K	2.8K	(2.7)S	(2.8)K
17	2.6F	2.7F	2.7F	2.7F	(2.8)J	(3.2)S	(3.5)S	3.2	3.0	2.9	2.8	(2.8)S
18	2.6F	(2.5)F	2.5F	2.5F	2.7F	3.0	3.0	2.7F	A	GK	2.5K	2.5K
19	(2.8)S	2.8	(2.6)S	2.6	2.8	2.9	3.1	3.0	(3.0)H	2.6	2.7	2.7
20	(2.6)S	2.5F	2.4F	2.5F	2.8F	3.0	2.9	2.9	2.9	2.7	2.6	2.7
21	2.6F	2.5F	2.7F	2.7F	2.8	(3.2)S	3.1	3.2	2.8	2.8	2.8	2.8
22	(2.6)S	2.6	2.6F	(2.7)S	2.7	2.8	3.0	2.8	2.8	2.7	(3.0)F	(2.9)S
23	(2.7)F	(2.8)S	2.8	2.7	2.6	2.8	3.1	3.0	2.8	2.8	2.8	(2.8)S
24	2.7	(2.8)S	2.9	(2.8)S	(2.8)S	3.0	3.3	3.2	(3.2)S	3.0	2.8	2.8
25	(2.9)S	2.8	(3.0)S	2.9	2.8	2.9	3.2	3.1	3.0	2.9	2.8	(2.8)S
26	2.6	(2.8)S	2.7	2.7	(2.8)S	2.8	3.2	3.2	3.0	2.8	2.8	(2.6)S
27	2.7	2.8	3.0	(2.7)S	2.7	3.0	3.1	(3.1)S	3.2	2.9	2.8	(2.8)S
28	(2.6)S	2.8	2.8	2.6	2.8	2.9	(3.1)S	(3.2)S	3.0	2.8	2.7	2.7
29	2.6	2.6	2.7	2.7	(2.9)S	2.9	(3.3)S	3.2	3.0	2.9	2.8	(2.8)S
30	2.6	2.6	2.6	2.8	2.7	3.2	(3.1)S	3.1	2.9	2.9	2.9	(2.8)S
31	2.6	(2.7)S	2.7	2.7	2.9	(2.9)S	3.2	3.0	3.0	2.9	2.8	(2.7)S
Median	2.7	2.8	2.7	2.8	3.1	3.1	3.0	2.9	2.9	2.8	2.8	2.8
Count	31	31	31	30	31	31	31	31	30	31	31	31

* SUPPLEMENTARY DATA FROM FT. BELVOIR,
LAT. 38°7'N., LONG. 77°10'W.

** SWEET-LO-MC TO 22.0 Mc IN 0.25 MIN

SWEEP-LO MC TO 22.0 Mc IN 0.25 MIN

Manual □ Automatic □

U. S. GOVERNMENT PRINTING OFFICE 1410-1253

TABLE 59
IONOSPHERIC DATA

(M3000)FI, (Unit)
 (Characteristics)
 Observed at Washington, D.C.

August 1949
 (Month)

Lat 39.0°N, Long 77.5°W

75°W Mean Time

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.
IONOSPHERIC DATA

National Bureau of Standards
 Scaled by: J.J.S., B.E.B.
 Calculated by: B.E.B., N.C.H., J.M.W.

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1																								
2																								
3																								
4																								
5																								
6																								
7																								
8																								
9																								
10																								
11																								
12																								
13																								
14																								
15																								
16																								
17																								
18																								
19																								
20																								
21																								
22																								
23																								
24																								
25																								
26																								
27																								
28																								
29																								
30																								
31																								
Median Count	-	3.2	3.4	3.6	3.7	3.5	3.6	3.5	3.4	3.5	3.6	3.5	3.5	3.6	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	
	10	10	10	23	24	24	30	27	26	26	26	23	23	23	24	24	23	23	24	24	24	24	24	

SUPPLEMENTARY DATA FROM FT. BELVOIR,
 LAT. 38.7°N, LONG. 77.1°W

*

Sweep 1.0 Mc to 25.0 Mc in .925 min
 Manual □ Automatic □
 U.S. GOVERNMENT PRINTING OFFICE 146-17259

(M1500)E
 (Characteristic)
 Observed at Washington, D.C.

 August[†], 1949
 (Month)
 Lot 39,00N, Long 77.5°W

 TABLE 60
 IONOSPHERIC DATA

Day	75°W Mean Time																								
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
1																									
2																									
3																									
4																									
5																									
6																									
7																									
8																									
9																									
10																									
11																									
12																									
13																									
14																									
15																									
16																									
17																									
18																									
19																									
20																									
21																									
22																									
23																									
24																									
25																									
26																									
27																									
28																									
29																									
30																									
31																									
Median	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Count	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	

 * SUPPLEMENTARY DATA FROM FT BELVOIR,
 LAT. 38°7'N, LONG. 77°1'W.

 Sweep-O-Lite Model 225.0 Mc in. 0.25 min
 Manual □ Automatic □

Table 61Ionospheric Storminess at Washington, D. C.August 1949

Day	Ionospheric character*		Principal storms		Geomagnetic character**	
	00-12 GCT	12-24 GCT	Beginning GCT	End GCT	00-12 GCT	12-24 GCT
1	1	3			1	2
2	2	1			3	3
3	5	6	0000	----	6	3
4	6	3	----	1200	5	4
5	1	1			3	4
6	1	2			2	2
7	1	4	1200	----	1	3
8	4	4	----	----	4	2
9	4	2	----	1300	2	2
10	4	5	0200	2400	2	2
11	2	1			1	1
12	1	2			2	1
13	1	2			2	2
14	2	3	2300	----	4	3
15	4	6	----	----	4	2
16	4	5	----	----	2	1
17	3	1	----	0400	2	3
18	3	5	1200	----	2	3
19	2	3	----	0200	2	2
20	2	1			3	3
21	2	1			2	2
22	2	3			2	2
23	1	2			2	1
24	1	0			1	0
25	0	1			0	1
26	1	3			1	1
27	0	2			3	2
28	1	2			2	1
29	2	1			2	2
30	2	1			2	1
31	2	1			1	2

*Ionosphere character figure (I-figure) for ionospheric storminess at Washington, D. C., during 12-hour period, on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

**Average for 12 hours of Cheltenham, Maryland, geomagnetic K-figures on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

----Dashes indicate continuing storm.

Table 62Sudden Ionosphere Disturbances Observed at Washington, D. C.August 1949

1949 Day	GCT		Location of transmitters	Relative intensity at minimum*	Other phenomena
	Beginning	End			
August 1	1952	2215	Ohio, D.C., New Brunswick	0.0	
2	1520	1540	Ohio, D.C., England	0.03	
5	2042	2110	Ohio, D.C., England	0.0	
6	1813	1845	Ohio, D.C.	0.1	
6	2251	2345	Ohio, D.C.	0.01	
16	1157	1220	Ohio, England	0.2	
16	1745	1820	Ohio, D.C., England, New Brunswick	0.3	Solar flare*** 1755
19	1843	1905	Ohio, D.C., England	0.1	Solar flare*** 1900
19	2110	2210	Ohio, D.C., England	0.0	Terr.mag.pulse** 2113-2210 Solar flare*** 2112
20	1200	1220	England	0.1	
20	1525	1600	Ohio, D.C., England	0.03	Terr.mag.pulse** 1525-1527 Solar flare*** 1530
22	1415	1440	Ohio, D.C.	0.2	Terr.mag.pulse** 1416-1418 Solar flare*** 1416
25	1543	1620	Ohio, D.C., England	0.05	Solar flare*** 1540
25	2025	2050	Ohio, D.C., England, New Brunswick	0.05	Solar flare*** 2015
31	1936	2020	Ohio, D.C., Canal Zone, England, New Brunswick	0.0	Solar flare*** 1940

*Ratio of received field intensity during SID to average field intensity before and after, for station WSKAL, 6080 kilocycles, 600 kilometers distant, for all SID except the following: Station GLH, 13525 kilocycles, received in New York, 5340 kilometers distant, was used for the SID on August 20.

** As observed on Cheltenham magnetogram of the United States Coast and Geodetic Survey.

*** Time of observation at McMath-Hulbert Observatory, Michigan.

Table 63

Sudden Ionosphere Disturbances Reported by Engineer-in-Chief, Cable and Wireless, Ltd., as Observed in England

1949 Day	GCT Beginning End	Receiving station	Location of transmitters	Other phenomena	1949 Day	GCT Beginning End	GCT Receiving station	Location of transmitters	Other phenomena
July 27	1227 1240	Brentwood	Barbados, Canary Is., Greece, Iran, Madagascar, Palestine, Portugal, Spain, Switzerland, Syria, Yugoslavia, Bahrain I., Barbados, Greece, India, Kenya, Palestine, Southern Rhodesia, Spain, Switzerland, Uruguay, U.S.S.R. ^o , Yugoslavia, Zanzibar	Terr. mag. pulse* 1222-1230	August 5	0805 ***	Brentwood	Afghanistan, Austria, Bahrain I., Belgian Congo, Canary Is., Eritrea, French Equatorial Africa, Greece, India, Iran, Kenya, Madagascar, Palestine, Portugal, Southern Rhodesia, Spain, Syria, Trans-Jordan, Turkey, U.S.S.R., Yugoslavia, Zanzibar	
28	1120 1140	Brentwood							
29	1415 1445	Brentwood	Bahrain I., Barbados, Belgium Congo, Bulgaria, Chile, Colombia, Eritrea, Greece, Iran, Kenya, Malta, Palestine, Portugal ^l , Southern Rhodesia, Spain, Switzerland, Syria, Uruguay, U.S.S.R., Venezuela, Yugoslavia, Zanzibar Argentina, Australia, Brazil, Canada, Ceylon, Gold Coast, New York, Union of S. Africa	Terr. mag. pulse* 1411-1430	5	0805	Somerston	Ceylon, China, Egypt, Gold Coast, India, New York, Nigeria, Union of S. Africa	
29	1415 1445	Somerston			5	1040	Brentwood	Bahamas, Belgian Congo, Bulgaria, Canary Is., Greece, India, Kenya, Malta, Palestine, Portugal, Southern Rhodesia, Spain, Switzerland, Syria, Trans-Jordan, Turkey, U.S.S.R., Yugoslavia	
30	0730 0750	Brentwood			16	1150	Brentwood	Austria, Bahrain I., Barbados, Belgian Congo, Canary Is., Chile, Greece, India, Iran, Kenya, Malta, Palestine, Portugal, Southern Rhodesia, Spain, Switzerland, Syria, Trans-Jordan, Uruguay, U.S.S.R., Yugoslavia, Zanzibar	
30	0730 0830	Somerston							
31	0845 0845	Brentwood							
31	1450 1615	Brentwood							
31	1515 1545	Somerston							
August 5	0720 0750	Brentwood							

*As observed on Cheltenham magnetogram of the United States Coast and Geodetic Survey.

**Time of observation at McMath-Hulbert Observatory, Michigan.

***Incomplete recovery of SID.

Table 64

Sudden Ionosphere Disturbances Reported by International Telephone
and Telegraph Corporation, as Observed at Platanos, Argentina

1949 Day	GCT		Location of transmitters	Other phenomena
	Beginning	End		
July 29	1420	1450	Bolivia, Brazil, Chile, Colombia, Cuba, Denmark, England, Germany, New York, Peru, Switzerland, Venezuela	Terr. mag. pulse* 1411-1430
	31	1510	Brazil, Chile, France, New York, Peru, Switzerland	Solar flare** 1505

*As observed on Cheltenham magnetogram of the United States Coast and Geodetic Survey.

**Time of observation at McMath-Hulbert Observatory, Michigan.

Table 65

Sudden Ionosphere Disturbances Reported by RCA Communications, Inc.,
as Observed at Point Reyes, California

1949 Day	GCT		Location of transmitters	Other phenomena
	Beginning	End		
August 6	2250	2400	Australia, Hawaii, Japan, Philippine Is.	
	19	2110	Australia, China, Hawaii, Japan, Philippine Is.	Solar flare* 2112
September 5	0110	0245	Australia, China, Chosen, Hawaii, Japan, Java, New York, Philippine Is.	
	9	0210	Australia, China, Japan, Java, Philippine Is.	
9	0052	0145	Australia, China, Chosen, Hawaii, Japan, Java, Philippine Is.	

*Time of observation at McMath-Hulbert Observatory, Michigan.

Note: Observers are invited to send to the CRPL information on times of beginning and end of sudden ionosphere disturbances for publication as above. Address letters to the Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

Table 66

Provisional Radio Propagation Quality Figures
 (Including Comparisons with CMFL Warnings and CRPL Probable Disturbed Period Forecasts)
July 1949

Day	North Atlantic						North Pacific						<u>Quality Figure Scale:</u>
	Quality figure	CRPL*	CRPL**	Forecast of probable disturbed periods	Geo-magnetic K _{Ch}	Quality figure	CRPL*	CRPL**	Forecast of probable disturbed periods	Geo-magnetic K _{Ch}	Quality figure	CRPL*	
	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	01-12 GCT 13-24 GCT	
1	6	5	X	2	2	6	7	X	2	2	2	1	1 - Useless
2	7	6		2	1	6	7		2	1	2	1	2 - Very poor
3	7	6		1	2	7	7		1	2	1	1	3 - Poor
4	7	7		1	1	6	6		1	1	1	1	4 - Poor to fair
5	7	6		1	1	6	6		1	1	1	1	5 - Fair
6	7	7		1	1	6	7		1	1	1	1	6 - Fair to good
7	7	6		2	2	7	7		2	2	2	2	7 - Good
8	5	7		2	3	7	7		2	3	2	3	8 - Very good
9	7	7		3	1	6	7		3	1	1	1	9 - Excellent
10	7	7		1	2	6	7		1	2	1	1	
11	7	7		1	1	6	7		1	1	1	1	
12	7	7		1	3	7	7		1	3	1	1	
13	5	6		3	2	6	6		3	2	1	1	
14	6	6		2	2	6	7		2	2	1	1	
15	6	7		0	1	6	7		0	1	1	1	
16	7	6		1	4	6	5		1	4	1	1	
17	6	5		3	2	7	6		3	2	1	1	
18	6	6		2	4	7	7		2	4	1	1	
19	5	5		4	3	6	7		4	3	1	1	
20	7	6		2	2	6	7		2	2	1	1	
21	7	6		1	2	6	7		1	2	1	1	
22	7	6		2	3	6	6		2	3	1	1	
23	7	6		3	3	6	6		3	3	1	1	
24	6	6		2	2	7	8		2	2	1	1	
25	6	5		3	3	6	7		3	3	1	1	
26	7	6		1	1	6	7		1	1	1	1	
27	7	6		0	0	7	7		0	0	1	1	
28	7	6		0	1	6	6		0	1	1	1	
29	7	6		1	2	6	7		1	2	1	1	
30	7	6		2	2	6	7		2	2	1	1	
31	6	5	X	2	2	7	7	X	2	2	1	1	
Scores:													
H		0	0			0	0						
M		0	0			0	0						
G		31	29			31	29						
(S)		0	2			0	0						
S		0	0			0	2						

*Broadcast on WWV, Washington, D.C. Times of warnings recorded to nearest half day as broadcast.

**In addition to dates marked X, the following was designated as a probable disturbed day on forecasts more than eight days in advance of said date: July 2.

Symbols:

X Warning given or probable disturbed date

H Quality 4 or worse on day or half day of warning

M Quality 4 or worse on day or half day of no warning

G Quality 5 or better on day of no warning

(S) Quality 5 on day of warning

S Quality 6 or better on day of warning

() Quality 4 or worse (disturbed)

Geomagnetic K_{Ch} on the standard scale of 0 to 9, 9 representing the greatest disturbance.

Table 67

American and Zürich Provisional Relative Sunspot NumbersAugust 1949

Date	R _A *	R _Z **	Date	R _A *	R _Z **
1	174	161	17	228	175
2	172	171	18	235	168
3	163	127	19	220	162
4	132	114	20	224	192
5	125	109	21	236	151
6	78	88	22	242	198
7	55	59	23	230	189
8	52	50	24	217	169
9	56	45	25	206	158
10	37	34	26	200	163
11	22	17	27	197	165
12	66	56	28	210	163
13	128	82	29	210	155
14	163	108	30	197	133
15	188	155	31	218	168
16	195	174	Mean:	163.7	130.9

*Combination of reports from 49 observers; see page 8.

**Dependent on observations at Zürich Observatory and its stations at Locarno and Arosa.

Table 68a

Coronal observations at Climax, Colorado (5303A), east limb

Table 69a

Coronal observations at Climax, Colorado (6374A), east limb

Table 68b

Coronal observations at Climax, Colorado (5303A), west limb

Date GGT	Degrees south of the solar equator															0°	Degrees north of the solar equator																															
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90														
1949	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X													
Aug. 1.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	6	7	7	7	9	10	10	11	10	11	12	15	14	10	7	4	2	-	-	-	-	-	-							
2.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	6	7	7	7	9	10	10	11	10	11	12	15	14	10	7	4	2	-	-	-	-	-	-	-						
3.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	8	12	12	13	13	12	13	14	14	15	19	17	13	10	5	-	-	-	-	-	-	-	-						
4.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	4	10	11	13	13	14	13	13	14	14	16	20	21	18	14	10	2	-	-	-	-	-	-	-						
5.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	8	12	14	15	13	18	14	8	9	11	14	15	13	12	9	3	-	-	-	-	-	-						
6.8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	3	4	11	13	15	15	13	11	10	7	7	12	14	13	13	11	1	-	-	-	-	-	-	-							
7.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	5	10	14	15	23	25	18	14	10	9	10	15	13	13	5	4	3	-	-	-	-	-	-							
8.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	5	8	11	15	18	17	11	9	9	10	12	13	9	5	1	-	-	-	-	-	-								
9.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	9	10	12	14	15	16	18	15	8	7	13	11	9	8	3	1	-	-	-	-	-	-							
10.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	4	9	11	13	12	13	17	26	22	15	10	9	6	3	-	-	-	-	-	-	-								
11.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	5	8	10	9	10	11	13	14	12	16	21	18	9	5	3	2	1	-	-	-	-	-	-						
12.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	10	11	13	11	13	14	13	12	15	13	9	9	8	6	2	-	-	-	-	-	-	-						
13.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	4	3	4	8	10	11	10	10	9	6	4	2	-	-	-	-	-	-	-	-									
14.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	4	6	8	9	9	10	13	14	13	14	11	10	12	9	8	7	6	7	5	3	1	-	-	-	-	-	-			
15.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	4	5	3	3	4	5	10	12	13	13	10	9	5	3	2	2	2	-	-	-	-	-	-	-					
16.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	3	2	2	4	6	14	12	12	10	9	10	14	13	11	10	8	4	3	4	3	2	-	-	-	-	-	-		
17.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	2	2	3	4	7	8	10	9	8	8	12	24	20	15	12	6	3	2	2	1	-	-	-	-	-	-			
18.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	3	3	2	2	4	5	5	5	8	11	14	14	10	5	2	-	-	-	-	-	-	-	-	-	-	-		
19.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	3	3	2	2	4	5	5	5	7	13	14	15	14	11	8	5	3	4	4	2	-	-	-	-	-	-		
20.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	4	5	5	6	10	9	9	5	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
21.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	4	5	7	8	9	10	7	12	18	11	13	12	9	4	2	1	-	-	-	-	-	-	-	-	-	-	-		
22.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	11	13	10	8	14	13	12	11	11	10	9	8	6	2	-	-	-	-	-	-	-	-	-	-	-	-	-		
23.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	4	4	5	6	10	11	13	13	14	14	14	15	11	5	3	2	1	-	-	-	-	-	-	-	-	-		
24.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	5	5	7	9	11	12	13	12	12	17	18	15	14	13	11	7	3	2	-	-	-	-	-	-	-	-	-		
25.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	4	8	9	9	8	9	13	14	14	12	10	5	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
26.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	5	7	10	13	20	12	8	10	20	30	31	32	20	12	8	5	3	2	-	-	-	-	-	-	-	-	-	-		
27.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	5	6	10	13	16	17	13	10	15	22	20	20	18	8	5	3	-	-	-	-	-	-	-	-	-	-	-			
29.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	8	10	10	10	9	11	10	11	12	13	11	6	3	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30.7	-	X	X	X	X	-	-	-	-	-	-	-	-	-	-	-	-	3	3	3	2	5	10	10	10	6	9	13	12	17	27	25	6	3	-	-	-	-	-	-	-	-	-	-	-	-		

Table 69b

Coronal observations at Climax, Colorado (6374A), west limb

Date GGT	Degrees south of the solar equator															0°	Degrees north of the solar equator																								
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90					
1949	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
Aug. 1.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	5	7	-	-	-	-	-	-	-	-	-	-	-	-	-				
2.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
3.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
4.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	2	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
6.8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	1	1	2	4	4	7	10	6	1	1	3	11	10	7	5	1	-	-	-	-	-	-	
7.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	2	5	4	2	1	1	1	1	1	1	1	-	-	-	-	-	-	-	
8.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	10	5	6	8	6	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	10	5	1	14	12	1	1	1	1														

Table 70a





Coronal observations at Climax, Colorado (6704A), east limb

Table 70b

Coronal observations at Climax, Colorado (6704A), west limb

GRAPHS OF IONOSPHERIC DATA

44






Fig. 5. BOSTON, MASSACHUSETTS
42.4°N, 71.2°W JULY 1949




Fig. 6. BOSTON, MASSACHUSETTS JULY 1949




Fig. 7. SAN FRANCISCO, CALIFORNIA
37.4°N, 122.2°W JULY 1949




Fig. 8. SAN FRANCISCO, CALIFORNIA JULY 1949




Fig. 9. WHITE SANDS, NEW MEXICO

32.3°N, 106.5°W

JULY 1949




Fig. 10. WHITE SANDS, NEW MEXICO

JULY 1949




Fig. 11. BATON ROUGE, LOUISIANA

30.5°N, 91.2°W

JULY 1949




Fig. 12. BATON ROUGE, LOUISIANA

JULY 1949




Fig. 13. OKINAWA I.
26.3°N, 127.7°E

JULY 1949




Fig. 14. OKINAWA I.

JULY 1949




Fig. 15. MAUI, HAWAII

20.8°N, 156.5°W

JULY 1949




















Fig. 16. MAUI, HAWAII


JULY 1949
















Fig. 33. OSLO, NORWAY
60.0°N, 11.0°E





MAY 1949

Fig. 34. OSLO, NORWAY
MAY 1949Fig. 35. WAKKANAI, JAPAN
45.4°N, 141.7°E

MAY 1949

Fig. 36. WAKKANAI, JAPAN
MAY 1949








Fig. 45. OKINAWA I.
26.3°N, 127.7°E MAY 1949




Fig. 46. OKINAWA I. MAY 1949




Fig. 47. BRISBANE, AUSTRALIA
27.5°S, 153.0°E MAY 1949
















Fig. 48. BRISBANE, AUSTRALIA MAY 1949










Fig. 61. MADRAS, INDIA
13.0°N, 80.2°E

APRIL 1949




Fig. 62. MADRAS, INDIA

APRIL 1949




Fig. 63. TIRUCHIRAPALLI, INDIA
10.8°N, 78.8°E

APRIL 1949




Fig. 64. TIRUCHIRAPALLI, INDIA

APRIL 1949




Fig. 65. FRIBOURG, GERMANY
48.1°N, 7.8°E

MARCH 1949




Fig. 66. FRIBOURG, GERMANY

MARCH 1949




Fig. 67. POITIERS, FRANCE
46.6°N, 2.0°W

MARCH 1949








Fig. 68. POITIERS, FRANCE

MARCH 1949






Fig. 73. WAKKANAI, JAPAN

45.4°N, 141.7°E

FEBRUARY 1949




Fig. 74. WAKKANAI, JAPAN

FEBRUARY 1949




Fig. 75. FUKAURA, JAPAN

40.6°N, 139.9°E

FEBRUARY 1949




Fig. 76. FUKAURA, JAPAN

FEBRUARY 1949




Fig. 77. SHIBATA, JAPAN
37.9°N, 139.3°E

FEBRUARY 1949




Fig. 78. SHIBATA, JAPAN

FEBRUARY 1949




Fig. 79. TOKYO, JAPAN

35.7°N, 139.5°E

FEBRUARY 1949












Fig. 80. TOKYO, JAPAN

FEBRUARY 1949








Fig. 89. FUKAURA, JAPAN
40.6°N, 139.9°E

JANUARY 1949




Fig. 90. FUKAURA, JAPAN

JANUARY 1949




Fig. 91. SHIBATA, JAPAN
37.9°N, 139.3°E

JANUARY 1949








Fig. 92. SHIBATA, JAPAN

JANUARY 1949



Index of Tables and Graphs of Ionospheric Datain CRPL-F61

	<u>Table page</u>	<u>Figure page</u>
Baton Rouge, Louisiana		
July 1949	11	46
Bombay, India		
April 1949	17	58
Boston, Massachusetts		
July 1949	10	45
Brisbane, Australia		
May 1949	15	55
Canberra, Australia		
May 1949	16	56
Christchurch, New Zealand		
May 1949	16	57
Delhi, India		
April 1949	17	58
Fribourg, Germany		
March 1949	18	60
February 1949	18	61
January 1949	20	64
Fukaura, Japan		
May 1949	14	53
February 1949	19	62
January 1949	21	66
Guam I.		
July 1949	12	48
Hobart, Tasmania		
May 1949	16	56
Huancayo, Peru		
July 1949	13	50
Johannesburg, Union of S. Africa		
June 1949	13	51
Madras, India		
April 1949	17	59
Maui, Hawaii		
July 1949	11	47
Okinawa I.		
July 1949	11	47
June 1949	13	51
May 1949	15	55
Oslo, Norway		
July 1949	10	44
June 1949	13	50
May 1949	14	52
Palmyra I.		
July 1949	12	49

Index (CRPL-F61, continued)

	<u>Table page</u>	<u>Figure page</u>
Poitiers, France		
April 1949	16	57
March 1949	18	60
February 1949	18	61
January 1949	20	65
San Francisco, California		
July 1949	10	45
San Juan, Puerto Rico		
July 1949	12	48
Shibata, Japan		
May 1949	14	53
February 1949	19	63
January 1949	21	66
Tiruchirapalli, India		
April 1949	17	59
Tokyo, Japan		
May 1949	15	54
February 1949	19	63
January 1949	21	67
Trinidad, British West Indies		
July 1949	12	49
Wakkanai, Japan		
May 1949	14	52
February 1949	19	62
January 1949	20	65
Washington, D. C.		
August 1949	10	44
White Sands, New Mexico		
July 1949	11	46
Yamakawa, Japan		
May 1949	15	54
February 1949	20	64
January 1949	21	67



CRPL and IRPL Reports

[A list of CRPL Section Reports is available from the Central Radio Propagation Laboratory upon request]

Daily:

Radio disturbance warnings, every half hour from broadcast station WWV of the National Bureau of Standards. Telephoned and telegraphed reports of ionospheric, solar, geomagnetic, and radio propagation data.

Weekly:

CRPL-J. Radio Propagation Forecast (of days most likely to be disturbed during following month).

Semimonthly:

CRPL-Ja. Semimonthly Frequency Revision Factors for CRPL Basic Radio Propagation Prediction Reports.

Monthly:

CRPL-D. Basic Radio Propagation Predictions—Three months in advance. (Dept. of the Army, TB 11-499-, monthly supplements to TM 11-499; Dept. of the Navy, DNC-13-1 (), monthly supplements to DNC-13-1.)

CRPL-F. Ionospheric Data.

Quarterly:

*IRPL-A. Recommended Frequency Bands for Ships and Aircraft in the Atlantic and Pacific.

*IRPL-H. Frequency Guide for Operating Personnel.

Circulars of the National Bureau of Standards:

NBS Circular 462. Ionospheric Radio Propagation.

NBS Circular 465. Instructions for the Use of Basic Radio Propagation Predictions.

Reports issued in past:

IRPL-C61. Report of the International Radio Propagation Conference, 17 April to 5 May 1944.

IRPL-G1 through G12. Correlation of D. F. Errors With Ionospheric Conditions.

IRPL-R. Nonscheduled reports:

R4. Methods Used by IRPL for the Prediction of Ionosphere Characteristics and Maximum Usable Frequencies.

R5. Criteria for Ionospheric Storminess.

R6. Experimental Studies of Ionospheric Propagation as Applied to the Loran System.

R7. Second Report on Experimental Studies of Ionospheric Propagation as Applied to the Loran System.

R9. An Automatic Instantaneous Indicator of Skip Distance and MUF.

R10. A Proposal for the Use of Rockets for the Study of the Ionosphere.

R11. A Nomographic Method for Both Prediction and Observation Correlation of Ionosphere Characteristics.

R12. Short Time Variations in Ionospheric Characteristics.

R14. A Graphical Method for Calculating Ground Reflection Coefficients.

R15. Predicted Limits for F2-layer Radio Transmission Throughout the Solar Cycle.

R17. Japanese Ionospheric Data—1943.

R18. Comparison of Geomagnetic Records and North Atlantic Radio Propagation Quality Figures—October 1943 Through May 1945.

R21. Notes on the Preparation of Skip-Distance and MUF Charts for Use by Direction-Finder Stations. (For distances out to 4000 km.)

R23. Solar-Cycle Data for Correlation with Radio Propagation Phenomena.

R24. Relations Between Band Width, Pulse Shape and Usefulness of Pulses in the Loran System.

R25. The Prediction of Solar Activity as a Basis for the Prediction of Radio Propagation Phenomena.

R26. The Ionosphere as a Measure of Solar Activity.

R27. Relationships Between Radio Propagation Disturbance and Central Meridian Passage of Sunspots Grouped by Distance From Center of Disc.

R30. Disturbance Rating in Values of IRPL Quality-Figure Scale from A. T. & T. Co. Transmission Disturbance Reports to Replace T. D. Figures as Reported.

R31. North Atlantic Radio Propagation Disturbances, October 1943 Through October 1945.

R33. Ionospheric Data on File at IRPL.

R34. The Interpretation of Recorded Values of fEs.

R35. Comparison of Percentage of Total Time of Second-Multiple Es Reflections and That of fEs in Excess of 3 Mc.

IRPL-T. Reports on tropospheric propagation:

T1. Radar operation and weather. (Superseded by JANP 101.)

T2. Radar coverage and weather. (Superseded by JANP 102.)

CRPL-T3. Tropospheric Propagation and Radio-Meteorology. (Reissue of Columbia Wave Propagation Group WPG-5.)

*Items bearing this symbol are distributed only by U. S. Navy. They are issued under one cover as the DNC-14 series.

