

National Bureau of Standards
Library, N.W. Bldg

Reference book in 10 06
taken from the Library.

APR 4 1956

PART B
SOLAR - GEOPHYSICAL DATA

ISSUED
MARCH 1956

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
CENTRAL RADIO PROPAGATION LABORATORY
BOULDER, COLORADO

SOLAR - GEOPHYSICAL DATA

CONTENTS

INTRODUCTION

Description of Tables and Graphs

I RELATIVE SUNSPOT NUMBERS

- (a) American and Zürich Daily Numbers
- (b) Graph of Sunspot Cycle

II SOLAR CENTERS OF ACTIVITY

- (a) Calcium Plage and Sunspot Regions
- (b) Coronal Line Emission Indices

III SOLAR FLARES

- (a) Optical Observations
- (b) Optical Observations (continued)
- (c) Ionospheric Effects

IV SOLAR RADIO WAVES

- (a) 167 Mc -- 3-hourly and Daily Flux (Boulder)
- (b) 460 Mc -- 3-hourly and Daily Flux (Boulder)
- (c) 167 Mc -- Outstanding Events (Boulder)
- (d) 460 Mc -- Outstanding Events (Boulder)

V GEOMAGNETIC ACTIVITY INDICES

- (a) C, Kp, Ap, and Selected Quiet and Disturbed Days
- (b) Chart of Kp by Solar Rotations

VI RADIO PROPAGATION QUALITY INDICES

North Atlantic:

- (a) CRPL Quality Figures and Forecasts
- (b) Graphs Comparing Forecast and Observed Quality
- (c,d) Graphs of Useful Frequency Ranges

North Pacific:

- (e) CRPL Quality Figures and Forecasts
- (f) Graphs Comparing Forecast and Observed Quality

SOLAR - GEOPHYSICAL DATA

INTRODUCTION

This monthly report series is intended to keep research workers abreast of the major particulars of solar activity and the associated ionospheric, radio propagation and other geophysical effects. It is made possible through the cooperation of many observatories, laboratories and agencies as recorded in the detailed description of the tables and graphs which follows. The Editor is Miss J. V. Lincoln.

I RELATIVE SUNSPOT NUMBERS

American and Zürich Daily Numbers -- The table lists (1) the daily American relative sunspot numbers, RA' , as compiled by the Solar Division of the American Association of Variable Star Observers, and (2) the provisional daily Zürich relative sunspot numbers, RZ , as communicated by the Swiss Federal Observatory. Because of the time required to collect and reduce the observations, RA' will normally appear one month later than RZ .

The relative sunspot number is an index of the activity of the entire visible disk. It is determined each day without reference to preceding days. Each isolated cluster of sunspots is termed a sunspot group and it may consist of one or a large number of distinct spots whose size can range from 10 or more square degrees of the solar surface down to the limit of resolution (e.g. $1/8$ square degrees). The relative sunspot number is defined as $R=K(10g+s)$, where g is the number of sunspot groups and s is the total number of distinct spots. The scale factor K (usually less than unity) depends on the observer and is intended to effect the conversion to the scale originated by Wolf. The observations for sunspot numbers are made by a rather small group of extraordinarily faithful observers, many of them amateurs, each with many years of experience. The counts are made visually with small, suitably protected telescopes.

Final values of RZ appear in the IAU Quarterly Bulletin on Solar Activity, the Journal of Geophysical Research and elsewhere. They usually differ slightly from the provisional values. The American numbers, RA' , are not revised.

Graph of Sunspot Cycle -- The graph illustrates the recent trend of Cycle 19 of the 11-year sunspot cycle and some predictions of the future level of activity. The customary "12-month" smoothed

index, \bar{R} , is used throughout, the data being final R_z numbers except for the current year. Predictions shown are those made for one year after the latest available datum by the method of A. G. McNish and J. V. Lincoln (Trans. Am. Geophys. Union, 30, 673-685, 1949) modified by the use of regression coefficients and mean cycle values recomputed for Cycles 8 through 18. Cycle 19 began April 1954, when the minimum \bar{R} of 3.4 was reached.

II SOLAR CENTERS OF ACTIVITY

Calcium Plage and Sunspot Regions -- The table gives particulars of the centers of activity visible on the solar disk during the preceding month. These are based on estimates made and reported on the day of observation and are therefore of limited reliability.

The table gives the heliographic coordinates of each center (taken as the calcium plage unless two or more significantly and individually active sunspot groups are included in an extended plage) in terms of the Greenwich date of passage of the sun's central meridian (CMP) and the latitude; the serial number of the plage as assigned by McMath-Hulbert Observatory; the serial number of the center in the previous solar rotation, if it is a persisting region; particulars of the plage at three times during its transit of the visible disk (first appearance, maximum development, last appearance): the date, the area, the central intensity; particulars of the associated sunspot group, if any, at analogous times: the date, the area, the spot count. The unit of area is a millionth of the area of a solar hemisphere; the central intensity of calcium plages is roughly estimated on a scale of 1=faint to 5=very bright.

Calcium plage data are available through the cooperation of the McMath-Hulbert Observatory of the University of Michigan and the Mt. Wilson Observatory. The sunspot data are compiled from reports from the U. S. Naval Observatory, Mt. Wilson Observatory, and from reports from Europe and Japan received through the daily Ursigram messages.

Coronal Line Emission Indices -- In the table are summarized solar coronal emission intensity indices for the green (Fe XIV at $\lambda 5303$) and red (Fe X at $\lambda 6374$) coronal lines. The indices are based on measurements made at 5° intervals around the periphery of the solar disk by the High Altitude Observatory at Climax, Colorado, and by Harvard University observers at Sacramento Peak (The USAF Upper Air Research Observatory at Sunspot, New Mexico, under contract AF 19(604)-146). The measurements are expressed as the number of millionths of an Angstrom of the continuum of the center of the solar disk (at the same wavelength as the line) that would contain the same energy as the observed coronal line. The indices have the following meanings:

G_6 = mean of six highest line intensities in quadrant for $\lambda 5303$.

R_6 = same for $\lambda 6374$.

G_1 = highest value of intensity in quadrant, for $\lambda 5303$.

R_1 = same for $\lambda 6374$.

The dates given in the table correspond to the approximate time of CMP of the longitude zone represented by the indices. The actual observations were made for the North East and South East quadrants 7 days before; for the South West and North West quadrants 7 days after the CMP date given.

To obtain rough measures of the integrated emission of the entire solar disk in either of the lines, assuming the coronal changes to be small in a half solar rotation, it is satisfactory to perform the following type of summation given in example for 15 October:

$$\left(\text{MEAN DISK EMISSION} \right)_{15 \text{ OCT}}^{\text{IN } \lambda 5303} = \frac{1}{N} \left[\sum_{15 \text{ OCT}}^{22 \text{ OCT}} \left\{ (G_6)_{\text{NE}} + (G_6)_{\text{SE}} \right\} + \sum_{8 \text{ OCT}}^{14 \text{ OCT}} \left\{ (G_6)_{\text{SW}} + (G_6)_{\text{NW}} \right\} \right]$$

where N is the number of indices entering the summation.

Such integrated disk indices as well as integrated whole-sun indices are computed for each day and are published quarterly in the "Solar Activity Summary" issued by the High Altitude Observatory at Boulder, Colorado. In the same reports are given maps of the intensity distribution of coronal emission derived from all available Climax and Sacramento Peak observations, as well as other information on solar activity, such as maps made from daily limb prominence surveys in $H\alpha$ and notes regarding the history of active regions on the solar disk.

Preliminary summaries of solar activity, prepared on a fast schedule, are issued Friday of each week from High Altitude Observatory in conjunction with CRPL and include solar activity through the preceding day. These are useful to groups needing information on the current status of activity on the visible solar disk, but are not recommended for research uses unless such a prompt schedule of reporting is essential. The same information is included in the subsequent quarterly reports, with extensive additions, corrections and evaluations.

III SOLAR FLARES

Optical Observations -- The table presents the preliminary record of solar flares as reported to the CRPL on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete data are published later in the Quarterly Bulletin of Solar Activity, I.A.U., in various observatory publications and elsewhere. The present listing serves to identify and roughly describe the phenomena observed.

Reporting directly to the CRPL are the following observatories: Mt. Wilson, McMath-Hulbert, U. S. Naval, Wendelstein, and Sacramento Peak. The remainder report through the URSIgram centers in Europe and Japan. Observations are in the light of the center of the H-alpha line unless noted otherwise. The reports from Sacramento Peak, New Mexico (communicated to CRPL by the High Altitude Observatory at Boulder) are from observations at the USAF Upper Air Research Observatory at Sunspot, New Mexico, by Harvard University observers, under contract AF 19(604)-146.

For each flare are listed the reporting observatory, date, times of beginning and ending of observing period, duration of flare (when known), total area in millionths of visible hemisphere, the McMath serial number of the region with which the flare is associated, the heliographic coordinates in degrees, the time of maximum phase, maximum intensity of flare, fractional area having nearly maximum brightness, and finally the flare importance on the IAU scale of 1- to 3+. A final column lists provisionally the occurrence of simultaneous ionospheric effects as observed on selected field strength recordings of distant high-frequency radio transmissions; a more nearly definitive list of these ionospheric effects, including particulars, appears in these reports after the lapse of a month (see below). All times are Universal Time (UT or GCT).

Ionospheric Effects -- SID (and GID--gradual ionospheric disturbances) may be detected in a number of ways: short wave fadeouts, enhancement of low frequency atmospherics, increases in cosmic absorption, and so forth. The table lists events that have been recognized on field strength recordings of distant high-frequency radio transmissions. Under a coordinated program, the staffs at the following ionospheric sounding stations contribute reports that are screened and synthesized at CRPL-Boulder: Puerto Rico, Ft. Belvoir, Va., and Anchorage, Alaska (CRPL Stations: PR, BE, AN); Huancayo, Peru, and College, Alaska (CRPL-Associated Laboratories: HU, CO); and White Sands, N. Mex., Adak, Alaska, and Okinawa (U. S. Signal Corps Stations: WS, AD, OK). McMath-Hulbert Observatory (MC) also contributes such reports. In addition, reports are volunteered by RCA Communications Inc.,

Marconi Wireless, Netherlands Postal and Telecommunications Services, Swedish Telecommunications, and others; these usually specify times of SID and the radio paths involved.

In the coordinated program, the abnormal fades of field strength not obviously ascribable to other causes, are described as short wave fadeouts with the following further classification:

- S-SWF: sudden drop-out and gradual recovery
- Slow S-SWF: drop-out taking 5 to 15 minutes and gradual recovery
- G-SWF: gradual disturbance; fade irregular in both drop-out and recovery.

When there is agreement among the various reporting stations on the time (UT) of an event, it is accepted as a widespread phenomenon and listed in the table.

The degree of confidence in identifying the event, a subjective estimate, is reported by the stations and this is summarized in an index of certainty that the event is widespread, ranging from 1 (possible) to 5 (definite). The times given in the table for the event are from the report of a station (underlined in table) that identified it with high confidence. The criteria for the subjective importance rating assigned by each station on a scale of 1- to 3+ include amplitude of the fade, duration and confidence; greater consideration is given to reports on paths near the subsolar point in arriving at the summary importance rating given in the table.

Note: The tables of SID observed at Washington included in CRPL F-reports prior to F-135 were restricted to events classed here as S-SWF.

IV SOLAR RADIO WAVES

The data on solar radio waves are from observations at 167 Mc and 460 Mc made at the Gunbarrel Hill (Boulder) station of the National Bureau of Standards. The half-width of the antenna lobe is appreciably greater than the solar disk. Polarization has not been determined. All times are in Universal Time (UT or GCT); when the observing period extends slightly into the next Greenwich day, the time scale is extended beyond 24 hours.

3-hourly and Daily Flux -- Flux is given in power units. These units are approximately 10^{-22} watt meter⁻²(c/s)⁻¹ for both polarizations together. They will be subject to a correction factor when gain measurements of the antenna have been made. The median flux is measured for every one-hour period that contains a usable calibration and at least thirty minutes of usable record. A three-hour value of flux is obtained by averaging the available one-hour medians (at least two required). A daily value of flux is obtained by averaging all available one-hour medians (at least 4 required). A dash indicates that insufficient measurements were made to meet the above requirements or that the records were not of usable quality. Parentheses indicate that the value is somewhat doubtful because of atmospheric noise or local interference.

The variability index, given for each three-hour interval, is on a scale 0 to 3 defined as follows:

0 - The instantaneous flux did not drop below one-half the median level or exceed twice the median level at any time.

1 - The instantaneous flux made from one to ten excursions outside the range described above.

2 - The instantaneous flux made from ten to one hundred excursions outside the range described above.

3 - The instantaneous flux made more than one hundred excursions outside the range described above.

For the purpose of the variability index, an excursion whose maximum intensity is M times the median level is counted as M excursions. A dash is used to indicate that measurements were made for less than one hour during the period. Parentheses surround variability indices which are in doubt because of atmospheric noise or local interference.

Outstanding Events -- A separate table lists the occurrences that are not adequately described by the three-hourly values of median flux and variability. These are classified in general accordance with the system described and illustrated by Dodson, Hedeman, and Owren (Ap. J. 118, 169, 1953). The categories of events are identified in the table by numbers, which do not necessarily indicate the magnitude of the event:

0 - Rise in base level -- A temporary increase in the continuum with duration of the order of tens of minutes to an hour.

1 - Series of bursts -- Bursts or groups of bursts, occurring intermittently over an interval of time of the order of minutes or hours. Such series of bursts are assigned as distinctive events only when they occur on a smooth record or show as a distinct change in the activity.

2 - Groups of bursts -- A cluster of bursts occurring in an interval of time of the order of minutes.

3 - Minor burst -- A burst of moderate or small amplitude, and duration of the order of one or two minutes.

4 - Minor burst and second part -- A double rise in flux in which the early rise is a minor burst.

5 - Noise storm ends -- A noise storm (see 6) which ceases at some time during the observing period.

6 - Noise storm -- A temporary increase in radiation characterized by numerous closely spaced bursts, by an increase in the continuum, or by both. Duration is of the order of hours or days.

7 - Noise storm begins -- The onset of a noise storm occurs at some time during the observing period.

8 - Major burst -- An outburst, or other burst of large amplitude and more than average duration. A major burst is usually complex, with a duration of the order of one to ten minutes.

9 - Major burst and second part -- A double rise in flux, the first part of which is a major burst. The second part may consist of a rise in base level, a group or series of bursts, or the onset of a noise storm.

Starting times and durations are enclosed in parentheses when they are limited by the period of observation. The maximum instantaneous flux (Inst. Flux) is measured from the sky level as are the hourly medians. The maximum smoothed flux (Smd. Flux) is that obtained by taking the difference of the maximum value of a smooth curve drawn through the outstanding occurrence with a smoothing period of 20 percent to 50 percent of the total duration, and the value of the interpolated hourly median at that same time had the event not occurred, both measured from the sky level.

V GEOMAGNETIC ACTIVITY INDICES

C, Kp, Ap, and Selected Quiet and Disturbed Days -- The data in the table are: (1) preliminary international character figures, C; (2) geomagnetic planetary three-hour range indices, Kp; (3) daily "equivalent amplitude," Ap; (4) magnetically selected quiet and disturbed days.

This table is made available by the Committee on Characterization of Magnetic Disturbance of IAGA, IUGG. The Meteorological Office, De Bilt, Holland collects the data from magnetic observatories distributed throughout the world, and compiles C and selected days. The Chairman of the Committee computes the planetary and equivalent amplitude indices. The same data are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm).

Kp is the mean standardized K-index from 12 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g. 5- is $4\frac{2}{3}$, 5o is $5\frac{0}{3}$, and 5+ is $5\frac{1}{3}$. This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of Kp has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948" of the Association of Terrestrial Magnetism and Electricity (IATME), International Union of Geodesy and Geophysics.

Ap is a daily index of magnetic activity on a linear scale rather than on the quasi-logarithmic scale of the K-indices. It is the average of the eight values of an intermediate 3-hourly index "ap," defined as one-half the average gamma range of the most disturbed of the three force components, in the three-hour interval at standard stations; in practice, ap is computed from the Kp for the 3-hour interval. The extreme range of the scale of Ap is 0 to 400. The method is described in IATME Bulletin No. 12h (for 1953) p. viii f. Values of Ap (like Kp and Cp) have been published for the Polar Year 1932/33 and for the years 1937 onwards.

The magnetically quiet and disturbed days are selected in accordance with the general outline in Terr. Mag. (predecessor to J. Geophys. Res.) 48, pp 219-227, December 1943. The method in current use calls for ranking the days of a month by their geomagnetic activity as determined from the following three criteria with equal weight: (1) the sum of the eight Kp's; (2) the sum of the squares of the eight Kp's; and (3) the greatest Kp.

Chart of Kp by Solar Rotations -- The graph of Kp by solar rotations is furnished through the courtesy of Dr. J. Bartels, Geophysikalisches Institute, Göttingen.

VI RADIO PROPAGATION QUALITY INDICES

One can take as the definition of a radio propagation quality index: the measure of the efficiency of a medium-powered radio circuit operated under ideal conditions in all respects, except for the variable effect of the ionosphere on the propagation of the transmitted signal. The indices given here are derived from monitoring and circuit performance reports, and are the nearest practical approximation to the ideal index of propagation quality.

Quality indices are usually expressed on a scale that ranges from one to nine. Indices of four or less are generally taken to represent significant disturbance. (Note that for geomagnetic K-indices, disturbance is represented by higher numbers.) The adjectival equivalents of the integral quality indices are as follows:

1 = useless	4 = poor-to-fair	7 = good
2 = very poor	5 = fair	8 = very good
3 = poor	6 = fair-to-good	9 = excellent

CRPL forecasts are expressed on the same scale. The tables summarizing the outcome of forecasts include categories P-Perfect; S-Satisfactory; U-Unsatisfactory; F-Failure. The following conventions apply:

P - forecast quality equal to observed	U - forecast quality two or more grades different from observed when <u>both</u> forecast and observed were ≥ 5 , or both ≤ 5
S - forecast quality one grade different from observed	F - other times when forecast quality two or more grades different from observed

Full discussion of the reliability of forecasts requires consideration of many factors besides the over-simplified summary given.

The quality figures represent a consensus of experience with radio propagation conditions. Since they are based entirely on monitoring or traffic reports, the reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often

be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality for reasons such as multipath or interference. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

North Atlantic Radio Path -- The CRPL quality figures, Qa, are compiled by the North Atlantic Radio Warning Service (NARWS), the CRPL forecasting center at Ft. Belvoir, Virginia, from radio traffic data for North Atlantic transmission paths closely approximating New York-to-London. These are reported to CRPL by the Canadian Defense Research Board, Canadian Broadcasting Company, and the following agencies of the U. S. Government:--Coast Guard, Navy, Army Signal Corps, U. S. Information Agency. Supplementing these data are CRPL monitoring, direction-finding observations and field strength measurements of North Atlantic transmissions made at Belvoir.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the original scale. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year, with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. The 6-hourly quality figure is the mean of the reports available for that period.

The 6-hourly quality figures are given in this table to the nearest one-third of a unit, e.g. 5.0 is 5 and 0/3; 5.5 is 4 and 2/3; 5.8 is 5 and 1/3. Other data included are:

(a) Whole-day radio quality indices, which are weighted averages of the four 6-hourly indices, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which seek to designate the days of significant disturbance or unusually quiet conditions.

(b) Short-term forecasts, issued every six hours by the North Atlantic Radio Warning Service. These are issued one hour before 00h, 06h, 12h, 18h, UT and are applicable to the period 1 to 7 hours ahead.

(c) Advance forecasts, issued twice weekly by the NARWS (CRPL-J reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole-day quality indices.

(d) Half-day averages of the geomagnetic K indices measured by the Cheltenham Magnetic Observatory of the U. S. Coast and Geodetic Survey.

A chart compares the short-term forecasts with Qa-figures. A second chart compares the outcome of advance forecasts (1 to 3 or 4 days ahead) with a type of "blind" forecast. For the latter, the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

Ranges of useful frequencies on the North Atlantic radio path are shown in a series of diagrams, one for each day. Time is the angular coordinate and radio frequency in Mc is the radius vector. The shaded area indicates the range of frequencies for which transmissions of quality 5 or greater were observed. The blacker the diagram, the quieter the day has been; a narrow strip indicates either high LUHF, low MUF or both. These diagrams are based on data reported to CRPL by the German Post Office through the Fernmeldetechnischen Zentralamtes, Darmstadt, Germany, being observations every one and a half hours of selected transmitters located in the eastern portion of North America.

Note: Beginning with data for September 1955, Qa has been determined from reports that are available within a few hours or at most within a few days, including for the first time, the CRPL observations. Therefore these are the indices by which the forecasters assess every day the conditions in the recent past. Over a period of several years, they have closely paralleled the former Qa indices which included CRPL observations and included three additional reports received after a considerable lag. Qa was first published to the nearest one-third of a unit at the same time.

North Pacific Radio Path -- The CRPL quality figures, Qp, are compiled by the North Pacific Radio Warning Service (NPRWS), the CRPL forecasting center at Anchorage, Alaska, from radio traffic data for moderately long transmission paths in the North Pacific equivalent to Seattle-to-Anchorage or Anchorage-to-Tokyo. These include reports to CRPL by the Alaskan Communications Service, Aeronautical Radio, Inc., U. S. Air Force and Civil Aeronautical Administration. In addition, there are CRPL monitoring, direction-finder observations and field strength measurements of suitable transmissions.

The original reports are on various scales and for various time intervals. The observations for each 9 hours or 24 hour period are averaged on the original scale. This average is compared with reports for the same period in the preceding two months and expressed as a deviation from the 3-month mean. The deviations are put on the 1 to 9 scale of quality which is assumed to have a standard deviation of 1.25 and a mean for the various periods as follows:

03-12 hours UT	5.33
09-18	5.33
18-03	6.00
00-24	5.67

The 9-hour and 24-hour indices Q_p are determined separately. Each index is a weighted mean where the CRPL observations have unit weight and the others are weighted by the correlation coefficient with the CRPL observations.


The table, analagous to that for Q_a , includes the 9-hourly quality figures; whole day quality figures; short term forecasts issued by NPRWS three times daily at 02^h, 09^h, and 18^h UT, applicable to the stated 9-hour periods; advance forecasts issued twice weekly by NPRWS (CRPL-Jp report); and half-day averages of geomagnetic K indices from Sitka.

The chart compares the outcome of advance forecasts, on the same basis as the similar chart for the North Atlantic Radio Path.

RELATIVE SUNSPOT NUMBERS

American Relative Sunspot Numbers	
January 1956	
Date	R_A
1	48
2	42
3	52
4	55
5	42
6	44
7	47
8	57
9	52
10	46
11	49
12	59
13	78
14	69
15	97
16	115
17	108
18	97
19	95
20	106
21	100
22	89
23	86
24	79
25	81
26	57
27	41
28	43
29	57
30	52
31	41
Mean:	67.2

Zürich Provisional Relative Sunspot Numbers	
February 1956	
Date	R_Z
1	40
2	50
3	60
4	57
5	26
6	34
7	29
8	25
9	31
10	29
11	56
12	80
13	90
14	142
15	168
16	224
17	237
18	270
19	246
20	260
21	208
22	186
23	177
24	156
25	149
26	140
27	122
28	132
29	132
Mean:	122.6



PREDICTED AND OBSERVED SUNSPOT NUMBERS

CALCIUM PLAGE AND SUNSPOT REGIONS

FEBRUARY 1956

CMP Feb. 1956	Lat.	McMath Plage Number	Return of Region	Calcium Plage Data			Sunspot Data		
				Date-Area-Intensity			Date-Area-Count		
				First seen	Maximum	Last seen	First seen	Maximum	Last seen
02.4	N23	3389 (1)*	New	03- 800-2	----	04- 300-1.5	03- xx -1	----	04- xx -1
04.0	N25	3390 (5)	3370	01- 200-1	02- 1800-2	05- 500-1	01- 60 -2	01- 60 -2	04- xx -1
05.1	S36	3386 (1)	New	01- 200-1	02- 700-1	10-1100-1	29- 50a-x	02- 200a-x	04- xx -2
05.2	S18	3387 (3)	3372	01- 500-1	02- 1000-3	10- 700-1	31-110 -1	02- 200a-x	06- 50a-x
06.5	N30	3388 (4)	3371	01- 2000-2	07- 5700-4	10- 4800-2.5	31-170 -1	01- 540 -5	10- 50a-x
07.0	S25	3391 (3)	3372	03- 1000-2	04- 1100-2	10- 400-1			
08.6	S15	3392 (2)	3373	03- 1200-2.5	----	10- 300-1			
09.0	N24	3394 (3)	3374	03- 2000-2	----	10- 1000-1	08- 10 -x	09- 110 -2	12- 50a-x
09.3	S27	3393 (2)	3375	03- 2000-2	----	14- 800-1			
09.8	N20	3395 (3)	3374	08- 700-2	13- 1000-2.5	15- 1000-2			
11.3	N22	3406 (3)	3374	15- 800-2.5	----	16- 1500-2			
12.3	N09	3408 (1)	New	18- 1000-3.5	----	18- 1000-3.5			
12.6	N17	3407 (1)	New	16- 700-1.5	16- 700-1.5	16- 700-1.5			
12.9	S32	3396 (3)	3378	10- 1300-2.5	10- 1300-2.5	15- 1000-1			
13.0	S17	3397 (1)	New	10- 1200-2	10- 1200-2	16- 900-2			
15.5	N30	3398 (2)	3379	10- 1900-2.5	19- 3500-2	20- 3000-1.5			
16.2	S22	3399 (5)	3380	10- 1000-1	18- 1300-2	21-1500-2	11- 10 -x	13- 130 -6	20- 60 -5
16.5	N42	3401 (1)	New	10- 1000-2.5	13- 4100-2	21- 600-2			
16.8	N21	3400 (2)	3379	10- 2000-3	19-18000-3.5	23- 9000-3	10-340 -1**	{ 18- 650 -6 14-2410 -28 21-1730 -15	24-870 -2**
18.9	S22	3403 (1)	New	14- 5000-3.5	18- 6300-3.5	25- 5000-3	13-290 -3	18-1660 -20	24-970 -2
20.0	N22	3404 (1)	New	15-10000-4	18-12600-3.5	26-10000-3	14-150 -1	20-1280 -35	26-580 -3
21.3	S22	3405 (1)	New	14- 2000-3	16- 3500-4	27- 1400-1	14-730 -1	26- 970 -1	27-440 -1
23.0	N23	3411 (1)	New	19- 500-2	25- 800-2	28-1000-1	23- 40 -4	23- 40 -4	23- 40 -4
24.2	N36	3409 (?)	3383	18- 900-1	19- 1000-2	21- 800-1			
25.7	S22	3410 (?)	3384	18- 1500-1.5	29- 5000-2	01- 3000-2	19- 70 -1	24- 220 -6	01- 10 -1
25.9	N23	3412 (?)	3385	19- 4000-3.5	29-13000-3	02-11400-2	19-290 -1	24- 970 -10	02-390 -7
28.5	S24	3413 (1)	New	20- 1000-2	25- 5800-3	05- 3000-2	21-390 -3	21- 390 -3	02- 20 -1

* Parenthetical value following McMath plage region number is age of region in number of rotations.

** First and last parts of the giant region.

CORONAL LINE EMISSION INDICES

FEBRUARY 1956

CMP Date 1956	North East Quadrant (observed 7 days earlier)				South East Quadrant (observed 7 days earlier)				South West Quadrant (observed 7 days later)				North West Quadrant (observed 7 days later)			
	G ₆	G ₁	R ₆	R ₁	G ₆	G ₁	R ₆	R ₁	G ₆	G ₁	R ₆	R ₁	G ₆	G ₁	R ₆	R ₁
Feb.																
1	25 ^a	35 ^a	7	10	31 ^a	42 ^a	9	9	X	X	X	X	X	X	X	X
2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
3	X	X	X	X	X	X	X	X	15	18	8	10	16	19	10	14
4	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
5	26 ^a	40 ^a	X	X	X	X	X	X	31	43	17	19	45	68	15	41
6	X	X	X	X	X	X	X	X	34	43	18	22	66	100	21	26
7	X	X	X	X	X	X	X	X	26	32	18	22	54	72	14	18
8	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
9	65	92	13	21	49	80	11	16	X	X	X	X	X	X	X	X
10	45	83	X	X	36	61	16	25	X	X	X	X	X	X	X	X
11	72	99	12	15	85	150	19	36	X	X	X	X	X	X	X	X
12	17	25	8	10	40	77	15	32	30	34	14	23	38	49	15	25
13	19	23	8	11	50	58	16	22	29	34	21	32	31	38	21	25
14	X	X	X	X	X	X	X	X	31	41	7	12	35	50	31	70
15	X	X	X	X	X	X	X	X	X	X	X	X	45	59	X	X
16	83*	109	20	25	X	X	X	X	X	X	X	X	X	X	X	X
17	74*	93	32	38	26	45	15	28	X	X	X	X	X	X	X	X
18	X	X	X	X	X	X	X	X	31	47	16	25	51*	79	28	46
19	57*	88	24	32	36*	60	23	39	X	X	X	X	X	X	X	X
20	70*	92	25	46	35*	60	21	34	27	43	20	36	65	120	25	36
21	47	72	17	23	18	28	18	22	22	31	25	45	34	62	20	30
22	X	X	X	X	X	X	X	X	18	20	15	22	33	42	12	19
23	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
24	X	X	X	X	X	X	X	X	38	88	17	36	61	83	16	28
25	X	X	X	X	X	X	X	X	44	96	24	39	91*	105	22	36
26	50	72	14	25	47	83	16	30	31	64	23	39	55	75	20	22
27	31	38	X	X	54	87	15	25	29 ^a	49 ^a	14 ^a	43	34 ^a	45 ^a	14 ^a	18 ^a
28	35	40	14	20	52	85	18	28	15 ^a	20 ^a	6 ^a	16 ^a	14 ^a	17 ^a	7 ^a	9 ^a
29	X	X	X	X	55	86	X	X	14	22	7	10	12	23	13	20

* Yellow line observed.
a Index computed from low weight data.

SOLAR FLARES

FEBRUARY 1956

Observatory	Date Feb. 1956	Time Observed		Dura- tion	Total Area	McMath Plage Region Number	Approx. Position		Time Max. Phase	Max. Int. Arb.	Rel. Area of Max. Tenths	Importance	Provis. Iono- spheric Effect
		Start UT	End UT				Lat.	Mer. Dist.					
{ S. Peak	10	2050	2119	29	117	3400	N19	E90	2113	25	8	1	S-SWF
{ McMath	10	2110	2140	30		3400	N20	E90				3	
{ S. Peak	13	1438	1558	80	285	3400	N18	E47	1450	30	5	1+	S-SWF
{ McMath	13	b1530	a1550	> 20		3400	N20	E40				2	
McMath	13	b1718	a1730	> 12		3400	N20	E40				1+	Slow S-SWF
McMath	16	b1750				3403?	S22	E30?				1?	
{ McMath	16	1805	1925	80		3400	N22	E10				2+	Slow S-SWF
{ S. Peak	16	b1815	a2015	>120	227	3400	N18	E08	a1858	25	6	1+	
Wendel.	17	b0750	0830	> 40		3400	N24	E09				1+	
Schau.	17	b0905	0917	> 12		3400	N22	E06				1	
Schau.	17	0906	0918	12		3400	N21	W07				1	
{ Wendel.	17	b1030	1138	> 68		3400	N21	W09				2-	S-SWF
{ Neder.	17	b1125				3400	N19	W02				2	
{ S. Peak	18	2020	a2033	> 13	130	3404	N18	E10	2027	16	2	1	
{ McMath	18	2028	2110	42		3404	N18	E12				1	
S. Peak	19	b1430	1657	> 87	270	3400	N23	W27	1445	30	5	1+	S-SWF
Mt. Wilson	19	b1806		< 10		3404	N25	E15				1	
{ McMath	19	b1819				3403	S20	W06				2	Slow S-SWF
{ S. Peak	19	1820	1845	25	176	3403	S25	W08	1828	25	4	1	
{ Mt. Wilson	19	b1822		< 10		3403	S25	W05				2	
McMath	20	b1405				3400	N24	W40				1	G-SWF
McMath	20	1817	1835	18		3400	N22	W30				1	G-SWF
{ McMath	20	1930	-----+	53		3403	S20	W20				2	S-SWF
{ McMath	20	1957	2023			3403	S22	W23				1	Slow S-SWF
{ S. Peak	20	1935	1955	20	139	3403	S22	W23	1940	30	8	1	
{ S. Peak	20	1958	2011	13	115	3403	S21	W24	2000	26	5	1	
McMath	21	1740	a1750	> 10		3403	S20	W30				1+	S-SWF
S. Peak	22	2137	2210	33	225	3404	N22	W41	2149	20	4	1+	
Tokyo	23	0100	0120	20		3412	N35	E45				1	
Tokyo	23	b0334	0414	> 40		3400	N25	W85				3	S-SWF
McMath	23	b1330				3403	S20	W60				1	
Tokyo	24	0143	0203	20		3412	N25	E35				1	
Tokyo	26	2305	2335	30		3412	N35	E05				1	
Wendel.	28	0715	0832	17		3419	S23	E73				1	
Wendel.	28	0832	0850	18		3419	S23	E72				1	

+ Second brightening of this flare at 1957 UT.

SOLAR FLARES

FEBRUARY 1956

Subflares noted as follows (Date, time (UT), region):

Feb. 10,	1620 (3400)	Feb. 15,	1610 (3400)*	Feb. 21,	1835 (3398)
	2123 (3400)	17,	2006 (3403)		1845 (3404)
11,	1533 (3400)		2124 (3404)		1910 (3412)
	a1539 (3400)	18,	b2042 (3400)*		b2125 (3398)
	1541 (3400)	20,	1444 (3400)	22,	1519 (3404)
	1720 (3400)		1454 (3404)		1620 (3404)
	2008 (3397)		1503 (3403)		1740 (3404)
12,	1500 (3400)		1516 (3400)		1924 (3404)
	1830 (3399)		1549 (3400)		2233 (3403)
	1933 (3400)		1631 (3404)	23,	1539 (3412)
	2056 (3403)		1935 (3404)		1601 (3403)
	2110 (3400)		1959 (3413)		b1705 (3412)
	2210 (3400)	21,	1457 (3400)		b1919 (3412)*
13,	1710 (3400)		1541 (3400)	25,	1520 (3418)
	1810 (3403)		1647 (3400)		1540 (3404)
	1915 (3403)		1657 (3403)		b1805 (3418)*
	2020 (3400)**		a1725 (3404)	26,	1645 (3404)*
	2129 (3400)		1730 (3403)	29,	b2005 (3420)*
	2149 (3404)		b1748 (3404)		
15,	1526 (3400)*		1830 (3412)		

* McMath observation; all others are Sac. Peak

** Observed by both McMath and Sac. Peak

IONOSPHERIC EFFECTS OF SOLAR FLARES

JANUARY 1956

Jan. 1956	Start UT	End UT	Type	Wide- spread Index	Importance	Observation Stations
3	0210	0220	G-SWF	2	1-	AN, <u>OK</u>
4	0900	0950	G-SWF	1	1	<u>OK</u>
7	1914	1930	G-SWF	2	1-	AN, <u>HU</u> , PR
13	1532	1545	G-SWF	2	1-	BE, <u>PR</u>
	1606	1627	G-SWF	3	1	BE, MC, <u>PR</u>
	1717	1726	G-SWF	3	1-	AN, <u>BE</u> , PR
14	0130	0225	G-SWF	1	2	<u>OK</u>
	1932	2007	S-SWF	5	1+	AN, <u>BE</u> , <u>HU</u> , MC, PR
15	1528	1540	Slow S-SWF	4	1+	<u>HU</u> , MC, <u>PR</u>
16	1342	1410	Slow S-SWF	3	1	<u>HU</u> , <u>PR</u> , NERA*
	1646	1720	G-SWF	3	1	AN, <u>HU</u> , MC, PR
17	0847	0922	G-SWF	2	2	NERA*, RCA**
	1642	1718	G-SWF	2	1-	AN, <u>BE</u> , <u>PR</u>
	2016	2042	G-SWF	3	1	AN, <u>HU</u> , <u>PR</u>
18	1423	1450	Slow S-SWF	5	1+	BE, <u>HU</u> , MC, PR
	1755	1816	Slow S-SWF	5	1+	BE, <u>HU</u> , MC, WS
19	0015	0050	S-SWF	4	2	<u>OK</u> , Japan ⁺
	0305	0320	S-SWF	5	2	AN, <u>OK</u> , Japan ⁺
	0558	0630	Slow S-SWF	1	1+	<u>OK</u>
	1556	1725	G-SWF	5	1	BE, <u>HU</u> , MC, PR, NERA*
	1907	1943	G-SWF	4	1+	BE, <u>HU</u> , MC, PR
	2127	2145	G-SWF	3	1	AN, <u>HU</u> , PR
	2245	2300	G-SWF	2	1	AN, <u>OK</u>
20	0405	0435	Slow S-SWF	1	2+	<u>OK</u>
21	1850	1912	G-SWF	3	1-	AN, <u>BE</u> , MC
23	1245	1335	G-SWF	2	1	<u>HU</u> , MC
	1450	1500	G-SWF	4	1	<u>HU</u> , MC, PR, NERA*
	1620	1705	G-SWF	5	2+	BE, <u>HU</u> , MC, <u>PR</u> , NERA*
24	0400	0430	S-SWF	1	1+	<u>OK</u>
	0540	0622	Slow S-SWF	1	1+	<u>OK</u>
25	1336	1405	S-SWF	2	1-	PR, NERA*
	1723	1734	G-SWF	2	1-	BE, MC
26	1648	1724	S-SWF	5	2	BE, <u>HU</u> , MC, PR, WS, NERA*

* Nederhorst den Berg, Netherlands.

** RCA Communications Inc. at Somerton and Brentwood, England.

+ Hiraiso Radio Wave Observatory, Japan.

SOLAR RADIO WAVES (BOULDER) -- 167 MC

3-HOURLY AND DAILY FLUX

FEBRUARY 1956

Feb. 1956	Flux				Daily	Variability				Daily	Observed Periods
	Hours UT					Hours UT					Hours UT
	12 15	15 18	18 21	21 24		12 15	15 18	18 21	21 24		
1											
2											
3											
4											
5											
6	No Scalable Records February 1 through 13.										
7											
8											
9											
10											
11											
12											
13											
14	--	22	25	31	25	--	2	2	2	2	1356-2421
15	--	37	67	--	58	--	2	3	3	3	1354-2150,2238-2422
16	--	66	>300	94	>160	--	2	2	2	2	1353-2423
17	--	72	58	45	60	--	3	2	2	3	1352-2423
18	--	16	21	81	37	--	2	2	2	2	1350-2424
19	--	>220	>280	130	>220	--	1	1	1	1	1349-2426
20	--	34	21	17	25	--	2	2	2	2	1348-2427
21	--	54	50	38	49	--	2	3	3	3	1346-2428
22	--	16	16	14	16	--	2	2	1	2	1345-2430
23	--	16	18	17	17	--	2	2	2	2	1344-2430
24	--	13	12	11	12	--	0	2	1	2	1342-2431
25	--	--	--	--	--	--	--	--	--	--	-----
26	--	12	11	10	11	--	1	1	(1)	(1)	1339-2433
27	--	--	11	10	11	--	1	2	2	2	1338-1606,1827-2433
28	--	15	15	--	15	--	2	2	2	2	1336-2434
29	--	13	13	23	16	--	1	2	3	3	1335-2436

SOLAR RADIO WAVES (BOULDER) -- 460 MC

3-HOURLY AND DAILY FLUX

FEBRUARY 1956

Feb. 1956	Flux					Variability					Observed Periods	
	Hours UT					Hours UT					Hours UT	
	12 15	15 18	18 21	21 24	Daily	12 15	15 18	18 21	21 24	Daily		
1	--	33	34	33	33	--	0	0	0	0	1410-2405	
2	--	34	35	36	35	0	0	1	0	1	1409-2406	
3	--	29	31	--	30	0	0	0	0	0	1408-2407	
4	--	--	29	31	30	0	0	0	(0)	(0)	1407-2408	
5	--	30	31	32	31	0	0	0	0	0	1406-2410	
6	--	31	33	37	34	0	(0)	0	1	1	1405-2411	
7	--	35	33	40	37	0	(0)	(0)	2	2	1404-2412	
8	--	32	--	33	32	(0)	0	1	(0)	1	1403-2413	
9	--	--	--	35	--	0	--	--	(0)	(0)	1401-1643, 2035-2414	
10	--	30	32	41	34	(0)	1	1	2	2	1400-1650, 1708-2416	
11	--	33	33	33	33	0	1	0	0	1	1359-2417	
12	--	32	34	34	33	0	0	1	1	1	1358-2418	
13	--	38	34	34	36	1	0	0	1	1	1357-2312, 2333-2420	
14	--	37	38	38	38	0	(0)	0	0	(0)	1356-2421	
15	--	46	--	--	48	1	0	0	1	1	1354-2010, 2157-2422	
16	--	(44)	(109)	(84)	(83)	0	1	2	2	2	1353-2423	
17	--	--	46	53	47	0	1	1	1	1	1352-1707, 1724-2423	
18	--	39	42	45	42	0	1	0	0	1	1350-2424	
19	--	46	49	47	47	2	2	1	0	2	1349-2426	
20	--	47	47	45	46	1	1	1	(0)	1	1348-2427	
21	--	44	43	42	43	0	1	0	0	1	1346-2428	
22	--	38	36	38	37	1	0	1	(0)	1	1345-2430	
23	--	41	40	39	40	(0)	(0)	1	(0)	1	1344-2203, 2213-2430	
24	--	37	37	38	37	0	0	1	0	1	1342-2431	
25	--	--	--	--	--	--	--	--	--	--		
26	--	--	35	35	35	1	0	0	0	1	1339-2433	
27	--	33	35	35	34	0	0	0	0	0	1338-2433	
28	--	36	37	37	36	0	0	0	(0)	(0)	1336-2213, 2250-2434	
29	--	35	38	44	39	0	0	0	2	2	1335-1847, 1909-2436	

OUTSTANDING EVENTS

FEBRUARY 1956

Feb. 1956	Type	Start UT	Duration Hrs:Mins	Maximum			Remarks
				Time UT	Inst. Flux	Smd. Flux	
14	6	(1356)	(10:25)	~2200	>360	18	Off scale
15	6	(1354)	(10:28)	~2100	>360	70	Off scale
16	6	(1353)	(04:11)	~1500	>300*	60	
16	9	1804	(07:19)	~1930	>620	>320	Off scale
17	6	(1352)	(10:31)	~1700	>1200	60	Off scale
18	6	(1350)	(06:29)	~1600	>900	5	
18	9	2019	(04:05)	2048	>540	100	
19	6	(1349)	(10:37)	~1800	>720	> 280	
19	8	1430	00:07	--	>1300	> 580	Off scale
20	6	(1348)	(10:39)	~1500	>440	23	
21	6	(1346)	(10:42)	~2130	>780	43	
22	6	(1345)	(10:45)	~2000	>320	4	
23	6	(1344)	(10:46)	~1800	>250	7	
24	2	1912	00:06	1917.5	>690	90	
24	3	2312.0	00:00.5	2312.2	130	--	
26	3	1348	00:00.5	1348.2	57	--	
26	8	1440	00:29	1450.4	67	28	
26	4	1941	(04:52)	1941.8	140	--	
27	1	(1338)	(02:28)	1420	78	--	
27	1	(1827)	(06:06)	2029	190	--	
28	6	(1336)	(10:58)	~1800	--	4	
28	3	1627.4	00:00.1	1627.4	>500	--	
28	8	1840.8	00:08.0	1844.0	>940	>360	
29	2	(1335)	(08:49)	1846.0	>290	--	
29	3	1956.1	00:01.8	1956.2	>340	44	
29	9	2224	(02:12)	2225	>1000	120	Off scale

*Most > signs indicate that flux value given was on scale but beyond the presently calibrated range of the recorders and that the actual value is greater.

OUTSTANDING EVENTS


FEBRUARY 1956

Feb. 1956	Type	Start UT	Duration Hrs:Mins	Maximum			Remarks
				Time UT	Inst. Flux	Smd. Flux	
2	3	1427.0	00:00.9	1427.2	~130	--	
6	2	2235.5	00:03.3	2238.7	120	8	
7	8	2128.0	00:05.3	2129.5	>2300	>1000	
10	6	(1400)	(07:16)	~1800	--	3	
10	8	2046.0	00:01.0	2046.7	>1200	--	
10	8	2101.3	00:01.0	2101.5	>1800	--	
10	9	2116	(03:00)	2117.3	>1800	16	
11	4	1534.9	00:03.9	1535.9	>1800	35	
11	3	1602.6	00:02.0	1603.1	>1100	35	
12	2	2111	00:18	2128	130	8	
13	9	1442	02:03	1448	460	5	
13	6	1645	(08:35)	~1700	--	4	
13	2	2026.8	00:01.7	2027.2	160	23	
13	3	2215.9	00:01.0	2216.2	130	--	
14	6	(1356)	(10:25)	~2200	--	9	
14	2	2255.8	00:03.6	2258.3	78	11	
15	6	(1354)	(10:28)	~1900	--	19	
16	6	(1353)	(04:05)	1525	--	14	
16	9	1758	(06:25)	~2020	420	100	
17	6	(1353)	(10:30)	~2100	--	22	
17	3	1402.1	00:01.7	1403.2	~700	--	
17	8	1646.3	00:03.6	1646.8	>1400	32	
18	6	(1350)	(10:34)	~2100	--	17	
19	9	1420	01:07	1436.3	>2200	320	
19	6	1527	(08:59)	~1500	--	19	
19	8	1836.6	00:08	1839.0	290	180	
20	6	(1348)	(10:39)	~1600	--	17	
20	8	1918.8	00:04.0	1920.3	450	140	
21	6	(1346)	(10:42)	~1700	--	13	
21	3	1706.0	00:01.6	1706.7	500	33	
22	1	1416	(10:14)	2058	>750	--	
23	8	1916	00:08	1916.2	130	8	
24	6	(1342)	(10:49)	~2200	--	8	
24	8	1912.0	00:04.5	1913.9	280	7	
24	8	2157	00:11	2205	100	16	
26	3	1416.4	00:00.4	1416.5	>700	--	
28	6	(1336)	(10:58)	~2100	--	8	
29	6	(1335)	(08:47)	~2000	--	10	
29	9	2222	(02:14)	2228.6	410	78	

GEOMAGNETIC ACTIVITY INDICES

JANUARY 1956

Jan. 1956	C	Values Kp								Sum	Ap	Final Selected Days
		Three-hour Gr. interval										
		1	2	3	4	5	6	7	8			
1	1.1	4-	3-	2-	3o	2+	4+	4o	5-	26+	20	Five Quiet
2	0.9	3+	3+	2o	2o	2+	3+	4-	4-	24-	15	
3	0.8	3o	3+	3-	3+	2-	3-	2+	4-	23-	14	8
4	1.2	3-	3+	3-	2+	2+	2+	5o	4-	24+	17	15
5	0.9	3+	3+	3-	2o	2o	2o	4-	2-	21-	12	16
6	1.0	4o	1+	1+	2o	4o	4o	3+	3-	23-	16	20
7	1.0	1+	1+	0o	2-	2+	4o	3-	3o	16+	10	26
8	0.2	2+	1+	1o	2o	1o	1o	2o	1-	11+	5	
9	0.9	2+	4-	2o	4-	4-	3+	2o	2-	22+	14	
10	1.4	0o	3+	4-	4-	4-	5+	5-	4o	28+	26	
11	1.6	6o	4+	5-	6o	4o	3+	5+	2o	36-	42	Five Disturbed
12	1.4	1o	2+	2+	4-	3-	4-	6-	4-	25o	21	
13	0.6	4o	3o	2o	1-	2o	1o	1+	0+	14+	9	
14	0.7	2-	3-	2o	4-	2o	2o	2-	2-	17+	9	11
15	0.0	0o	0+	1o	2-	2-	1o	0+	1+	7+	4	18
												19
16	0.2	3o	1+	1+	2o	2-	1-	2-	1o	13-	6	24
17	0.3	1-	2-	2o	2-	2+	1-	1+	3o	13+	7	28
18	1.7	5o	5+	4+	4o	4o	4o	5-	6-	37o	40	
19	1.3	6-	4o	4+	4-	5o	4-	4-	3-	33-	32	
20	0.1	2o	2-	2o	2-	1o	1-	1-	1o	11-	5	
21	0.8	2o	0o	0o	1-	1-	3+	3-	4+	14-	9	Ten Quiet
22	1.4	6-	6-	4+	4-	2-	1o	4o	1+	27+	29	
23	1.2	1o	3-	3-	1+	2o	3o	4o	5o	22-	16	
24	1.5	3o	5o	4-	5-	5-	4o	6-	5o	36-	38	7
25	0.9	6-	4-	4-	2-	2-	1o	3-	2-	22-	18	8
												13
26	0.0	0o	1-	1o	1-	1-	1-	1-	1+	6-	3	14
27	1.6	2-	1+	2+	4o	4+	5-	6o	3+	28-	27	15
28	1.6	5o	6+	5-	4o	4-	3+	4o	3+	34+	37	16
29	1.0	3-	3-	3-	3o	4+	4-	4-	2+	25o	17	17
30	1.0	4-	3+	2+	2o	3-	4o	2+	4-	24o	16	20
31	0.9	3o	3+	3-	4-	3+	3+	3-	4-	26-	17	21
Mean:	0.94									Me n:	15	26

PLANETARY MAGNETIC
THREE-HOUR-RANGE INDICES

Kp till 1956 January 31

(Ks from Wingst and Göttingen till 1956 February 20)



NORTH ATLANTIC

JANUARY 1956

Jan. 1956	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day index	Advance forecasts (J-reports) for whole day; issued in advance by:			Geomag- netic K _{Ch}	
	00 to 06	06 to 12	12 to 18	18 to 24	00	06	12	18		1-4 days	4-7 days	8-25 days	Half Day (1) (2)	
1	7-	6+	7o	7-	7	6	7	6	7-	7	7		2	3
2	6+	6+	7o	6+	6	6	7	7	7-	7	7		3	3
3	7-	6+	7o	7o	6	6	7	7	7-	7	7		3	2
4	7o	6+	7o	6+	7	7	7	7	7-	7	7		2	3
5	6+	6+	7+	6+	7	6	7	7	7-	7	7		3	2
6	6o	6o	7o	6+	6	7	7	6	7-	7	7		2	3
7	6+	6-	7-	7o	6	6	7	6	7-	7	7		1	3
8	7-	6-	7+	7-	7	6	7	7	7-	7	7		1	1
9	6o	6+	7o	7-	7	6	7	6	7-	7	7		3	2
10	6-	6+	7-	6-	6	6	7	6	6o	6	7		2	(4)
11	4+	5-	7-	6-	5	5	6	6	5+	6	7		(5)	3
12	5o	4+	6o	5o	5	5	6	6	5-	6	7		2	3
13	5o	5+	7-	7o	5	5	7	6	6+	6	6		3	1
14	6+	6-	7+	6+	6	6	7	7	7-	6	6		2	1
15	6o	6-	7o	7o	6	6	7	7	7-	7	6		1	1
16	7-	6+	7+	7-	6	6	7	7	7-	7	6		1	1
17	6+	6+	6+	7-	6	6	7	6	7-	7	7		1	1
18	5+	6-	7-	6o	6	5	6	6	6o	7	7		(4)	(4)
19	6-	6o	6o	6-	4	5	7	6	6-	7	7		(4)	3
20	5o	6-	7-	6+	5	6	6	6	6o	6	6		2	1
21	7-	6+	7o	7-	6	7	7	7	7-	6	6		0	3
22	5-	6-	7-	7o	6	5	6	6	6+	6	6		(4)	2
23	7-	6+	7o	6o	6	6	6	6	7-	6	7		2	3
24	6+	5+	6+	5o	5	6	6	6	6-	6	7		(4)	(4)
25	6-	6o	7-	6+	6	6	6	6	6+	6	7		3	1
26	6o	6o	7-	7o	6	6	7	7	7-	6	7		1	1
27	6+	6+	7+	6o	6	6	7	6	7-	7	6		2	(4)
28	5o	6o	7-	6o	5	4	6	6	6o	7	6		(4)	3
29	5+	6-	7-	6-	6	6	7	6	6o	7	6		2	3
30	5-	5-	7-	5+	5	6	7	6	6-	7	6		2	2
31	6-	6-	7o	7-	5	6	7	6	6+	6	7		3	3
Score: Quiet Periods					P	18	21	22	17		20	18		
					S	11	8	9	14		11	11		
					U	0	0	0	0		0	2		
					F	1	1	0	0		0	0		
Disturbed Periods					P	0	0	0	0		0	0		
					S	1	1	0	0		0	0		
					U	0	0	0	0		0	0		
					F	0	0	0	0		0	0		

() represent disturbed values.



CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS NORTH ATLANTIC

JANUARY 1956

— Short-term forecast

o Quality figure









| Range of reports











USEFUL FREQUENCY RANGES -- NORTH ATLANTIC PATH

JANUARY 1956


DATA ONLY EVERY THIRD HOUR
UNTIL JAN. 2, 0600UT



DATA NOT RECEIVED FROM JAN. 9, 0900UT TO JAN. 26, 0300UT



JANUARY 1956



Adapted from Observations by Deutschen Bundespost

CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS

NORTH PACIFIC

JANUARY 1956

Jan. 1956	North Pacific 9-hourly quality figures			Short-term fore- casts issued at			Whole day index	Advance forecasts (Jp reports) for whole day; issued in advance by:			Geomag- netic K _{Si}	
	03 to 12	09 to 18	18 to 03	02	09	18		1-4 days	4-7 days	8-25 days	Half day (1) (2)	
1	6	6	6	5	5	6	6	6	6		2	3
2	5	5	6	6	5	6	6	6	6		1	3
3	6	6	7	5	6	7	7	6	6		2	2
4	6	5	6	6	6	6	6	6	6		3	3
5	5	5	6	6	6	7	6	5	6		2	2
6	5	5	6	5	5	6	6	5	6		1	(4)
7	5	5	6	6	5	6	6	6	6		1	3
8	5	6	6	5	5	6	5	6	6		2	1
9	6	6	6	5	5	6	6	6	6		3	3
10	6	5	5	6	6	6	6	6	6		3	(5)
11	3	2	5	4	4	5	(3)	6	5		(5)	(5)
12	4	4	4	4	4	5	(4)	5	5		2	(4)
13	3	3	5	5	4	5	(3)	5	6		1	1
14	6	6	5	4	5	6	6	6	4		2	1
15	5	6	6	5	5	6	5	6	4		1	1
16	4	5	5	5	5	6	5	5	5		1	1
17	6	6	5	5	6	6	6	5	5		1	2
18	5	5	5	6	6	6	5	5	6		(4)	(5)
19	5	4	6	5	5	4	(4)	5	6		(4)	(4)
20	5	5	6	5	5	5	6	6	4		2	1
21	5	5	6	6	5	5	5	4	4		0	2
22	6	6	6	6	4	7	6	4	4		(4)	2
23	6	5	4	7	6	7	5	5	5		1	3
24	4	4	4	6	5	5	(3)	5	5		(4)	(5)
25	4	5	5	4	5	6	(4)	6	3		(4)	2
26	5	4	5	5	5	6	5	6	4		0	0
27	5	6	4	4	5	6	(4)	6	5		2	(4)
28	5	5	6	5	5	6	5	5	5		(5)	3
29	5	5	5	6	5	6	5	5	5		2	(4)
30	5	5	5	6	5	6	5	6	6		2	3
31	5	5	6	6	5	6	5	6	6		3	3
Score: Quiet Periods				P	10	13	13	13	12			
				S	14	11	13	10	9			
				U	0	0	0	0	0			
				F	1	1	1	1	3			
Disturbed Periods				P	2	1	0	0	0			
				S	2	4	2	2	3			
				U	1	1	0	2	2			
				F	1	0	2	3	2			

() represent disturbed values.

CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS

NORTH PACIFIC

JANUARY 1956

OUTCOME OF ADVANCE FORECASTS (1 TO 4 DAYS AHEAD)

