

Reference should not be  
made from this

PART B  
SOLAR - GEOPHYSICAL DATA

ISSUED  
APRIL 1958

U. S. DEPARTMENT OF COMMERCE  
NATIONAL BUREAU OF STANDARDS  
CENTRAL RADIO PROPAGATION LABORATORY  
BOULDER, COLORADO



## SOLAR - GEOPHYSICAL DATA

### CONTENTS

#### INTRODUCTION

Description of Tables and Graphs

#### I DAILY SOLAR INDICES

- (a) Relative Sunspot Numbers and 2800 Mc Solar Flux
- (b) Graph of Sunspot Cycle

#### II SOLAR CENTERS OF ACTIVITY

- (a) Calcium Plage and Sunspot Regions
- (b) Coronal Line Emission Indices

#### III SOLAR FLARES

- (a-k) Optical Observations
  - (l) Flare Patrol Observations
  - (m) Subflares
  - (n) Ionospheric Effects

#### IV SOLAR RADIO WAVES

- (a,b) 2800 Mc -- Outstanding Occurrences (Ottawa)
- (c) 2800 Mc -- Times of Observations (Ottawa)
- (d) 200 Mc -- Daily Data (Cornell) March 1958
- (e) 200 Mc -- Outstanding Occurrences (Cornell) March 1958
- (f) 167 Mc -- Daily Data (Boulder) February 1958
- (g,h) 167 Mc -- Outstanding Occurrences (Boulder) February 1958
- (i) 470 Mc -- Daily Data (Boulder) February 1958
- (j) 470 Mc -- Outstanding Occurrences (Boulder) February 1958

#### V GEOMAGNETIC ACTIVITY INDICES

- (a) C, Kp, Ap, and Selected Quiet and Disturbed Days
- (b) Charts of Kp by Solar Rotations

#### VI RADIO PROPAGATION QUALITY INDICES

##### North Atlantic:

- (a) CRPL Quality Figures and Forecasts
- (b) Graphs Comparing Forecast and Observed Quality
- (c,d) Graphs of Useful Frequency Ranges

##### North Pacific:

- (e) CRPL Quality Figures and Forecasts
- (f) Graphs Comparing Forecast and Observed Quality

#### VII ALERT PERIODS AND SPECIAL WORLD INTERVALS

- (a) IGY World Warning Agency Decisions for Alerts and SWI



# SOLAR - GEOPHYSICAL DATA

## INTRODUCTION

This monthly report series is intended to keep research workers abreast of the major particulars of solar activity and the associated ionospheric, radio propagation and other geophysical effects. It is made possible through the cooperation of many observatories, laboratories and agencies as recorded in the detailed description of the tables and graphs which follows. The report is edited by Miss J. V. Lincoln of the Sun-Earth Relationships Section.

### I DAILY SOLAR INDICES

Relative Sunspot Numbers -- The table includes (1) the daily American relative sunspot numbers,  $R_A'$ , as compiled by the Solar Division of the American Association of Variable Star Observers, and (2) the provisional daily Zürich relative sunspot numbers,  $R_Z$ , as communicated by the Swiss Federal Observatory. Because of the time required to collect and reduce the observations,  $R_A'$  will normally appear one month later than  $R_Z$ .

The relative sunspot number is an index of the activity of the entire visible disk. It is determined each day without reference to preceding days. Each isolated cluster of sunspots is termed a sunspot group and it may consist of one or a large number of distinct spots whose size can range from 10 or more square degrees of the solar surface down to the limit of resolution (e.g.  $1/8$  square degrees). The relative sunspot number is defined as  $R=K(10g+s)$ , where  $g$  is the number of sunspot groups and  $s$  is the total number of distinct spots. The scale factor  $K$  (usually less than unity) depends on the observer and is intended to effect the conversion to the scale originated by Wolf. The observations for sunspot numbers are made by a rather small group of extraordinarily faithful observers, many of them amateurs, each with many years of experience. The counts are made visually with small, suitably protected telescopes.

Final values of  $R_Z$  appear in the IAU Quarterly Bulletin on Solar Activity, the Journal of Geophysical Research and elsewhere. They usually differ slightly from the provisional values. The American numbers,  $R_A'$ , are not revised.

Solar Flux Values, 2800 Mc -- The table also lists the daily values of solar flux at 2800 Mc recorded in watts/ $M^2$ /cycle/second bandwidth ( $\times 10^{-22}$ ) in two polarizations by the National Research Council at Ottawa, Canada. These solar radio noise indices are being published in accordance with CCIR Report 25 that a basic solar index for ionospheric propagation should be measured objectively and "preferably refer to a property of the sun such as radiation flux which has direct physical relationship to the ionosphere."

Graph of Sunspot Cycle -- The graph illustrates the recent trend of Cycle 19 of the 11-year sunspot cycle and some predictions of the future level of activity. The customary "12-month" smoothed index,  $R$ , is used throughout, the data being final  $R_Z$  numbers except for the current year. Predictions shown are those made for one year after the latest available datum by the method of A. G. McNish and J. V. Lincoln (Trans. Am. Geophys. Union, 30, 673-685, 1949) modified by the use of regression coefficients and mean cycle values recomputed for Cycles 8 through 18. Cycle 19 began April 1954, when the minimum  $R$  of 3.4 was reached.

## II SOLAR CENTERS OF ACTIVITY

Calcium Plage and Sunspot Regions -- The table gives particulars of the centers of activity visible on the solar disk during the preceding month. These are based on estimates made and reported on the day of observation and are therefore of limited reliability.

The table gives the heliographic coordinates of each center (taken as the calcium plage unless two or more significantly and individually active sunspot groups are included in an extended plage) in terms of the Greenwich date of passage of the sun's central meridian (CMP) and the latitude; the serial number of the plage as assigned by McMath-Hulbert Observatory; the serial number of the center in the previous solar rotation, if it is a persisting region; particulars of the plage at CMP: area, central intensity; a summary of the development of the plage during the current transit of the disk, where  $b$  = born on disk,  $l$  = passed to or from invisible hemisphere,  $d$  = died on disk, and  $/$  = increasing,  $-$  = stable,  $\backslash$  = decreasing; and age in solar rotations; particulars of the associated sunspot group, if any, at CMP: area and spot count and the summary of development during the current disk transit, similar to the above. The unit of area is a millionth of the area of a solar hemisphere; the central intensity of calcium plages is roughly estimated on a scale of 1 = faint to 5 = very bright.

Calcium plage data are available through the cooperation of the McMath-Hulbert Observatory of the University of Michigan and the Mt. Wilson Observatory. The sunspot data are compiled from reports from the U. S. Naval Observatory, Mt. Wilson Observatory, and from reports from Europe and Japan received through the daily Ursigram messages.

Coronal Line Emission Indices -- In the table are summarized solar coronal emission intensity indices for the green (Fe XIV at  $\lambda 5303$ ) and red (Fe X at  $\lambda 6374$ ) coronal lines. The indices are based on measurements made at  $5^\circ$  intervals around the periphery of the solar disk by the High Altitude Observatory at Climax, Colorado, and by Harvard University observers at Sacramento Peak (The USAF Upper Air Research Observatory at Sunspot, New Mexico, under contract AF 19(604)-146). The measurements are expressed as the number of millionths of



an Angstrom of the continuum of the center of the solar disk (at the same wavelength as the line) that would contain the same energy as the observed coronal line. The indices have the following meanings:

$G_6$  = mean of six highest line intensities in quadrant for  $\lambda 5303$ .

$R_6$  = same for  $\lambda 6374$ .

$G_1$  = highest value of intensity in quadrant, for  $\lambda 5303$ .

$R_1$  = same for  $\lambda 6374$ .

The dates given in the table correspond to the approximate time of CMP of the longitude zone represented by the indices. The actual observations were made for the North East and South East quadrants 7 days before; for the South West and North West quadrants 7 days after the CMP date given.

To obtain rough measures of the integrated emission of the entire solar disk in either of the lines, assuming the coronal changes to be small in a half solar rotation, it is satisfactory to perform the following type of summation given in example for 15 October:

$$\left( \text{MEAN DISK EMISSION} \right)_{\text{IN } \lambda 5303} \Big|_{15 \text{ OCT}} = \frac{1}{N} \left[ \sum_{15 \text{ OCT}}^{22 \text{ OCT}} \left\{ (G_6)_{\text{NE}} + (G_6)_{\text{SE}} \right\} + \sum_{8 \text{ OCT}}^{14 \text{ OCT}} \left\{ (G_6)_{\text{SW}} + (G_6)_{\text{NW}} \right\} \right]$$

where N is the number of indices entering the summation.

Such integrated disk indices as well as integrated whole-sun indices are computed for each day and are published quarterly in the "Solar Activity Summary" issued by the High Altitude Observatory at Boulder, Colorado. In the same reports are given maps of the intensity distribution of coronal emission derived from all available Climax and Sacramento Peak observations, as well as other information on solar activity, such as maps made from daily limb prominence surveys in  $H\alpha$  and notes regarding the history of active regions on the solar disk.

Preliminary summaries of solar activity, prepared on a fast schedule, are issued Friday of each week from High Altitude Observatory in conjunction with CRPL and include solar activity through the preceding day. These are useful to groups needing information on the current status of activity on the visible solar disk, but are not recommended for research uses unless such a prompt schedule of reporting is essential. The same information is included in the subsequent quarterly reports, with extensive additions, corrections and evaluations.

### III SOLAR FLARES

Optical Observations -- The table presents the preliminary record of solar flares as reported to the CRPL on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete data are published later in the Quarterly Bulletin on Solar Activity, I.A.U., in various observatory publications and elsewhere. The present listing serves to identify and roughly describe the phenomena observed.

Reporting directly to the CRPL are the following observatories: McMath-Hulbert, Wendelstein, Sacramento Peak, Mitaka and Swedish Astrophysical Station on Capri. The remainder report through the URSIgram centers or are available through the IGY World Data Center for Solar Activity in Boulder. Observations are in the light of the center of the H-alpha line unless noted otherwise. The reports from Sacramento Peak, New Mexico (communicated to CRPL by the High Altitude Observatory at Boulder) are from observations at the USAF Upper Air Research Observatory at Sunspot, New Mexico, by Harvard University observers, under contract AF 19(604)-146.

For each flare are listed the reporting observatory, the date, beginning and ending times, time of maximum phase, the heliographic coordinates in degrees, McMath serial number of the region, duration, the flare importance on the IAU scale of 1- to 3+, observing conditions where 1 means poor, 2 fair and 3 good, time of measurement for tabulated width of H $\alpha$  or tabulated area, measured (i.e. projected) maximum area in square degrees, corrected maximum area in square degrees which equals measured area times secant  $h$  where  $h$  is the heliocentric angle, maximum effective line-width in H $\alpha$  expressed in Angstroms, and maximum intensity of H $\alpha$  expressed in per cent of the continuous spectrum. The following symbols are used in the table:

D = Greater than  
E = Less than

F = Approximately  
G = Plus

A final column lists provisionally the occurrence of simultaneous ionospheric effects as observed on selected field-strength recordings of distant high-frequency radio transmissions; a more nearly definitive list of these ionospheric effects, including particulars, appears in these reports after the lapse of a month (see below). All times are Universal Time (UT or GCT). Subflares (importance 1-) are listed by date, time of beginning and their heliographic coordinates. A graph presents intervals for which there were no patrols for flare observations from the observatories whose complete data are published in the table.

Ionospheric Effects -- SID (and GID--gradual ionospheric disturbances) may be detected in a number of ways: short wave fadeouts, enhancement of low frequency atmospherics, increases in cosmic absorption, and so forth. The table lists events that have been recognized on field-strength recordings of distant high-frequency radio transmissions.



Under a coordinated program, the staffs at the following ionospheric sounding stations contribute reports that are screened and synthesized at CRPL-Boulder: Puerto Rico, Ft. Belvoir, Va., and Anchorage, Alaska (CRPL Stations: PR, BE, AN); Huancaayo, Peru, and College, Alaska (CRPL-Associated Laboratories: HU, CO); and White Sands, N. Mex., Adak, Alaska, and Okinawa (U.S. Signal Corps Stations: WS, AD, OK). McMath-Hulbert Observatory (MC) also contributes such reports. In addition, reports are volunteered by RCA Communications Inc., Marconi Wireless, Netherlands Postal and Telecommunications Services, Swedish Telecommunications, and others; these usually specify times of SID and the radio paths involved.

In the coordinated program, the abnormal fades of field strength not obviously ascribable to other causes, are described as short wave fadeouts with the following further classification:

- S-SWF: sudden drop-out and gradual recovery
- Slow S-SWF: drop-out taking 5 to 15 minutes and gradual recovery
- G-SWF: gradual disturbance; fade irregular in both drop-out and recovery.

When there is agreement among the various reporting stations on the time (UT) of an event, it is accepted as a widespread phenomenon and listed in the table.

The degree of confidence in identifying the event, a subjective estimate, is reported by the stations and this is summarized in an index of certainty that the event is widespread, ranging from 1 (possible) to 5 (definite). The times given in the table for the event are from the report of a station (underlined in table) that identified it with high confidence. The criteria for the subjective importance rating assigned by each station on a scale of 1- to 3+ include amplitude of the fade, duration and confidence; greater consideration is given to reports on paths near the subsolar point in arriving at the summary importance rating given in the table.

Note: The tables of SID observed at Washington included in CRPL F-reports prior to F-135 were restricted to events classed here as S-SWF.

## IV SOLAR RADIO WAVES

### 2800 Mc Observations

The data on solar radio wave events made in Ottawa, Canada by the Radio and Electrical Engineering Division of the National Research Council (A. E. Covington) at 2800 Mc (10-cm emission) are presented. Near local noon (about 1700 UT) the sensitivity of the radiometer is determined and a mean flux for the whole day calculated. These values are given in a tabular form (see table I-1) in units of  $10^{-22}$  watts/M<sup>2</sup>/c/s. Burst phenomena are measured above this level and are given in terms especially suitable for the variations

observed on this frequency. The basis for the classifications is described by Covington - J.R. Astro. Soc. Can. 45, 49, 1951 and Dodson, Hedeman and Covington, Ap. J. 119, 541, 1954. A modification in terminology with a view to simplification has been introduced and consists essentially of the omission of the descriptive word "Single" from the "Single-Simple" and "Single-Complex" classes; in designating the "Single", "Single-Simple" and "Rise and Fall" bursts into a single classification designated as "Simple Bursts" with an appropriate type number; in the addition of the letter "f" to indicate that the burst deviates from the basic pattern by the presence of one or more small fluctuations in intensity; and by the addition of the letter "A" to indicate that the event has another smaller duration event superimposed upon it.

### Simple Burst

Any single burst which rises to one maximum and then decreases to the pre-burst level.

1 - Simple 1 -- Simple burst, type 1 (formerly "single"). Bursts of intensity less than 7 1/2 flux units and duration less than 7 1/2 minutes.

2 - Simple 2 -- Simple burst, type 2 (formerly "single-simple"). Bursts of impulsive nature with intensity greater than 7 1/2 flux units.

3 - Simple 3 -- Simple burst, type 3 (formerly "rise and fall"). Bursts of moderate intensity with duration greater than 7 1/2 minutes.

4 - Post-burst increase -- Postburst level is greater than the preburst level. The gradual return to normal flux may require as long as several hours.

5 - Absorption following burst (negative post).

6 - Complex -- (formerly "single-complex"). A single burst which shows two or more comparable maxima before the activity has declined to zero.

7 - Period of irregular activity or fluctuations -- Series of overlapping bursts of moderate intensity and duration.

8 - Group -- Series of single isolated bursts occurring in succession with intensity between the events equal to the level before and after the group.

9 - Precursor -- A small increase of intensity occurring before a larger increase.

## Great Burst


Infrequently occurring bursts of great intensity, often of complicated structure.

## Letter "A"

Indicates that this event has another event superimposed upon it.

## Letter "f"

Indicates that the basic form of the event is modified by secondary fluctuations.



## 200 Mc Observations

Data on solar radio waves made at Cornell University, Ithaca, N.Y. (Marshall Cohen) on 201.5 Mc are presented. All times are in Universal Time (UT or GCT). The antenna is linearly polarized and has a pattern appreciably broader than the solar disk. Flux is reported in units of  $10^{-22}$  watts/m<sup>2</sup>/cps and the tabulated numbers are twice the values observed in the one linear component.

Tables of flux and outstanding occurrences are given in general according to the systems used for the NBS 170 Mc and 450 Mc data.

## 170 Mc and 450 Mc Observations

Data on solar radio emission at the nominal frequencies of 170 Mc and 450 Mc recorded at the Gunbarrel Hill (Boulder) station of the National Bureau of Standards (R.S. Lawrence) are presented. The half width of the antenna lobe is appreciably greater than the solar disk. Polarization is not determined, but the dipole is oriented E-W. All times are in Universal Time (UT or GCT).

3-Hourly and Daily Flux Density and Variability -- Flux density is given in power units. These units are approximately  $10^{-22}$  watts meter<sup>-2</sup>(c/s)<sup>-1</sup> for both polarizations together. They will be subject to a correction factor when gain measurements of the antenna have been made. The median flux is measured for every one-hour period having at least thirty minutes of usable record and an applicable gain calibration. A three-hour value of flux is obtained by averaging the available one-hour medians (at least two required). A daily value of flux is obtained by averaging all available one-hour medians (at least four required). A dash indicates that insufficient measurements were made to meet the above requirements or that the records were not of usable quality. Flux values may be followed by the qualifying symbols D, S, and X defined subsequently.

The variability index, given for each three-hour interval, is on a scale 0 to 3 defined as follows:

0 - The instantaneous flux did not drop below one-half the median level or exceed twice the median level at any time.

1 - The instantaneous flux made from one to ten excursions



outside the range described above.

2 - The instantaneous flux made from ten to one hundred excursions outside the range described above.

3 - The instantaneous flux made more than one hundred excursions outside the range described above.

For the purpose of the variability index, an excursion whose maximum intensity is  $M$  times the median level is counted as  $M$  excursions. The variability index is omitted if measurements were made for less than one hour during the period. The variability for the day is the mean of the three-hourly values. The letter  $S$  follows variability indices which are in doubt because of atmospheric or local interference.

The observing periods are given in U. T. to the nearest 1/10 hour and they usually extend into the next Greenwich day.

Outstanding Occurrences -- A separate table lists the occurrences which are not adequately described by the three-hourly values of flux density and variability. Two classifications are given: (1) A system in general accord with that described and illustrated by Dodson, Hedeman, and Owren (Ap. J. 118, 169, 1953) and (2) the system described in the IGY Solar Activity Instruction Manual, prepared by the Radio Emission editor of the I.A.U. Quarterly Bulletin on Solar Activity.

In system (1) the occurrences are identified by numbers which do not necessarily indicate the magnitude of the event, as follows:

0 - Rise in base level -- A temporary increase in the continuum with duration of the order of tens of minutes to an hour.

1 - Series of bursts -- Bursts or groups of bursts, occurring intermittently over an interval of time of the order of minutes or hours. Such series of bursts are assigned as distinctive events only when they occur on a smooth record or show as a distinct change in the activity.

2 - Groups of bursts -- A cluster of bursts occurring in an interval of time of the order of minutes.

3 - Minor burst -- A burst of moderate or small amplitude, and duration of the order of one or two minutes.

4 - Minor burst and second part -- A double rise in flux in which the early rise is a minor burst.




6 - Noise storm -- A temporary increase in radiation characterized by numerous closely spaced bursts, by an increase in the continuum, or by both. Duration is of the order of hours or days.

7 - Noise storm begins -- The onset of a noise storm occurs at some time during the observing period.


8 - Major burst -- An outburst, or other burst of large amplitude and more than average duration. A major burst is usually complex, with a duration of the order of one to ten minutes.

9A, 9B, or 9 - Major burst and second part or large event without distinct first and second parts -- If there is a double rise in flux, the first part, a major burst, is listed as 9A and the second part as 9B. The second part may consist of a rise in base level, a group or series of bursts, a noise storm. A major increase in flux with duration greater than ten minutes but without distinct first and second parts, is listed simply as 9.


O-RISE IN BASE LEVEL




I - SERIES




2 - GROUP




3 - MINOR




4 - MINOR+




7-ONSET OF NOISE STORM




6-NOISE STORM IN PROGRESS



8 - MAJOR



9-MAJOR +



In system (2) combinations of the following letters are used to describe some distinctive characteristics of the recorded disturbances:

- S = simple rise and fall of intensity,
- C = complex variation of intensity,
- A = appears to be part of general activity,
- D = distinct from (i.e. apparently superimposed upon) the general background,
- M = multiple peaks separated by relatively long periods of quietness,
- F = multiple peaks separated by relatively short periods of quietness,
- E = sudden commencement or rise of activity.

Starting and maximum times are read to the nearest 1/10 minute if they are very definite and otherwise to the nearest minute. If the duration is less than five minutes, it is given to the nearest 1/10 minute; otherwise to the nearest minute (see also qualifying symbols below).

Maximum flux densities are given in units of  $10^{-22}$  watts meter<sup>-2</sup>(c/s)<sup>-1</sup>. The instantaneous maximum flux density is the highest peak in the disturbance measured above the sky level. The smoothed maximum flux density is the maximum value of a smooth curve drawn through the outstanding occurrence with a smoothing period of 20 to 50 percent of the total duration; it is measured above the estimated level in the absence of the disturbance. The intention is that (smoothed maximum) x (duration) should give a measure of the energy radiated in the disturbance.

A dash indicates missing or insignificant data. Observations are interrupted during the period from 26 to 29 minutes after each hour for calibrations. Observing periods are given in the Daily Data tables. The following qualifying symbols are used:

- B - Event in progress before observations began.
- D - Greater than.
- I - Event apparently continued during an interruption of the observations. The period of the interruption may be given in the remarks.
- N - See footnotes.
- X - Measurement is uncertain or doubtful.
- S - Measurement may be influenced by interference or atmospherics.

## V GEOMAGNETIC ACTIVITY INDICES

C, Kp, Ap, and Selected Quiet and Disturbed Days -- The data in the table are: (1) preliminary international character figures, C; (2) geomagnetic planetary three-hour range indices, Kp; (3) daily "equivalent amplitude," Ap; (4) magnetically selected quiet and disturbed days.

This table is made available by the Committee on Characterization of Magnetic Disturbance of IAGA, IUGG. The Meteorological Office, De Bilt, Holland collects the data from magnetic observatories distributed throughout the world, and compiles C and selected days. The Chairman of the Committee computes the planetary and equivalent amplitude indices. The same data are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm).

Kp is the mean standardized K-index from 12 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g. 5- is  $4\frac{2}{3}$ , 5o is  $5\frac{0}{3}$ , and 5+ is  $5\frac{1}{3}$ . This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of Kp has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948" of the Association of Terrestrial Magnetism and Electricity (IATME), International Union of Geodesy and Geophysics.

Ap is a daily index of magnetic activity on a linear scale rather than on the quasi-logarithmic scale of the K-indices. It is the average of the eight values of an intermediate 3-hourly index "ap," defined as one-half the average gamma range of the most disturbed of the three force components, in the three-hour interval at standard stations; in practice, ap is computed from the Kp for the 3-hour interval. The extreme range of the scale of Ap is 0 to 400. The method is described in IATME Bulletin No. 12h (for 1953) p. viii f. Values of Ap (like Kp and Cp) have been published for the Polar Year 1932/33 and for the years 1937 onwards.

The magnetically quiet and disturbed days are selected in accordance with the general outline in Terr. Mag. (predecessor to J. Geophys. Res.) 48, pp 219-227, December 1943. The method in current use calls for ranking the days of a month by their geomagnetic activity as determined from the following three criteria with equal weight: (1) the sum of the eight Kp's; (2) the sum of the squares of the eight Kp's; and (3) the greatest Kp.

Chart of Kp by Solar Rotations -- The graph of Kp by solar rotations is furnished through the courtesy of Dr. J. Bartels, Geophysikalisches Institute, Göttingen.

## VI RADIO PROPAGATION QUALITY INDICES

One can take as the definition of a radio propagation quality index: the measure of the efficiency of a medium-powered radio circuit operated under ideal conditions in all respects, except for the variable effect of the ionosphere on the propagation of the transmitted signal. The indices given here are derived from monitoring and circuit performance reports, and are the nearest practical approximation to the ideal index of propagation quality.

Quality indices are usually expressed on a scale that ranges from one to nine. Indices of four or less are generally taken to represent significant disturbance. (Note that for geomagnetic K-indices, disturbance is represented by higher numbers.) The adjectival equivalents of the integral quality indices are as follows:

1 = useless	4 = poor-to-fair	7 = good
2 = very poor	5 = fair	8 = very good
3 = poor	6 = fair-to-good	9 = excellent

CRPL forecasts are expressed on the same scale. The tables summarizing the outcome of forecasts include categories P-Perfect; S-Satisfactory; U-Unsatisfactory; F-Failure. The following conventions apply:

P - forecast quality equal to observed	U - forecast quality two or more grades different from observed when <u>both</u> forecast and observed were $> 5$ , or both $< 5$
S - forecast quality one grade different from observed	F - other times when forecast quality two or more grades different from observed

Full discussion of the reliability of forecasts requires consideration of many factors besides the over-simplified summary given.

The quality figures represent a consensus of experience with radio propagation conditions. Since they are based entirely on monitoring or traffic reports, the reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often



be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality for reasons such as multipath or interference. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

North Atlantic Radio Path -- The CRPL quality figures,  $Q_a$ , are compiled by the North Atlantic Radio Warning Service (NARWS), the CRPL forecasting center at Ft. Belvoir, Virginia, from radio traffic data for North Atlantic transmission paths closely approximating New York-to-London. These are reported to CRPL by the Canadian Defense Research Board, Canadian Broadcasting Corporation, and the following agencies of the U. S. Government:--Coast Guard, Navy, Army Signal Corps, U. S. Information Agency. Supplementing these data are CRPL monitoring, direction-finding observations and field-strength measurements of North Atlantic transmissions made at Belvoir.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the original scale. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year, with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. The 6-hourly quality figure is the mean of the reports available for that period.

The 6-hourly quality figures are given in this table to the nearest one-third of a unit, e.g. 5.0 is 5 and 0/3; 5- is 4 and 2/3; 5+ is 5 and 1/3. Other data included are:

(a) Whole-day radio quality indices, which are weighted averages of the four 6-hourly indices, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which seek to designate the days of significant disturbance or unusually quiet conditions.

(b) Short-term forecasts, issued every six hours by the North Atlantic Radio Warning Service. These are issued one hour before 00<sup>h</sup>, 06<sup>h</sup>, 12<sup>h</sup>, 18<sup>h</sup>, UT and are applicable to the period 1 to 7 hours ahead.

(c) Advance forecasts, issued twice weekly by the NARWS (CRPL-J reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole-day quality indices.



(d) Half-day averages of the geomagnetic K indices measured by the Fredericksburg Magnetic Observatory of the U. S. Coast and Geodetic Survey.

A chart compares the short-term forecasts with Qa-figures. A second chart compares the outcome of advance forecasts (1 to 3 or 4 days ahead) with a type of "blind" forecast. For the latter, the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

Ranges of useful frequencies on the North Atlantic radio path are shown in a series of diagrams, one for each day. The shaded area indicates the range of frequencies for which transmissions of quality 5 or greater were observed. The blacker the diagram, the quieter the day has been; a narrow strip indicates either high LUHF, low MUF, or both. These diagrams are based on data reported to CRPL by the German Post Office through the Fernmeldetechnischen Zentralamt, Darmstadt, Germany, being observations every one and a half hours of selected transmitters located in the eastern portion of North America. Since January 6, 1958 the transmitters monitored are restricted to those located north of 39° latitude. The magnetic activity index,  $A_F$ , from Fredericksburg, Va., is also given for each day.

Note: Beginning with data for September 1955, Qa has been determined from reports that are available within a few hours or at most within a few days, including for the first time, the CRPL observations. Therefore these are the indices by which the forecasters assess every day the conditions in the recent past. Over a period of several years, they have closely paralleled the former Qa indices which excluded CRPL observations and included three additional reports received after a considerable lag. Qa was first published to the nearest one-third of a unit at the same time.

North Pacific Radio Path -- The CRPL quality figures, Qp, are compiled by the North Pacific Radio Warning Service (NPRWS), the CRPL forecasting center at Anchorage, Alaska, from radio traffic data for moderately long transmission paths in the North Pacific equivalent to Seattle-to-Anchorage or Anchorage-to-Tokyo. These include reports to CRPL by the Alaska Communications System, Aeronautical Radio, Inc., U. S. Air Force and Civil Aeronautical Administration. In addition, there are CRPL monitoring, direction finder observations and field strength measurements of suitable transmissions.

The original reports are on various scales and for various time intervals. The observations for each 8 hours or 24 hour period are averaged on the original scale. This average is compared with reports for the same period in the preceding two months and expressed

as a deviation from the 3-month mean. The deviations are put on the 1 to 9 scale of quality which is assumed to have a standard deviation of 1.25 and a mean for the various periods as follows:

03-10 hours UT	5.33
11-18	5.33
19-02	6.00
00-24	5.67

The 8-hour and 24-hour indices  $Q_p$  are determined separately. Each index is a weighted mean where the CRPL observations have unit weight and the others are weighted by the correlation coefficient with the CRPL observations.

The table, analagous to that for  $Q_a$ , includes the 8-hourly quality figures; whole day quality figures; short-term forecasts issued by NPRWS three times daily at 02<sup>h</sup>, 10<sup>h</sup>, and 18<sup>h</sup> UT, applicable to the stated 8-hour periods; advance forecasts issued twice weekly by NPRWS (CRPL-Jp report); and half-day averages of geomagnetic K indices from Sitka.

The chart compares the outcome of advance forecasts, on the same basis as the similar chart for the North Atlantic Radio Path.

Note: Beginning with November 1956 the short-term forecast formerly made at 0900 UT was changed to 1000 UT. The North Pacific quality figures used for evaluation are now 8-hourly rather than 9-hourly.

## VII ALERT PERIODS AND SPECIAL WORLD INTERVALS


A table gives the Alert Periods and Special World Intervals (SWI) as designated by the IGY World Warning Agency at Ft. Belvoir, Va. For each day of the Alert or SWI are given the number of flares of importance two or greater reported promptly to the IGY World Warning Agency and the magnetic activity index  $A_{Be}$  observed at the IGY World Warning Agency.

# DAILY SOLAR INDICES

Feb. 1958	American Relative Sunspot Numbers $R_A'$
1	154
2	143
3	158
4	159
5	176
6	119
7	160
8	131
9	155
10	117
11	160
12	143
13	128
14	130
15	143
16	162
17	145
18	121
19	103
20	156
21	159
22	171
23	129
24	156
25	163
26	139
27	108
28	85
Mean:	141.9

Mar. 1958	Zürich Provisional Relative Sunspot Numbers $R_Z$	Daily Values Solar Flux at 2800 Mc, Ottawa, Canada Flux
1	109	195
2	90	209
3	140	223
4	185	232
5	203	233
6	215	251
7	220	256
8	187	251
9	177	255
10	181	242
11	168	235
12	156	232
13	145	238
14	158	227
15	165	217
16	155	214
17	164	208
18	162	210
19	155	220
20	154	232
21	156	224
22	163	266
23	187	268
24	204	274
25	180	258
26	194	284
27	226	302
28	292	295
29	302	332
30	338	344
31	342	338
Mean:	189.4	250.5







## CALCIUM PLAGE AND SUNSPOT REGIONS

MARCH 1958

CMP Mar. 1958	Lat	McMath Plage Number	Return of Region	Calcium Plage Data			Sunspot Data		
				CMP Values Area Int.		History, Age	CMP Values Area Count		History
01.6	S12	4442	*	(5000)	(3)	$l \text{ --- } l$ 5	70	3	$b \text{ --- } l$
06.2	N24	4443	4399	1800	3	$l \text{ --- } l$ 2	660	20	$b \text{ --- } l$
07.3	N32	4444	New	2500	3	$l \text{ --- } l$ 1	820	16	$l \text{ --- } l$
07.7	S15	4445	<del>4400</del>	8000	2.5	$l \text{ --- } l$ 5	2210	24	$l \text{ --- } l$
08.8	N21	4446	4405	4000	2.5	$l \text{ --- } l$ 3	70	2	$l \text{ --- } l$
09.7	S23	4447	4400	1300	2	$l \text{ --- } l$ 5			
10.5	N25	4450	4411	1000	1.5	$l \text{ --- } l$ 3			
10.6	N10	4448	4412	1000	2	$l \text{ --- } l$ 2			
11.9	N14	4449	4410	8300	3	$l \text{ --- } l$ 3	1060	12	$l \text{ --- } l$
12.2	S12	**	New	600	2.5	$b \text{ --- } l$ 1			
12.5	N41	4454	New	200	2	$b \text{ --- } d$ 1			
13.4	N25	4452	4410	1000	2	$l \text{ --- } l$ 3	50	2	$b \text{ --- } d$
13.5	N14	4453	New	2300	3	$l \text{ --- } l$ 1	850	6	$l \text{ --- } l$
15.6	S26	4455	4414	500	1.5	$l \text{ --- } d$ 8			
16.7	N08	4462	New	300	1.5	$b \text{ --- } d$ 1			
17.6	N27	4468	+	400	1	$b \text{ --- } l$ 1			
17.7	S20	4457	4422	900	1.5	$l \text{ --- } l$ 5			
18.2	N13	4456	New	6200	3	$l \text{ --- } l$ 1	880	20	$l \text{ --- } l$
19.1	S04	4472	New	500	1	$b \text{ --- } l$ 1			
20.5	N36	4460	New	1500	3	$l \text{ --- } l$ 1	200	7	$l \text{ --- } l$
20.7	N20	4461	4424	400	1	$l \text{ --- } d$ 4	(10)	(1)	$l \text{ --- } d$
20.8	N08	4463	4430	400	1.5	$l \text{ --- } d$ 2			
21.0	S18	4459	++	2200	2.5	$l \text{ --- } l$ 3	140	1	$l \text{ --- } l$
22.3	S06	4466	New	300	1	$l \text{ --- } d$ 1			
22.4	S20	4473	4427	1100	1	$l \text{ --- } l$ 3			
22.6	N22	4465	New	4000	3	$l \text{ --- } l$ 1	1440	20	$l \text{ --- } l$
23.7	N11	4467	New	1200	2.5	$l \text{ --- } l$ 1	170	9	$l \text{ --- } l$
24.2	S18	4470	4428	2000	2	$l \text{ --- } l$ 4			
24.9	N26	4469	New	2200	3	$l \text{ --- } l$ 1	600	27	$l \text{ --- } l$
26.6	N18	4474	New	2300	3.5	$l \text{ --- } l$ 1	200	2	$l \text{ --- } l$
28.2	S24	4479	New	800	2	$b \text{ --- } l$ 1			
28.3	N20	4475	New	1600	3	$l \text{ --- } l$ 1	190	10	$l \text{ --- } d$
28.5	S12	4476	New	12,000	3.5	$l \text{ --- } l$ 1	2090	37	$l \text{ --- } l$
28.9	N31	4477	4435	500	2.5	$l \text{ --- } l$ 2			
29.1	N09	4482	New	700	2.5	$b \text{ --- } l$ 1	80	4	$b \text{ --- } d$
30.2	S22	4478	4438	2400	2.5	$l \text{ --- } l$ 2	1720	11	$l \text{ --- } l$
30.5	N06	4491	New	(200)	(1.5)	$b \text{ --- } d$ 1			

COMMERCE - STANDARDS - BOULDER

\* 4393 and 4394.

\*\* 4451 (4458).

+ In position of 4417.

++ 4431 and 4426.

CORONAL LINE EMISSION INDICES

MARCH 1958

CMP Mar. 1958	North East Quadrant (observed 7 days earlier)				South East Quadrant (observed 7 days earlier)				South West Quadrant (observed 7 days later)				North West Quadrant (observed 7 days later)			
	G6	G1	R6	R1	G6	G1	R6	R1	G6	G1	R6	R1	G6	G1	R6	R1
1	117	147	42	109	106	140	17	30	83	108	x	x	87	116	x	x
2	81	100	x	x	76	93	x	x	x	x	x	x	x	x	x	x
3	x	x	x	x	x	x	x	x	45	59	20	32	79	104	19	42
4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
6	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
7	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
8	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
9	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
10	147	205	42	68	93	118	21	32	79	96	x	x	176	222	x	x
11	111	136	x	x	68	84	x	x	x	x	x	x	x	x	x	x
12	123	178	30	50	53	70	12	14	x	x	x	x	x	x	x	x
13	103	128	x	x	44	52	x	x	x	x	x	x	x	x	x	x
14	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
15	81	96	x	x	37	44	x	x	x	x	x	x	x	x	x	x
16	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
17	123	169	18	32	63	84	23	42	x	x	x	x	x	x	x	x
18	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
19	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
20	x	x	x	x	x	x	x	x	106a	139a	57a	150a	102a	108a	10a	12a
21	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
22	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
23	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
24	125	238	20	30	135	216	x	x	84	92	18	36	129	194	31	42
25	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
26	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
27	x	x	x	x	x	x	x	x	174	260	x	x	169	230	38	90
28	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
29	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
30	x	x	x	x	x	x	x	x	97	118	x	x	59	72	x	x
31	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x

\* = yellow line observed.  
a = index computed from low weight data.  
x = no observations.

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION			DURA- TION — MINUTES	IM- PORT- ANCE	OBS. COND.	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT
		START	END	APPROX. LAT.	APPROX. LONG.	MAX. PHASE				TIME — UT	MCAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	
WENDEL	01 0911	0938		S09 W48	4436		27	36			34.00		S-SWF
SIMEIZ	01 0912	1007		S10 W46	4436		55	3				4.10	S-SWF
SCHAUINS	01 0916	0950		S12 W42	4436		34	3					
ONOREJOV	01 0918	0922		S13 W45	4436		4	16	0918				
WENDEL	01 0925	0938		S16 W80	4428		13	1			3.00	5.00	
ONOREJOV	01 0929	0933		S17 E78	4445		4	16	0929				
WENDEL	01 1007	1036		S19 W80	4428		29	1			3.00		
ZURICH	01 1346	1348		S16 E77	4445		2	1	1346		2.00		
ZURICH	01 1408	1412		N15 W31	4434		4	1	1408		1.00		
AROSA	01 1527	1539		S12 W56	4436		12	1					
UCCLE	02 0820	E 0838	0	N32 W22	4435		18	0	0820	3.40			
CAPRI S	03 1008	E 1042	0	S14 E59	4445		34	0	1026	5.00	10.00		
NIZAMIAH	03 1013	E 1048		S19 E60	4445		35	0	1018	9.72	18.93	3.60	
ARCETRI	03 1015	E 1100		S12 E58	4445		45	0					
R O HERST	03 1016	E 1115	1022	S17 E62	4445		59	0	1021	7.10	14.80	6.70	250
STOCKHOLM	03 1031	E 1036	0	S17 E63	4445		5	0					
KODAIKUN	03 1040	E 1407	0	S15 E54	4445		121	0	1040	2.00	4.40	2.00	
CAPRI S	03 1206	E 1411	0	S18 E65	4445		131	0	1209	1.47	3.11		72
USNRL	03 1812	E 1832	D	S20 E61	4445		20	0	1825	2.50			
CLIMAX	03 1812	E 1832	D	S21 E70	4445		16	1	2346	1.90	2.40		
HAWAII	03 2340	E 2356		S14 E40	4445		50	1					
USNRL	04 1320	E 1410	1322	S25 W45	4445		26	0	1322	1.13	1.59	1.00	106
USNRL	04 1724	E 1750		S17 E40	4445		17	0	1725	2.60	3.30		
HUANCAYO	04 2208	E 2225	D	S17 E33	4445		15	0					
ZURICH	05 0908	E 0923		S17 E29	4445		15	0	908		4.00		
ONOREJOV	05 0956	E 1018		S22 E28	4445		22	0	1007		3.00	2.30	
ZURICH	05 1001	E 1018		S19 E25	4445		17	1	1001		4.00		
WENDEL	05 1004	E 1135		S21 E27	4445		91	0					
ONOREJOV	05 1021	E 1037	1022	S22 E30	4445		16	1	1022			2.50	
AROSA	05 1024	E 1035		S21 E28	4445		11	1			6.00		
WENDEL	05 1307	E 1333	0	S18 E29	4445		26	0			4.00		
ZURICH	05 1313	E 1335		S17 E26	4445		22	1	1313				
AROSA	05 1315	E 1330		S19 E28	4445		15	0					
ONOREJOV	05 1322	E 1342	0	S18 E28	4445		20	0	1327	1.92	3.19	2.20	
USNRL	05 1413	E 1411	0	S20 W54	4442		266	0	1414	2.94	3.40	1.00	71
USNRL	05 1635	E 1721		S20 E28	4445		26	1	1642	4.52	11.03		
GOTTAWA	05 1647	E 1710	1656	N24 W58	4435		33	0	1656	1.70	5.45	102	
USNRL	05 1702	E 1720		N32 W65	4435		21	2	1649	1.70	5.45		
USNRL	05 1649	E 1720		N32 W65	4435		21	2	1649	1.70	5.45		
USNRL	05 1702	E 1720		N32 W65	4435		21	2	1703	2.50	2.94	1.00	63
MT WILSON	05 1720	E 1728		S14 E30	4445		8	1	1703				
USNRL	05 1721	E 1730		S15 E28	4445		9	1	1723	.90	1.03	108	
GOTTAWA	05 1721	E 1740		S13 E29	4445		19	1	1724	3.36	3.86		
SAC PEAK	05 1820	E 1842		S15 E29	4445		12	1					
USNRL	05 1902	E 2100	0	S08 W57	4442		118	0	1902	2.60	4.54	1.00	18
USNRL	05 2047	E 2100	0	S19 E23	4445		13	0	2051	1.13	1.22		107
CLIMAX	05 2311	E 2341	2328	S09 E15	4445		30	0	2328	4.90			
ARCETRI	06 0820	E 0844	0	S20 W63	4441		24	0	0820	1.30	2.80		
ARCETRI	06 0846	E 0931	E	S17 E19	4445		16		0846	2.50	2.70		
UCCLE	06 0931	E 0931	E	N24 W50	4443				0846			PAGE	1

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION		DUR.	IM.	OBS. COND.	TIME	MEASUREMENTS		PROVISIONAL IONOSPHERIC EFFECT
		START	END	APPROX. LAT.	APPROX. LONG.	MINUTES	FOR. TRACE			AREA Sq. Deg.	CONTR. AREA Sq. Deg.	
CAPRI S	06 1246 E	1330 D		S14 W61	4442	44 0	1	3	1246	1.00	22.0	
AROSA	06 1511	1519		S20 E12	4445	8	1					
AROSA	06 1538	1547		S20 E12	4445	9	1		2023	5.00		
CLIMAX	06 2018	2030 0		S22 E13	4445	12 0	1	1	2050	1.02	2.78	78
USNRL	06 2050	2107 0		S08 W71	4442	17 D	1					
MITAKA	07 0525 E	0537 D		N22 E23	4446	12 0	16	1	0530	5.67	7.06	149
UCCLE	07 0828	0900	0829	N11 E85	4449	32	16	2	0829	2.00	4.00	
UCCLE	07 1024	1156		N11 E86	4449	86	2	2	1024	1.50	3.40	
CAPRI S	07 1030	1156		N14 E72	4449	40 0	2	2	1054	2.00	6.80	
R O HERST	07 1043 E	1050 0	1043 E	N10 E72	4449	7 0	1	1	1043	.60	2.40	
STOCKHOLM	07 1100 E	1140 0		N10 E70	4449	40 0	2	1				
NEGERHORST	07 1105 E	1200		N10 E72	4449	55 0	2	1				
MEUDON	07 1110 E			N12 E68	4449		1					
UCCLE	07 1203 E	1212 D		N11 E86	4449	9 D	3	2	1115	6.00	13.00	
ONDREJOV	07 1239 E	1319		N08 E69	4449	40 D	16	2	1208	1.99	5.71	65
USNRL	07 1335 E	1307		N06 E70	4449	32 0	16	1	1242	3.56	3.64	116
USNRL	07 1326	1336		S18 E01	4445	12	1	3	1246	1.57	1.61	117
USNRL	07 1453	1536		S18 W01	4445	43	1	3	1330	1.24	3.40	68
USNRL	07 1815	1845		N08 E85	4449	30	1	3	1455	1.81	1.83	103
HUANCAYO	07 1947 E	2024	2001	S18 W03	4445	37 0	1	1	1815			
MITAKA	07 2033	2110 0	2033	S15 W80	4442	37 0	1	1				
MITAKA	08 0530 E	0542		N25 W22	4443	12 0	1	1	0534	1.84	2.26	96
MITAKA	08 0557 E	0631 0		N12 E56	4449	25 0	1	1	0602	3.30	3.92	120
ATHENS	08 0851	0935		S21 W31	4445	42	16	4		.40	5.40	
ATHENS	08 0914	0922		S11 W20	4442	7	2	4				
AROSA	08 1051	1059		N32 W08	4445	8	1					
AROSA	08 1720	1755	1727	S18 W14	4445	35	2	2	6.60			30
SAC PEAK	08 2158	2215 D		N33 W17	4444	17 D	1	1	2208	1.60	3.00	
HAWAII	08 2159	2249		N12 E52	4449	50	1	1	2209	2.20		
CLIMAX	08 2200	2217	2202	N05 E54	4449	17	1	2	2340	2.50		20
SAC PEAK	08 2336	2348 0	2340	N32 W20	4444	12 0	1	1	2346	2.70		
CLIMAX	08 2352 D	2344		N17 E57	4449	16 0	1	1	2337	7.57	4.90	
HAWAII	08 2337 E	2356 0	2337	N09 E48	4449	21 0	26	1		16.20		
MITAKA	09 0210 E	0221		N24 W35	4443	11 0	16	1	0216	7.57	11.50	122
MITAKA	09 0443 E	0514		N32 W23	4444	31 0	1	1	0453	1.84	2.11	113
MITAKA	09 0452 E	0505 0		S15 W19	4445	13 0	1	1	0453	1.84	2.47	100
MITAKA	09 0421 E	0428 D		N10 E42	4445	7 0	16	1	0621	3.71	5.68	115
WENDEL	09 0757 E	0804		N19 E45	4449	27 0	16	2		6.00		
UCCLE	09 0900	0927	0920	N32 W58	4449	55	1	2	0920	3.40	3.00	
UCCLE	09 0902	0937		N12 E50	4453	13	16	1	0922	2.20	3.00	
UCCLE	09 0933	0946	0936	N22 W50	4443	42	2	1	0938	3.40	5.40	
UCCLE	09 0946	1032	0956	N14 E20	4453	42	2	4	0956	4.50	5.40	
UCCLE	09 0955	1025	1004	N34 W34	4444	30	1	4	1004	2.20	2.00	
UCCLE	09 1031	1038	1033	N12 E50	4449	5 0	16	3	1033	3.40	5.10	
ONDREJOV	09 1033 E			N09 E44	4449	5 0	16	3	1034	4.00	4.40	
UCCLE	09 1056	1104	1059	N12 E45	4449	7 0	16	2	1059			
ONDREJOV	09 1057 E	1104		N11 E40	4449	7 0	16	3	1059			
ONDREJOV	09 1120 E	1135	1122	N12 E51	4453	15 0	16	3	1122			2

COMETICE - STANBARD - BOULDER

SOLAR FLARES  
MARCH 1958

OBSERVATORY	DATE MAR. 1958	OBSERVED TIME		LOCATION		DURA- TION MINUTES	DE- FOR- TANCE	OBS. COND.	TIME — U T	MEASUREMENTS		MAX. WIDTH INT. %	PROVISIONAL IONOSPHERIC EFFECT
		START	END	AFFIX.	M-MATR LAT. LONG. REGION					N-CAL AREA Sq. Deg.	COIL. AREA Sq. Deg.		
ONDREJOV	09	1145 E	1154 O		N13 E49 4453	9 D	1	3	1345	1.58	2.76	80	S-SMF
USNRL	09	1217 E	1341		N15 E55 4453	84 D	1	2	1341	2.30		16	
SAC PEAK	09	1447	1507		S21 W35 4445	20	1	2	1455	1.91	2.37	25	
OTTAWA	09	1454	1500		S19 W35 4445	6	1	2	1455	6.80			
SAC PEAK	09	1540	1740 U		N34 W32 4444	120 D	2	2	1648	6.38	10.00		S-SMF
OTTAWA	09	1542	1709		N32 W32 4444	87 D	2	2	1645	2.71	4.33	1.00	S-SMF
USNRL	09	1543	1642 D		N35 W30 4444	59 D	16	2	1645	2.50		18	S-SMF
SAC PEAK	09	1957	2030		N11 E37 4449	33	1	2	0013	1.84	2.98	2.45	134
MITAKA	10	0007 E	0031		N32 W38 4444	24 D	1	2	0013	5.67	6.73	1.91	87
MITAKA	10	0111 E	0131		N33 W39 4444	20 D	1	1	0134	1.84	1.81	1.00	100
MITAKA	10	0134 E	0140 O		N33 W39 4444	6 D	1	1	0154	5.67	2.87	1.99	128
MITAKA	10	0148 E	0155		N33 W33 4444	7 D	1	1	0212	3.90	4.40	2.22	149
RODAIKNL	10	0208	0241 O		N11 E35 4453	33 D	16	3	0213	1.80	2.00	154	Slow S-SMF G-SMF
ATHENS	10	0210 E	0217 D		N34 W36 4444	7 D	1	3	1316	1.39	2.45	4.50	S-SMF G-SMF
CAPRI S	10	0709 E	0743		N08 E24 4449	34 D	16	1	1411	1.16	2.12		S-SMF G-SMF
NEDERHORST	10	1316 E	1350 O		N37 W35 4444	10	2	1	1412	2.00		1.00	104
OTTAWA	10	1335	1345		N31 W41 4444	9 D	1	2	2050	3.06	4.50	20	G-SMF
ONDREJOV	10	1408	1420 D		N32 W42 4444	8	1	3	0034	3.30	3.50		G-SMF
OTTAWA	10	1710	1718		N34 W46 4444	11	1	3	0833	4.00	6.00		G-SMF
CLIMAX	10	1955	2006		S11 W50 4445	54	2	2	0902	1.50	3.00		G-SMF
CLIMAX	10	2024	2128		S11 W50 4445	26 D	2	2	0923	2.00	2.20	2.60	
HAWAII	10	2026	2032		S13 W48 4445	6 D	16	2	0923	2.78	5.93	1.35	
USNRL	10	2028	2128 D		S10 W50 4445	1 D	1	2	2358	7.63	7.95	2.29	227
SAC PEAK	10	2040 E	2041 O		N11 E02 4449	12 D	1	1	0044	4.89	4.70	1.90	120
HAWAII	11	0030 E	0042 D		N29 W85 4443	5	0	1	0044	4.70	4.89	1.90	120
UCCLE	11	0802	0906		N10 W04 4449	45 D	1	2	0044	4.70	4.89	1.90	120
UCCLE	11	0902	0927 D		N14 E18 4453	16 D	16	4	0944	3.40	4.70		
ONDREJOV	11	0918	0929 O		N15 E70 4456	4	16	4	0944	3.40	4.70		
MITAKA	11	2352 E	2400		N12 E15 4453	16 D	2	4	0945	5.60	5.10		
MITAKA	12	0024	0233		N15 E03 4449	12 D	1	4	1113	3.40	5.10		
MITAKA	12	0027 E	0032 D		N15 E70 4456	32 D	16	4	1131	3.40	5.10		
MITAKA	12	0043 E	0128 D		N32 W80 4444	4	16	4	1158	5.60	8.90		
UCCLE	12	0943 E	0947		N23 W50 4446	25 D	2	4	1137	4.50	5.60		
UCCLE	12	1127	1159		N12 E15 4453	22 D	1	4	1148	3.40	5.10		
UCCLE	12	1128	1200		S20 W69 4445	55 D	26	2	1441	2.04	5.50	2.00	162
UCCLE	12	1132	1137		N10 E78 4455	16 D	1	1	0034	4.89	1.22	3.23	
USNRL	12	1436 E	1531		N00 E90 4461	12 D	1	1	0034	1.00		2.40	
MCNATH	12	1440 E			N19 W01 4453	44	1	1	2220	4.00	4.50		
MITAKA	13	0018 E	0034 O		N10 W08 4453	34 D	1	1	0251	2.78	2.89	2.10	107
MITAKA	13	0020	0032 D									3	
ONDREJOV	13	1053 E	1058										
HAWAII	13	2216	2300										
MITAKA	14	0246 E	0320										



# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME			LOCATION			DUR- TION — MINUTES	IM- POR- TANCE	OBS. COND.	MEASUREMENTS				PROVISIONAL LONGPERIOD EFFECT	
		START	END	MAX. PHASE	APPROX.		PLACE REGION				TIME — UT	MEAS. AREA Sq. Deg.	CORE AREA Sq. Deg.	MAX WIDTH H <sub>0</sub>		MAX INT. %.
					LAT.	LONG.										
NIZAMIAH MITAKA AROSA AROSA ZURICH ZURICH AROSA UCCLE AROSA UCCLE UCCLE UCCLE ZURICH R O HERST ONOREJOV	14	0442	0458	0446	N09 W11	4453	16	1	3	0446	3.04	3.22	1.60	1.84	131	Slow S-SNF
	14	0456 E	0512	0500	N11 W12	4453	16	0	1	0502	11.40	11.80	1.60	1.84	131	
	14	0846	0910		N08 W18	4449	24	1								
	14	0920	0941		N08 W19	4449	21	2								
	14	0948 E	1100	0	N11 W13	4453	72	2	1	948		7.00				
	14	1000 E	1029		N10 W25	4449	29	1	4	1018						
	14	1013	1036		N06 W26	4449	23	1	3							
	14	1013	1100	1021	N10 W15	4453	47	16	4	1021	4.50					
	14	1023	1043		N08 W20	4449	20	1	3	1143	2.20					
	14	1136	1200	1143	N10 W15	4453	24	1	2	1320		6.00				
	14	1301 E	1325 D		N11 W15	4453	24	0	2	1507	4.50					
	14	1504 E	1530	1507	N11 W90	4446	26	0	1	1507						
	14	1518 E	1541	1541	S23 W80	4445	23	0	1	1518			2.20			
	15	0822	0850	0831	N12 W25	4453	28	16	4	0831	4.00	4.40				
15	0834	0841	0835	N37 E58	4460	7	1	4	0835	1.00	2.00					
15	1010 E	1028		N10 W25	4453	18	0	2	1010	3.00	3.00					
15	1030	1033	1030	N12 W25	4453	3	1	4	1030	2.00	2.00					
15	1208	1235	1219	N36 E65	4460	27	1	2								
15	1342	1347	1344	N13 W27	4453	5	1	2	1344	3.50	3.70					
15	1345	1346		N10 W25	4453	1	1	2	1345	1.00	1.00					
15	1541 E	1547		N11 W27	4453	6	0	1	1541			2.20				
15	1542 E	1547	1542	N13 W27	4453	5	1	2	1542	2.00	2.30					
16	0839	0850	0841	N13 W41	4453	11	2	4	0841	4.50	6.30					
16	0840	0848	0842	N14 E32	4456	8	16	4	0842	3.60	4.50					
16	0840 E	0905		N17 E56	4461	25	0	2	840							
16	0851	0930	0918	N15 E30	4456	39	16	4	0918	4.50	5.60					
16	0857	0902		N13 E29	4456	5	1	2	857							
16	0950	0955		N27 W90	4446	5	1	1								
16	1358	1412		N12 E25	4456	14	1	1								
17	0438	0453	0444	N10 E18	4456	15	16	3	0444	4.86	5.34					
17	0804 E	0817		N15 W71	4449	13	0	2	804		2.00					
17	0804 E	0909	0	N20 E71	4465	65	0	2	804		6.00					
17	0819 E	0825		N25 E85	4465	6	0	3								
17	0938 E	0948	0	N23 E70	4465	10	0	1	0938	4.60	2.10					
17	1006 E	1116	0	N23 E77	4465	70	0	1			3.00					
17	1008	1054		N10 E15	4456	46	16	1			7.00					
17	1008	1120		N10 E16	4456	72	2	4	1025	6.00	6.00					
17	1011	1144	1025	N11 E15	4456	33	16	3	1030	3.65	4.00					
17	1027 E	1043	1030	N10 E18	4456	16	0	1			4.00					
17	1153 E	1220		N23 E77	4465	27	0	1			5.00					
17	1215	1338		N35 E32	4460	23	0	1			3.00					
17	1314	1336	0	N23 E73	4463	22	0	1			3.50					
17	1515 E	1535	1515	N22 E68	4462	20	0	3	1515	1.50	1.50					
17	1516	1559	1518	N23 E70	4462	43	1	2	1518	1.81	1.81					
17	1516 E	1550		N13 E10	4456	34	0	1	1518	1.02	1.09					
18	0804	1125	0835	N22 E60	4465	20	1	4	0835	6.80	6.80					
18	0805	0824	0	N18 E35	4465	19	0	4	0810	3.40	4.20					
18	0810	0834	0825	N09 W70	4453	24	1	4	0825	2.80	2.80					
18	0922	0930	0927	N13 W80	4453	8	1	4	0927	2.20	2.20					
													PAGE		4	

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED		LOCATION		DURA- TION — MINUTES	IM- PAC- TANCE	OBS. COND.	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT
		START	END	APPROX. LAT.	APPROX. LONG.				TIME — UT	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	
UCCLLE	18	1145	1201 D	N22 E60	4465	16 D	16	4	1155	3.40		G-SNF
HUANCAYD	18	1905 E	1955 D	N22 E53	4465	50 D	16	2				
MITAKA	19	0300 E	0332 D	N23 E48	4465	32 D	16	1	0321	13.40	25.90	120
ATHENS	19	0730 E	0750 D	N24 E48	4465	20 D	1	3		.60	2.40	
UCCLLE	19	0956 E		N24 E48	4465		1	1				
UCCLLE	19	0956 E		N24 E48	4465		1	1				
MEUDON	19	1022	1140	N14 W13	4456	78	1	1			4.00	
UCCLLE	19	1027 E	1032 D	N12 W13	4456	5 D	1	1	1028	2.10	2.10	
CAPRI S	19	1045 E	1125 D	N11 W11	4456	40 D	1	2	1050	2.40	2.50	
ARDSA	19	1114 E	1120	N24 E39	4465	6 D	1	1				
UCCLLE	19	1114 E	1140	N12 W15	4456	26 D	2	1				
UCCLLE	19	1117 E	1135 D	N12 W11	4456	18 D	16	1	1125	4.50	4.50	
MT WILSON	19	1910	1931	N14 W18	4456	21	1	1				
MT WILSON	19	2137	2152	N22 E40	4465	15	1					
ONDRE JUV	20	0656 E	0706	N24 E56	4469	10 D	1	2	0700	7.90		S-SNF
UCCLLE	20	0723	0802	N22 E35	4469	59	2	4	0752	5.10	5.40	
UCCLLE	20	0728	0851	N21 E32	4465	3	2	3	0759	5.80	5.40	
UCCLLE	20	0728	0737	N21 E32	4469	152	1	4	0726	3.20	3.40	
UCCLLE	20	0825	0848	N18 W58	4456	11	16	3	0730	3.20	5.80	
UCCLLE	20	0828	0900	N22 E58	4453	32	2	4	0828	5.60		
UCCLLE	20	0850	0920	N20 E52	4465	32	1	4	0838	2.20		
UCCLLE	20	0905	0955	N20 E52	4465	50	1	4	0855	6.20		
UCCLLE	20	0907	0957	N23 E55	4465	50	16	4	0920	3.40		
UCCLLE	20	0959	1030	N23 E43	4469	31	16	2				
MEUDON	20	1127 E	1144 D	N23 E35	4465	17	2	2				
NEDEHORST	20	1259	1330	N25 E28	4465	31	2					
MEUDON	20	1305 E	1320	N23 E29	4465	15	2					
MEUDON	20	1314	1344	N18 W30	4452	30	1					
UCCLLE	20	1324 E	1356	N23 E35	4465	32 D	16	3		4.50		
UCCLLE	20	1324 E	1356	N20 W32	4456	32 D	16	2		4.10		24
SAC PEAK	20	1445	1545	N22 E26	4445	60	16	2				
SAC PEAK	20	1452	1550	N25 E28	4465	58	16	2				
UCCLLE	20	1455 E	1552 D	N24 E29	4465	42 D	2	1	1517	5.00	6.70	S-SNF
UCCLLE	20	1510 E	1538 D	N23 E29	4465	3 D	16	2		4.50		
UCCLLE	20	1535 E	1930	N23 E33	4465	65	1	2		2.70		15
SAC PEAK	20	1825	2130	N23 E25	4465	65	2	2		3.80		24
SAC PEAK	20	2025	2130	N22 E23	4465	65	2	2		3.80		16
SAC PEAK	20	2230	2315 D	N22 E22	4465	45 D	1	2				
UCCLLE	21	1019	1050	N17 E21	4465	31	2	2	1022	5.50	5.50	
KODAIKUN	21	1021 E	1040 D	N20 E20	4465	19 D	16	2	1030	3.90	4.00	S-SNF
R D EDIN	21	1027 E	1044	N22 E17	4465	17 D	2	1	1031	6.00	7.20	154
NIZANIAH	21	1040 E		N E	4465		1	1		2.00	1.80	
CAPRI S	21	1613	1432 D	N23 E14	4465	19 D	1	1	1415	2.00	2.20	S-SNF
SAC PEAK	21	1850	1937 D	N22 E12	4465	47 D	16	3		4.10		24
HAWAII	21	1852	1920	N23 E07	4465	28	16	3	1902	4.10	4.80	Slow S-SNF
UCCLLE	21	1900 E	1925	N22 E12	4465	25 D	2					
UCCLLE	22	0925 E	1129	N18 W60	4456	124 D	1	2	0944	2.20	2.60	
UCCLLE	22	1123 E	1155	N22 E05	4465	32 D	16	2	1130	4.50		G-SNF
MEUDON	22	1125	1150	N20 E07	4465	25	1					

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE M.D.Y.	OBSERVED UNIVERSAL TIME		LOCATION		DURA- TION MINUTES	IN- FOR- TANCE	OBS. COND.	MEASUREMENTS				PROVINCIAL LONGITUDINAL EFFECT		
		START	END	APPROX. LAT.	MER. DIST. REGION				PLAGE	TIME U T	MEAS. AREA Sq. Deg.	CORR. Sq. Deg.		MAX WIDTH In	MAX INT. %
MEUDON	22 1958	1147	1201	S20 E90	4478	14	1								
MEUDON	22 1210	1230		S12 E90	4476	20	1								
WENDEL	23 0844	0857		N13 W68	4456	13	1								
UCCLE	23 0950	1200	1005	S15 E80	4476	130	0	4	1005	29.00	3.00				
CAPRI S	23 0950	1211		S14 E75	4476	141	36		1028	8.00	64.00				
WENDEL	23 0950	1358	0	S14 E76	4476	248	0	3			25.00				
MOSCOW	23 0951	1102	0	S25 E67	4476	71	0								
SCHAUINS	23 0957	1137	0	S13 E78	4476	100	0								
ZURICH	23 1005	1055		S12 E76	4476	50	3	2							S-SHF
MEUDON	23 1009	E 1119		S09 E80	4476	70	3								
UCCLE	23 1012	1200	0	S12 E85	4476	108	D	4	1018	5.47	10.61	3.60			
NIZAMIAH	23 1016	E 1027	D	S18 E60	4476	11	D	1							
WENDEL	23 1056	E 1329		S16 E73	4476	153	D								
AROSA	23 1105	E 1200	D	S14 E83	4476	55	D	2							
ZURICH	23 1215	E 1319	0	S14 E73	4476	64	0	1	1215	10.00	10.00				
USNRL	23 1216	E 1227		S13 E75	4476	11	0	2	1218	2.04	6.51	2.00	85		
ZURICH	23 1227	E 1252		N23 W12	4465	25	1	1	1227		2.00				
NEDERHORST	23 1258	E 1415		S20 E88	4478	77	D								
NEDERHORST	23 1258	E 1415		S12 E90	4476	77	D								
HAWAII	23 1826	E 1838		N12 W85	4456	12	1	2	1828	1.50	3.10				S-SHF
HAWAII	24 0048	E 0052	D	N23 W01	4469	4	D	1	0048	2.10	2.40				
AROSA	24 0655	E 0703		N17 W26	4465	8	D								
UCCLE	24 0713	E 0726		N17 W26	4465	13	1								
UCCLE	24 0717	E 0731		N20 W26	4465	19	1	4	0720	2.20	3.10				
UCCLE	24 0731	E 0740	0734	S17 E72	4476	11	16	4	0734	2.20	4.80				
UCCLE	24 0745	E 0756	0746	S17 E72	4476	35	1	4	0744	3.40	3.40				
UCCLE	24 0748	E 0823	0759	N21 W27	4465	38	1	4	0759	3.40					
WENDEL	24 0749	E 0827		N20 W25	4465	35	1								
ZURICH	24 0758	E 0825		N20 W26	4465	27	D	2	758	3.00	3.00				S-SHF
UCCLE	24 0758	E 0803		S22 E88	4478	7	2	4	758	6.80	4.00				
AROSA	24 0758	E 0834		S16 E65	4476	36	0	2							
ZURICH	24 0800	E 0817		S17 W27	4465	17	0	4							
WENDEL	24 0810	E 0831		S17 E72	4476	21	1	4	0822	2.20	3.00				
UCCLE	24 0811	E 0830		S17 E72	4476	15	1								
AROSA	24 0813	E 0828		S16 E66	4476	15	1	4		2.20	3.00				
UCCLE	24 0917	E 0925		S17 E72	4476	8	1	4		2.20	3.00				
ZURICH	24 0953	E 1020	1014	S16 E64	4476	27	1	2	953						
WENDEL	24 0954	E 1014	0	S17 E64	4476	20	D								
UCCLE	24 0955	E 1007		S17 E72	4476	12	1	4	1047	2.20	3.00				
UCCLE	24 1045	E 1056		S22 E88	4478	11	2	2	1052	3.40	6.80				
OMOREJOV	24 1051	E 1055		S21 E81	4478	4	D	2	1114		2.20				
OMOREJOV	24 1110	E 1119	0	S16 E57	4476	9	D	2			2.50				
AROSA	24 1110	E 1124		S16 E55	4476	14	D	1							
WENDEL	24 1110	E 1140		S15 E57	4476	30	D	1							
UCCLE	24 1111	E 1122		S15 E61	4476	11	16	4	1117	3.40	4.30				
UCCLE	24 1138	E 1200		N20 W30	4465	18	1	4	1145	2.20	3.00				
WENDEL	24 1139	E 1157		N17 W29	4465	4	0								
AROSA	24 1238	E 1242	0	N19 W28	4465	27	0	4							
WENDEL	24 1311	E 1338	D	N17 W30	4465	9	1								
AROSA	24 1313	E 1322		N17 W30	4465	9	1	4	1400	1.20	2.40				
UCCLE	24 1355	E 1404		S22 E80	4478	9	1								
WENDEL	24 1607	E 1632	0	S17 E60	4476	25	0	1							G-SHF
														PAGE 6	

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION		DURATION — MINUTES	RM TRACE	ONE COND.	MEASUREMENTS				PROVISIONAL IONOSPHERIC EFFECT		
		START	END	APPROX. LAT.	APPROX. LONG.				TIME — UT	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	MAX WIDTH H <sub>o</sub>		MAX DIF. %	
MT WILSON	23	1614 E	1643	S13 E56	4476	8	1	4	1638	3.40	4.10		15	G-SNF G-SNF Slow S-SNF	
	24	1655	1737	N20 W32	4465	3 D	1	2		3.00	3.30				
	24	1734 E	2028	S14 E58	4476	16	1	2	2016	1.80	3.30				
	24	2208	2224	N25 W16	4465	16	1	3	2210	3.30	4.00				
	24	2306	0006	S18 E60	4476	60	16	2	2320	3.10	5.90				
NIZANIAH	25	0347	0354	S13 E51	4476	7	1	2	0349	1.52	2.38	1.60	154	S-SNF S-SNF S-SNF	
	25	0529 E	0555	N18 E52	4474	26	0	2	0530	7.25	8.74	2.30			
	25	0558 E	0647 D	N15 E25	4474	9	0	1	0538	5.50	6.60	2.00			
	25	0557 E	0626	S13 E51	4476	29	0	2	0603	4.86	7.62	2.50			
	25	0605 E	0622	S18 E50	4476	17	0	2	0605			2.60			
	25	0823	0828	N20 W40	4465	5	1	3	0825			2.20			
	25	0823	0841	N18 W37	4465	18	1	1			3.00				
	25	0830 E	0841	N20 W41	4465	11	0	1	830		1.00				
	25	0835 E	0844	N17 W39	4465	9	0	1				2.60			
	25	0848	0851	S24 E60	4478	3	1	3	0850						
	25	0848	0853	S25 E60	4478	5	1	1							
	25	0859	0905	S25 E63	4478	6	1	1	859		1.00				
	25	1032	1045	S15 E43	4476	13	1	1							
	25	1048	1101	N40 W90	4460	13	1	3							
	25	1056 E	1118	S09 E49	4476	22	0	1			3.00				
	25	1124 E	1127	N23 W09	4469	26	0	1			3.00				
	25	1124 E	1127	S18 E63	4478	3	0	1	3	1125		7.00			
	25	1151	1252	N37 W68	4460	62	16	2							
	25	1202	1214	N35 W85	4460	12	0	1	2						
	25	1411	1435	N19 W42	4465	24	0	1	2	1415	2.00	5.00			G-SNF
	25	1414	1428	N21 W44	4465	14	1	2	2	1418	1.80	2.66	2.20		
	25	1418 E	1430	N19 W42	4465	12	0	1	2	1418	1.80	2.66	2.20		
	25	1418 E	1556	S15 E49	4476	67	16	1	3	1420		6.00			
	25	1449	1605	S14 E46	4476	71	16	1	3	1503	1.91	2.78	2.40		
	25	1454	1605	S15 E47	4476	42	0	1	3	1458					
25	1457 E	1538	S13 E46	4476	58	0	2	3							
25	1502 E	1600	S14 E47	4476	13	0	1	1	1510	1.80	3.20			Slow S-SNF	
25	1509 E	1522	S15 E50	4476	20	2	1	1	1510	1.80	3.20				
25	1513 E	1533	S15 E47	4476	46	0	2	2	1520	5.00	7.00				
25	1519 E	1605 U	S15 E44	4476	17	1	2	2	1520	2.20	5.20			Slow S-SNF	
25	1708	1725	S17 E47	4476	17	1	2	2		3.30					
25	1715 E														
25	1818		1830	N37 W75	4460	12	1	2							
HAWAII	26	0036 E	0040 D	N21 W50	4465	4	0	1	0036	2.10	3.90				
	26	0750 E	0755 D	N22 E12	4474	5	0	1							
	26	1652	1712	N18 E00	4474	20	1	1							
	26	1742	1755	S19 E32	4476	13	1	1							
	26	1754	1805	S08 E06	4476	11	1	1							
	26	1815	1827	N10 W48	4467	12	1	1							
	26	1930	2106	S29 W01	4476	20	1	4	1945	1.80	1.96				
	26	2046	2049	S05 E31	4476	20	1	1	2049	2.26	2.66				
OTTAWA	26	2142	2210 D	S17 E28	4476	43	0	3	2155	1.80	2.08				
	26	2347	2340	S18 E29	4476	50	2	2	2340	10.20	6.20				
	26	2350		S18 E28	4476	50	2	2		5.40	6.20				
ISAC PEAK	26												25	S-SNF	
	26														
NIZANIAH	27	0537 E	0549	N24 W56	4465	12	0	1	0542	1.22	2.79	1.70	7		

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED TIME		MAX. PHASE	LOCATION		DURA- TION — MINUTES	IM- POR- TANCE	OBS. COND.	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT	
		START	END		APPROX. LAT. — MER. DIST.	APPROX. MAGN. — REGION				TIME — U.T.	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	MAX. WIDTH He	MAX. INT. %
ZURICH	27	0823	0850		S19 E25	4476	27	1	2	823		2.00		
AROSA	27	0835	0843		N22 W67	4465	8 D	1						
ZURICH	27	0835	0846		N23 W54	4465	11	1	2	835		4.20		
WENDEL	27	0905	0917		S24 E43	4478	12	1				4.00		
WENDEL	27	0907	0920 D	0908	S30 E34	4478	13 D	1				3.00		
WENDEL	27	1016	1026		N22 W65	4465	10 D	1				4.00		
UCCLE	27	1029	1040	1033	N16 W09	4474	11 D	1	4	1033	1.50	3.00		
WENDEL	27	1030	1044		N15 W07	4474	14	1					2.10	
ONDREJOV	27	1031	1035		N24 W41	4469	4	1	3	1032			2.10	
ONDREJOV	27	1037	1047		N15 W09	4474	10	1	3	1040				
WENDEL	27	1042	1116		N22 W69	4465	34	26				14.00		
AROSA	27	1045	1108		N22 W69	4465	23	1						
UCCLE	27	1045	1113		N25 W80	4465	28	2	4		3.00	6.40		
ONDREJOV	27	1046	1107		N22 W62	4465	21 D	16	3	1054			3.30	
UCCLE	27	1104	1109		N22 W72	4465	5 D	16	3		2.20	4.40		
UCCLE	27	1110	1115		S14 E80	4480	5	1	3	1112		4.00	2.00	
WENDEL	27	1159	1212		S23 W56	4470	13	1						
NEDERHORST	27	1201	1210		N23 W57	4465	9 D	1						
ONDREJOV	27	1201	1211	1202	N24 W58	4465	10	16	3	1202		4.20		
OTTAWA	27	1318	1335	1322	S18 E22	4476	17	16	3	1322	3.31	3.65	4.20	
ONDREJOV	27	1319	1326	1321	S18 E23	4476	7	1	3	1321			2.10	
WENDEL	27	1319	1332		S18 E21	4476	13	1				4.00		
ONDREJOV	27	1417	1428	1419	N23 W66	4465	11	1	3	1419			2.00	
ONDREJOV	27	1510	1515 D		S23 E38	4478	5 D	1	3					
ONDREJOV	27	1510	1517		S28 E39	4478	7 D	1	3	1512			1.90	
SAC PEAK	27	1535	1705	1557 U	S16 E22	4476	90	26	2		10.20	4.93		26
OTTAWA	27	1536	1710		S17 E23	4476	94	2	3	1555	4.47			
ONDREJOV	27	1537	1620 D	1552	S15 E23	4476	53 D	2	3	1552			3.00	
ONDREJOV	27	1545	1616		S19 E30	4476	31 D	2	3	1547			2.60	
OTTAWA	27	1702	1726	1705	S23 E37	4478	24	16	3	1705	2.32	3.01		
SAC PEAK	27	1937	1945	1940	N22 W79	4465	8	1	2		2.50			16
OTTAWA	27	1938	1941	1941	N19 W75	4465	25	16	3	1941	2.32			
SAC PEAK	27	2147	2212	2157 U	N27 W78	4465	25	2	2		7.50			17
MAWATI	27	2150	2210	2156	N28 W85	4465	20	26	1	2156	8.70			
AROSA	28	0645	0700		S07 W06	4476	15 D	1						
UCCLE	28	0807	0817	0814	S25 E31	4478	12 D	1	2	0805	5.10	5.60		
AROSA	28	0807	0819		S24 E23	4478	12	1						
ZURICH	28	0807	0832		S25 E30	4478	23	1	3	809		2.00		
WENDEL	28	0812	0833		S23 E27	4478	21 D	16				5.00		
ZURICH	28	0842	0903		S24 E23	4478	21	1	3	842		3.00		
ZURICH	28	0843	0906		S22 E29	4478	23	1	3	843		1.00		
AROSA	28	0843	0855		S24 E23	4478	10 D	1				6.80		
UCCLE	28	0912	0920 D	0921	N20 W88	4465	24 D	2	4	0921	3.40	6.00		
WENDEL	28	0916	0930 D		S12 E50	4465	15 D	16						
AROSA	28	0955	1015		S12 E50	4476	20 D	1						
AROSA	28	1000	1025		S12 E54	4476	20	1						
UCCLE	28	1012	1035	1016	S12 E54	4476	23 D	1	2	1016	3.40			
AROSA	28	1012	1100 D	1043	S23 E52	4478	48 D	2	2	1043	6.80			
STOCKHOLM	28	1032	1104		S22 E52	4478	34	26						
UCCLE	28	1033	1114		S22 E58	4478	18	2	2	1038	7.90	4.60		
UCCLE	28	1034	1052	1038	S25 E50	4478	20 D	16	2	1052	3.80	6.00		
CAPRI S	28	1035	1055 D		S23 E24	4478	29 D	2	3	1045				
ZURICH	28	1045	1114 D		S24 E20	4478	29 D	2					PAGE 8	



# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED		LOCATION			DURATION — MINUTES	IN- POUN- TANCE	OBS. COND.	MEASUREMENTS				PROVISIONAL LONGPERIODIC EFFECT	
		START	END	APPROX. LAT.	MOON REL. DIST.	PHASE REGION				TIME U.T.	MEAS. AREA Sq. Deg.	CORR. Sq. Deg.	MAX. WIDTH H <sub>g</sub>		MAX. INT. %
ZURICH WENDEL AROSA UCCLE WENDEL UCCLE USNRL AROSA UCCLE WENDEL SAC PEAK USNRL	Mar. 1958	28 1045	E 1114 O	S24 E28	4478	29 0	1	3	1045		3.00			S-SWF	
		28 1145	E 1155	S17 W59	4469	10 0	1					3.00			
		28 1151	1237	S14 E12	4476	46	1								
		28 1153	1226	S15 E19	4476	23	2	2	1211	5.60					
		28 1156	1220	S15 E13	4476	53	1					4.00			
		28 1200	1233 O	S20 E20	4478	33 0	1	2	1218	3.40					
		28 1224	1437	S15 E12	4476	13 0	16	2	1227	3.96		1.00	106		
		28 1304	1312 O	N32 E90	4486	8 0	1					4.40			
		28 1428	1437	N20 W88	4465	9	16	3	1430	2.20					
		28 1429	1439	N22 W85	4465	10	1					3.00			
		28 1510	1535	N27 W90	4465	25	1	2	1548	1.13		1.00	14		
		28 1547	1622	S06 W12	4476	35	1	2							
MCMAH WENDEL OTAWA USNRL OTAWA USNRL OTAWA MCMAH USNRL MCMAH OTAWA USNRL	Mar. 1958	28 1550	E 1608	S04 W15	4476	20 0	2						G-SWF		
		28 1552	E 1608	S07 W11	4476	16 0	1								
		28 1707	E 1822	S15 E10	4476	73 0	26	3	1715	7.37					
		28 1709	E 1822	S15 E08	4476	102	16	2	1714	5.20		2.00		143	
		28 1722	1904	S15 E08	4476	72	16	2	1724	3.17				124	
		28 1725	1820	N15 W25	4474	55	16	3	1738	4.18					
		28 1735	E 1813	N20 W20	4474	37	1	2	1737	2.26		1.00		80	
		28 1736		S15 E10	4476	43	1								
		28 1833	1916	S15 E10	4476	43	1	2	1838	1.82					
		28 1834	1922	N21 W50	4463	48	2	2	1838	1.98					
		28 2042	2131	S22 E20	4478	49	2	4	2049	5.68					
		28 2044	2120 O	S23 E23	4478	36 0	16	2	2045	3.39					
SAC PEAK USNRL SAC PEAK SAC PEAK MITAKA MITAKA AROSA AROSA AROSA WENDEL MEUNEL ZURICH ZURICH ZURICH ZURICH AROSA MEUNEL WENDEL ZURICH CAPRI S UCCLE STOCKHOLM ONOREJOV AROSA UCCLE ONOREJOV WENDEL ONOREJOV AROSA ONOREJOV	Mar. 1958	28 2054	E 2055 D	S25 E20	4478	1 0	2	1			5.30		S-SWF		
		28 2054	E 2055 D	S25 E20	4478	1 0	2	1				2.70			20
		28 2227	2240	N24 W40	4465	18	1	1				2.90			20
		28 2237	2308	S14 E03	4476	31	1	1							
		29 0208	0215 O	S14 E01	4476	7 0	1	1	0208	2.78	2.81	2.36		149	
		29 0244	E 0250 O	N21 W90	4469	6 0	1	1	0244	1.86		3.75			
		29 0640	E 0652 O	N33 E90	4484	12 0	1								
		29 0648	0703	S15 E58	4480	15	1								
		29 0652	0708	N24 W48	4469	16	1								
		29 0730	0736 O	N25 W58	4469	6 0	16					5.00			
		29 0735	E 0748 O	N27 W60	4469	13 0	16					10.80			
		29 0755	E 0806	S09 W24	4476	11 0	1	2	755	2.80					
29 0755	E 0812	S22 E18	4476	17 0	1	2	755	2.00							
29 0759	E 0816 O	N25 W67	4469	17 0	16	2	759	4.90							
29 0853	E 0909 O	S17 W03	4476	16 0	1	2	853	1.00							
29 0912	1012	N22 W30	4474	60	2										
29 0915	1010	N19 W30	4474	55	16						6.00				
29 0919	1008 O	N22 W30	4474	49 0	16						7.00				
29 0920	1018 O	N18 W31	4474	58 0	2	2	922	6.00							
29 0921	E 0955 O	N21 W29	4474	34 0	1	2	941	3.00							
29 0921	1001	N25 W30	4474	40	16	3	934	5.20							
29 0950	E 1019 O	N20 W30	4474	29 0	1										
29 0953	E 1028 O	N19 W30	4474	35 0	1	2	1001				2.50				
29 1031	1043	S06 W04	4476	12	1										
29 1032	1040	S06 W02	4476	8	1	2					2.00				
29 1033	E 1040	S10 W03	4476	7 0	1	3	1035				2.70				
29 1034	1047	S06 E00	4476	13	1	1									
29 1055	1100	N33 E70	4484	5	1						4.00				
29 1217	E 1227	S25 E19	4478	10 0	1	3	1222				2.80	PAGE 9			

# SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION		DATA TION — MINUTES	IN- FOR- TANCE	OBS. COND.	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT
		START	END	APPROX. LAT.	APPROX. LONG.				TIME U T	MEAS. SOL. WAVE	CONTR. AREA SOL. WAVE	
WENDEL	29 1218	1235	1343	S23 E15	4478	17	1	3	1343	2.38	3.00	S-SWF
OTTAWA	29 1339	1405	1346	N36 E76	4484	26	16	3	1346		11.20	
ONDRÉJOV	29 1340	1405	1346	N36 E73	4484	17 D	26	3			6.00	
WENDEL	29 1343	1407		N33 E80	4484	20	16					S-SWF
WENDEL	29 1345	1421 D		N32 E78	4484	24 D	16					
WENDEL	29 1345	1421 D		N32 E78	4484	24 D	16					
ONDRÉJOV	29 1353	1408	1357	S17 E82	4483	15	1	3	1357	1.57	5.00	S-SWF
OTTAWA	29 1356	1408	1357	S18 E83	4483	18 D	16	3	1357			
SAC PEAK	29 1401	1421	1401 E	S18 E82	4483	15 D	1	1	1409	2.50	2.16	
WENDEL	29 1403	1421 D		S18 E82	4483	15	1	3	1409	1.22	3.00	S-SWF
WENDEL	29 1410	1416	1413	S15 E29	4480	14 D	1	3	1413		3.50	
ONDRÉJOV	29 1439	1446	1446	S15 E29	4480	14 D	1	3	1413		2.20	
OTTAWA	29 1448	1502		S15 W00	4476	5 D	1	1	1448	1.62		S-SWF
USNRL	29 1449	1507	1450	S25 W67	4489	14 D	1	2	1450	.79	2.83	
MCNATH	29 1450	1507	1450	N27 W70	4469	18	1	3	1451		3.00	
ONDRÉJOV	29 1507	1512		N23 W72	4476	8 D	1	3	1510		2.00	S-SWF
ONDRÉJOV	29 1507	1512		S10 W09	4476	5 D	1	3	1510			
SAC PEAK	29 1547	1637	1607	S17 W11	4476	50	1	2	1557	2.04	2.11	
USNRL	29 1552	1617 D	1557	S15 W10	4476	53	1	2	1554		2.50	S-SWF
ONDRÉJOV	29 1820	1910	1822	S23 E08	4478	25 D	1	3	1822	3.90	2.70	
SAC PEAK	29 1822	1840	1822	S32 E05	4478	18	1	1	1822	2.50	5.00	
HAWAII	29 1825			S11 W05	4476	2	16	1	2134	2.50	5.00	S-SWF
MCNATH	29 2132	2134 D		N38 E87	4484	2	16	1	2134	3.60		
SAC PEAK	29 2133	2150	2133	N36 E85	4484	17 D	16	1	0029	.97	1.00	
MITAKA	30 0021	0046	0022	S08 W13	4476	25 D	1	1	0109		8.00	S-SWF
MITAKA	30 0102	0109 D		N36 E67	4484	7 D	2	1	0150	1.84	1.90	
MITAKA	30 0149	0200 D		S08 W14	4476	11 D	2	1	0156	1.84	2.75	
MITAKA	30 0152	0203	0156	N36 E66	4484	11 D	16	1	0216	.89	2.85	S-SWF
MITAKA	30 0221	0229	0217	N36 E66	4484	14 D	16	1	0225	3.71	3.78	
MITAKA	30 0401	0415 D	0408	S08 W07	4476	14 D	16	1	0408	1.84	2.76	
MITAKA	30 0456	0509 D	0458	N36 E65	4484	13 D	16	1	0459	1.84	5.88	S-SWF
ATHENS	30 0606	0628		N36 E62	4484	22	1	3		1.00	3.50	
ATHENS	30 0655	0702		N34 E67	4484	7	1	3		.60	2.10	
ATHENS	30 0656	0718		N36 E62	4484	22	1	3				S-SWF
AROSA	30 0755	0806		N14 W47	4474	11 D	1	4	0815	2.80	4.00	
WENDEL	30 0755	0812	0815	N14 W47	4474	17 D	1	4			3.10	
USCCLL	30 0811	0827	0815	S18 E43	4480	16 D	1	4				S-SWF
MOSCOW	30 0817	0846 D		N30 E60	4484	29 D	26					
AROSA	30 0820	0830		S15 W10	4476	10	1					
WENDEL	30 0842	0851	0844	N34 E69	4484	9	16	4	0844	4.50	5.00	S-SWF
USCCLL	30 0842	0853		N34 E69	4484	11	2	4			7.60	
AROSA	30 0843	0850		N34 E66	4484	7	1	4	0915	5.60		
USCCLL	30 0850	1000	0915	S23 W12	4478	70 D	2	4				S-SWF
SCHAUNINS	30 0915	1330		S14 W21	4476	255 D	2	4				
ZURICH	30 0938	0958		N36 E69	4484	20 D	1	3	938		4.00	
ZURICH	30 0941	0944	0941	N23 W68	4469	3	1	3	941		2.00	S-SWF
WENDEL	30 0944	1106		S12 W16	4476	82	16	3			14.00	
WENDEL	30 0944	1230		S16 W19	4476	166	26	3	959		5.00	
USCCLL	30 0945	1042	0959	S18 W20	4476	57	1	4	0957	6.80		S-SWF
USCCLL	30 0945	1421	0957	S17 W22	4476	276	26	4		PAGE	10	

III

## SOLAR FLARES

MARCH 1958

OBSERVATORY	DATE Mar. 1958	OBSERVED UNIVERSAL TIME		LOCATION			DURA- TION MINUTES	IN- TER- FER- ENCE	OBS. COND.	MEASUREMENTS				PROVISIONAL IONOSPHERIC EFFECT
		START	END	MAX. PHASE	APPROX. LAT.	APPROX. LONG.				TIME U T	AREA Sq. Deg.	CORR. Sq. Deg.	MAX. WIDTH Mm	MAX. INT. %
ZURICH	30	0952	1042	1002	S11 W18	4476	50	1	3	1002	3.00			
AROSA	30	1000	1100		S18 W21	4476	60	2						
UCCLE	30	1007	1054	1020	S12 W20	4476	47	2	4	1020	5.60			
CAPRI S	30	1010	1032	D	S17 W19	4476	22	2	1	1015	2.40			
STOCKHOLM	30	1011	1020	D	S13 W15	4476	9	2						
MEUDON	30	1017	1030	D	S20 W20	4476	13	2						
MEUDON	30	1017	1030	D	S20 W20	4476	13	2						
WENDEL	30	1042	1100		S13 W13	4476	18	1						
UCCLE	30	1113	1118	1115	S25 E04	4478	5	1	4	1115	2.20			
UCCLE	30	1114	1134		S36 E80	4484	20	1	4	1121	2.20			
CAPRI S	30	1120	1205	D	S18 W21	4476	45	1	4	1155	2.20			
MEUDON	30	1148	1210	D	S20 W20	4476	22	1						
UCCLE	30	1249	1316	1250	N23 W90	4469	27	1	4	1250	2.20	4.50		
MEUDON	30	1300	1351	D	S20 W20	4476	51	1						
UCCLE	30	1422	1429	1425	S08 W16	4476	7	1	4	1425	2.20			
UCCLE	30	1455	1504	1457	S20 W07	4478	9	1	4	1457	2.20			
UCCLE	30	1533	1621	1544	N36 E80	4484	48	2	2	1544	3.40	5.80		
WENDEL	30	1540	1602		N33 E61	4484	25	1						
CAPRI S	30	1547	1557	D	N35 E65	4484	17	1	2	1546	.80			
WENDEL	30	1614	1626		N34 E65	4484	12	1						
MITAKA	30	1623	E		N38 E60	4484	4	1	1	1624	.99			
SAC PEAK	30	1740	1827	U	S16 E37	4480	47	1	2		3.60			16
SAC PEAK	30	1747	1800	1755	N35 E64	4484	13	1	2		2.40			16
SAC PEAK	30	2007	2117	U	N23 W51	4474	70	2	2		5.10			17
HAWAII	30	2010	2112	2016	N20 W50	4474	62	2	1	2016	3.50	6.40		
HAWAII	30	2036	E	2040	S07 W21	4476	4	1	1	2036	2.00			
SCHAUINS	30	2114	E	0129	S15 E09	4476	25	2						
HAWAII	30	2114	E	0129	S07 E60	4484	18	2	1	2158	2.60	6.40		
HAWAII	30	2130	E	2208	S07 E60	4484	18	2	1	2309	5.00			
HAWAII	30	2340	E	2350	S07 E60	4476	16	1						
MITAKA	30	2347	E	2353	S08 W51	4476	6	1	1	2351	9.43	11.40	1.83	146
SYDNEY	31	0005	0025	0015	S15 W20	4476	20	1						
MITAKA	31	0008	0036	0012	S13 W24	4476	28	2	1	0008	8.56	9.33	3.34	227
HAWAII	31	0014	E	0020	S16 W27	4476	6	2	1	0018	7.20	8.10		
HAWAII	31	0028	E	0032	N38 E55	4484	4	2	1	0032	2.70	6.70		
MITAKA	31	0028	E	0030	N35 E54	4484	10	2	1	0031	3.80	8.60	7.65	149
SYDNEY	31	0040	E	0110	S08 W22	4476	30	1						
MITAKA	31	0049	E	0113	S07 W24	4476	24	2	1	0052	9.43	10.00	4.32	278
HAWAII	31	0050	E	0106	D	4476	16	2	1	0052	6.00	6.50		
MITAKA	31	0119	E	0124	S07 W25	4476	5	2	1	0120	1.84	2.95	2.99	120
MITAKA	31	0146	E	0200	S07 W25	4476	14	2	1	0149	1.86	1.99	2.37	149
MITAKA	31	0231	E	0244	S15 W21	4476	13	2	1	0233	3.80	8.60	3.00	125
SYDNEY	31	0314	0345	0320	N35 E53	4484	31	1						
MITAKA	31	0423	E	0427	S05 W50	4476	4	1	1	0423	2.78	9.73	2.93	96
MITAKA	31	0535	0555	D	N23 W74	4469	4	2	1	0545	.89	3.12	2.29	
ATHENS	31	0650	0704		N25 W72	4469	14	1	3		.80			
MITAKA	31	0856	E	0904	S19 E28	4480	8	2	3	0856	2.43	2.81	1.60	
MITAKA	31	0901	E	0914	S18 E27	4480	13	2	1					
STOCKHOLM	31	1015	E	1025	N20 E61	4485	10	2						
AROSA	31	1016	1032		S09 W27	4476	6	1	1					
UCCLE	31	1119	1140		N23 W90	4469	21	1	2					
UCCLE	31	1120	1131		N22 E66	4485	11	1	2					
UCCLE	31	1143	E	1146	S18 E27	4480	3	2	2	1146	2.10	2.30		
UCCLE	31	1244	1448	1445	S07 W69	4476	4	1	2	1445	2.00	4.00		
UCCLE	31	1244	1534	1528	S14 W41	4476	17	1	4	1528	1.70	2.00		
AROSA	31	1325	E	1328	S15 W40	4476	10	2						
SAC PEAK	31	1332	E	1335	S12 W42	4476	13	1	3					
SAC PEAK	31	1332	E	1335	S12 W42	4476	13	1	3					
SAC PEAK	31	1332	E	1335	S12 W42	4476	13	1	3					
SAC PEAK	31	2305	E	2332	D	4483	27	2	1					

COMMERCIAL - STANDARDS - BOULDER

SAC PEAK: ALL VALUES IN MAX. INT. COLUMN ARE ARBITRARY UNITS (0-40),  
NOT PERCENT OF CONTINUOUS SPECTRUM.


E - LESS THAN  
D - GREATER THAN  
U - APPROXIMATE  
+ - PLUS  
- - MINUS

ANACAPRI SWEDISH  
KODAIKANAL  
KRASNAYA PAKHRA  
ROYAL OBSERVATORY, EDINBURGH  
GREENWICH ROYAL OBSERVATORY, HERSTMONCEUX  
SAC PEAK  
SCHAUINSLAND  
UNITED STATES NAVAL RESEARCH LABORATORY

CAPRI S  
KODAIKANAL  
KRASNAYA  
R O EDIN  
R O HERST  
SAC PEAK  
SCHAUINS  
USNR.

## INTERVALS OF NO FLARE PATROL OBSERVATIONS

MARCH 1958



COMMERCE - STANDARDS - BOULDER

Anacapri (Swedish)  
 Arcetri  
 Arosa  
 Athens  
 Climax  
 Greenwich Royal Observatory,  
 Herstmonceux

Hawaii  
 Huancayo  
 Kodaikanal  
 Meudon  
 Mitaka  
 Nizamia  
 Ondrejov

Ottawa  
 Royal Observatory, Edinburgh  
 Sacramento Peak  
 Uccle  
 U. S. Naval Research Laboratory  
 Zurich



## SUBFLARES NOTED AS FOLLOWS: DATE - UNIVERSAL TIME - COORDINATES

FEBRUARY 1958

WENDEL	01	0848	E	S06	W03	USNRL	08	1407	S12	W42	UCCLE	17	1055	S17	E44		
UCCLE	01	0901		S09	E67	CLIMAX	08	1625	S19	E18	UCCLE	17	1139	N12	W63		
UCCLE	01	0914		S31	E20	USNRL	08	1628	S20	E13	SAC PEAK	17	1600	N13	W57		
WENDEL	01	0945	E	S06	W02	SAC PEAK	08	1650	E	S21	E11	SAC PEAK	17	1637	N12	W58	
UCCLE	01	0959		S30	W55	CLIMAX	08	1738	S18	E10	SAC PEAK	17	1855	N07	W80		
WENDEL	01	1042	E	S30	W51	MC MATH	08	1745	S18	E08	SAC PEAK	17	2150	N09	W61		
SAC PEAK	01	1935		N27	W33	SAC PEAK	08	1927	S12	W45	SAC PEAK	17	2225	S25	E60		
SAC PEAK	01	2140		S21	E90	SAC PEAK	08	2042	E	S19	E07						
SAC PEAK	01	2217		S21	E90												
						* ATHENS	09	0700	E	S11	W48	UCCLE	18	1219	S12	W09	
ATHENS	02	0808	E	S05	W14	* ATHENS	09	0753	S11	W51	UCCLE	18	1414	E	S25	W25	
WENDEL	02	1212	E	S06	W18	UCCLE	09	0857	N08	W72	MC MATH	18	1438	N07	W02		
WENDEL	02	1245	E	S06	W18	UCCLE	09	0905	E	S17	W14	MC MATH	18	1540	S22	W32	
USNRL	02	1444		S19	W90	SAC PEAK	09	1515	U	S15	E08	SAC PEAK	18	1542	S10	E54	
SAC PEAK	02	1612		S12	E77	ZURICH	09	1530	E	S13	E07	SAC PEAK	18	1542	S26	W27	
SAC PEAK	02	1615		S06	W24	SAC PEAK	09	1710	S14	W56	* USNRL	18	1621	S15	W05		
SAC PEAK	02	1640		S12	E74	HAWAII	09	2054	S17	W58	CLIMAX	18	1624	S12	W02		
SAC PEAK	02	1855		S13	E74	HAWAII	09	2118	S17	E07	SAC PEAK	18	1922	S28	W23		
CLIMAX	02	1856		S11	E76	SAC PEAK	09	2142	N19	E09							
						HAWAII	09	2144	N20	E05	UCCLE	19	1259	N11	W16		
ATHENS	03	0729		S12	E33	SAC PEAK	09	2157	S12	W54	SAC PEAK	19	1507	E	N10	W18	
UCCLE	03	1007		S12	E34	HAWAII	09	2200	S17	W57	SAC PEAK	19	1555	N10	W89		
UCCLE	03	1042		S12	E33						SAC PEAK	19	1620	N21	W90		
UCCLE	03	1103		S12	E34	WENDEL	10	0834	E	S22	W08	USNRL	19	1622	N20	W90	
UCCLE	03	1111		S20	E85	UCCLE	10	0908	E	S16	W00	USNRL	19	1843	N09	W22	
* R O HERST	03	1207	E	S11	E31	UCCLE	10	0917	S13	W69	SAC PEAK	19	1900	N21	W90		
UCCLE	03	1248		S27	W85	UCCLE	10	1012	N14	E35	USNRL	19	1901	N19	W90		
UCCLE	03	1251		S12	E34	* MEUDON	10	1129	E	S14	W60	* SAC PEAK	19	1912	S15	W21	
UCCLE	03	1303		S11	E70	WENDEL	10	1205	E	S16	W01	* HAWAII	19	2010	E	N21	E10
UCCLE	03	1311		S17	E82	UCCLE	10	1205	S20	W37	USNRL	19	2012	N20	W90		
UCCLE	03	1437	E	S12	E32	UCCLE	10	1256	S21	W11	CLIMAX	20	1822	N11	E21		
CAPRI S	03	1449	E	S12	E33	UCCLE	10	1433	S16	W03	USNRL	20	1851	S24	E10		
* SAC PEAK	03	1530		S20	E78	USNRL	10	1443	E	S17	W05	USNRL	20	1908	S12	E16	
SAC PEAK	03	1540		S10	E58	SAC PEAK	10	1450	E	S14	W03	USNRL	20	1954	S15	W09	
SAC PEAK	03	1540		S28	W80	USNRL	10	1509	N17	E42							
SAC PEAK	03	1648		S11	E28	SAC PEAK	10	1510	N15	E43	ATHENS	21	0738	S25	E02		
CLIMAX	03	1850		S12	E63	* SAC PEAK	10	1540	S21	W13	ATHENS	21	0853	N11	E13		
* CLIMAX	03	1855		S22	E79	UCCLE	10	1541	E	N08	E35	CAPRI S	21	1453	E	S22	W01
* SAC PEAK	03	1856	E	S20	E77	USNRL	10	1542	S22	W13	CLIMAX	21	1533	S15	E09		
* SAC PEAK	03	2015		S20	E77	SAC PEAK	10	1900	S16	W23	USNRL	21	1535	S13	E09		
* SAC PEAK	03	2039		S12	E28	USNRL	10	1901	S18	W23	CLIMAX	21	1607	S13	E04		
* SAC PEAK	03	2200	U	N19	E57	* HAWAII	10	1904	S16	W65	USNRL	21	1608	S13	E05		
SAC PEAK	03	2230	E	S12	E25	HAWAII	10	2326	S23	W18	* USNRL	21	1646	S13	E05		
SAC PEAK	03	2230		N24	W58						USNRL	21	1710	S04	W20		
						ATHENS	11	0745	S21	W18	USNRL	21	1846	S12	E02		
* CAPRI S	04	0855	E	S11	E51	* ATHENS	11	0749	S18	W27	USNRL	21	2046	S12	E02		
* USNRL	04	1324		S11	E16	ATHENS	11	0753	N11	E08							
USNRL	04	1337		S10	E18	ATHENS	11	0824	S16	W39	ATHENS	23	0645	E	N20	W32	
USNRL	04	1425		N13	E05	UCCLE	11	0952	E	S20	W48	* CLIMAX	23	2049	S11	W04	
USNRL	04	1429		S09	E15	USNRL	11	1319	S23	W25							
USNRL	04	1457		S10	E16	USNRL	11	1615	N12	E20	ATHENS	24	0723	N04	W54		
SAC PEAK	04	1510	E	S13	E14	CLIMAX	11	1620	E	N12	E18	ATHENS	24	0727	N13	E16	
USNRL	04	1512		S11	E15	USNRL	11	1658	N22	E18	ATHENS	24	0730	S34	E80		
USNRL	04	1518		N06	W10	USNRL	11	1753	N12	E18	USNRL	24	1227	E	S19	W26	
SAC PEAK	04	1617		S14	E67	USNRL	11	1834	E	S22	W44	USNRL	24	1256	E	S13	E12
SAC PEAK	04	1712		S13	E13	HUANCAYO	11	1855	S19	W42	USNRL	24	1306	S25	W11		
CLIMAX	04	1721		S13	E14	USNRL	11	1900	S22	W45	USNRL	24	1323	S05	W60		
SAC PEAK	04	1725		S15	E19	* HUANCAYO	11	1940	S19	W23	USNRL	24	1346	S20	W43		
CLIMAX	04	1852		N19	E44						USNRL	24	1446	S22	W41		
USNRL	04	1853		N21	E45	ATHENS	12	0637	E	S15	W28	USNRL	24	1842	S06	E38	
CLIMAX	04	1942		S12	E44	WENDEL	12	1113	E	S12	W62	SAC PEAK	24	1845	S04	E38	
						WENDEL	12	1127	E	S16	W47	USNRL	24	1901	S05	E08	
ATHENS	05	0714		N19	E37	CAPRI S	12	1211	E	N12	E18	USNRL	24	1905	S16	W36	
ATHENS	05	0721		S12	E05	SAC PEAK	12	1550	N11	E05	USNRL	24	1906	N15	E39		
OTTAWA	05	1517	E	S07	E34	USNRL	12	1551	N11	E07	SAC PEAK	24	1912	N13	E39		
CLIMAX	05	1819		S19	E29	USNRL	12	1600	N10	E05	SAC PEAK	24	1945	U	S15	E07	
CLIMAX	05	1903		S10	W28	SAC PEAK	12	1632	N14	E04	USNRL	24	1948	E	S14	E08	
CLIMAX	05	1932		S19	E27	CLIMAX	12	1633	N14	E03							
CLIMAX	05	1946		S13	E19	CLIMAX	12	1645	N09	E03							
						SAC PEAK	12	1645	N08	E03	WENDEL	25	1123	E	S07	W34	
ATHENS	06	0840	E	S12	W08	USNRL	12	1646	N09	E05	WENDEL	25	1127	E	S11	E01	
* CAPRI S	06	1153	E	S10	W10	USNRL	12	1719	N25	E05	WENDEL	25	1138	E	S24	E66	
* R O EDIN	06	1154		S11	W14	CLIMAX	12	1737	N11	E03	USNRL	25	1317	S24	W49		
OTTAWA	06	1447	E	S10	W11	SAC PEAK	12	1745	N10	E05	USNRL	25	1434	S18	W48		
SAC PEAK	06	1640		S16	E22	USNRL	12	1748	N11	E04	* SAC PEAK	25	1630	S12	W52		
OTTAWA	06	1641	E	S16	E22	USNRL	12	1753	S10	W90	MC MATH	25	2119	S22	W60		
SAC PEAK	06	1650		S13	W08	USNRL	12	1757	S10	W60	USNRL	25	2120	S23	W60		
* SAC PEAK	06	1652		N30	W90	USNRL	12	1851	S12	W90							
OTTAWA	06	1652	E	S13	W07	SAC PEAK	12	2127	N07	E07	UCCLE	26	1439	E	S27	W48	
USNRL	06	1725	E	S18	E37	SAC PEAK	12	2127	N22	E69	SAC PEAK	26	2100	S12	W16		
OTTAWA	06	1746	E	S18	E29												
SAC PEAK	06	1802		S11	W12	UCCLE	13	0946	N09	W05	R O HERST	27	1226	E	S12	W21	
USNRL	06	1803		S11	W12	UCCLE	13	1017	N15	W06	SAC PEAK	27	1637	S14	W26		
OTTAWA	06	1803	E	S11	W11	ZURICH	13	1446	N07	W08	* SAC PEAK	27	1745	E	N11	W28	
SAC PEAK	06	2100		S10	W13	USNRL	13	1830	N24	E01	SAC PEAK	27	1925	N12	W32		
											SAC PEAK	27	1935	S21	W80		
SAC PEAK	07	1502	E	S21	E23	UCCLE	14	0942	N25	W65	SAC PEAK	27	1955	N34	E15		
SAC PEAK	07	1502		N30	W90	UCCLE	14	1130	N13	W17	SAC PEAK	27	2025	S13	W27		
SAC PEAK	07	1700		N30	W90	CAPRI S	14	1259	E	N11	W13	SAC PEAK	27	2107	S22	E90	
SAC PEAK	07	1710		S21	E26	SAC PEAK	14	1642	N17	W12	SAC PEAK	27	2137	S14	W28		
SAC PEAK	07	1710		S10	W26	USNRL	14	1644	N17	W12	SAC PEAK	27	2200	S13	W29		
SAC PEAK	07	1815		S21	E26	SAC PEAK	14	1737	N11	W24	SAC PEAK	27	2245	S14	W29		
SAC PEAK	07	1837		N17	E90												
SAC PEAK	07	1905		N11	W28	ATHENS	15	0713	S26	W72	SAC PEAK	28	1730	S13	W38		
SAC PEAK	07	1925		N15	E71	UCCLE	15	0923	S16	W75	SAC PEAK	28	1820	S16	E90		
SAC PEAK	07	1925		S21	E26	WENDEL	15	1347	E	S12	W33	SAC PEAK	28	2040	S16	E90	
SAC PEAK	07	1945		S14	E18	SAC PEAK	15	1742	N12	W37	SAC PEAK	28	2130	S14	W38		
HUANCAYO	07	2000		S13	E16	SAC PEAK	15	1807	N13	W35	SAC PEAK	28	2250	S10	W44		
SAC PEAK	07	2150		S21	E25	SAC PEAK	15	1920	N12	W37							
SAC PEAK	07	2225	E														



## IONOSPHERIC EFFECTS OF SOLAR FLARES

(SHORT-WAVE RADIO FADEOUTS)

FEBRUARY 1958

Feb. 1958	Start UT	End UT	Type	Wide Spread Index	Import- tance	Observation Stations	Known Flare, UT CRPL-F 163 B
2	1707	1745	Slow S-SWF	5	2+	<u>BE</u> , CR, HU, MC, PR, WS	
3	1547	1550	Slow S-SWF	2	1+	HU, <u>PR</u>	1522
5	1035	1125	Slow S-SWF	5	2+	<u>HU</u> , <u>JU</u>	
6	0053	0105	S-SWF	3	1	AD, <u>CA</u>	
6	1658	1718	G-SWF	4	1+	AN, HU, MC, <u>PR</u>	1652E
6	1724	1752	Slow S-SWF	5	2-	BE, <u>CR</u> , HU, MC, PR, WS, CW*	
8	0406	0425	S-SWF	4	1+	CA, <u>OK</u> , TO, CW+	*
8	1112	1130	S-SWF	3	2	KU, NE, <u>PU</u>	*
8	1755	1850	G-SWF	4	3-	BE, HU, <u>MC</u> , PR, WS	1740
9	0210	0235	Slow S-SWF	5	2	AD, <u>CA</u> , <u>OK</u> , TO	0207
9	0558	0610	S-SWF	1	1+	<u>KO</u>	0548E
9	0654	0739	S-SWF	4	3	<u>KO</u> , NE	0658
9	0843	0902	S-SWF	4	2	<u>KO</u> , KU	0837E
9	1332	1418	S-SWF	5	3	BE, HU, MC, NE, PR, <u>PU</u>	1330
9	1421	1436	S-SWF	5	2	BE, HU, MC, <u>NE</u> , PR, <u>PU</u>	1415
9	1935	1957	G-SWF	3	1+	<u>HU</u> , MC, PR	
9	2124	2144	Slow S-SWF	5	1	<u>CA</u> , HU, <u>TO</u> , WS	2108
10	1325	1400	S-SWF	5	3	BE, DA, HU, MC, NE, <u>PR</u> , <u>PU</u>	1323
10	1903	1950	S-SWF	4	3	BE, HU, <u>MC</u> , PR, WS	1900
11	0809	0837	S-SWF	5	3	JU, KO, ND, <u>PU</u>	0820
11	1322	1335	S-SWF	5	1	HU, <u>KU</u>	1319E
11	1345	1410	S-SWF	5	3	HU, <u>JU</u> , <u>PR</u>	1342
12	1750	1840	G-SWF	4	3-	BE, CR, HU, <u>WS</u>	
12	1840	1930	S-SWF	4	3-	BE CR, HU, PR, <u>WS</u>	1839
15	1628	1740	G-SWF	4	1+	AN, BE, <u>MC</u> , WS	
15	1955	2023	G-SWF	4	2	<u>HU</u> , MC, PR, WS	1957
17	1218	1255	S-SWF	2	1+	<u>BE</u> , PR	*
19	1630	1715	G-SWF	3	2	<u>HU</u> , PR	1630
25	0505	0541	S-SWF	1	1	<u>OK</u>	0445
25	2008	2040	Slow S-SWF	4	2	<u>BE</u> , MC, PR, WS	1954
26	0432	0508	G-SWF	3	2+	<u>KO</u> , OK, TO	0449E
26	0540	0636	Slow S-SWF	4	2+	<u>KO</u> , OK, TO, CW+	0547
27	0318	0415	Slow S-SWF	3	1+	AD, <u>OK</u>	*
27	1153	1307	S-SWF	1	3	<u>PU</u>	1155
27	1420	1501	G-SWF	4	2-	<u>HU</u> , MC, PR, WS	

COMMERCE - STANDARDS - BOULDER

\* No known flare patrol at this time.

CA = Canberra, Australia.

CR = Cornell University, N.Y.

DA = Darmstadt, G.F.R.

HH = Heinrich Hertz Institute, Berlin.

JU = Juhlesruh, G.D.R.

KO = Kodaikanal.

KU = Kuhlungsborn

NE = Nederhorst den Berg, Netherlands.

PU = Prague, Czech.

SW = Enkoping, Sweden.

TO = Hiraio Radio Wave Observatory, Japan.

ZU = Zurich, Switzerland.

CW\* = Barbadoes.

CW+ = Hong Kong

SOLAR RADIO EMISSION  
OUTSTANDING OCCURRENCES

OTTAWA

MARCH 1958

2800 MC

Mar. 1958	Type*	Start UT Hrs:Mins	Duration Hrs:Mins	Maximum		Remarks
				Time UT Hrs:Mins	Peak Flux	
3	1 Simple 1 f	21 47	1	21 47.8	5	
4	1 Simple 1	15 32.5	4	15 33.5	6	
4	2 Simple 2	16 04.5	2.5	16 05.5	28	
4	1 Simple 1 f	17 23	5	17 24.5	6	
5	3 Simple 3 A	16 36	35	indet.	9	
8	Group (2)	16 44.2	4.2			
1	Simple 1	16 44.2	2	16 44.9	6	
1	Simple 1	16 48.2	0.2	16 48.3	6	
5	1 Simple 1	17 20.5	1.5	17 21	3	
5	1 Simple 1	20 57.5	4	20 59.5	6	
6	2 Simple 2	22 28.8	2	22 29.3	16	
7	6 Complex	18 14	2.5	18 15	90	
8	6 Complex f	13 26	5	13 26.8	88	
8	3 Simple 3 A	17 22	1 40	17 43	13	
8	Group (2)	17 22.4	9.6			
2	Simple 2	17 22.4	2.5	17 23.4	26	
1	Simple 1	17 28	4	17 29.5	7	
6	Complex	18 00	6	18 02.4	11	
6	Complex	18 55	5.5	18 56.3	7	
8	2 Simple 2	20 59.2	2.5	21 01	9	
9	2 Simple 2	15 43	8	15 45.9	85	
4	Post Increase A		5 15		22	
2	Simple 2	20 07.3	2	20 07.9	9	
10	6 Complex	13 15.2	9	13 16.1	51	
10	1 Simple 1	18 25.2	1.8	18 25.8	6	
10	3 Simple 3 A	20 24	1 30	20 35	10	
6	Complex f	20 28	7	20 32.5	72	
2	Simple 2	21 31.8	1.4	21 32.4	13	
11	3 Simple 3 A	15 00	> 35	indet.	13	
6	Complex	15 12.6	10	15 16.6	50	
12	3 Simple 3 A	14 28	47	14 43	9	
2	Simple 2	14 37	2.5	14 38	33	
12	2 Simple 2	17 02.4	1.3	17 02.8	10	
12	1 Simple 1	20 42.5	1.5	20 43.1	6	
13	6 Complex	13 10.5	4	13 13	6	
13	1 Simple 1	16 20.3	1	16 20.8	6	
13	2 Simple 2	22 16.5	4	22 17.2	10	
14	9 Precursor	14 53	5.5		13	
6	Complex	14 58.5	13	15 01	210	
4	Post Increase		2 45		40	
15	2 Simple 2	18 19.8	1.5	18 20.3	9	
15	1 Simple 1	19 07.5	4	19 09	6	
15	1 Simple 1	21 11.5	1	21 12	6	
16	1 Simple 1	14 10	2.5	14 11.2	3	
16	1 Simple 1	15 33	3	15 34.5	2	
19	2 Simple 2	17 27.5	2.5	17 28.5	13	
19	2 Simple 2	19 09.5	5	19 11	37	
4	Post Increase		35		6	
19	2 Simple 2 f	21 07	9	21 09.5	14	
4	Post Increase		30			
20	8 Group (2)	13 04	16.5			
2	Simple 2	13 04	5	13 04.7	350	
2	Simple 2	13 17	3.5	13 18.2	14	
20	3 Simple 3 A	14 54	40	14 59	16	
2	Simple 2	14 54.6	2.5	14 55.2	32	
20	3 Simple 3	18 50	45	19 02	7	
20	3 Simple 3 f	20 47	15	20 52	7	
20	1 Simple 1	21 41.5	2.5	21 42.5	3	
21	1 Simple 1	13 11	1.5	13 11.7	7	
21	3 Simple 3 f	18 55	40	18 59.3	18	
21	3 Simple 3 f	21 15	35	21 17.2	8	
22	2 Simple 2	12 07	1.5	12 07.4	12	
22	3 Simple 3 f A	18 04	3	18 23	26	
2	Simple 2 f	18 42.2	15	18 44	160	
23	3 Simple 3 A	b11 15	> 9	indet.	45*	*estimated (in sunrise)
7	Period Irreg. Activity	b11 15	> 4 35	11 34	300*	
2	Simple 2	18 26.2	4	18 27.2	20	
24	2 Simple 2	11 38.5	1.5	11 39	30	

SOLAR RADIO EMISSION  
OUTSTANDING OCCURRENCES

OTTAWA

MARCH 1958

2800 MC

Mar. 1958	Type*	Start UT Hrs:Mins	Duration Hrs:Mins	Maximum		Remarks
				Time UT Hrs:Mins	Peak Flux	
25	2 Simple 2	14 13.7	2	14 14.3	46	
25	3 Simple 3 f	14 53.5	40	14 58.5	7	
25	1 Simple 1 f	18 17.3	1	18 17.8	6	
26	2 Simple 2	12 55	2.5	12 55.8	30	
26	2 Simple 2	13 28.5	4	13 29	100	
27	2 Simple 2	11 59.8	14	12 00.6	470	
27	8 Group (2)	13 20	5			
	1 Simple 1	13 20	1	13 20.3	5	
	2 Simple 2	13 24	1	13 24.5	11	
27	2 Simple 2 f	15 05	2	15 06	85	
27	6 Complex	15 43.8	11	15 46.4	220	
	4 Post Increase A		3 10		50	
	6 Complex f	17 01	6	17 03	162	
	4 Post Increase		25		12	
27	1 Simple 1	19 38	3	19 38.5	7	
27	1 Simple 1	21 04.8	1	21 05.1	6	
27	2 Simple 2	21 48.8	6	21 49.7	93	
27	2 Simple 2	23 04	3	23 05	60	
28	7 Period Irreg. Activity	11 47	40	11 58.5	16	In sunset osc.
28	8 Group (2)	15 46.5	10.5			
	2 Simple 2	15 46.5	3	15 47.9	9	
	1 Simple 1	15 52	5	15 54.5	4	
28	9 Precursor f	17 02.5	6		7	
	2 Simple 2 f	17 08.5	14	17 11.5	575	
	4 Post Increase A		2 20		60	
	2 Simple 2	18 35	3.5	18 35.7	100	
28	2 Simple 2 f	20 23.3	5	20 25	9	
28	6 Complex	20 43	33	20 45.1	520	
28	3 Simple 3 f	21 25	40	indet.	24	
29	7 Period Irreg. Activity	12 05	35	12 22.9	53	
29	1 Simple 1	13 02	1	13 02.5	6	
29	2 Simple 2 f	13 40.5	10	13 42	310	
29	2 Simple 2	14 08.6	2.5	14 09	38	
29	1 Simple 1	14 34.4	0.3	14 34.5	7	
29	6 Complex	14 47.1	4	14 49.1	42	
29	1 Simple 1	15 29.5	1	15 30	4	
29	3 Simple 3 A	15 36	1 10	15 56	22	
	1 Simple 1	16 27	3	16 28.5	7	
29	1 Simple 1	16 52	1	16 52.5	7	
29	2 Simple 2	18 20.5	12.5	18 21.8	1400	
	4 Post Increase		1 30		34	
29	8 Group (4)	21 17.5	20.4			
	1 Simple 1	21 17.5	1	21 17.9	7	
	1 Simple 1	21 24.8	2	21 25.7	6	
	6 Complex	21 29.2	3.5	21 31.4	220	
	2 Simple 2	21 36.4	1.5	21 36.8	12	
30	1 Simple 1	12 16.5	5	12 18.5	6	
30	8 Group (2)	14 22.2	6.3			
	2 Simple 2	14 22.2	1	14 22.4	9	
	2 Simple 2	14 26	2.5	14 26.8	52	
30	3 Simple 3	14 57	30	15 01	7	
30	2 Simple 2	15 39.3	2	15 40	58	
30	6 Complex	15 50.5	2.5	15 51.8	12	
30	2 Simple 2	15 59	3	16 00.5	18	
30	1 Simple 1	17 12.2	0.7	17 12.4	7	
30	2 Simple 2	17 20.4	1.5	17 20.8	24	
30	3 Simple 3 A f	17 45	1 45	indet.	17	
	8 Group (4)	17 49	23.8			
	2 Simple 2	17 49	3	17 49.5	44	
	2 Simple 2	17 55.3	3.5	17 56.2	71	
	6 Complex	18 03	1.5	18 04	20	
	1 Simple 1	18 10.8	2	18 11.1	7	
	2 Simple 2	18 59	1.5	18 59.5	8	
	2 Simple 2	19 08	1.5	19 08.3	22	
30	3 Simple 3 f A	19 55	1 25	indet.	15	
	Simple 2	20 57.3	2	20 57.8	23	
30	2 Simple 2 f	21 57.5	5	21 58.3	22	
31	1 Simple 1	12 56	1.5	12 56.5	7	
31	2 Simple 2	14 40.5	2	14 41	42	
31	2 Simple 2	16 51.8	1.5	16 52.2	23	
31	2 Simple 2	17 29.2	1.5	17 29.6	14	
31	7 Period Irreg. Activity	19 30	45	19 43	10	

## OTTAWA

2800 MC

HOURS OF OBSERVATIONS: JANUARY, FEBRUARY, MARCH 1958

OBSERVING PERIOD: January 1300 UT - 2120 UT (approx.)  
 February 1250 UT - 2200 UT (approx.)  
 March 1155 UT - 2245 UT (approx.)

with the following exceptions:

## (1) Records obscured by interference:-

Jan. 6	1535 - 1545	1710 - 1800	1815 - 1840
9	2000 - 2050		
10	1950 - 2020		
Feb. 4	1815 - 1845		
6	2055 - 2100		
15	1930 - 2005		
23	1555 - 1620	1640 - 1700	
26	2010 - 2025		
Mar. 10	1840 - 1850		
12	1940 - 1950	2100 - 2110	
13	1835 - 1850	1910 - 1920	2050 - 2115
14	1925 - 2000	2020 - 2025	
16	1835 - 1855		
17	1840 - 1850	1915 - 1940	2000 - 2020
18	1720 - 1730		
22	1630 - 1640		
23	1730 - 1745	1755 - 1810	
24	1630 - 1700		
25	1830 - 1845		
28	1800 - 1815		
31	1750 - 1820		

## (2) No observations:

Jan. 30	1600 - 1615	1630 - 1645	
Feb. 3	1705 - 1720		
5	1600 - 1615		
6	1520 - 1530		
7	1505 - 1525	1550 - 1600	1610 - 1625
20	1635 - 1645	1830 - 1850	
21	1605 - 1620		
22	1650 - 1715		
23	1625 - 1640		
24	1605 - 1620		
28	1450 - 1910		
Mar. 1	1605 - 1615		
21	1620 - 1635		

## SOLAR RADIO EMISSION

## DAILY DATA

MARCH 1958

CORNELL

200 MC

Mar. 1958	Flux Density $10^{-22} \text{w m}^{-2} (\text{c/s})^{-1}$			Variability 0 to 3			Observing Periods	
	Hours UT			Hours UT			Hours UT	
	12 15	15 18	18 21	12 15	15 18	18 21		
1	[[26	22]	--	[[2	2]	-	1355-1700	
2	[[14	14]	--	[[0	0]	-	1335-1715	
3	[[62	36	18	[[1	1	0	1350-2100	
4	[[13	15	15	[[0	1	1	1335-2110	
5	[[14	14	15	[[1	0	1	1340-2100	
6	[[16	19	19	[[1	1	1	1340-2100	
7	[[24	25	27	[[1	2	3	1340-2100	
8	[[36	40]	--	[[2	2]	-	1340-1700	
9	[[44	48]	--	[[2	1]	-	1310-1700	
10	[[54	61	46	[[1	1	1	1340-2100	
11	[[24	27	35	[[1	1	2	1330-2105	
12	[[35	40	36	[[2	2	1	1335-2100	
13	[[19	22	22	[[1	1	2	1330-2100	
14	[[12	13	12	[[1	1	1	1345-2100	
15	[[14	14]	--	[[0	0]	-	1315-1700	
16	[[12	12]	--	[[0	0]	-	1330-1700	
17	[[12	13	13	[[1	0	0	1345-2105	
18	[[16	13	13	[[1	0	0	1345-1450, 1520-2100	
19	[[12	21	17	[[3	3	2	1340-2100	
20	[[64	90	111	[[3	3	3	1330-2105	
21	[[60	47	39	[[1	2	2	1335-2110	
22	[[23	25	22	[[2	2	2	1345-2045	
23	[[34	28]	--	[[2	1]	-	1325-1700	
24	--	16	17	-	1	1	1520-2105	
25	[[32	33	34	[[1	1	1	1345-2100	
26	[[35	40	45	[[1	1	1	1350-1625, 1740-2100	
27	[[46	54	53	[[1	1	1	1330-2130	
28	[[52	44	42	[[1	1	1	1405-2100	
29	[[52	54]	--	[[3	3]	-	1300-1715	
30	[[58	100]	--	[[3	3]	-	1255-1700	
31	[[20	21	24	[[2	1	2	1340-2105	

COMMERCE - STANDARDS - BOULDER

[ = first hour missing.

[[ = first two hours missing.

] = last hour missing.

]] = last two hours missing.



SOLAR RADIO EMISSION  
OUTSTANDING OCCURRENCES

CORNELL

MARCH 1958

200 MC

Mar. 1958	Type Ap. J	Start UT	Time of Maximum	Duration Minutes	Type IAU	Max. Flux Density $10^{-22} \text{ W m}^{-2} (\text{c/s})^{-1}$		Remarks
						Inst.	Smooth	
3	0	1404		146	F			
	2	1754		1	CD	110	58	
	3	1803.5		.5	CD	>204	>142	
5	8	1732.5		2.5	CD	> 51	> 34	off-scale 1733-33.5, 1734-34.5 UT
	3	1843		.5	CD	> 51	> 32	
7	7	1558			E			
	8	1606		2	CA	> 58	> 26	off-scale 1607.5-08 UT
10	0	1517.5		50	SA			
	3	1642.5		.5	CD	>224	>104	off-scale
	3	2019.5		.5	CD	>204	>110	
11	2	1726.5		3	CA	> 54	> 18	
	2	1948		14	CA	> 54	> 16	
12	3	1906.5		1.5	CA	>233	>135	off-scale 1907.5 UT
13	0	1510		70	CA			
	0	1832		91	F			
14	8	1457		20	ECD	> 54	> 40	off-scale 1504.5 UT
	8	1940.5		10	ECD	> 52	> 37	off-scale 1945, 1946, 1948-50 UT
21	8	1750		2.5	CA	>217	>115	off-scale 1750-50.5 UT
	8	1940		5.5	CD	>224	>132	off-scale 1941-41.5, 1942-42.5, 1943, 1944.5 1945 UT
	2	2045		15	F			
22	0	1602		69	F	78	39	
24	8	1635		6	ECD	>190	>146	off-scale 1637-39 UT
	2	1722		12	E			
25	8	1413.5		2	ECD	>204	>109	off-scale
	3	1418.5		2.5	CD	>204	>115	off-scale 1419-19.5 UT
	3	1423	1423.5	1	CD	156	86	
	8	1817		1	CD	>196	>106	off-scale 1817.5-18 UT
	3	2008		2	CD	200	121	
26	3	1528		.5	CD	>204	>121	
28	7,4	1736.5		92	E			
	8	1836.5		3.5	CD	>240	> 84	off-scale
	2	2023		5.5	F			
	3	2023	2023.5	1.5	CD	>240	>134	
29	8	1631.5		2	ECA	>204	>104	
30	7	1318		92	F			
	0	1524.5		107.5	E			
31	7	1935		93	E			

## SOLAR RADIO EMISSION

## DAILY DATA

FEBRUARY 1958

BOULDER

167 MC

Feb. 1958	Flux Density $10^{-22} \text{ W m}^{-2} (\text{c/s})^{-1}$						Variability 0 to 3						Observing Periods	
	Hours UT					Day	Hours UT					Day	Hours UT	
	0 3	12 15	15 18	18 21	21 24		0 3	12 15	15 18	18 21	21 24			
1	-	-	-	-	18	-	-	-	0	1S	0	0	19.8-24.1	
2	-	-	20	19	19	20	-	-	1	0	0	0	14.2-24.1	
3	-	-	22	19	19	20	-	-	2	1S	1S	1S	14.2-24.1	
4	-	-	21	20	25	22	-	-	2	2	2	2	14.2-24.2	
5	-	-	39	39	56	43	-	-	3	2	3	3	14.2-16.4, 16.8-24.2	
6	-	-	141	224	205	188	-	-	1	1S	2	1S	14.7-24.2	
7	-	-	769	754	579	716	-	-	1	2	2S	2	14.1-24.2	
8	-	-	584	448	349	474	-	-	1	0	1S	1	14.1-24.2	
9	-	-	103	63	903	288	-	-	1	1	2	1	14.1-24.3	
10	-	-	147	38	30	76	-	-	1S	2S	2S	2S	14.0-24.3	
11	-	-	19	18	18	19	-	-	1	1S	1S	1S	14.0-24.3	
12	-	-	15	18	55	26	-	-	2	1S	2S	2S	14.0-24.3	
13	-	-	23	24	24	24	-	-	1S	2	2S	2S	14.0-24.3	
14	-	-	21	22	20	21	-	-	2	2S	2	2	13.9-24.3	
15	-	-	17	17	18	17	-	-	1S	0S	0S	0S	13.9-24.3	
16	-	-	17	18	19	18	-	-	1	1	2S	1	14.3-24.3	
17	-	-	20	19	19	19	-	-	1S	1S	0S	1S	13.9-24.3	
18	-	-	20	18	20	19	-	-	2S	2S	2S	2S	13.8-23.3	
19	-	-	17	17	23	19	-	-	2S	2S	2S	2S	14.2-24.4	
20	-	-	19	23	18	20	-	-	2S	2S	2S	2S	13.8-24.4	
21	-	-	18	18	18	18	-	-	1S	0S	0S	0S	13.8-24.5	
22	-	-	25	24	21	23	-	-	2S	1S	2S	2S	13.8-24.5	
23	-	-	53	32	45	43	-	-	3	2S	2S	2S	13.8-24.5	
24	-	-	108	109	106	108	-	-	2S	2	2S	2S	13.8-24.5	
25	-	-	187	252	273	237	-	-	2S	2	2	2	13.7-24.5	
26	-	-	457	431	326	405	-	-	2	2	2S	2	13.7-24.5	
27	-	-	321	348	315	328	-	-	2	2	2	2	14.3-24.6	
28	-	-	222	174	136	177	-	-	1	1S	2S	1S	13.6-24.6	

COMMERCE - STANDARDS - BOULDER

SOLAR RADIO EMISSION  
OUTSTANDING OCCURRENCES

BOULDER

FEBRUARY 1958

167 MC

Feb. 1958	Type Ap.J	Start UT	Time of Maximum	Duration Minutes	Type IAU	Max. Flux Density $10^{-22} \text{ W m}^{-2} (\text{c/s})^{-1}$		Remarks
						Inst.	Smooth	
1	2	2334	2338.3	06 D	CD	120	-	B2258.9, 2323.8
2	1	1742	1748	13	MF	37	-	
3	1	1410 B	1448.4	595 D	CD	520 D	-	
3	3	1643.3	1643.3	01	ECD	500 D	-	
3	3	1723.1	1723.4	00.6	ECD	440 D	-	B(groups)1939, 2253
4	1	1410 B	1449	400 D	MF	1500 D	-	B1423.7, 1434.4, 1532.4
4	2	1509.9	1514.9	05.3	CD	1500 D	450 D	
4	2	1710.7	1714.1	14.8	ECD	1300 D	140	
4	6	2050	2308.9	200 D	CD	640 D	7	I 1626-1648
5	6	1410 B	1720	600 D	CD	1400 D	34	LB1706, 1750, 1923.7
5	3	1931.4	1931.9	00.9	CD	1000 D	-	N2
6	6	1440 B	2102.2	570 D	CD	1800 D	210	B1950.4, LB2336.0
7	6	1405 B	1711.5	605 D	CD	1800 D	790	N3
8	6	1405 B	1637.0	605 D	CD	1000 D	570	N4
9	6	1405 B	1431.0	430 D	CD	1100 D	100	
9	9	2115	I	175 D	CD	1900 D	910 D	N5, I 2215-2231
10	6	1400 B	2405.2	615 D	CD	1200 D	130	LB1415.6, B1437.1, 1520.4
10	8	1910	1912.2	03	ECD	1700 D	750 D	
10	2	2336	2336.9	02	CD	920 D	540 D	B2344, LB2414.5
11	1	1400 B	1449.2	615 D	F	280	-	S, B1408, 1509.6
12	3	1751.8	1752.0	00.8	ESD	830 D	-	B1555.1
12	1	1756	1758.5	264 D	F	120	-	
12	9	2220	2331.8	120 I	CD	1600 D	600 D	
13	1	1400 B	1712.5	620 D	MF	430 D	-	B(groups)2012, 2318
14	6	1430	1803.8	590 D	CD	830 D	5	

COMMERCE - STANDARDS - BOULDER

- Notes: 1. Interference may obscure or be mistaken for solar events. Relatively small events are not reported.
2. February 5, Bursts 1950.0, 2151.2, 2156.2, 2338.3, 2404.1.
3. February 7, Bursts 1905.4, 2003.8, Large Bursts 2153.2, 2316.7, 2336.9.
4. February 8, Bursts 1433.0, 2152.8, 2307.1, 2341.3, Large Bursts 1638.4, 2144.0.
5. February 9, Group of large bursts 1414-1420, Bursts 1708.3, 1852.9.

SOLAR RADIO EMISSION  
OUTSTANDING OCCURRENCES

BOULDER

FEBRUARY 1958

167 MC

Feb. 1958	Type Ap.J	Start UT	Time of Maximum	Duration Minutes	Type IAU	Max. Flux Density $10^{-22} \text{ w m}^{-2} (\text{c/s})^{-1}$		Remarks
						Inst.	Smooth	
14	3	1436.4	1436.4	00.3	ESD	1100 D	-	B2121.8, 2319.6, LB2341.8
14	3	2130.2	2130.2	00.7	ESD	680 D	-	
16	1	1420 B	2019.4	600 D	MF	93	-	
16	3	2248.8	2249.2	01.1	ECD	200	-	
17	1	1355 B	1801.8	625 D	MF	73	-	S
18	1	1350 B	2102.1	567 D	MF	750 D	-	S, B1403.4, 1409.0, 2112.3
18	8	1538	1539.9	04	ECD	340	100	
18	8	1619	1622.9	07 I	ECD	250	49	S, B2038.2, 2133.5, 2333.6
19	1	1410 B	1633.2	410 D	MF	170	-	
19	6	2100	2253	205 D	CD	310	8	
20	1	1350 B	1521.0	190 D	MF	200	-	S
20	6	1700	2118.5	445 D	CD	500 D	9	S, LB1907.0
21	1	1345 B	1721.9	645 D	MF	110	-	S
22	6	1345 B	1437.8	645 D	CD	210	8	S, B1708.6
23	6	1345 B	1355.1	645 D	CD	770 D	36	N6
24	6	1345 B	-	645 D	CD	1200 D	110	N7
25	6	1340 B	1800 X	650 D	CD	1300 D	270	
26	6	1340 B	1600 X	650 D	CD	1200 D	440	
27	6	1415 B	2100 X	620 D	CD	1200 D	340	
28	6	1335 B	1420.9	660 D	CD	950 D	200	

COMMERCE - STANDARDS - BOULDER

- Notes: 6. February 23, Large bursts 1357, 1426.4, 1550.8, 1555.4, 1642.1.  
 7. February 24, Two large bursts occurred at 1914.0, 1941.8, either of which could be considered the maximum. Other large bursts 1749.9, 1916.6.

SOLAR RADIO EMISSION

DAILY DATA  
FEBRUARY 1958

BOULDER

470 MC

Feb. 1958	Flux Density $10^{-22} \text{ w m}^{-2} (\text{c/s})^{-1}$						Variability 0 to 3						Observing Periods
	Hours UT					Day	Hours UT					Day	Hours UT
	0 3	12 15	15 18	18 21	21 24		0 3	12 15	15 18	18 21	21 24		
1	-	-	88	88	95	90	-	-	0	0	0	0	14.2-24.1
2	-	-	81	80	80	80	-	-	0	0	0	0	14.2-24.1
3	-	-	81	80	80	80	-	-	0	0	OS	0	14.2-24.1
4	-	-	80	80	81	81	-	-	1	0	0	0	14.2-24.2
5	-	-	81	81	82	81	-	-	0	1	1	1	14.2-24.2
6	-	-	82	82	82	82	-	-	0	0	OS	0	14.9-24.2
7	-	-	82	82	82	82	-	-	0	1	2	1	14.1-24.2
8	-	-	81	81	81	81	-	-	2	0	0	1	14.1-24.2
9	-	-	81	81	161	101	-	-	0	0	3	2	14.1-24.3
10	-	-	81	81	81	81	-	-	0	2	1	1	14.1-24.3
11	-	-	81	81	81	81	-	-	0	0	0	0	14.0-24.3
12	-	-	80	80	81	80	-	-	0	0	0	0	14.0-24.3
13	-	-	81	81	81	81	-	-	OS	0	OS	OS	14.0-24.3
14	-	-	80	81	81	81	-	-	0	0	0	0	13.9-24.3
15	-	-	81	81	81	81	-	-	0	0	0	0	13.9-24.3
16	-	-	80	81	80	80	-	-	0	0	0	0	14.4-24.3
17	-	-	80	80	80	80	-	-	0	0	0	0	13.9-24.4
18	-	-	81	80	80	80	-	-	0	0	0	0	13.8-24.4
19	-	-	81	80	-	81	-	-	0	OS	OS	OS	13.8-24.4
20	-	-	-	80	80	80	-	-	OS	OS	OS	OS	13.8-21.0, 21.5-24.4
21	-	-	81	80	81	81	-	-	OS	OS	OS	OS	13.8-24.5
22	-	-	81	80	81	81	-	-	OS	OS	OS	OS	13.8-24.5
23	-	-	81	81	81	81	-	-	0	OS	0	0	13.8-24.5
24	-	-	81	81	81	81	-	-	OS	OS	OS	OS	13.8-24.5
25	-	-	81	-	81	81	-	-	OS	-	OS	OS	13.7-16.9, 22.0-24.5
26	-	-	81	81	81	81	-	-	OS	OS	OS	OS	13.7-24.5
27	-	-	81	81	81	81	-	-	0	OS	18	OS	13.7-15.1, 16.2-24.5
28	-	-	81	81	81	81	-	-	OS	OS	OS	OS	13.6-24.5



SOLAR RADIO EMISSION  
OUTSTANDING OCCURRENCES

BOULDER

FEBRUARY 1958

470 MC

Feb. 1958	Type Ap.J	Start UT	Time of Maximum	Duration Minutes	Type IAU	Max. Flux Density $10^{-22} \text{ w m}^{-2} (\text{c/s})^{-1}$		Remarks
						Inst.	Smooth	
4	1	1410 B	1514.9	595 D	MF	200	-	N2,N3
5	1	1410 B	2308.1	595 D	MF	270	-	
5	2	1929 B	1929.4	0.6 I	CD	970	-	
6	1	1453 B	1803.4	557 D	F	150	-	N4
7	1	1405 B	2007.8	605 D	MF	560	-	
7	2	2102.1	2102.6	02.2	ECD	980	120	
7	3	2331.7	2331.8	00.2	ECD	440	-	N5
8	2	1632	1638.6	08	CD	810	90	
9	1	1405 B	1419.9	427 D	MF	590	-	
9	9	2112	2204.3	133	CD	2300 D	230	
10	3	1859.4	1859.5	00.5	ECD	2500 D	-	
10	8	1904	1906.5	21	CD	280	10	
10	1	2115	2242.4	105	MF	130	-	
10	8	2200	2202.2	03.7	ECD	210	100	
12	0	2229 B	2238.4	15 I	CD	130	-	
24	1	1345 B	2155.8	645 D	MF	170	-	
27	2	2200.6	2205.7	05.2	CD	490	-	S, N6

COMMERCE - STANDARDS - BOULDER

- Notes: 1. Interference may occasionally obscure or be mistaken for solar events.  
 2. February 3, small burst at 1648.7.  
 3. February 4, burst at 1434.3.  
 4. February 7, small groups of bursts at 2253 and 2336.  
 5. February 8, large burst 1414.3.  
 6. February 25, Probable type "1" or "MF" all day

## GEOMAGNETIC ACTIVITY INDICES

FEBRUARY 1958

Feb. 1958	C	Values Kp								Sum	Ap	Final Selected Days	
		Three hour Gr. interval											
		1	2	3	4	5	6	7	8				
1	0.4	3-	1-	2+	2+	3o	1+	2-	3-	17-	9	Five Quiet	
2	0.4	3-	2+	3-	3o	2+	1+	2-	0+	16+	9		
3	0.2	1-	1+	1o	1+	3o	1+	2o	2o	13-	6		
4	1.1	3-	0+	1o	3o	4-	3+	5-	4-	22+	17		3
5	1.4	3+	4+	4-	4o	5-	4-	5-	5-	33o	30		15
6	1.3	4o	4+	4+	4+	4+	5-	3o	4+	33+	30	24	
7	1.2	4o	4-	4o	4o	3+	4o	4o	4o	31o	25	25	
8	1.2	4-	5-	4o	4o	4-	4-	4o	4o	32-	27	26	
9	1.0	4o	4o	2o	3-	3-	3+	4o	3o	26-	18		
10	1.2	4o	4-	2-	2o	2+	5o	5-	4+	28-	24		
11	2.0	9o	8+	9-	8+	8o	5+	6o	6o	60-	199	Five Disturbed	
12	1.8	6o	6-	6+	5+	4o	5-	6o	4+	42+	59		
13	1.0	4-	2-	3+	4-	4+	4-	3o	2+	26-	18		
14	1.2	4o	5o	4o	3o	3+	4+	2+	2o	28o	23		6
15	0.5	2-	1o	1o	2+	2+	3-	2+	3-	16o	8		11
16	0.9	2+	1-	1+	3+	3o	4o	3+	3o	21o	14	12	
17	1.3	4o	5-	4o	4-	5-	4o	5-	4o	34-	31	17	
18	1.3	4o	5o	4+	4+	4o	3+	5o	4-	34-	32	18	
19	1.1	4+	4-	5-	4-	3+	3o	4o	4-	30+	25		
20	1.2	4-	4o	3o	4o	4o	3o	5-	5-	31o	26		
21	1.3	3+	5-	4+	4-	4+	3o	5-	5o	33o	31	Ten Quiet	
22	1.1	5-	4o	4-	4o	4-	3o	4-	4o	31-	25		
23	0.9	4+	4o	4o	4-	3o	3+	3o	3-	28o	21		
24	0.3	2+	3-	2o	3-	3-	2-	2-	1-	16+	8		1
25	0.2	1-	2o	3-	2o	1-	1-	1+	1-	11-	5		2
26	0.2	2o	2o	3-	2o	1+	1o	2-	1+	14o	7	3	
27	0.5	3-	1o	1o	1o	2-	2o	3+	4-	16+	10	15	
28	0.8	2o	2+	3+	4o	4+	3-	2+	2o	23o	15	16	
												24	
												25	
												26	
												27	
												28	
Mean:	0.96									Mean:	27		



CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS  
NORTH ATLANTIC  
FEBRUARY 1958

Feb. 1958	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day index	Advance forecasts (J-reports) for whole day; issued in advance by:			Geomag- netic K <sub>Fr</sub>	
	00 to 06	06 to 12	12 to 18	18 to 24	00	06	12	18		1-4 days	4-7 days	8-25 days	Half Day (1) (2)	
1	7-	7-	7o	7o	7	7	7	7	7o	7	7		2	2
2	6+	7-	7o	7o	7	7	7	7	7-	7	7		3	1
3	7o	7-	7o	7+	7	7	7	7	7o	7	7		1	2
4	7o	7-	7o	6o	7	7	7	7	7-	7	7		2	3
5	5+	5o	7o	6o	6	5	6	6	6-	7	7		(4)	(4)
6	6-	6-	7-	6o	6	6	7	6	6o	7	7		(4)	(4)
7	6o	6-	7-	6o	6	6	7	6	6+	6	7		3	3
8	6+	7-	7-	6+	6	6	7	7	7-	6	7		(4)	(4)
9	6+	6+	7-	6o	6	7	7	7	6+	6	7		3	3
10	7-	7-	6+	5-	7	7	7	6	6o	6	7		3	3
11	3-	1+	3o	3+	6	2	3	3	(3-)	6	6		(9)	(6)
12	3+	4o	6+	6-	3	2	5	5	(4+)	5	7		(5)	(5)
13	6+	6+	7-	6+	5	6	7	6	6+	4	7		3	3
14	6+	6+	7-	6+	6	6	6	6	6+	6	4		3	2
15	6+	7-	7o	7-	6	6	6	6	7-	6	6		1	2
16	7-	7-	7-	7o	6	6	7	6	7-	6	6		1	3
17	7-	6+	7-	6-	6	6	6	6	6+	7	6		(4)	(4)
18	5+	5o	6+	6-	5	5	6	5	6-	6	7		(4)	3
19	6o	6o	7-	6+	5	6	6	6	6+	6	7		(4)	3
20	6-	6+	7o	6+	6	6	6	6	6+	6	7		3	3
21	6-	6o	7o	6o	6	5	6	6	6+	6	6		(4)	3
22	6-	6+	7o	6+	6	5	7	6	6+	6	6		3	3
23	6-	6o	7o	7-	6	6	7	6	6+	6	6		(4)	3
24	6o	7-	7o	7o	6	6	7	7	7-	6	6		2	2
25	7o	7-	7+	7+	7	7	7	7	7o	7	6		2	1
26	7o	7o	7o	7o	7	7	7	7	7o	7	6		2	2
27	7o	7o	7o	7o	7	7	7	7	7o	7	6		1	3
28	7-	7-	7o	7o	7	7	7	7	7-	7	7		3	3
Score: Quiet Periods														
					P	20	19	18	18					
					S	6	7	9	9					
					U	0	0	0	0					
					F	0	0	0	0					
Disturbed Periods														
					P	1	0	1	1					
					S	0	1	0	0					
					U	0	1	0	0					
					F	1	0	0	0					

( ) represent disturbed values.



## CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS


## NORTH ATLANTIC

FEBRUARY 1958

— Short-term forecast


| Range of reports

o Quality figure



## OUTCOME OF ADVANCED FORECASTS



1 TO 4 DAYS AHEAD







# USEFUL FREQUENCY RANGES -- NORTH ATLANTIC PATH

FEBRUARY 1958



FEBRUARY 1958



COMMERCE - STANDARDS - BOULDER






CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS  
NORTH PACIFIC  
FEBRUARY 1958

OUTCOME OF ADVANCED FORECASTS

1 TO 4 DAYS AHEAD



## ALERT PERIODS AND SPECIAL WORLD INTERVALS

Alert Issued Ends 1600 UT 1600 UT	SWI Issued Ends 0001 UT 2400 UT	A <sub>Be</sub> On Days of Alert Period (SWI Underlined)	Number of Flares of IMP $\geq$ 2 Reported Promptly on Days of Alert Period
1958			
Mar 02-Mar 07	Mar 05-Mar 05	09-21-27- <u>31</u> -31-20	0-1-0-0-0-1
Mar 12-Mar 13		44- <u>38</u>	2-0
Mar 14-Mar 16	Mar 15-Mar 15	16- <u>24</u> -19	1-0-0
Mar 20-Apr 01	Mar 23-Mar 25	23-33-22- <u>16</u> - <u>16</u> - <u>24</u> -18-	8-3-0-3-2-7-1-
	Mar 30-Mar 31	-16-12-10- <u>27</u> - <u>14</u> -20	-6-9-2-9-2-1

COMMERCE - STANDARDS - BOULDER





